ERRATUM TO: DEFORMATION QUANTIZATION IN ALGEBRAIC GEOMETRY

AMNON YEKUTIELI

ABSTRACT. This note contains a correction of the *proofs* of the main results of the paper [A. Yekutieli, Deformation quantization in algebraic geometry, Adv. Math. **198** (2005), 383-432]. The results are correct as originally stated.

0. INTRODUCTION

This note contains a correction of the *proofs* of the main results of [Ye1], namely Theorems 0.1 and 0.2. The results are correct as originally stated.

The mistake in my original proofs was discovered Michel Van den Bergh, and I thank him for calling my attention to it. The way to fix the proofs is essentially contained in his paper [VdB].

Let me begin by explaining the mistake. As can be seen in Example 0.1 below, the mistake itself is of a rather elementary nature, but it was obscured by the complicated context.

Suppose \mathbb{K} is a field of characteristic 0, and X is a smooth separated *n*-dimensional scheme over \mathbb{K} . Recall that the coordinate bundle Coor X is an infinite dimensional bundle over X, with free action by the group $\operatorname{GL}_{n,\mathbb{K}}$. The quotient bundle is by definition

$$\operatorname{LCC} X := \operatorname{Coor} X/\operatorname{GL}_{n,\mathbb{K}},$$

and the projection π_{gl} : Coor $X \to LCC X$ is a $GL_{n,\mathbb{K}}$ -torsor.

The erroneous (implicit) assertion in [Ye1] is that the de Rham complexes satisfy

$$(\pi_{\mathrm{gl}*}\,\Omega_{\mathrm{Coor}\,X})^{\mathrm{GL}_n(\mathbb{K})} = \Omega_{\mathrm{LCC}\,X}.$$

From that I deduced (incorrectly, top of page 424) that the Maurer-Cartan form ω_{MC} is a global section of the sheaf

$$\Omega^1_{\operatorname{LCC} X} \widehat{\otimes}_{\mathcal{O}_{\operatorname{LCC} X}} \pi^{\widehat{\ast}}_{\operatorname{lcc}}(\mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{T}^0_{\operatorname{poly},X}).$$

(This false, as can be seen from [VdB, Lemma 6.5.1]). This led to many incorrect formulas in [Ye1, Section 7].

The correct thing to do is to work with the infinitesimal action of the Lie algebra $\mathfrak{g} := \mathfrak{gl}_n(\mathbb{K})$. For $v \in \mathfrak{g}$ one has the contraction (inner derivative) ι_v , which is a degree -1 derivation of the de Rham complex $\pi_{\mathrm{gl}*} \Omega_{\mathrm{Coor} X}$. Recall that the Lie derivative is $L_v := d \circ \iota_v + \iota_v \circ d$. A local section $\omega \in \pi_{\mathrm{gl}*} \Omega_{\mathrm{Coor} X}$ is said to be \mathfrak{g} -invariant if $\iota_v(\omega) = L_v(\omega) = 0$ for all $v \in \mathfrak{g}$. According to [VdB, Lemma 9.2.3] one has

$$(\pi_{\mathrm{gl}*}\,\Omega_{\mathrm{Coor}\,X})^{\mathfrak{g}} = \Omega_{\mathrm{LCC}\,X}.$$

1

Date: 13 August 2007.

AMNON YEKUTIELI

It is worthwhile to note that in my incorrect proof there was no need to invoke Kontsevich's property (P5) from [Ko]. The correct proof does require property (P5) – cf. [VdB, Lemma 9.2.1].

Example 0.1. Here is a simplified example. Suppose G is the affine algebraic group $\operatorname{GL}_{1,\mathbb{K}} = \operatorname{Spec} \mathbb{K}[t, t^{-1}]$, and X is the variety G, with regular left action. The group of rational points is $G(\mathbb{K}) = \mathbb{K}^{\times}$. The action of G on X is free, the invariant ring is $\mathcal{O}(X)^{G(\mathbb{K})} = \mathbb{K}$, and the quotient is $X/G = \operatorname{Spec} \mathbb{K}$. For the de Rham complex

$$\Omega(X) = \mathcal{O}(X) \oplus \Omega^{1}(X) = \mathbb{K}[t, t^{-1}] \oplus \mathbb{K}[t, t^{-1}] \cdot \mathrm{d}t$$

we have $\Omega(X)^{G(\mathbb{K})} \neq \mathbb{K}$, since it contains $t^{-1}dt$. But for the infinitesimal action of the Lie algebra $\mathfrak{g} := \mathfrak{gl}_1(\mathbb{K})$ it is easy to see that $\Omega(X)^{\mathfrak{g}} = \mathbb{K}$.

After some deliberation I decided that the best way to present the erratum is by completely rewriting [Ye1, Section 7]. This is Section 1 below. Section 2 contains some additional minor corrections to [Ye1].

1. The Global L_{∞} Quasi-isomorphism

This is a revised version of [Ye1, Section 7]. In this section we prove the main results of the paper [Ye1], namely Theorem 0.1 (which is repeated here as Corollary 1.19), and Theorem 0.2 (which is repeated here, with more details, as Theorem 1.2). Throughout \mathbb{K} is a field containing \mathbb{R} , and X is a smooth irreducible separated *n*-dimensional scheme over \mathbb{K} . We use all notation, definitions and results of [Ye1, Sections 1-6] freely. However the bibliography references relate to the list at the end of this note.

Suppose $U = \{U_0, \ldots, U_m\}$ is an open covering of the scheme X, consisting of affine open sets, each admitting an étale coordinate system, namely an étale morphism $U_i \to \mathbf{A}_{\mathbb{K}}^n$. For every *i* let $\sigma_i : U_i \to \text{LCC } X$ be the corresponding section of $\pi_{\text{lcc}} : \text{LCC } X \to X$, and let $\boldsymbol{\sigma}$ be the resulting simplicial section (see [Ye1, Theorem 6.5]).

Let \mathcal{M} be a bounded below complex of quasi-coherent \mathcal{O}_X -modules. The mixed resolution $\operatorname{Mix}_{U}(\mathcal{M})$ was defined in [Ye1, Section 6]. For any integer *i* let

$$G^{i} \operatorname{Mix}_{U}(\mathcal{M}) := \bigoplus_{j=i}^{\infty} \operatorname{Mix}_{U}^{j}(\mathcal{M}),$$

so $\{G^i \operatorname{Mix}_{U}(\mathcal{M})\}_{i \in \mathbb{Z}}$ is a descending filtration of $\operatorname{Mix}_{U}(\mathcal{M})$ by subcomplexes, with $G^i \operatorname{Mix}_{U}(\mathcal{M}) = \operatorname{Mix}_{U}(\mathcal{M})$ for $i \leq 0$, and $\bigcap_i G^i \operatorname{Mix}_{U}(\mathcal{M}) = 0$. Let

$$\operatorname{gr}_{\mathrm{G}}^{i}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{M}) := \operatorname{G}^{i}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{M}) / \operatorname{G}^{i+1}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{M})$$

and $\operatorname{gr}_{\operatorname{G}}\operatorname{Mix}_{U}(\mathcal{M}) := \bigoplus_{i} \operatorname{gr}_{\operatorname{G}}^{i}\operatorname{Mix}_{U}(\mathcal{M}).$

By [Ye1, Proposition 6.3], if \mathcal{G}_X is either $\mathcal{T}_{\text{poly},X}$ or $\mathcal{D}_{\text{poly},X}$, then $\text{Mix}_U(\mathcal{G}_X)$ is a sheaf of DG Lie algebras on X, and the inclusion

$$\eta_{\mathcal{G}}: \mathcal{G}_X \to \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{G}_X)$$

is a DG Lie algebra quasi-isomorphism.

Note that if $\phi : \operatorname{Mix}_{U}(\mathcal{M}) \to \operatorname{Mix}_{U}(\mathcal{N})$ is a homomorphism of complexes that respects the filtration $\{\mathrm{G}^{i} \operatorname{Mix}_{U}\}$, then there exists an induced homomorphism of complexes

$$\operatorname{gr}_{\operatorname{G}}(\phi) : \operatorname{gr}_{\operatorname{G}}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{M}) \to \operatorname{gr}_{\operatorname{G}}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{N}).$$

 $\mathbf{2}$

Suppose \mathcal{G} and \mathcal{H} are sheaves of DG Lie algebras on a topological space Y. An \mathcal{L}_{∞} morphism $\Psi : \mathcal{G} \to \mathcal{H}$ is a sequence of sheaf morphisms $\psi_j : \prod^j \mathcal{G} \to \mathcal{H}$, such that for every open set $V \subset Y$ the sequence $\{\Gamma(V, \psi_j)\}_{j \geq 1}$ is an \mathcal{L}_{∞} morphism $\Gamma(V, \mathcal{G}) \to \Gamma(V, \mathcal{H})$. If $\psi_1 : \mathcal{G} \to \mathcal{H}$ is a quasi-isomorphism then Ψ is called an \mathcal{L}_{∞} quasi-isomorphism.

Recall that there is a canonical quasi-isomorphism of complexes of \mathcal{O}_X -modules

(1.1)
$$\mathcal{U}_1: \mathcal{T}_{\mathrm{poly},X} \to \mathcal{D}_{\mathrm{poly},X}.$$

According to [Ye2, Theorem 4.17], the induced homomorphism

$$\operatorname{gr}_{G}(\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{U}_{1})) : \operatorname{gr}_{G}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\operatorname{poly},X}) \to \operatorname{gr}_{G}\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X})$$

is a quasi-isomorphism.

Theorem 1.2. Let X be an irreducible smooth separated \mathbb{K} -scheme. Let $U = \{U_0, \ldots, U_m\}$ be an open covering of X consisting of affine open sets, each admitting an étale coordinate system, and let σ be the associated simplicial section of the bundle LCC $X \to X$. Then there is an induced L_{∞} quasi-isomorphism

$$\Psi_{\boldsymbol{\sigma}} = \{\Psi_{\boldsymbol{\sigma};j}\}_{j\geq 1} : \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\operatorname{poly},X}) \to \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X}).$$

The homomorphism $\Psi_{\sigma,1}$ respects the filtration {Gⁱ Mix_U}, and

$$\operatorname{gr}_{G}(\Psi_{\boldsymbol{\sigma};1}) = \operatorname{gr}_{G}(\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{U}_{1})).$$

Proof. Let Y be some K-scheme, and denote by \mathbb{K}_Y the constant sheaf. For any p we view Ω_Y^p as a discrete inv \mathbb{K}_Y -module, and we put on $\Omega_Y = \bigoplus_{p \in \mathbb{N}} \Omega_Y^p$ direct sum dir-inv structure. So Ω_Y is a discrete (and hence complete) DG algebra in Dir Inv Mod \mathbb{K}_Y .

We shall abbreviate $\mathcal{A} := \Omega_{\operatorname{Coor} X}$, so that $\mathcal{A}^0 = \mathcal{O}_{\operatorname{Coor} X}$ etc. As explained above, \mathcal{A} is a DG algebra in Dir Inv Mod $\mathbb{K}_{\operatorname{Coor} X}$, with discrete (but not trivial) dir-inv module structure.

There are sheaves of DG Lie algebras $\mathcal{A} \otimes \mathcal{T}_{poly}(\mathbb{K}[[t]])$ and $\mathcal{A} \otimes \mathcal{D}_{poly}(\mathbb{K}[[t]])$ on the scheme Coor X. The differentials are $d_{for} = d \otimes 1$ and $d_{for} + 1 \otimes d_{\mathcal{D}}$ respectively. As explained just prior to [Ye1, Theorem 3.16], \mathcal{U} extends to a continuous \mathcal{A} multilinear L_{∞} morphism

$$\mathcal{U}_{\mathcal{A}} = \{\mathcal{U}_{\mathcal{A};j}\}_{j\geq 1} : \mathcal{A} \ \widehat{\otimes} \ \mathcal{T}_{\mathrm{poly}}(\mathbb{K}[[t]]) o \mathcal{A} \ \widehat{\otimes} \ \mathcal{D}_{\mathrm{poly}}(\mathbb{K}[[t]])$$

of sheaves of DG Lie algebras on $\operatorname{Coor} X$.

The MC form $\omega := \omega_{\mathrm{MC}}$ is a global section of $\mathcal{A}^1 \widehat{\otimes} \mathcal{T}^0_{\mathrm{poly}}(\mathbb{K}[[t]])$ satisfying the MC equation in the DG Lie algebra $\mathcal{A} \widehat{\otimes} \mathcal{T}_{\mathrm{poly}}(\mathbb{K}[[t]])$. See [Ye1, Proposition 5.9]. According to [Ye1, Theorem 3.16], the global section $\omega' := \mathcal{U}_{\mathcal{A};1}(\omega) \in$ $\mathcal{A}^1 \widehat{\otimes} \mathcal{D}^0_{\mathrm{poly}}(\mathbb{K}[[t]])$ is a solution of the MC equation in the DG Lie algebra $\mathcal{A} \widehat{\otimes} \mathcal{D}_{\mathrm{poly}}(\mathbb{K}[[t]])$, and there is a continuous \mathcal{A} -multilinear \mathcal{L}_{∞} morphism

$$\mathcal{U}_{\mathcal{A},\omega} = \{\mathcal{U}_{\mathcal{A},\omega;j}\}_{j\geq 1} : \left(\mathcal{A}\,\widehat{\otimes}\,\mathcal{T}_{\mathrm{poly}}(\mathbb{K}[[t]])\right)_{\omega} \to \left(\mathcal{A}\,\widehat{\otimes}\,\mathcal{D}_{\mathrm{poly}}(\mathbb{K}[[t]])\right)_{\omega'}$$

between the twisted DG Lie algebras. The formula is

(1.3)
$$\mathcal{U}_{\mathcal{A},\omega;j}(\gamma_1\cdots\gamma_j) = \sum_{k\geq 0} \frac{1}{(j+k)!} \mathcal{U}_{\mathcal{A};j+k}(\omega^k\cdot\gamma_1\cdots\gamma_j)$$

for $\gamma_1, \ldots, \gamma_j \in \mathcal{A} \widehat{\otimes} \mathcal{T}_{\text{poly}}(\mathbb{K}[[t]])$. The two twisted DG Lie algebras have differentials $d_{\text{for}} + ad(\omega)$ and $d_{\text{for}} + ad(\omega') + \mathbf{1} \otimes d_{\mathcal{D}}$ respectively. This sum in (1.3) is actually finite, the number of nonzero terms in it depending on the bidegree of $\gamma_1 \cdots \gamma_j$. Indeed, if $\gamma_1 \cdots \gamma_j \in \mathcal{A}^q \widehat{\otimes} \mathcal{T}^p_{\text{poly}}(\mathbb{K}[[t]])$, then

(1.4)
$$\mathcal{U}_{\mathcal{A};j+k}(\omega^k \cdot \gamma_1 \cdots \gamma_j) \in \mathcal{A}^{q+k} \widehat{\otimes} \mathcal{D}^{p+1-j-k}_{\text{poly}}(\mathbb{K}[[t]]),$$

which is is zero for k > p - j + 2; see proof of [Ye2, Theorem 3.23].

By [Ye1, Theorem 5.6] (the universal Taylor expansions) there are canonical isomorphisms of graded Lie algebras in Dir $\mathsf{Inv} \mathsf{Mod} \mathbb{K}_{\mathrm{Coor} X}$

(1.5)
$$\mathcal{A} \widehat{\otimes} \mathcal{T}_{\text{poly}}(\mathbb{K}[[t]]) \cong \mathcal{A} \widehat{\otimes}_{\mathcal{A}^0} \pi^*_{\text{coor}}(\mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{T}_{\text{poly},X})$$

and

(1.6)
$$\mathcal{A} \widehat{\otimes} \mathcal{D}_{\text{poly}}(\mathbb{K}[[t]]) \cong \mathcal{A} \widehat{\otimes}_{\mathcal{A}^0} \pi^{\widehat{\ast}}_{\text{coor}}(\mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{D}_{\text{poly},X}).$$

[Ye1, Proposition 5.8] tells us that

$$\mathbf{d}_{\mathrm{for}} + \mathrm{ad}(\omega) = \nabla_{\mathcal{P}}$$

under these identifications. Therefore

(1.7)
$$\mathcal{U}_{\mathcal{A},\omega} = \{\mathcal{U}_{\mathcal{A},\omega;j}\}_{j\geq 1} : \mathcal{A}\widehat{\otimes}_{\mathcal{A}^0} \pi^*_{\text{coor}}(\mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{T}_{\text{poly},X}) \\ \to \mathcal{A}\widehat{\otimes}_{\mathcal{A}^0} \pi^*_{\text{coor}}(\mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{D}_{\text{poly},X})$$

is a continuous \mathcal{A} -multilinear L_{∞} morphism between these DG Lie algebras, whose differentials are $\nabla_{\mathcal{P}}$ and $\nabla_{\mathcal{P}} + \mathbf{1} \otimes d_{\mathcal{D}}$ respectively. As in the proof of [Ye1, Theorem 5.6], under the identifications (1.5) and (1.6) we have the equality

(1.8)
$$\mathcal{U}_{\mathcal{A};1} = \mathbf{1} \otimes \pi^*_{\mathrm{coor}}(\mathbf{1} \otimes \mathcal{U}_1),$$

i.e. it is the pullback of the map (1.1).

Let us filter the DG algebra \mathcal{A} by the descending filtration $\{G^i\mathcal{A}\}_{i\in\mathbb{Z}}$, where $G^i\mathcal{A} := \bigoplus_{j=i}^{\infty} \mathcal{A}^i$. The DG Lie algebras appearing in equation (1.7) inherit this filtration. From formulas (1.3) and (1.4) we see that the homomorphism of complexes $\mathcal{U}_{\mathcal{A},\omega;1}$ respects the filtration, and from (1.8) we see that

$$\operatorname{gr}_{\mathrm{G}}(\mathcal{U}_{\mathcal{A},\omega;1}) = \operatorname{gr}_{\mathrm{G}}(\mathcal{U}_{\mathcal{A};1}) = \mathbf{1} \otimes \pi_{\operatorname{coor}}^{\widehat{*}}(\mathbf{1} \otimes \mathcal{U}_{1}).$$

Let $n := \dim X$. As noted earlier, the action of $\mathfrak{g} := \mathfrak{gl}_n(\mathbb{K})$ gives

$$(\pi_{\mathrm{gl}*} \mathcal{A})^{\mathfrak{g}} = (\pi_{\mathrm{gl}*} \Omega_{\mathrm{Coor} X})^{\mathfrak{g}} = \Omega_{\mathrm{LCC} X}.$$

According to [VdB, Lemma 9.2.1], the L_{∞} morphism $\mathcal{U}_{\mathcal{A},\omega}$ commutes with the action of the Lie algebra \mathfrak{g} . Therefore $\mathcal{U}_{\mathcal{A},\omega}$ descends (i.e. restricts) to a continuous Ω_{LCCX} -multilinear L_{∞} morphism

(1.9)
$$\begin{array}{c} \mathcal{U}_{\mathcal{A},\omega}^{\mathfrak{g}}:\Omega_{\mathrm{LCC}\,X}\,\widehat{\otimes}_{\mathcal{O}_{\mathrm{LCC}\,X}}\,\pi_{\mathrm{lcc}}^{*}(\mathcal{P}_{X}\otimes_{\mathcal{O}_{X}}\mathcal{T}_{\mathrm{poly},X}) \\ \to \,\Omega_{\mathrm{LCC}\,X}\,\widehat{\otimes}_{\mathcal{O}_{\mathrm{LCC}\,X}}\,\pi_{\mathrm{lcc}}^{*}(\mathcal{P}_{X}\otimes_{\mathcal{O}_{X}}\mathcal{D}_{\mathrm{poly},X}). \end{array}$$

The DG Lie algebras in formula (1.9) also have filtrations $\{G^j\}_{j\in\mathbb{Z}}$, the homomorphism $\mathcal{U}_{\mathcal{A},\omega;1}^{\mathfrak{g}}$ respects this filtration, and we now have

(1.10)
$$\operatorname{gr}_{G}(\mathcal{U}_{\mathcal{A},\omega;1}^{\mathfrak{g}}) = \operatorname{gr}_{G}(\mathcal{U}_{\mathcal{A};1}^{\mathfrak{g}}) = \mathbf{1} \otimes \pi_{\operatorname{lcc}}^{*}(\mathbf{1} \otimes \mathcal{U}_{1}).$$

According to [Ye1, Theorem 6.4] there are induced operators

$$\Psi_{\boldsymbol{\sigma};j} := \boldsymbol{\sigma}^*(\mathcal{U}_{\mathcal{A},\omega;j}^{\mathfrak{g}}) : \prod^{J} \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\operatorname{poly},X}) \to \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X})$$

for $j \geq 1$. The L_{∞} identities in [Ye1, Definition 3.7], when applied to the L_{∞} morphism $\mathcal{U}^{\mathfrak{g}}_{\mathcal{A},\omega}$, are of the form considered in [Ye1, Theorem 6.4(iii)]. Therefore these

identities are preserved by $\boldsymbol{\sigma}^*$, and we conclude that the sequence $\Psi_{\boldsymbol{\sigma}} = \{\Psi_{\boldsymbol{\sigma},j}\}_{j=1}^{\infty}$ is an L_{∞} morphism. Furthermore, $\Psi_{\boldsymbol{\sigma};1}$ respects the filtration $\{\mathbf{G}^i \operatorname{Mix}_{\boldsymbol{U}}\}$, and from (1.10) we get

(1.11)
$$\operatorname{gr}_{G}(\Psi_{\boldsymbol{\sigma};1}) = \operatorname{gr}_{G}(\boldsymbol{\sigma}^{*}(\mathcal{U}_{\mathcal{A};1}^{\mathfrak{g}})) = \operatorname{gr}_{G}(\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{U}_{1})).$$

According to [Ye2, Theorem 4.17] the homomorphism $\operatorname{gr}_{G}(\operatorname{Mix}_{U}(\mathcal{U}_{1}))$ is a quasiisomorphism. Since the complexes $\operatorname{Mix}_{U}(\mathcal{T}_{\operatorname{poly},X})$ and $\operatorname{Mix}_{U}(\mathcal{D}_{\operatorname{poly},X})$ are bounded below, and the filtration is nonnegative and exhaustive, it follows that $\Psi_{\sigma;1}$ is also a quasi-isomorphism.

Corollary 1.12. Taking global sections in Theorem 1.2 we get an L_∞ quasi-isomorphism

$$\Gamma(X, \Psi_{\boldsymbol{\sigma}}) = \{ \Gamma(X, \Psi_{\boldsymbol{\sigma};j}) \}_{j \ge 1} : \Gamma(X, \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\operatorname{poly},X})) \to \Gamma(X, \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X})).$$

Proof. Theorem 1.2 tells us that $\Psi_{\sigma,1}$ is a quasi-isomorphisms of complexes of sheaves. By [Ye1, Theorem 6.2] it follows that

$$\Gamma(X, \Psi_{\boldsymbol{\sigma};1}) : \Gamma(X, \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\operatorname{poly},X})) \to \Gamma(X, \operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X}))$$

is a quasi-isomorphism.

Corollary 1.13. The data (U, σ) induces a bijection

$$\mathrm{MC}(\Psi_{\boldsymbol{\sigma}}) : \mathrm{MC}\Big(\Gamma\big(X, \mathrm{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\mathrm{poly},X})\big)[[\hbar]]^+\Big) \\ \xrightarrow{\simeq} \mathrm{MC}\Big(\Gamma\big(X, \mathrm{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\mathrm{poly},X})\big)[[\hbar]]^+\Big).$$

Proof. Use Corollary 1.12 and [Ye1, Corollary 3.10].

Recall that $\mathcal{T}_{\text{poly}}(X) = \Gamma(X, \mathcal{T}_{\text{poly},X})$ and $\mathcal{D}_{\text{poly}}^{\text{nor}}(X) = \Gamma(X, \mathcal{D}_{\text{poly},X}^{\text{nor}})$; and the latter is the DG Lie algebra of global poly differential operators that vanish if one of their arguments is 1.

Suppose $f : X' \to X$ is an étale morphism. According to [Ye2, Prposition 4.6] there are DG Lie algebra homomorphisms $f^* : \mathcal{T}_{\text{poly}}(X) \to \mathcal{T}_{\text{poly}}(X')$ and $f^* : \mathcal{D}_{\text{poly}}^{\text{nor}}(X) \to \mathcal{D}_{\text{poly}}^{\text{nor}}(X')$. These homomorphisms extend to formal coefficients, and we get functions

$$\operatorname{MC}(f^*) : \operatorname{MC}(\mathcal{T}_{\operatorname{poly}}(X)[[\hbar]]^+) \to \operatorname{MC}(\mathcal{T}_{\operatorname{poly}}(X')[[\hbar]]^+)$$

etc.

One says that X is a \mathcal{D} -affine variety if $\mathrm{H}^q(X, \mathcal{M}) = 0$ for every quasi-coherent left \mathcal{D}_X -module \mathcal{M} and every q > 0.

Theorem 1.14. Let X be an irreducible smooth separated \mathbb{K} -scheme. Assume X is \mathcal{D} -affine. Then there is a canonical function

$$Q: \mathrm{MC}(\mathcal{T}_{\mathrm{poly}}(X)[[\hbar]]^+) \to \mathrm{MC}(\mathcal{D}_{\mathrm{poly}}^{\mathrm{nor}}(X)[[\hbar]]^+)$$

called the quantization map. It has the following properties:

- (i) The function Q preserves first order terms.
- (ii) The function Q respects étale morphisms. Namely if X' is another D-affine scheme, with quantization map Q', and if f : X' → X is an étale morphism, then

$$Q' \circ \mathrm{MC}(f^*) = \mathrm{MC}(f^*) \circ Q.$$

(iii) If X is affine, then Q is bijective.

AMNON YEKUTIELI

(iv) The function Q is characterized as follows. Choose an open covering U = $\{U_0, \ldots, U_m\}$ of X consisting of affine open sets, each admitting an étale coordinate system. Let σ be the associated simplicial section of the bundle LCC $X \to X$. Then there is a commutative diagram

$$\begin{array}{ccc}
\operatorname{MC}(\mathcal{T}_{\operatorname{poly}}(X)[[\hbar]]^+) & \xrightarrow{Q} & \operatorname{MC}(\mathcal{D}_{\operatorname{poly}}^{\operatorname{nor}}(X)[[\hbar]]^+) \\
\operatorname{MC}(\eta_{\mathcal{T}}) & & \operatorname{MC}(\eta_{\mathcal{D}})
\end{array}$$

 $\mathrm{MC}\big(\Gamma(X,\mathrm{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\mathrm{poly},X}))[[\hbar]]^+\big) \xrightarrow{\mathrm{MC}(\Psi_{\boldsymbol{\sigma}})} \mathrm{MC}\big(\Gamma(X,\mathrm{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\mathrm{poly},X}))[[\hbar]]^+\big)$

in which the arrows $MC(\Psi_{\sigma})$ and $MC(\eta_{\mathcal{D}})$ are bijections. Here Ψ_{σ} is the L_{∞} quasi-isomorphism from Theorem 1.2, and $\eta_{\mathcal{T}}, \eta_{\mathcal{D}}$ are the inclusions of DG Lie algebras.

Let's elaborate a bit on the statement above. It says that to any MC solution $\alpha = \sum_{j=1}^{\infty} \alpha_j \hbar^j \in \mathcal{T}^1_{\text{poly}}(X)[[\hbar]]^+$ there corresponds an MC solution $\beta =$ $\sum_{j=1}^{\infty} \beta_j \hbar^j \in \mathcal{D}_{poly}^{nor,1}(X)[[\hbar]]^+$. The element $\beta = Q(\alpha)$ is uniquely determined up to gauge equivalence by the group $\exp(\mathcal{D}_{poly}^{nor,0}(X)[[\hbar]]^+)$. Given any local sections $f, g \in \mathcal{O}_X$ one has

(1.15)
$$\frac{1}{2}(\beta_1(f,g) - \beta_1(g,f)) = \alpha_1(f,g) \in \mathcal{O}_X.$$

The quantization map Q can be calculated (at least in theory) using the collection of sections σ and the universal formulas for deformation in [Ye1, Theorem 3.13].

We'll need a lemma before proving the theorem.

Lemma 1.16. Let $f, g \in \mathcal{O}_X = \mathcal{D}_{\text{poly},X}^{-1}$ be local sections.

(1) For any $\beta \in \operatorname{Mix}^0_{U}(\mathcal{D}^1_{\operatorname{poly},X})$ one has

 $[[\beta, f], q] = \beta(q, f) - \beta(f, q) \in \operatorname{Mix}^{0}_{U}(\mathcal{O}_{X}).$

- (2) For any $\beta \in \operatorname{Mix}^{1}_{U}(\mathcal{D}^{0}_{\operatorname{poly},X}) \oplus \operatorname{Mix}^{2}_{U}(\mathcal{D}^{-1}_{\operatorname{poly},X})$ one has $[[\beta, f], g] = 0$. (3) Let $\gamma \in \operatorname{Mix}_{U}(\mathcal{D}_{\operatorname{poly},X})^{0}$, and define $\beta := (\operatorname{d_{mix}} + \operatorname{d_{\mathcal{D}}})(\gamma)$. Then $[[\beta, f], g] = 0$.

Proof. (1) [Ye1, Proposition 6.3] implies that the embedding ([Ye1, (6.1)]:

$$\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\operatorname{poly},X}) \\ \subset \bigoplus_{p,q,r} \prod_{j \in \mathbb{N}} \prod_{\boldsymbol{i} \in \boldsymbol{\Delta}_{j}^{m}} g_{\boldsymbol{i}*} g_{\boldsymbol{i}}^{-1} \left(\Omega^{q}(\boldsymbol{\Delta}_{\mathbb{K}}^{j}) \widehat{\otimes} \left(\Omega_{X}^{p} \otimes_{\mathcal{O}_{X}} \mathcal{P}_{X} \otimes_{\mathcal{O}_{X}} \mathcal{D}_{\operatorname{poly},X}^{r} \right) \right)$$

is a DG Lie algebra homomorphism. So by continuity we might as well assume that $\beta = aD$ with $a \in \Omega^0_X = \mathcal{O}_X$ and $D \in \mathcal{D}^1_{\text{poly},X}$. Moreover, since the Lie bracket of $\Omega_X \otimes_{\mathcal{O}_X} \mathcal{P}_X \otimes_{\mathcal{O}_X} \mathcal{D}_{\mathrm{poly},X}$ is Ω_X -bilinear, we may assume that a = 1, i.e. $\beta = D$. Now the assertion is clear from the definition of the Gerstenhaber Lie bracket, see [Ko, Section 3.4.2].

(2) Applying the same reduction as above, but with $D \in \mathcal{D}^r_{\text{poly}, X}$ and $r \in \{0, -1\}$, we get $[[D, f], g] \in \mathcal{D}_{\text{poly}, X}^{r-2} = 0.$

(3) By part (2) it suffices to show that $[[\beta, f], g] = 0$ for $\beta := d_{\mathcal{D}}(\gamma)$ and $\gamma \in$ $\operatorname{Mix}^0_{U}(\mathcal{D}^0_{\operatorname{poly},X})$. As explained above we may further assume that $\gamma = D \in \mathcal{D}^0_{\operatorname{poly},X}$. Now the formulas for $d_{\mathcal{D}}$ and [-,-] in [Ko, Section 3.4.2] imply that $[[\mathrm{d}_{\mathcal{D}}(D), f], g] = 0.$

 $\mathbf{6}$

7

Proof of Theorem 1.14. Step 1. Take an open covering U as in property (iv). Since the sheaves $\mathcal{D}_{\text{poly},X}^{\text{nor},p}$ are quasi-coherent left \mathcal{D}_X -modules, it follows that $\mathrm{H}^q(X, \mathcal{D}_{\text{poly},X}^{\text{nor},p}) = 0$ for all p and all q > 0. Therefore $\Gamma(X, \mathcal{D}_{\text{poly},X}^{\text{nor}}) = \mathrm{R}\Gamma(X, \mathcal{D}_{\text{poly},X}^{\text{nor}})$ in the derived category $\mathsf{D}(\mathsf{Mod}\,\mathbb{K})$. Now by [Ye1, Theorem 3.12] the inclusion $\mathcal{D}_{\text{poly},X}^{\text{nor}} \to \mathcal{D}_{\text{poly},X}$ is a quasi-isomorphism, and by [Ye1, Theorem 6.2(1)] the inclusion $\mathcal{D}_{\text{poly},X} \to \mathrm{Mix}_U(\mathcal{D}_{\text{poly},X})$ is a quasi-isomorphism. According to [Ye1, Theorem 6.2(2)] we have $\Gamma(X, \mathrm{Mix}_U(\mathcal{D}_{\text{poly},X})) = \mathrm{R}\Gamma(X, \mathrm{Mix}_U(\mathcal{D}_{\text{poly},X}))$. The conclusion is that

(1.17)
$$\mathcal{D}_{\text{poly}}^{\text{nor}}(X) = \Gamma(X, \mathcal{D}_{\text{poly}, X}^{\text{nor}}) \to \Gamma(X, \text{Mix}_{U}(\mathcal{D}_{\text{poly}, X}))$$

is a quasi-isomorphism of complexes of K-modules. But in view of [Ye1, Proposition 6.3], this is in fact a quasi-isomorphism of DG Lie algebras.

From (1.17) we deduce that

$$\eta_{\mathcal{D}}: \mathcal{D}_{\text{poly}}^{\text{nor}}(X)[[\hbar]]^+ \to \Gamma(X, \text{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\text{poly},X}))[[\hbar]]^+$$

is a quasi-isomorphism of DG Lie algebras. Using [Ye1, Corollary 3.10] we see that $MC(\eta_D)$ is bijective. Therefore the diagram in property (iv) defines Q uniquely.

According to Corollary 1.13, the arrow marked $MC(\Psi_{\sigma})$ is a bijection. So we have established property (iv), except for the independence of the open covering.

Step 2. The left vertical arrow comes from the DG Lie algebra homomorphism

$$\eta_{\mathcal{T}}: \mathcal{T}_{\text{poly}}(X)[[\hbar]]^+ \to \Gamma(X, \operatorname{Mix}_{U}(\mathcal{T}_{\operatorname{poly},X}))[[\hbar]]^+,$$

which is a quasi-isomorphism when $\mathrm{H}^{q}(X, \mathcal{T}^{p}_{\mathrm{poly}, X}) = 0$ for all p and all q > 0. So in case X is affine, the quantization map Q is bijective. This establishes property (iii).

Step 3. Now suppose $U' = \{U'_0, \ldots, U'_{m'}\}$ is another such affine open covering of X, with sections $\sigma'_i : U'_i \to \text{LCC } X$. Without loss of generality we may assume that $m' \ge m$, and that $U'_i = U_i$ and $\sigma'_i = \sigma_i$ for all $i \le m$. There is a morphism of simplicial schemes $f : U \to U'$, that is an open and closed embedding. Correspondingly there is a commutative diagram

$$\begin{array}{ccc} \operatorname{MC}(\mathcal{T}_{\operatorname{poly}}(X)[[\hbar]]^{+}) & \xrightarrow{Q} & \operatorname{MC}(\mathcal{D}_{\operatorname{poly}}^{\operatorname{nor}}(X)[[\hbar]]^{+}) \\ & & \operatorname{MC}(\eta_{\mathcal{T}}) \downarrow & & \operatorname{MC}(\eta_{\mathcal{D}}) \downarrow \\ \end{array} \\ \operatorname{MC}(\Gamma(X,\operatorname{Mix}_{U'}(\mathcal{T}_{\operatorname{poly},X}))[[\hbar]]^{+}) & \xrightarrow{\operatorname{MC}(\Psi_{\sigma'})} & \operatorname{MC}(\Gamma(X,\operatorname{Mix}_{U'}(\mathcal{D}_{\operatorname{poly},X}))[[\hbar]]^{+}) \\ & & \operatorname{MC}(f^{*}) \downarrow & & & \operatorname{MC}(f^{*}) \downarrow \\ \end{array} \\ \operatorname{MC}(\Gamma(X,\operatorname{Mix}_{U}(\mathcal{T}_{\operatorname{poly},X}))[[\hbar]]^{+}) & \xrightarrow{\operatorname{MC}(\Psi_{\sigma})} & \operatorname{MC}(\Gamma(X,\operatorname{Mix}_{U}(\mathcal{D}_{\operatorname{poly},X}))[[\hbar]]^{+}), \end{array}$$

where the vertical arrows on the right are bijections. We conclude that Q is independent of U and σ . This concludes the proof of property (iv).

Step 4. Suppose $f : X' \to X$ is an étale morphism. Then we can choose an affine open covering U' of X' that refines U in the obvious sense. Each of the open sets U'_i inherits an étale coordinate system, and hence a section $\sigma'_i : U'_i \to \text{LCC } X'$. We

get a commutative diagram

$$\begin{array}{ccc} \mathrm{MC}\big(\Gamma(X,\mathrm{Mix}_{\boldsymbol{U}}(\mathcal{T}_{\mathrm{poly},X}))[[\hbar]]^+\big) & \xrightarrow{\mathrm{MC}(\Psi_{\boldsymbol{\sigma}})} & \mathrm{MC}\big(\Gamma(X,\mathrm{Mix}_{\boldsymbol{U}}(\mathcal{D}_{\mathrm{poly},X}))[[\hbar]]^+\big) \\ & & & & \\ \mathrm{MC}(f^*) \Big\downarrow & & & \\ \mathrm{MC}\big(\Gamma(X',\mathrm{Mix}_{\boldsymbol{U}'}(\mathcal{T}_{\mathrm{poly},X'}))[[\hbar]]^+\big) & \xrightarrow{\mathrm{MC}(\Psi_{\boldsymbol{\sigma}'})} & \mathrm{MC}\big(\Gamma(X',\mathrm{Mix}_{\boldsymbol{U}'}(\mathcal{D}_{\mathrm{poly},X'}))[[\hbar]]^+\big), \end{array}$$

This proves property (ii).

Step 5. Finally we must show that Q preserves first order terms, i.e. property (i). Let

$$\alpha = \sum_{j=1}^{\infty} \alpha_j \hbar^j \in \mathcal{T}_{\text{poly}}(X)^1[[\hbar]]^+$$

be an MC solution, and let

$$\beta = \sum_{j=1}^{\infty} \beta_j \hbar^j \in \mathcal{D}_{\text{poly}}^{\text{nor}}(X)^1[[\hbar]]^+$$

be an MC solution such that $\beta = Q(\alpha)$ modulo gauge equivalence. This means that there exists some

$$\gamma = \sum_{k \ge 1} \gamma_k \hbar^k \in \Gamma(X, \operatorname{Mix}_U(\mathcal{D}_{\operatorname{poly}, X}))^0[[\hbar]]^+$$

such that

$$\sum_{j\geq 1} \frac{1}{j!} \Psi_{\boldsymbol{\sigma};j}(\alpha^j) = \exp(\mathrm{af})(\exp(\gamma))(\beta),$$

with notation as in [Ye1, Lemma 3.2]. Cf. [Ye1, Theorem 3.8]. In the first order term (i.e. the coefficient of \hbar^1) of this equation we have

(1.18)
$$\Psi_{\boldsymbol{\sigma};1}(\alpha_1) = \beta_1 - (\mathbf{d}_{\min} + \mathbf{d}_{\mathcal{D}})(\gamma_1);$$

see [Ye1, equation (3.3)].

In order to apply Lemma 1.16(2), we are interested in in the component of $\Psi_{\sigma;1}(\alpha_1)$ living in the summand $\operatorname{Mix}^0_U(\mathcal{D}^1_{\operatorname{poly},X})$. But this is exactly

$$\operatorname{gr}_{G}^{0}(\Psi_{\sigma;1})(\alpha_{1}) \in \operatorname{gr}_{G}^{0}\operatorname{Mix}_{U}(\mathcal{D}_{\operatorname{poly},X}^{1}) = \operatorname{Mix}_{U}^{0}(\mathcal{D}_{\operatorname{poly},X}^{1})$$

Since according to Theorem 1.2 we have

$$\operatorname{gr}_{G}^{0}(\Psi_{\boldsymbol{\sigma};1}) = \operatorname{gr}_{G}^{0}(\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{U}_{1})),$$

it follows that the component we are interested in is

$$\operatorname{gr}_{G}^{0}(\operatorname{Mix}_{\boldsymbol{U}}(\mathcal{U}_{1}))(\alpha_{1}) = \mathcal{U}_{1}(\alpha_{1}).$$

Now take any two local sections $f, g \in \mathcal{O}_X$. Using Lemma 1.16 we get

$$\begin{split} [\Psi_{\sigma,1}(\alpha_1), f], g] &= [[\mathcal{U}_1(\alpha_1), f], g] = \mathcal{U}_1(\alpha_1)(g, f) - \mathcal{U}_1(\alpha_1)(f, g) = -2\alpha_1(f, g), \\ [[\beta_1, f], g] &= \beta_1(g, f) - \beta_1(f, g) \end{split}$$

and

$$[[(\mathbf{d}_{\min} + \mathbf{d}_{\mathcal{D}})(\gamma_1), f], g] = 0.$$

Combining these equations with equation (1.18) we see that equation (1.15) indeed holds. So the proof is done.

9

Corollary 1.19. Let X be an irreducible smooth separated \mathbb{K} -scheme. Assume X is \mathcal{D} -affine. Then the quantization map Q of Theorem 1.14 may be interpreted as a canonical function

$$Q: \frac{\{\text{formal Poisson structures on } X\}}{\text{gauge equivalence}} \to \frac{\{\text{deformation quantizations of } \mathcal{O}_X\}}{\text{gauge equivalence}}$$

The quantization map Q preserves first order terms, and commutes with étale morphisms $f: X' \to X$. If X is affine then Q is bijective.

Proof. By definition the left side is $\mathrm{MC}(\mathcal{T}_{\mathrm{poly}}(X)[[\hbar]]^+)$. On the other hand, according to [Ye1, Theorem 1.13] every deformation quantization of \mathcal{O}_X can be trivialized globally, and by [Ye1, Proposition 1.14] any gauge equivalence between globally trivialized deformation quantizations is a global gauge equivalence. Hence the right side is $\mathrm{MC}(\mathcal{D}_{\mathrm{poly}}^{\mathrm{nor}}(X)[[\hbar]]^+)$.

2. Miscellaneous Errors

Here is a list of minor errors in the paper [Ye1].

(1) Section 3, bottom of page 395: the formula should be

$$\operatorname{af}(\gamma)(\omega) := [\gamma, \omega] - \operatorname{d}(\gamma) = \operatorname{ad}(\gamma)(\omega) - \operatorname{d}(\gamma) \in \mathfrak{m} \widehat{\otimes} \mathfrak{g}^1,$$

(2) Definition 5.2, page 411: the formula should be

$$\nabla_{\mathcal{P}}: \mathcal{P}_X \to \Omega^1_X \otimes_{\mathcal{O}_X} \mathcal{P}_X.$$

References

- [CKTB] A. Cattaneo, B. Keller, C. Torossian and A. Bruguieres, "Déformation, Quantification, Théory de Lie", Panoramas et Synthèses 20 (2005), Soc. Math. France.
- [Ko] M. Kontsevich, Deformation quantization of Poisson manifolds, Lett. Math. Phys. 66 (2003), no. 3, 157-216.
- [VdB] M. Van den Bergh, On global deformation quantization in the algebraic case, J. Algebra 315 (2007), 326-395.
- [Ye1] A. Yekutieli, Deformation Quantization in Algebraic Geometry, Adv. Math. 198 (2005), 383-432.
- [Ye2] A. Yekutieli, Mixed Resolutions and Simplicial Sections, to appear in Israel J. Math. Eprint math.AG/0502206 at http://arxiv.org.
- [Ye3] A. Yekutieli, Continuous and Twisted L-infinity Morphisms, J. Pure Appl. Algebra 207 (2006), 575-606.
- [Ye4] A. Yekutieli, An Averaging Process for Unipotent Group Actions, Representation Theory 10 (2006), 147-157.

A. YEKUTIELI: DEPARTMENT OF MATHEMATICS BEN GURION UNIVERSITY, BE'ER SHEVA 84105, ISRAEL

E-mail address: amyekut@math.bgu.ac.il