

# Variational quantum Monte Carlo simulations with tensor-network states

A. W. Sandvik<sup>1,2,3</sup> and G. Vidal<sup>4</sup>

<sup>1</sup>*Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215*

<sup>2</sup>*Center for Theoretical Sciences, National Taiwan University, Taipei, Taiwan 106*

<sup>3</sup>*Department of Physics, National Taiwan University, Taipei, Taiwan 106*

<sup>4</sup>*School of Physical Sciences, The University of Queensland, QLD 4072, Australia*

(Dated: February 1, 2019)

We show that the formalism of tensor-network states, such as the matrix product states (MPS), can be used as a basis for variational quantum Monte Carlo simulations. Using a stochastic optimization method, we demonstrate the potential of this approach by explicit MPS calculations for the transverse Ising chain with up to  $N = 256$  spins at criticality, using periodic boundary conditions and  $D \times D$  matrices with  $D$  up to 32. The computational cost of our scheme formally scales as  $ND^3$ , whereas standard MPS approaches and the related density matrix renormalization group method scale as  $ND^5$  and  $ND^6$ , respectively, for periodic systems.

PACS numbers: 03.75.Mn, 03.75.Hh, 05.30.Fk, 05.10.Ln

Devising unbiased computational methods for correlated quantum many-body systems remains one of the greatest challenges in theoretical physics. Considerable progress has been made in recent years. Quantum Monte Carlo (QMC) methods with efficient loop-cluster updates [1, 2, 3] now enable simulations of certain classes of spin and boson hamiltonians on very large lattices—up to  $10^4$  sites essentially in the ground state and considerably more at elevated temperatures. Modern projector QMC methods [4] can also access large lattices. Both approaches are already contributing significantly to forefront areas of condensed matter physics, e.g., studies of exotic quantum phase transitions in antiferromagnets [5]. However, due to “sign problems” [6, 7], most fermion systems in more than one dimension and spin models with frustrated interactions are intractable to QMC simulations. The density matrix renormalization group (DMRG) method [8, 9], on the other hand, can produce essentially exact results for one-dimensional fermion systems and frustrated spins, including systems of a few coupled chains (ladders) [10]. These calculations are often restricted to open boundary conditions, however, which some times can be problematic. A more severe limitation is the exponential scaling in the computational complexity for systems with two or more dimensions [11].

The underlying reason for the problems with DMRG in higher dimensions has recently been identified as the inability of matrix product states (MPS), which are produced by the DMRG method [12], to properly account for entanglement in dimensions higher than one [13]. In order to overcome this limitation, a generalization of the MPS was proposed—the projected-entangled pair states (PEPS) [14]. These states are based on tensor-product networks, which can be contracted using an approximate scheme which has been argued to be adequate for typical many-body hamiltonians, even at quantum-critical points. While this approach is very promising, practical applications are still hampered by the severe increase of

the computational effort with the size  $D$  of the tensors in two dimensions. The scaling is typically  $\sim D^{10}$ , and calculations are therefore currently restricted to very small  $D \sim 2 - 5$  [15, 16]. Developing schemes with a more favorable  $D$  scaling is therefore a high priority.

In principle MPS and PEPS can be used in variational QMC calculations. Sampling the physical states, instead of contracting the tensor network over those indices, formally reduces the scaling in  $D$  [17]. In practice, it is not clear how much can be achieved this way, however. An efficient method is required to optimize tensors with hundreds or thousands of independent parameters, based on noisy Monte Carlo estimates of the energy and its derivatives. In this Letter we demonstrate that such a program is actually feasible. We develop a method based on a stochastic optimization scheme [18] which requires only the first derivatives of the energy with respect to the parameters (the tensor elements). Here we focus on MPS for simplicity, but the scheme can be applied to more generic tensor networks, e.g., PEPS, as well. We test the method on the Ising chain in a transverse external field, defined by the hamiltonian,

$$H = - \sum_{i=1}^L (\sigma_i^z \sigma_{i+1}^z + h \sigma_i^x), \quad (1)$$

where  $\sigma^x$  and  $\sigma^z$  are the standard Pauli matrices. This system undergoes a quantum phase transition from a ground state with long-range Ising order in the  $z$  direction for  $h < 1$  to a state with disordered  $z$  components when  $h > 1$ . We here consider exclusively the computationally most challenging  $h = 1$  critical point.

For a periodic chain, a translationally invariant matrix-product state with momentum  $k = 0$  is of the form [12]

$$|\Psi\rangle = \sum_{\{s\}} \text{Tr}\{A(s_1)A(s_2)\cdots A(s_N)\} |s_1, s_2, \dots, s_N\rangle, \quad (2)$$

where the spins  $s_i = \pm 1$  are the eigenvalues of  $\sigma_i^z$  and  $A(\pm 1)$  are two  $D \times D$  matrices (for a non-translationally invariant system the matrices would be site dependent). We here take the matrices to be real and symmetric, which corresponds to a state which is symmetric with respect to the reflection  $s_i \rightarrow s_{N-i+1}$ . The ground state should also be invariant with respect to spin inversion;  $s_i \rightarrow -s_i$  for all  $i$ . A sufficient condition for this is that  $A(-1)$  and  $A(1)$  are related to each other by a transformation with some matrix  $U$  such that  $U^{-1}A(1)U = A(-1)$  and  $U^{-1}A(-1)U = A(1)$ , which implies  $U^2 = I$  (the identity matrix). For simplicity, and because of indications that a greater flexibility of the matrices is advantageous for the optimization, we here only enforce the weaker condition that  $A(1)$  and  $A(-1)$  have identical eigenvalues, using a scheme to be discussed below.

Our goal is to find the matrix elements  $a_{ij}^s$ ,  $s = \pm 1$ , that minimize the energy expectation value  $E = \langle H \rangle$  of the MPS (2). Using the notation  $W(S)$  for the wave function coefficient for state  $|S\rangle = |s_1, \dots, s_N\rangle$ ,

$$W(S) = \text{Tr}\{A(s_1)A(s_2)\cdots A(s_N)\}, \quad (3)$$

the energy, for given matrices  $A(\pm 1)$ , can be written in the form appropriate for Monte Carlo sampling;

$$E = \frac{1}{Z} \sum_S W^2(S) E(S), \quad Z = \sum_S W^2(S), \quad (4)$$

where  $E(S)$  is the estimator

$$E(S) = \sum_{S'} \frac{W(S')}{W(S)} \langle S' | H | S \rangle. \quad (5)$$

Here the sum includes  $S' = S$  and the  $N$  states obtained by flipping one spin in  $|S\rangle$ . The energy can be evaluated using importance sampling of the spin configurations according to the weight  $W^2(S)$ ;  $E = \langle E(S) \rangle_{W^2}$ . Our scheme also requires the derivatives of the energy with respect to the matrix elements. These are given by

$$\frac{\partial E}{\partial a_{ij}^s} = 2\langle \Delta_{ij}^s(S) E(S) \rangle - 2\langle \Delta_{ij}^s(S) \rangle \langle E(S) \rangle, \quad (6)$$

where we have defined

$$\Delta_{ij}^s = \frac{1}{W(S)} \frac{\partial W(S)}{\partial a_{ij}^s}. \quad (7)$$

Introducing the matrices

$$B(m) = A(s_{m+1}) \cdots A(s_N) A(s_1) \cdots A(s_{m-1}), \quad (8)$$

the derivative of the weight (3) is

$$\frac{\partial W(S)}{\partial a_{ij}^s} = \frac{1}{1 + \delta_{ij}} \sum_{m=1}^N [B_{ij}(m) + B_{ji}(m)] \delta_{s, s_m}. \quad (9)$$

To sample the states over which (5) and (6) are averaged, we generate successive spin configurations from a stored  $S$  by flipping a single spin at a time;  $s_m \rightarrow -s_m$ . To indicate that spin  $m$  has been flipped we denote the new configuration  $S'_m$ . Visiting the spins sequentially;  $m = 1, 2, \dots, N$ , we flip them according to the Metropolis probability;  $P_{\text{flip}} = \min[W^2(S'_m)/W^2(S), 1]$ . To calculate this, we use the cyclic property of the trace and write the new coefficient as  $W(S'_m) = \text{Tr}\{A(-s_m)B(m)\}$ . Further, we write the matrix  $B(m)$  in Eq. (8) as a product of left and right matrices  $B(m) = L(m+1)R(m-1)$ , where  $L(m) = A(s_m) \cdots A(s_N)$  and  $R(m) = A(s_1) \cdots A(s_m)$ . Formally we also define  $L(N+1) = R(0) = I$ . Before starting the updating process we calculate and store the left matrices  $L(2), \dots, L(N)$ , based on the initial spin configuration (random or from a previous run). Having access to these matrices, each successive spin-flip attempt requires only one matrix multiplication, and another multiplication advances the right matrix;  $R(m) = R(m-1)A(s_m)$ . Since  $L(m)$  is no longer needed at this stage we store  $R(m)$  in its place for future use.

Diagonal quantities, like the Ising part of the energy,  $E_z = -\sum_i \sigma_i^z \sigma_{i+1}^z$ , can be simply obtained by averaging the appropriate spin correlations in the stored state  $|S\rangle$ . To calculate off-diagonal quantities, ratios  $W(S')/W(S)$  are needed. After a full sweep of spin updates, all the matrices  $R(m)$  have been generated and stored. We can use them to speedily measure the off-diagonal energy  $E_x = h \sum_i \langle \sigma_i^+ + \sigma_i^- \rangle$ , the estimator of which is

$$E_x(S) = h \sum_{m=1}^N \frac{W(S'_m)}{W(S)}, \quad (10)$$

as well as the derivatives (9). To evaluate the sums, we now traverse the system from  $m = N$  to 1, and in the process generate the left matrices  $L(m)$  and store them in the place of  $R(m)$ . Once this process is completed we again have what we need to carry out an updating sweep in the manner described above. A full updating sweep, including measurements, thus requires  $4N$  matrix multiplications (plus operations which have a lower scaling in  $D$ ), giving a formal scaling  $ND^3$  of the algorithm.

Carrying out successive simulations with fixed matrices  $A(\pm 1)$ , the energy and derivatives obtained on the basis of some number  $F$  of spin-flip sweeps (referred to as one simulation *bin*) are used to update the matrix elements with  $j \geq i$  according to (and subsequently  $a_{ji}^s = a_{ij}^s$ ) [18]:

$$a_{ij}^s \rightarrow a_{ij}^s - \delta(k) \cdot r_{ij}^s \cdot \text{sign}(\partial E / \partial a_{ij}^s). \quad (11)$$

Here  $r_{ij}^s \in [0, 1]$  is random and  $\delta(k)$  is the maximum change, which decreases as a function of a counter  $k = 0, 1, \dots$ . Thus, instead of moving in the direction of the approximately evaluated gradient, as in standard stochastic optimization [19, 20], each parameter is changed independently, using the “correct” sign but with a random and




FIG. 1: (Color online) Main panel: Convergence of the energy per site of a 16-site system at  $h = 1$ , using  $D = 8$  and starting from random matrices. The cooling parameters were  $Q = 0.9$ ,  $\delta_0 = 0.05$ ,  $G_0 = 10$ ,  $F_0 = 100$ . Inset: The later stages of the simulation on a more detailed scale, and a comparison with a run which started from a converged  $D = 6$  calculation (lower curve; red symbols); here  $\delta_0 = 0.005$ ,  $G_0 = 5$ ,  $F_0 = 50$ .

well bounded magnitude for the step. This results in a very stable optimization ideally suited for problems with large numbers of parameters. For the gradual reduction of  $\delta$ , we here use a geometric form;  $\delta = \delta_0 Q^k$ , with, typically,  $Q = 0.9 - 0.95$ , but other forms also work well, e.g.,  $\delta = \delta_0 k^{-\alpha}$ , with  $\alpha \in [1/2, 1]$ . For each  $k$ , we complete a number,  $G$ , of bins, each followed by updates of the matrix elements. The number of sweeps per bin,  $F$ , as well as  $G$  are increased with  $k$ . The rationale behind increasing  $F(k)$  is that, as we approach the energy minimum, the derivatives will become smaller and require more sampling in order not to be dominated by noise [21]. Increasing  $G$  leads effectively to a slower “cooling” rate. We typically use a linear dependence in both cases;  $F = F_0 k$ ,  $G = G_0 k$ . We output the energy and its statistical error computed on the basis of the  $G$  bins before each increment of  $k$ . Since  $F$  and  $G$  increase with  $k$ , the error bars will decrease. For a sufficiently long run, if the cooling is slow enough, the calculated  $E_D(k)$  should approach the optimal energy for a given matrix size  $D$ .

As we already mentioned, we wish to enforce the property that  $A(-1)$  and  $A(1)$  have the same eigenvalues. We do this after each adjustment of the matrix elements, by diagonalizing both matrices and averaging their eigenvalues. The averaged diagonal matrix is then transformed back using the diagonalizing matrices for the original  $A(\pm 1)$ . If we do not carry out this diagonalization step we still in practice do obtain matrices with approximately equal eigenvalue spectra. However, enforcing this condition exactly seems to have favorable effects on the ability of the optimization method to quickly converge to a spin-inversion invariant ground state. We normalize the matrices so that the largest element  $|a_{ij}^s| = 1$ .

It should be noted that the optimal matrices are not unique—there is a huge degeneracy in terms of simul-

TABLE I: Results for the critical transverse Ising chain of size  $N$  obtained with the variational MPS QMC approach (with the matrix size  $D$  shown) and the SSE method [22]. Statistical errors in the last displayed digits are indicated; in the case of the MPS energy the statistical errors are less than  $10^{-7}$ .

$N$	$D$	$E/N$ (MPS)	$E/N$ (SSE)	$M^2$ (MPS)	$M^2$ (SSE)
16	12	-1.2752872	-1.275297(9)	0.52236(3)	0.52231(2)
32	16	-1.2737510	-1.273763(7)	0.4408(2)	0.44082(3)
64	20	-1.2733673	-1.273367(6)	0.3720(6)	0.37344(4)
128	24	-1.2732714	-1.273277(7)	0.3120(9)	0.31286(8)
256	32	-1.2732469	-1.273235(5)	0.261(2)	0.26324(9)

taneous transformations of  $A(\pm 1)$  that leave the trace invariant. This may also be an advantage in the optimization, as we are not trying to locate a point, but only reach some large hypersurface in parameter space.

In Fig. 1 we show an example of the convergence of the optimization for a 16-site chain, using  $D = 8$  and starting from random  $A(\pm 1)$ . The initial maximum parameter shift was  $\delta_0 = 0.05$ . We compare with a run which started from matrices resulting from a calculation with  $D = 6$  (with the new matrix elements in the larger matrices generated at random in the range  $[-\delta_0, \delta_0]$ ), which allows for a smaller initial step  $\delta_0 = 0.005$ . The latter calculation produces a marginally lower energy, showing that the cooling rate in the former case was slightly too fast—cooling slower we obtain consistent results.

It is useful to start the optimization for some  $N$  and  $D$  from  $A(\pm 1)$  previously obtained for a smaller  $N$  and the same  $D$ , or the same  $N$  and smaller  $D$ . Another good strategy is to first do a short run with a large  $\delta_0 \approx 0.1$  to achieve convergence only approximately, and then to restart the calculation with a smaller  $\delta_0$  [but much larger than the smallest  $\delta(k)$  reached previously].

We do not claim that the cooling protocol presented above is optimal; further improvements could potentially lead to considerable efficiency gains. However, even as it stands now the scheme performs remarkably well.

When  $D$  is sufficiently large for almost exactly reproducing the true ground state energy, we observe a strong reduction of the variance in the energy  $E_D(k)$  as it converges (i.e., beyond what would be expected solely due to the increasing sampling times  $F$  and  $G$ ). For  $N = 8$  we can easily reproduce the exact energy to a relative precision of  $10^{-9}$  with  $D = 4$ . For larger  $N$ , we obtain excellent agreement with high-precision QMC results (using a stochastic series expansion, SSE, cluster method [22] at ultra-low temperatures). In Table I we summarize results for the energy as well as the squared magnetization;

$$M^2 = \left( \frac{1}{N} \sum_{i=1}^N \sigma_i^z \right)^2. \quad (12)$$

We show results for the largest  $D$  considered for each  $N$ ,




FIG. 2: (Color online) Relative error (deviation from our best converged results) in the energy and squared magnetization as a function of the matrix dimension  $D$ .

which are  $D \rightarrow \infty$  converged to within statistical errors.

The convergence with  $D$  is illustrated in Fig. 2, using as “exact” values for  $E$  and  $M^2$  either  $D$ -converged MPS or SSE results. In the case of the energy, a desired relative accuracy requires a  $D$  which eventually approaches a constant for large  $N$ . The squared magnetization is directly related to the long-distance physics, however, and our results are consistent with the expectation that  $D$  has to grow as some power,  $D \sim N^\alpha$ , to achieve a given relative accuracy. From Fig. 2 we obtain, roughly,  $\alpha \approx 0.65$ .

We have also carried out simulations with general non-symmetric matrices. In order to strictly enforce the lattice reflection and spin-inversion symmetries, we then use a wave function with a trace of four different matrix products related to each other by these symmetries, i.e.,

$$W(S) = \text{Tr}\{P(S) + P(S_R) + P(-S) + P(-S_R)\}, \quad (13)$$

where  $P(S) = A(s_1) \cdots A(s_N)$ , and  $-S$  and  $S_R$  are obtained by, respectively, spin-inverting and reflecting the configuration  $S$ . We find that this wave function has a lower optimum energy for given  $D$  than one with the single product  $P(S)$ . For small  $D$ , the energy is also better than for the symmetric matrices discussed above. However, the computational effort is higher by a factor of 4, and the convergence of the optimization scheme appears to be worse. In practice the pay-off of the larger number of parameters for given  $D$  is therefore questionable.

We have shown that it is feasible to stochastically optimize MPS with relatively large matrices. This demonstration makes it clear that the variational QMC approach can be successfully combined with the versatility of tensor-network states, for a sign-problem free and systematically refinable (through the tensor dimension  $D$ ) generic many-body method. The scaling with the matrix size  $D$  in the case of MPS for periodic chains is for-

mally reduced from  $D^5$  to  $D^3$ , and similar reductions are possible with tensor networks in higher dimensions [17]. There may of course be some further non-obvious  $D$  dependence in the convergence properties of the sampling and optimization schemes. For the Ising system studied here, critical slowing-down adversely affects the actual scaling in  $N$ . It should be noted, however, that other MPS schemes, as well as DMRG, also have convergence issues beyond the formal scaling in  $N$  and  $D$ .

At the late stages of completing this work we became aware of Ref. [23], where a different QMC approach is proposed in the same spirit and applied to PEPS.

AWS would like to thank Y.-J. Kao for stimulating discussions. This work was supported by the NSF under grant No. DMR-0513930 (AWS) and by the Australian Research Council grant No. FF0668731 (GV).

- 
- [1] H. G. Evertz, *Adv. Phys.* **52**, 1 (2003).
  - [2] N. V. Prokof'ev, B. V. Svistunov, and I. S. Tupitsyn, *Zh. Eks. Teor. Fiz.* **114**, 570 (1998) [*JETP* **87**, 311 (1998)].
  - [3] O. F. Syljuåsen and A. W. Sandvik, *Phys. Rev. E* **66**, 046701 (2002).
  - [4] A. W. Sandvik, *Phys. Rev. Lett.* **95**, 207203 (2005).
  - [5] A. W. Sandvik, *Phys. Rev. Lett.* **98**, 227202 (2007); R. G. Melko and R. K. Kaul, arXiv:0707.2961; K. Harada, N. Kawashima, and M. Troyer, arXiv:cond-mat/0608446.
  - [6] E. Y. Loh, J. E. Gubernatis, R. T. Scalettar, S. R. White, D. J. Scalapino, and R. L. Sugar, *Phys. Rev. B* **41**, 9301 (1990).
  - [7] P. Henelius and A. W. Sandvik, *Phys. Rev. B* **62**, 1102 (2000).
  - [8] S. R. White, *Phys. Rev. Lett.* **69**, 2863 (1992).
  - [9] U. Schollwöck, *Rev. Mod. Phys.* **77**, 259 (2005).
  - [10] S. R. White and D. J. Scalapino, *Phys. Rev. Lett.* **91**, 136403 (2003).
  - [11] S. Liang and H. Pang, *Phys. Rev. B* **49**, 9214 (1994).
  - [12] S. Östlund and S. Rommer, *Phys. Rev. Lett.* **75**, 3537 (1995).
  - [13] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, *Phys. Rev. Lett.* **90**, 227902 (2003).
  - [14] F. Verstraete and J. I. Cirac, arXiv:cond-mat/0407066.
  - [15] V. Murg, F. Verstraete, and J. I. Cirac, *Phys. Rev. A* **75**, 033605 (2007).
  - [16] J. Jordan, R. Orús, G. Vidal, F. Verstraete, and J. I. Cirac, arXiv:cond-mat/0703788.
  - [17] The scaling with MPS is reduced from  $D^3$  and  $D^5$  for open and periodic boundaries, respectively, to  $D^2$  and  $D^3$ . For PEPS with open boundaries the scaling goes from  $D^8 \tilde{D}^2 \approx D^{12}$  [15] to  $D^4 D'^2 + D^2 D'^3 \approx D^6$ , where  $\tilde{D} \approx D^2$  and  $D' \approx D$  are the ranks of the boundary MPS employed in the contraction. A reduction of several powers of  $D$  is also achieved in the case of PEPS with cylinder and torus boundary conditions.
  - [18] J. Lou and A. W. Sandvik, arXiv:cond-mat/0605034 (accepted for publication in *Phys. Rev. B*).
  - [19] H. Robbins and S. Monro, *Ann. Math. Stat.* **22**, 400 (1951); J. C. Spall, in *Wiley Encyclopedia of Electrical and Electronics Engineering*, Vol. 20, Edited by J. G.

- Webster (Wiley, 1999).
- [20] A. Harju, B. Barbiellini, S. Siljamäki, R. M. Nieminen, and G. Ortiz, Phys. Rev. Lett. **79**, 1173 (1997).
  - [21] Although the power of stochastic optimization partially lies in favorable effects of noise [19], there is clearly some limit beyond which too many errors in the signs of the derivatives will be detrimental.
  - [22] A. W. Sandvik, Phys. Rev. E **68**, 056701 (2003).
  - [23] N. Schuch, M. M. Wolf, F. Verstraete, and J. I. Cirac, arXiv:0708.1567.