

# QUANTUM $\mathfrak{gl}_\infty$ , INFINITE $q$ -SCHUR ALGEBRAS AND THEIR REPRESENTATIONS

JIE DU AND QIANG FU<sup>†</sup>

**ABSTRACT.** In this paper, we investigate the structure and representations of the quantum group  $\mathbf{U}(\infty) = \mathbf{U}_v(\mathfrak{gl}_\infty)$ . We will present a realization for  $\mathbf{U}(\infty)$ , following Beilinson–Lusztig–MacPherson (BLM) [1], and show that the natural algebra homomorphism  $\zeta_r$  from  $\mathbf{U}(\infty)$  to the infinite  $q$ -Schur algebra  $\mathcal{S}(\infty, r)$  is not surjective for any  $r \geq 1$ . We will give a BLM type realization for the image  $\mathbf{U}(\infty, r) := \zeta_r(\mathbf{U}(\infty))$  and discuss its presentation in terms of generators and relations. We further construct a certain completion algebra  $\widehat{\mathcal{K}}^\dagger(\infty)$  so that  $\zeta_r$  can be extended to an algebra epimorphism  $\tilde{\zeta}_r : \widehat{\mathcal{K}}^\dagger(\infty) \rightarrow \mathcal{S}(\infty, r)$ . Finally we will investigate the representation theory of  $\mathbf{U}(\infty)$ , especially the polynomial representations of  $\mathbf{U}(\infty)$ .

*Dedicated to Professor Leonard L. Scott on the occasion of his 65th birthday*

## 1. INTRODUCTION

The Lie algebra  $\mathfrak{gl}_\infty$  of infinite matrices and its extension  $A_\infty$  are interesting topics in the theory of infinite dimensional Lie algebras. Certain highest weight representations of these algebras have important applications in finding solutions of a large class of nonlinear equations (see, e.g., [3]) and in the representations theories of Heisenberg algebras, the Virasoro algebra and other Kac–Moody algebras (see, e.g., [21, 22, 32, 33]). The quantum versions of the corresponding universal enveloping algebras have also been studied in [24, 34, 35]. It should be noted that the structure and representations of quantum  $\mathfrak{gl}_\infty$  have various connections with the study of Lie superalgebras  $\mathfrak{gl}(m|n)$  (see, e.g., [2]) and the study of quantum affine  $\mathfrak{gl}_n$  (see [17, §6], [31, §2], [8] and [30]).

In this paper, we will investigate the quantum group  $\mathbf{U}(\infty) = \mathbf{U}_v(\mathfrak{gl}_\infty)$  and its representations through a series of its quotient algebras  $\mathbf{U}(\infty, r)$  and the infinite quantum Schur algebras  $\mathcal{S}(\infty, r)$ . The main idea is to extend the approach developed in [1] for quantum  $\mathfrak{gl}_n$  to quantum  $\mathfrak{gl}_\infty$ . Thus, we obtain a realization for  $\mathbf{U}(\infty)$  and an explicit description of the algebra homomorphism  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathbf{U}(\infty, r)$ . This in turn gives rise to a presentation and various useful bases for  $\mathbf{U}(\infty, r)$ . We will also prove that  $\mathbf{U}(\infty, r)$  is a proper subalgebra of  $\mathcal{S}(\infty, r)$ . This fact shows that the classical Schur–Weyl duality fails in this case. Finally, we investigate the ‘polynomial’ representation theory and classify all irreducible polynomial representations for  $\mathbf{U}(\infty)$ . We expect that this work will have further applications to quantum affine  $\mathfrak{gl}_n$ .

We organize the paper as follows. We recall the definition of quantum  $\mathfrak{gl}_\eta$  and  $q$ -Schur algebras  $\mathcal{S}(\eta, r)$  at any consecutive segment  $\eta$  of  $\mathbb{Z}$  in §2. In §3, we use the  $\eta$ -step flag variety to define the algebra  $\mathcal{K}(\eta, r)$  and discuss its stabilization property developed in [1] in a context suitable for infinite  $\eta$ . At the end of §3, we will focus on the infinite case  $\eta = \mathbb{Z}$ . First, the stabilization property allows us to define an algebra  $\mathcal{K}(\infty)$  over  $\mathbb{Q}(v)$  whose completion  $\widehat{\mathcal{K}}(\infty)$  contains a subalgebra  $\mathbf{V}(\infty)$  which is isomorphic to  $\mathbf{U}(\infty)$ . Second, there is a algebra homomorphism  $\xi_r : \mathbf{V}(\infty) \rightarrow \widehat{\mathcal{K}}(\infty, r)$

<sup>†</sup>Corresponding author.

Supported by the Australian Research Council (Grant: DP 0665124) and partially by the National Natural Science Foundation of China (10601037 & 10671142). The paper was written while the second author was visiting the University of New South Wales.

with image  $\mathbf{V}(\infty, r)$ . A Drinfeld–Jimbo type presentation for  $\mathbf{V}(\infty, r)$  will be given in §4. In §5, we will establish isomorphisms  $\mathcal{S}(\infty, r) \cong \widehat{\mathcal{K}}^\dagger(\infty, r)$  and between  $\mathbf{V}(\infty, r)$  and the homomorphic image  $\mathbf{U}(\infty, r)$  of the natural homomorphism  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathcal{S}(\infty, r)$ . Thus, we conclude that  $\zeta_r$  is not surjective for any  $r \geq 1$ . In §6, by identifying  $\mathcal{K}(\infty)$  with the modified quantum group  $\dot{\mathbf{U}}(\infty)$ , we derive an algebra epimorphism  $\zeta_r : \mathcal{K}(\infty) \rightarrow \mathcal{K}(\infty, r)$  and hence, extend the map  $\zeta_r$  to an epimorphism from  $\widehat{\mathcal{K}}^\dagger(\infty)$  to  $\mathcal{S}(\infty, r)$ . In the last three sections, we investigate the representation theory of  $\mathbf{U}(\infty)$ . The highest weight representations of  $\mathbf{U}(\infty)$  is studied in §7, and the polynomial representations of  $\mathbf{U}(\infty)$  is in §9 as an application of the representation theory of  $\mathcal{S}(\infty, r)$  investigated in §8.

Throughout the paper, we will encounter several (associative) algebras  $\mathcal{A}$  over a commutative ring  $\mathcal{R}$  without the identity element, but with many orthogonal idempotents  $e_i$ ,  $i \in I$ , such that  $\mathcal{A} = \bigoplus_{i, i' \in I} e_i \mathcal{A} e_{i'}$  and, for all  $i, i' \in I$ ,  $e_i \mathcal{A} e_{i'}$  are free over  $\mathcal{R}$ . Clearly, the index set  $I$  must be an infinite set, since if  $I$  was a finite set, then  $\sum_{i \in I} e_i$  would be the identity element of  $\mathcal{A}$ .

Choose an  $\mathcal{R}$ -basis  $\mathcal{B} = \{a_j\}_{j \in J}$  such that  $e_i \mathcal{B} e_{i'} = \{e_i a_j e_{i'}\}_{j \in J} \setminus \{0\}$  is a basis for  $e_i \mathcal{A} e_{i'}$  for all  $i, i' \in I$ , and  $\mathcal{B} = \cup_{i, i' \in I} e_i \mathcal{B} e_{i'}$ . We further assume that  $e_i \in \mathcal{B}$  for all  $i \in I$ . It is clear that, for any  $j \in J$ , there exist unique  $i_j, i'_j \in I$  such that  $e_{i_j} a_j = a_j$  and  $a_j e_{i'_j} = a_j$ . We will write  $ro(j) := i_j$  and  $co(j) := i'_j$  for all  $j \in J$ .

For a formal infinite linear combination  $f = \sum_{j \in J} f_j a_j$  with  $f_j \in \mathcal{R}$ , let, for any  $i \in I$ ,  $J(e_i, f) := \{j \in J \mid f_j \neq 0, e_i a_j = a_j\} = \{j \in J \mid f_j \neq 0, ro(j) = i\}$  and  $J(f, e_i) := \{j \in J \mid f_j \neq 0, a_j e_i = a_j\} = \{j \in J \mid f_j \neq 0, co(j) = i\}$ . One can easily check the following lemma.

**Lemma 1.1.** *Let  $\mathfrak{L}$  be the set<sup>1</sup> of formal (possibly infinite) linear combinations of  $\mathcal{B}$ , and let*

$$\begin{aligned} \dagger \widehat{\mathcal{A}} &:= \{f \in \mathfrak{L} \mid \forall i \in I, |J(e_i, f)| < \infty\}, & \widehat{\mathcal{A}}^\dagger &:= \{f \in \mathfrak{L} \mid \forall i \in I, |J(f, e_i)| < \infty\}, \\ \widehat{\mathcal{A}} &:= \{f \in \mathfrak{L} \mid \forall i, i' \in I, |J(e_i, f)| < \infty, |J(f, e_{i'})| < \infty\} = \dagger \widehat{\mathcal{A}} \cap \widehat{\mathcal{A}}^\dagger. \end{aligned}$$

We define the product of two elements  $\sum_{s \in J} f_s a_s, \sum_{t \in J} g_t a_t$  in  $\widehat{\mathcal{A}}^\dagger$  (resp.,  $\dagger \widehat{\mathcal{A}}$ ) to be  $\sum_{s, t \in J} f_s g_t a_s a_t$  where  $a_s a_t$  is the product in  $\mathcal{A}$ . Then  $\dagger \widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}^\dagger$  become associative algebras with identity  $1 = \sum_{i \in I} e_i$  and  $\widehat{\mathcal{A}}$  is a subalgebra with the same identity.

The algebras  $\dagger \widehat{\mathcal{A}}, \widehat{\mathcal{A}}^\dagger$  and  $\widehat{\mathcal{A}}$  are called the *completion algebras* of  $\mathcal{A}$ .

Note that, if  $\mathcal{A}$  admits an algebra antiautomorphism  $f$  satisfying  $f(e_i) = e_i$  for all  $i \in I$ , then  $\dagger \widehat{\mathcal{A}} \cong (\widehat{\mathcal{A}}^\dagger)^{\text{op}}$ , the algebras with the same underlying space as  $\widehat{\mathcal{A}}^\dagger$  but opposite multiplication.

**Some notations and conventions.** A consecutive segment  $\eta$  of  $\mathbb{Z}$  is either a finite interval of the form  $[m, n] := \{i \in \mathbb{Z} \mid m \leq i \leq n\}$  or an infinite interval of the form  $(-\infty, m], [n, +\infty)$  and  $(-\infty, \infty) = \mathbb{Z}$ , where  $m, n \in \mathbb{Z}$ . Let

$$\eta^\dagger = \begin{cases} \eta \setminus \{\eta_{\max}\}, & \text{if there is a maximum element } \eta_{\max} \text{ in } \eta; \\ \eta, & \text{otherwise.} \end{cases}$$

<sup>1</sup>We may identify the set  $\mathfrak{L}$  with the direct product  $\prod_{j \in J} \mathcal{R} a_j$ .

Let  $M_\eta(\mathbb{Z})$  (resp.  $\mathbb{Z}^\eta$ ) be the set of all matrices  $(a_{i,j})_{i,j \in \eta}$  (resp. all sequences  $(a_i)_{i \in \eta}$ ) over  $\mathbb{Z}$  with *finite support* if  $\eta$  is infinite. We will use the following index sets throughout the paper.

$$\begin{aligned}
 \tilde{\Xi}(\eta) &= \{(a_{ij}) \in M_\eta(\mathbb{Z}) \mid a_{ij} \geq 0 \ \forall i \neq j \text{ in } \eta\} = \Xi^-(\eta) + \tilde{\Xi}^0(\eta) + \Xi^+(\eta), \\
 \Xi(\eta) &= \{(a_{ij}) \in M_\eta(\mathbb{Z}) \mid a_{ij} \geq 0 \ \forall i, j \in \eta\} = \Xi^-(\eta) + \Xi^0(\eta) + \Xi^+(\eta), \\
 \Xi^\pm(\eta) &= \{(a_{ij}) \in M_\eta(\mathbb{Z}) \mid a_{ii} = 0 \ \forall i \in \eta\} = \Xi^-(\eta) + \Xi^+(\eta), \\
 \Xi(\eta, r) &= \{A \in \Xi(\eta) \mid \sigma(A) := \sum_{i,j \in \eta} a_{ij} = r\}, \quad \text{and} \\
 \Lambda(\eta, r) &= \{\lambda \in \mathbb{N}^\eta \mid \sigma(\lambda) := \sum_{i \in \eta} \lambda_i = r\}
 \end{aligned}
 \tag{1.1.1}$$

where  $\Xi^+(\eta)$  (resp.,  $\Xi^-(\eta)$ ,  $\tilde{\Xi}^0(\eta)$ ) is the subset of  $\tilde{\Xi}(\eta)$  consisting of those matrices  $(a_{ij})$  with  $a_{ij} = 0$  for all  $i \geq j$  (resp.,  $i \leq j$ ,  $i \neq j$ ),  $\Xi(\eta) = \tilde{\Xi}^0(\eta) \cap \Xi(\eta)$ , and  $\mathbb{N}^\eta \subset \mathbb{Z}^\eta$ . The sum  $\Xi^-(\eta) + \tilde{\Xi}^0(\eta) + \Xi^+(\eta)$  indicates the following matrix decomposition: for  $A \in \tilde{\Xi}(\eta)$ ,  $A = A^+ + A^0 + A^-$  with  $A^+ \in \Xi^+(\eta)$ ,  $A^- \in \Xi^-(\eta)$ ,  $A^0 \in \tilde{\Xi}^0(\eta)$ . We also write  $A^\pm = A^+ + A^- \in \Xi^\pm(\eta)$ . Also, for  $A = (a_{ij}) \in M_\eta(\mathbb{N})$ , define

$$\deg(A) := \sum_{i,j \in \eta} |j - i| a_{ij}.
 \tag{1.1.2}$$

Throughout the paper,  $v$  denotes an indeterminate and  $\mathbb{Q}(v)$  denotes the fraction field of the integral Laurent polynomial ring  $\mathcal{Z} := \mathbb{Z}[v, v^{-1}]$ .

## 2. QUANTUM $\mathfrak{gl}_\eta$ AND $q$ -SCHUR ALGEBRAS AT $\eta$

Let  $\eta$  be a fixed consecutive segment of  $\mathbb{Z}$ .

**Definition 2.1.** The quantum  $\mathfrak{gl}_\eta$  over  $\mathbb{Q}(v)$  is the  $\mathbb{Q}(v)$ -algebra  $\mathbf{U}(\eta) := \mathbf{U}_v(\mathfrak{gl}_\eta)$  presented by generators

$$E_i, F_i \quad (i \in \eta^{-1}), \quad K_j, K_j^{-1} \quad (j \in \eta)$$

and relations

- (a)  $K_i K_j = K_j K_i$ ,  $K_i K_i^{-1} = 1$ ;
- (b)  $K_i E_j = v^{\delta_{i,j} - \delta_{i,j+1}} E_j K_i$ ;
- (c)  $K_i F_j = v^{\delta_{i,j+1} - \delta_{i,j}} F_j K_i$ ;
- (d)  $E_i E_j = E_j E_i$ ,  $F_i F_j = F_j F_i$  when  $|i - j| > 1$ ;
- (e)  $E_i F_j - F_j E_i = \delta_{i,j} \frac{\tilde{K}_i - \tilde{K}_i^{-1}}{v - v^{-1}}$ , where  $\tilde{K}_i = K_i K_{i+1}^{-1}$ ;
- (f)  $E_i^2 E_j - (v + v^{-1}) E_i E_j E_i + E_j E_i^2 = 0$  when  $|i - j| = 1$ ;
- (g)  $F_i^2 F_j - (v + v^{-1}) F_i F_j F_i + F_j F_i^2 = 0$  when  $|i - j| = 1$ .

The algebra  $\mathbf{U}(\eta)$  is a Hopf algebra with comultiplication  $\Delta$  defined on generators by  $\Delta(E_i) = E_i \otimes \tilde{K}_i + 1 \otimes E_i$ ,  $\Delta(F_i) = F_i \otimes 1 + \tilde{K}_i^{-1} \otimes F_i$ ,  $\Delta(K_j) = K_j \otimes K_j$ . For notational simplicity, we set

$$\mathbf{U}(\eta) = \begin{cases} \mathbf{U}(n) = \mathbf{U}(\mathfrak{gl}_n), & \text{if } \eta = [1, n]; \\ \mathbf{U}(\infty) = \mathbf{U}(\mathfrak{gl}_\infty), & \text{if } \eta = (-\infty, +\infty). \end{cases}$$

Let  $\mathbf{U}^+(\eta)$  (resp.,  $\mathbf{U}^-(\eta)$ ,  $\mathbf{U}^0(\eta)$ ) be the subalgebra of  $\mathbf{U}(\eta)$  generated by the  $E_i$  (resp.,  $F_i$ ,  $K_j^{\pm 1}$ ). The subalgebras  $\mathbf{U}^+(\eta)$  and  $\mathbf{U}^-(\eta)$  are both  $\mathbb{N}$ -graded in terms of the degrees of monomials in the  $E_i$  and  $F_i$ . For monomials  $M$  in the  $E_i$  and  $M'$  in the  $F_i$ , and an element  $h \in \mathbf{U}^0(\eta)$ , write  $\deg(MhM') = \deg(M) + \deg(M')$ . Note that  $\deg$  does *not* define an algebra grading on

$\mathbf{U}(\eta)$ . However, if  $\Pi(\eta) = \{\alpha_j := e_j - e_{j+1} \mid j \in \eta^{-1}\}$  denotes the set of simple roots, where  $e_i = (\dots, 0, 1, 0 \dots) \in \mathbb{Z}^\eta$ , then there is an algebra grading over the root lattice  $\mathbb{Z}\Pi(\eta)$ ,

$$(2.1.1) \quad \mathbf{U}(\eta) = \bigoplus_{\nu \in \mathbb{Z}\Pi(\eta)} \mathbf{U}(\eta)_\nu$$

defined by the conditions  $\mathbf{U}(\eta)_{\nu'} \mathbf{U}(\eta)_{\nu''} \subseteq \mathbf{U}(\eta)_{\nu'+\nu''}$ ,  $K^{\mathbf{j}} \in \mathbf{U}(\eta)_0$ ,  $E_i \in \mathbf{U}(\eta)_{\alpha_i}$ ,  $F_i \in \mathbf{U}(\eta)_{-\alpha_i}$  for all  $\nu', \nu'' \in \mathbb{Z}\Pi(\eta)$ ,  $i \in \eta^{-1}$  and  $\mathbf{j} \in \mathbb{Z}^\eta$ .

Let  $[m]^! = [1][2] \cdots [m]$  where  $[t] = \frac{v^t - v^{-t}}{v - v^{-1}} \in \mathcal{Z}$ . The Lusztig  $\mathcal{Z}$ -form is the  $\mathcal{Z}$ -subalgebra  $U(\eta)$  of  $\mathbf{U}(\eta)$  generated by the elements  $E_i^{(m)} = \frac{E_i^m}{[m]^!}$ ,  $F_i^{(m)} = \frac{F_i^m}{[m]^!}$ ,  $K_j$  and

$$(2.1.2) \quad \begin{bmatrix} K_j; c \\ t \end{bmatrix} = \prod_{s=1}^t \frac{K_j v^{c-s+1} - K_j^{-1} v^{-c+s-1}}{v^s - v^{-s}},$$

for all  $i \in \eta^{-1}$ ,  $j \in \eta$ ,  $m, t \in \mathbb{N}$  and  $c \in \mathbb{Z}$ . Let  $U^+(\eta)$  (resp.,  $U^-(\eta)$ ,  $U^0(\eta)$ ) be the subalgebra of  $U(\eta)$  generated by the  $E_i^{(m)}$  (resp.,  $F_i^{(m)}$ ,  $K_j^{\pm 1}$  and  $\begin{bmatrix} K_j; c \\ t \end{bmatrix}$ ).

For each  $A \in \Xi^\pm(\eta)$  and  $\mathbf{j} = (j_i)_{i \in \mathbb{Z}} \in \mathbb{Z}^\eta$ , choose  $m, n \in \eta$  such that  $m \leq n$  and  $A \in \Xi^\pm([m, n])$  and let

$$E^{(A^+)} = M_n M_{n-1} \cdots M_{m+1}, \quad F^{(A^-)} = M'_{m+1} M'_{m+2} \cdots M'_n \quad \text{and} \quad K^{\mathbf{j}} = \prod_{i \in \mathbb{Z}} K_i^{j_i},$$

where

$$M_j = E_{j-1}^{(a_{j-1,j})} (E_{j-2}^{(a_{j-2,j})} E_{j-1}^{(a_{j-2,j})}) \cdots (E_m^{(a_{m,j})} E_{m+1}^{(a_{m,j})}) \cdots E_{j-1}^{(a_{m,j})},$$

and

$$M'_j = (F_{j-1}^{(a_{j,m})} \cdots F_{m+1}^{(a_{j,m})} F_m^{(a_{j,m})}) \cdots (F_{j-1}^{(a_{j,j-2})} F_{j-2}^{(a_{j,j-2})}) F_{j-1}^{(a_{j,j-1})}.$$

It is clear that  $E^{(A^+)}$  and  $F^{(A^-)}$  are independent of the selection of  $m, n$ . Note that we have, for  $A \in \Xi^\pm(\eta)$ ,  $\deg(E^{(A^+)}) = \deg(A^+)$  and  $\deg(F^{(A^-)}) = \deg(A^-)$ .

For a finite  $\eta$ , the following result is due to Lusztig (see, e.g., [26, 2.14], [1, 5.7] and [11]). The infinite case follows from the algebra isomorphism

$$\mathbf{U}(\infty) \cong \lim_{\substack{\longrightarrow \\ n}} \mathbf{U}([-n, n]),$$

induced from the natural embeddings  $\mathbf{U}([-n, n]) \subseteq \mathbf{U}([-n-1, n+1]) \subseteq \mathbf{U}(\infty)$  for all  $n \geq 0$ .

**Proposition 2.2.** (1) *The set  $\{E^{(A^+)} K^{\mathbf{j}} F^{(A^-)} \mid A \in \Xi^\pm(\eta), \mathbf{j} \in \mathbb{Z}^\eta\}$  forms a  $\mathbb{Q}(v)$ -basis for  $\mathbf{U}(\eta)$ .*

(2) *The set  $\left\{ E^{(A^+)} \prod_{i \in \eta} K_i^{\delta_i} \begin{bmatrix} K_i; 0 \\ t_i \end{bmatrix} F^{(A^-)} \mid A \in \Xi^\pm(\eta), \mathbf{t} \in \mathbb{N}^\eta, \delta_i \in \{0, 1\} \text{ for } i \in \eta \right\}$  forms a  $\mathcal{Z}$ -basis for  $U(\eta)$ .*

We now extend the definition of  $q$ -Schur algebras (or quantum Schur algebras). Let  $\Omega_\eta$  be a free  $\mathcal{Z}$ -module with basis  $\{\omega_i\}_{i \in \eta}$ . Let  $\mathbf{\Omega}_\eta = \Omega_\eta \otimes \mathbb{Q}(v)$ . Then  $\mathbf{U}(\eta)$  acts naturally on  $\mathbf{\Omega}_\eta$  by  $K_a \omega_b = v^{\delta_{a,b}} \omega_b$  ( $a, b \in \eta$ ),  $E_a \omega_b = \delta_{a+1,b} \omega_a$  and  $F_a \omega_b = \delta_{a,b} \omega_{a+1}$  ( $a \in \eta^{-1}, b \in \eta$ ). The tensor space  $\mathbf{\Omega}_\eta^{\otimes r}$  is a  $\mathbf{U}(\eta)$ -module via the comultiplication  $\Delta$  on  $\mathbf{U}(\eta)$ . Further, restriction gives the  $U(\eta)$ -module  $\Omega_\eta^{\otimes r}$ .

Let  $\mathcal{H}$  be the Hecke algebra over  $\mathcal{Z}$  associated with the symmetric group  $\mathfrak{S}_r$ , and let  $\mathcal{H} = \mathcal{H} \otimes_{\mathcal{Z}} \mathbb{Q}(v)$ . Thus,  $\mathcal{H}$  as a  $\mathcal{Z}$ -algebra has a basis  $\{\mathcal{T}_w\}_{w \in \mathfrak{S}_r}$  subject the relations: for all  $w \in \mathfrak{S}_r$  and  $s \in S := \{(i, i+1) \mid 1 \leq i \leq r-1\}$

$$\mathcal{T}_s \mathcal{T}_w = \begin{cases} \mathcal{T}_{sw}, & \text{if } \ell(sw) > \ell(w); \\ (v - v^{-1})\mathcal{T}_w + \mathcal{T}_{sw}, & \text{if } \ell(sw) < \ell(w), \end{cases}$$

where  $\ell$  is the length function on  $\mathfrak{S}_r$  with respect to  $S$ .

The Hecke algebra  $\mathcal{H}$  acts on  $\Omega_\eta^{\otimes r}$  on the right by ‘‘place permutations’’ via

$$(\omega_{i_1} \cdots \omega_{i_r}) \mathcal{T}_{(j,j+1)} = \begin{cases} \omega_{i_1} \cdots \omega_{i_{j+1}} \omega_{i_j} \cdots \omega_{i_r}, & \text{if } i_j < i_{j+1}; \\ v \omega_{i_1} \cdots \omega_{i_r}, & \text{if } i_j = i_{j+1}; \\ (v - v^{-1}) \omega_{i_1} \cdots \omega_{i_r} + \omega_{i_1} \cdots \omega_{i_{j+1}} \omega_{i_j} \cdots \omega_{i_r}, & \text{if } i_j > i_{j+1}; \end{cases}$$

cf. [5, (14.6.4)]. The endomorphism algebras

$$\mathcal{S}(\eta, r) := \text{End}_{\mathcal{H}}(\Omega_\eta^{\otimes r}), \quad \mathcal{S}(\eta, r) := \text{End}_{\mathcal{H}}(\Omega_\eta^{\otimes r})$$

are called  $q$ -Schur algebras at  $(\eta, r)$ . Note that  $\mathcal{S}(n, r) := \mathcal{S}(\eta, r)$  for  $\eta = [1, n]$  is usually called the  $q$ -Schur algebra of bidegree  $(n, r)$ . In general, if  $n = |\eta|$  is finite,  $\mathcal{S}(\eta, r) \cong \mathcal{S}(n, r)$  and  $\mathcal{S}(\eta, r) \cong \mathcal{S}(\eta, r) \otimes_{\mathbb{Z}} \mathbb{Q}(v)$ . If  $\eta = \mathbb{Z}$ ,  $\mathcal{S}(\infty, r) := \mathcal{S}(\eta, r)$  is called an *infinite  $q$ -Schur algebra*. We shall use similar notations for their integral versions.

Since the  $\mathcal{H}$ -action commutes with the action of  $\mathbf{U}(\eta)$  (see, e.g., [11], [2, §2-d]), we obtain algebra homomorphisms

$$(2.2.1) \quad \zeta_r : \mathbf{U}(\eta) \longrightarrow \mathcal{S}(\eta, r), \quad \zeta_r|_{U(\eta)} : U(\eta) \longrightarrow \mathcal{S}(\eta, r).$$

Let  $\mathbf{U}(\eta, r) = \text{Im}(\zeta_r)$  and  $U(\eta, r) = \text{Im}(\zeta_r|_{U(\eta)})$ . It is known that both maps are surjective if  $\eta$  is finite. However, this is not the case when  $\eta$  is infinite; see 5.4 below. Thus, for  $\eta = (-\infty, \infty)$  both  $\mathbf{U}(\infty, r)$  and  $U(\infty, r)$  are proper subalgebras. We will investigate the relationship between  $\mathbf{U}(\infty, r)$  and  $\mathcal{S}(\infty, r)$  in §5 via the BLM type realizations for  $\mathbf{U}(\infty)$  and  $\mathbf{U}(\infty, r)$  discussed in the next section.

### 3. THE BLM REALIZATION $\mathbf{V}(\infty)$ OF $\mathbf{U}(\infty)$ AND ITS QUOTIENTS $\mathbf{V}(\infty, r)$

In this section, we review the geometric construction of the  $q$ -Schur algebra and extend it to the infinite case.

Let  $V$  be a vector space of dimension  $r$  over a field  $k$ . An  $\eta$ -step flag is a collection  $\mathfrak{f} = (V_i)_{i \in \eta}$  of subspaces of  $V$  such that  $V_i \subseteq V_{i+1}$  for all  $i \in \eta^\dagger$  and  $\bigcup_{i \in \eta} V_i = V$  and  $V_i = 0$  for  $i \ll 0$  if  $\eta$  has no minimum element. If  $\eta$  has a minimum element  $\eta_{\min}$ , let  $V_{\eta_{\min}-1} = 0$ .

Let  $\mathcal{F}$  be the set of  $\eta$ -step flags. The group  $G := GL(V)$  acts naturally on  $\mathcal{F}$ , and hence, diagonally on  $\mathcal{F} \times \mathcal{F}$ . For  $(\mathfrak{f}, \mathfrak{f}') \in \mathcal{F} \times \mathcal{F}$ , where  $\mathfrak{f} = (V_i)_{i \in \eta}$  and  $\mathfrak{f}' = (V'_i)_{i \in \eta}$ , we let  $a_{i,j} = \dim(V_{i-1} + (V_i \cap V'_j)) - \dim(V_{i-1} + (V_i \cap V'_{j-1}))$ . Then the map  $(\mathfrak{f}, \mathfrak{f}') \mapsto (a_{i,j})$  induces a bijection from the set of  $G$ -orbits on  $\mathcal{F} \times \mathcal{F}$  to the set  $\Xi(\eta, r)$ . Let  $\mathcal{O}_A \subseteq \mathcal{F} \times \mathcal{F}$  be the  $G$ -orbit corresponding to the matrix  $A \in \Xi(\eta, r)$ . When  $k$  is a finite field of  $q$ -elements, every orbit  $\mathcal{O}_A$  is a finite set. Thus, for any  $A, B, C \in \Xi(\eta, r)$  and any fixed  $(\mathfrak{f}_1, \mathfrak{f}_2) \in \mathcal{O}_C$ , the number

$$g_{A,B,C;q} := \{\mathfrak{f} \in \mathcal{F} \mid (\mathfrak{f}_1, \mathfrak{f}) \in \mathcal{O}_A, (\mathfrak{f}, \mathfrak{f}_2) \in \mathcal{O}_B\}$$

is independent of the selection of  $(\mathfrak{f}_1, \mathfrak{f}_2)$ . It is well-known that there exists a polynomial  $g_{A,B,C} \in \mathbb{Z}[v^2]$  such that  $g_{A,B,C}|_{v^2=q} = g_{A,B,C;q}$  for any  $q$ .

Let  $\mathcal{K}(\eta, r)$  be the free  $\mathbb{Z}$ -module with basis  $\{e_A \mid A \in \Xi(\eta, r)\}$ . By [1, 1.2] there is a associative  $\mathbb{Z}$ -algebra structure on  $\mathcal{K}(\eta, r)$  with multiplication  $e_A \cdot e_B = \sum_{C \in \Xi(\eta, r)} g_{A,B,C} e_C$ . Note that, if  $\eta$  is finite, then  $\mathcal{K}(\eta, r)$  is isomorphic to the  $q$ -Schur algebra  $\mathcal{S}(|\eta|, r)$  (see, e.g., [11]). However, when  $\eta$  is infinite, the algebra  $\mathcal{K}(\eta, r)$  has no identity element. We further consider the basis  $\{[A]\}_{A \in \Xi(\eta, r)}$  for  $\mathcal{K}(\eta, r)$ , where  $[A] := v^{-d_A} e_A$  with  $d_A = \sum_{i \geq k, j < l} a_{ij} a_{kl}$ .

For a finite segment  $\eta$  of  $\mathbb{Z}$ , let  $I = I_\eta \in \Xi(\eta)$  be the identity matrix of size  $|\eta|$ . Consider  $A \in \tilde{\Xi}([m_0, n_0])$ . For any  $a, m, n \in \mathbb{Z}$  with  $m \leq m_0$  and  $n_0 \leq n$ , let  ${}_a(A^{[m,n]})$  denote the matrix in  $\Xi([m, n])$  obtained from  $A$  by adding 0s at  $(i, j)$  positions with  $m \leq i < m_0$  or  $n_0 < j \leq n$ , and let

${}_a(A^{[m,n]}) = A^{[m,n]} + aI \in \tilde{\Xi}([m,n])$ . Clearly, when  $a$  is large enough,  ${}_a(A^{[m,n]}) \in \Xi([m,n])$ . By [1] one can easily show the following generalized version of the *stabilization property*.

**Theorem 3.1.** *Let  $\mathcal{Z}_1$  be the subring of  $\mathbb{Q}(v)[v']$  generated by  $\prod_{1 \leq i \leq t} \frac{v^{-2(a-i)}v'^2-1}{v^{-2i}-1}$  and  $v^j$  with  $a \in \mathbb{Z}$ ,  $t \geq 1$  and  $j \in \mathbb{Z}$ . Let  $\eta$  be a fixed consecutive segment of  $\mathbb{Z}$ . For any  $A, B, C \in \tilde{\Xi}(\eta)$  with  $co(A) = ro(B)$ , there exist finitely many nonzero elements  $f_{A,B,C}(v, v') \in \mathcal{Z}_1$  and an integer  $a_0 \geq 1$  such that, if  $A, B, C \in \tilde{\Xi}([m_0, n_0])$  for some  $m_0 \leq n_0$  in  $\mathbb{Z}$  with  $[m_0, n_0] \subseteq \eta$ , then*

$$[{}_a(A^{[m,n]})] \cdot [{}_a(B^{[m,n]})] = \sum_{C \in \tilde{\Xi}([m_0, n_0])} f_{A,B,C}(v, v^{-a}) [{}_a(C^{[m,n]})]$$

for all  $m \leq m_0$ ,  $n \geq n_0$  with  $[m, n] \subseteq \eta$ , and  $a \geq a_0$ .

We now use the polynomials  $f_{A,B,C}(v, v')$  given in 3.1 to define a new associative algebra.

For  $C = (c_{i,j}) \in M_\eta(\mathbb{Z})$  let  $ro(C) = (\sum_j c_{i,j})_{i \in \eta} \in \mathbb{Z}^\eta$  and  $co(C) = (\sum_i c_{i,j})_{j \in \eta} \in \mathbb{Z}^\eta$ . Consider the free  $\mathcal{Z}_1$ -module with basis  $\{A \mid A \in \tilde{\Xi}(\eta)\}$ . Define a multiplication on this module by linearly extending the products on basis elements:

$$(3.1.1) \quad A \cdot B := \begin{cases} \sum_{C \in \tilde{\Xi}(\eta)} f_{A,B,C}(v, v') C, & \text{if } co(A) = ro(B) \\ 0 & \text{otherwise} \end{cases}$$

Then we get an associative algebra over  $\mathcal{Z}_1$  which has no identity element.

By specializing  $v' = 1$ , we get an associative  $\mathcal{Z}$ -algebra  $\mathcal{K}(\eta)$  with basis  $[A] := A \otimes 1$  ( $A \in \tilde{\Xi}(\eta)$ ) in which the product  $[A] \cdot [B]$  is given by  $\sum_{C \in \tilde{\Xi}(\eta)} f_{A,B,C}(v, 1)[C]$ , if  $co(A) = ro(B)$ , and it is zero, otherwise.

Let  $\mathcal{K}(\eta) = \mathcal{K}(\eta) \otimes_{\mathcal{Z}} \mathbb{Q}(v)$  and  $\mathcal{K}(\eta, r) = \mathcal{K}(\eta, r) \otimes_{\mathcal{Z}} \mathbb{Q}(v)$ . The set  $\{[\text{diag}(\lambda)] \mid \lambda \in \mathbb{Z}^\eta\}$  (resp.,  $\{[\text{diag}(\lambda)] \mid \lambda \in \Lambda(\eta, r)\}$ ) is a set of orthogonal idempotents of  $\mathcal{K}(\eta)$  (resp.,  $\mathcal{K}(\eta, r)$ ). By 1.1 we may construct the completion algebra  $\hat{\mathcal{K}}(\eta)$  (resp.,  $\hat{\mathcal{K}}(\eta, r)$ ) of  $\mathcal{K}(\eta)$  (resp.,  $\mathcal{K}(\eta, r)$ ). The element  $\sum_{\lambda \in \mathbb{Z}^\eta} [\text{diag}(\lambda)]$  (resp.,  $\sum_{\lambda \in \Lambda(\eta, r)} [\text{diag}(\lambda)]$ ) is the identity element of  $\hat{\mathcal{K}}(\eta)$  (resp.,  $\hat{\mathcal{K}}(\eta, r)$ ). Note that if  $\eta$  is a finite set, then we have  $\mathcal{K}(\eta, r) = \hat{\mathcal{K}}(\eta, r)$ .

Given  $r > 0$ ,  $A \in \Xi^\pm(\eta)$  and  $\mathbf{j} \in \mathbb{Z}^\eta$ , we define

$$(3.1.2) \quad \begin{aligned} A(\mathbf{j}, r) &= A(\mathbf{j}, r)_\eta = \sum_{\lambda \in \Lambda(\eta, r - \sigma(A))} v^{\lambda \cdot \mathbf{j}} [A + \text{diag}(\lambda)] \in \hat{\mathcal{K}}(\eta, r), \\ A(\mathbf{j}) &= A(\mathbf{j})_\eta = \sum_{\lambda \in \mathbb{Z}^\eta} v^{\lambda \cdot \mathbf{j}} [A + \text{diag}(\lambda)] \in \hat{\mathcal{K}}(\eta). \end{aligned}$$

where  $\lambda \cdot \mathbf{j} = \sum_{i \in \mathbb{Z}} \lambda_i j_i < \infty$ . (Note that these summations depend on  $\eta$ .) In particular,

$$\begin{aligned} A(\mathbf{0}) &= \sum_{\lambda \in \mathbb{Z}^\eta} [A + \text{diag}(\lambda)], \text{ for the zero vector } \mathbf{0} \in \mathbb{Z}^\eta, \\ 0(\mathbf{j}) &= \sum_{\lambda \in \mathbb{Z}^\eta} v^{\lambda \cdot \mathbf{j}} [\text{diag}(\lambda)], \text{ for the zero matrix } 0 \in \Xi^\pm(\eta). \end{aligned}$$

Thus, (3.1.1) implies the following.

**Lemma 3.2.** *Let  $\lambda, \mu \in \mathbb{Z}^\eta$ . For  $i \in \eta^{-1}$  we have*

$$\begin{aligned} E_{i,i+1}(\mathbf{0})[\text{diag}(\lambda)] &= [E_{i,i+1} + \text{diag}(\lambda - \mathbf{e}_{i+1})] = [\text{diag}(\lambda + \alpha_i)] E_{i,i+1}(\mathbf{0}) \\ E_{i+1,i}(\mathbf{0})[\text{diag}(\lambda)] &= [E_{i+1,i} + \text{diag}(\lambda - \mathbf{e}_i)] = [\text{diag}(\lambda - \alpha_i)] E_{i+1,i}(\mathbf{0}). \end{aligned}$$

Let  $\mathbf{V}(\eta)$  (resp.  $\mathbf{V}(\eta, r)$ ) be the subspace of  $\widehat{\mathcal{K}}(\eta)$  (resp.  $\widehat{\mathcal{K}}(\eta, r)$ ) spanned by the elements  $A(\mathbf{j})$  (resp.,  $A(\mathbf{j}, r)$ ) for  $A \in \Xi^\pm(\eta)$  and  $\mathbf{j} \in \mathbb{Z}^\eta$ . It is clear that the elements  $A(\mathbf{j})$  ( $A \in \Xi^\pm(\eta)$ ,  $\mathbf{j} \in \mathbb{Z}^\eta$ ) form a  $\mathbb{Q}(v)$ -basis of  $\mathbf{V}(\eta)$ . There are similar bases for  $\mathbf{V}(\eta, r)$ ; see [12] and Corollary 4.8 below.

When  $\eta = [m, n]$  is *finite*, it is known from [1, 5.7] that  $\mathbf{V}([m, n])$  is a subalgebra of  $\widehat{\mathcal{K}}([m, n])$  and there is an isomorphism

$$(3.2.1) \quad \mathbf{V}([m, n]) \cong \mathbf{U}([m, n]) \quad \text{via} \quad E_h \mapsto E_{h, h+1}(\mathbf{0}), \quad K^{\mathbf{j}} \mapsto 0(\mathbf{j}), \quad F_h \mapsto E_{h+1, h}(\mathbf{0}).$$

Moreover,  $\mathcal{K}([m, n], r) = \mathbf{V}([m, n], r) = \widehat{\mathcal{K}}([m, n], r)$  is a  $q$ -Schur algebra in this case.

When  $\eta$  is *infinite*, we will see below that the isomorphism  $\mathbf{U}(\eta) \cong \mathbf{V}(\eta)$  continues to hold, but the equalities  $\mathcal{K}([m, n], r) = \mathbf{V}([m, n], r) = \widehat{\mathcal{K}}([m, n], r)$  are replaced by a chain relation  $\mathcal{K}(\eta, r) \subseteq \mathbf{V}(\eta, r) \subseteq \widehat{\mathcal{K}}(\eta, r)$ ; see 3.3(2) and 3.6(1) below. Moreover,  $\mathbf{V}(\eta, r)$  is isomorphic to  $\mathbf{U}(\eta, r)$ .

For simplicity, we will only consider the case when  $\eta = \mathbb{Z}$  in the sequel.

**Theorem 3.3.** (1)  $\mathbf{V}(\infty)$  is a subalgebra of  $\widehat{\mathcal{K}}(\infty)$ . Moreover, the algebra  $\mathbf{U}(\infty)$  is isomorphic to  $\mathbf{V}(\infty)$  by sending  $E_h$  to  $E_{h, h+1}(\mathbf{0})$ ,  $F_h$  to  $E_{h+1, h}(\mathbf{0})$ , and  $K^{\mathbf{j}}$  to  $0(\mathbf{j})$  where  $h \in \mathbb{Z}$  and  $\mathbf{j} \in \mathbb{Z}^\infty$ .

(2) There is an algebra homomorphism, the truncation map,

$$(3.3.1) \quad \xi_r : \mathbf{V}(\infty) \longrightarrow \widehat{\mathcal{K}}(\infty, r)$$

by sending  $A(\mathbf{j})$  to  $A(\mathbf{j}, r)$  for any  $A \in \Xi^\pm(\infty)$  and  $\mathbf{j} \in \mathbb{Z}^\infty$ . In particular, we have  $\mathbf{V}(\infty, r) = \xi_r(\mathbf{V}(\infty))$  is a subalgebra of  $\widehat{\mathcal{K}}(\infty, r)$ .

*Proof.* Let  $f$  be the injective linear map from  $\mathbf{V}([-n, n])$  to  $\mathbf{V}(\infty)$  by sending  $A(\mathbf{j})$  to  $A(\mathbf{j})_\infty$  for any  $A \in \Xi^\pm([-n, n])$  and  $\mathbf{j} \in \mathbb{Z}^{[-n, n]}$ . Let  $X := \{E_{h, h+1}(\mathbf{0}), E_{h+1, h}(\mathbf{0}) \mid -n \leq h < n, \mathbf{j} \in \mathbb{Z}^{[-n, n]}\}$ . Since the formulas similar to [1, 5.3] hold in  $\widehat{\mathcal{K}}(\infty)$ , we have  $f(xA(\mathbf{j})) = f(x)f(A(\mathbf{j}))$  for any  $x \in X$  and  $A \in \Xi^\pm([-n, n])$  and  $\mathbf{j} \in \mathbb{Z}^{[-n, n]}$ . Since the algebra  $\mathbf{V}([-n, n])$  is generated by  $X$ ; see (3.2.1), it follows that  $f$  is an algebra homomorphism. Thus,  $\mathbf{V}(\infty)$  is the direct limit of  $\mathbf{V}([-n, n])$  and, hence, is a subalgebra of  $\widehat{\mathcal{K}}(\infty)$ , proving (1). The statement (2) can be proved similarly.  $\square$

We shall see in Proposition 5.3 below that  $\widehat{\mathcal{K}}(\infty, r)$  identifies with a subalgebra of  $\mathcal{S}(\infty, r)$ . Thus, identifying  $\mathbf{U}(\infty)$  with  $\mathbf{V}(\infty)$  yields the fact that the maps  $\xi_r$  in (3.3.1) and  $\zeta_r$  in (2.2.1) are identical.

The quotient algebras  $\mathbf{V}(\infty, r)$  share many properties with the usual  $q$ -Schur algebras. We first prove that  $\mathcal{K}(\infty, r)$  is a subalgebra of  $\mathbf{V}(\infty, r)$ .

For  $i \in \mathbb{Z}$ , let

$$\mathbf{k}_i = 0(\mathbf{e}_i, r), \quad \mathbf{e}_i = E_{i, i+1}(\mathbf{0}, r) \quad \text{and} \quad \mathbf{f}_i = E_{i+1, i}(\mathbf{0}, r),$$

and, for any  $\mathbf{t} \in \mathbb{N}^\infty$  and  $\mathbf{j} \in \mathbb{Z}^\infty$ , let

$$\mathbf{k}(\mathbf{t}) = \prod_{i \in \mathbb{Z}} [\mathbf{k}_i; t_i]^!, \quad \mathbf{k}_{\mathbf{t}} = \prod_{i \in \mathbb{Z}} \begin{bmatrix} \mathbf{k}_i; 0 \\ t_i \end{bmatrix} \quad \text{and} \quad \mathbf{k}^{\mathbf{j}} = \prod_{i \in \mathbb{Z}} \mathbf{k}_i^{j_i}$$

where  $[x; t]^! = (x-1)(x-v) \cdots (x-v^{t-1})$ ,  $[x; 0]^! = 1$  and  $\begin{bmatrix} \mathbf{k}_i; 0 \\ t_i \end{bmatrix}$  is defined as in (2.1.2). Since  $\mathbf{t}$  has finite support, the products are well-defined. By Theorem 3.3,  $\mathbf{e}_i, \mathbf{k}_i, \mathbf{f}_i$  ( $i \in \mathbb{Z}$ ) generate  $\mathbf{V}(\infty, r)$ . One can easily get the following result by direct calculation; cf. [7], [14].

**Lemma 3.4.** (1) We have  $\mathbf{k}(\mathbf{t}) = 0$  for all  $\mathbf{t} \in \mathbb{N}^\infty$  with  $\sigma(\mathbf{t}) > r$ .

(2) For any  $\mathbf{t} \in \mathbb{N}^\infty$ , we have  $\mathbf{k}_{\mathbf{t}} = \begin{cases} 0, & \text{if } \sigma(\mathbf{t}) > r; \\ [\text{diag}(\mathbf{t})], & \text{if } \sigma(\mathbf{t}) = r. \end{cases}$  In particular, if  $\lambda \in \Lambda(\infty, r)$ , then

$$[\text{diag}(\lambda)] = \mathbf{k}_\lambda \in \mathbf{V}(\infty, r) \quad \text{and hence} \quad 1 = \sum_{\lambda \in \Lambda(\infty, r)} \mathbf{k}_\lambda.$$

(3) For any  $\lambda \in \Lambda(\infty, r)$ , we have  $\mathbf{k}_i \mathbf{k}_\lambda = v^{\lambda_i} \mathbf{k}_\lambda$  and  $\begin{bmatrix} \mathbf{k}_i; c \\ t \end{bmatrix} \mathbf{k}_\lambda = \begin{bmatrix} \lambda_i + c \\ t \end{bmatrix}$ .

One can easily check the following commutator relations by direct calculation; cf. [7, 3.4] and [14, 4.8,4.9]. Let  $\leq$  be the partial order on  $\Lambda(\eta, r)$  defined by setting, for  $\lambda, \mu \in \Lambda(\eta, r)$ ,

$$\lambda \leq \mu \text{ if and only if } \lambda_i \leq \mu_i \text{ for all } i \in \eta.$$

For  $A \in \tilde{\Xi}(\infty)$  let

$$(3.4.1) \quad \sigma(A) = (\sigma_i(A))_{i \in \mathbb{Z}}, \text{ where } \sigma_i(A) = a_{i,i} + \sum_{j < i} (a_{i,j} + a_{j,i}).$$

**Proposition 3.5.** *Let  $A \in \Xi^\pm(\infty)$  and  $\lambda \in \Lambda(\infty, r)$ .*

- (1) *If  $\lambda \geq \sigma(A^+)$ , then  $\mathbf{e}^{(A^+)} \mathbf{k}_\lambda = \mathbf{k}_{\lambda - \text{co}(A^+) + \text{ro}(A^+)} \mathbf{e}^{(A^+)}$ ; otherwise,  $\mathbf{e}^{(A^+)} \mathbf{k}_\lambda = 0$ .*
- (2) *If  $\lambda \geq \sigma(A^-)$ , then  $\mathbf{f}^{(A^+)} \mathbf{k}_\lambda = \mathbf{k}_{\lambda + \text{co}(A^-) - \text{ro}(A^-)} \mathbf{f}^{(A^-)}$ ; otherwise,  $\mathbf{f}^{(A^+)} \mathbf{k}_\lambda = 0$ .*

For every  $A = (a_{s,t}) \in \tilde{\Xi}(\eta)$  and  $i \neq j$ , let  $\sigma_{i,j}(A) = \sum_{s \leq i; t \geq j} a_{s,t}$  for  $i < j$ , and  $\sigma_{i,j}(A) = \sum_{s \geq i; t \leq j} a_{s,t}$  for  $i > j$ . Define  $B \preceq A$  if and only if  $\sigma_{i,j}(B) \leq \sigma_{i,j}(A)$  for all  $i, j \in \eta$ ,  $i \neq j$ . Put  $B \prec A$  if  $B \preceq A$  and  $\sigma_{i,j}(B) < \sigma_{i,j}(A)$  for some  $i \neq j$ . For  $A \in \Xi(\infty)$  with  $\nu = \sigma(A)$ , let

$$\mathbf{m}^{(A)} = \mathbf{e}^{(A^+)} \mathbf{k}_\nu \mathbf{f}^{(A^-)}.$$

In general, for  $A \in \Xi^\pm(\infty)$  and  $\lambda \in \Lambda(\infty, r)$ , let  $\mathbf{m}^{(A,\lambda)} = \mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)}$ . Since the formula similar to [1, 5.5(c)] hold in  $\mathbf{V}(\infty, r)$ , the following result can be proved similar to the proof of [14, 5.5 and 5.6].

**Theorem 3.6.** (1) *For any  $A \in \Xi(\infty, r)$ , we have the following equation in  $\mathbf{V}(\infty, r)$ .*

$$(3.6.1) \quad \mathbf{m}^{(A)} = [A] + \sum_{B \in \Xi(\infty, r), B \prec A} f_{B,A} [B] \quad (\text{a finite sum})$$

where  $f_{B,A} \in \mathcal{Z}$ . Hence, the elements  $\mathbf{m}^{(A)}$ ,  $A \in \Xi(\infty, r)$ , form a  $\mathcal{Z}$ -basis of  $\mathcal{K}(\infty, r)$ , and  $\mathcal{K}(\infty, r) \subseteq \mathbf{V}(\infty, r)$ .

(2) *Suppose  $\mathbf{m}^{(A,\lambda)} \neq 0$  for some  $A \in \Xi^\pm(\infty)$  and  $\lambda \in \Lambda(\infty, r)$ . If there exists  $D \in \Xi^0(\infty)$  such that  $\text{co}(A + D) = \lambda + \text{co}(A^-) - \text{ro}(A^-)$ , then  $\mathbf{m}^{(A,\lambda)} = \mathbf{m}^{(A+D)}$ . Otherwise,*

$$(3.6.2) \quad \mathbf{m}^{(A,\lambda)} = \sum_{B \in \Xi(\infty, r), B \prec A} f'_{B,A} \mathbf{m}^{(B)} \quad (\text{a finite sum}).$$

where  $f'_{B,A} \in \mathbb{Q}(v)$ .

We end this section by showing that Borel subalgebras of (finite dimensional)  $q$ -Schur algebras are natural subalgebras of  $\mathbf{V}(\infty, r)$ , a fact which plays a crucial role in the determination of a presentation for  $\mathbf{V}(\infty, r)$ .

Let  $\mathbf{V}^{\geq 0}([-n, n], r)$  (resp.  $\mathbf{V}^{\leq 0}([-n, n], r)$ ) be the subalgebras of  $\mathbf{V}([-n, n], r)$  generated by  $\mathbf{e}'_i = E_{i, i+1}(\mathbf{0}, r)_{[-n, n]}$  (resp.  $\mathbf{f}'_i = E_{i+1, i}(\mathbf{0}, r)_{[-n, n]}$ ) and  $\mathbf{k}'_j = 0(e_j, r)_{[-n, n]}$  with  $i \in [-n, n-1]$  and  $j \in [-n, n]$ . For  $\lambda \in \Lambda([-n, n], r)$ , let

$$\mathbf{k}'_\lambda = \prod_{i=-n}^n \begin{bmatrix} \mathbf{k}'_i; 0 \\ \lambda_i \end{bmatrix} \in \mathbf{V}^{\geq 0}([-n, n], r) \cap \mathbf{V}^{\leq 0}([-n, n], r)$$

where  $\mathbf{k}'_n = v^r \mathbf{k}'_{-n}^{-1} \cdots \mathbf{k}'_{n-1}^{-1}$ .

**Proposition 3.7.** *For  $n \geq 1$  there are algebra monomorphisms (sending 1 to 1)*

$$\begin{aligned} \varphi_n^{\geq 0} : \mathbf{V}^{\geq 0}([-n, n], r) &\rightarrow \mathbf{V}(\infty, r) & \text{satisfying} & \quad \mathbf{e}'_i \mapsto \mathbf{e}_i, \mathbf{k}'_i \mapsto \mathbf{k}_i \quad (-n \leq i \leq n-1) \\ \varphi_n^{\leq 0} : \mathbf{V}^{\leq 0}([-n, n], r) &\rightarrow \mathbf{V}(\infty, r) & \text{satisfying} & \quad \mathbf{f}'_i \mapsto \mathbf{f}_i, \mathbf{k}'_i \mapsto \mathbf{k}_i \quad (-n \leq i \leq n-1). \end{aligned}$$



Moreover, for any  $\lambda \in \Lambda([-n, n], r)$ , we have  $\varphi_n^{\geq 0}(\mathbf{k}'_\lambda) = \varphi_n^{\leq 0}(\mathbf{k}'_\lambda) = \mathbf{h}_{\lambda, n}$ , where

$$(3.7.1) \quad \mathbf{h}_{\lambda, n} = \sum_{\substack{\mu \in \Lambda(\infty, r) \\ \mu_{-n} = \lambda_{-n}, \dots, \mu_{n-1} = \lambda_{n-1}}} [\text{diag}(\mu)].$$

*Proof.* By [14, 8.1] there is an algebra homomorphism  $\varphi_n^{\geq 0} : \mathbf{V}^{\geq 0}([-n, n], r) \rightarrow \mathbf{V}(\infty, r)$  mapping  $\mathbf{e}'_i$  to  $\mathbf{e}_i$  and  $\mathbf{k}'_i$  to  $\mathbf{k}_i$  for  $-n \leq i \leq n-1$ . By (3.6.1) and [14, 8.2] we have  $\varphi_n^{\geq 0}$  is injective. The proof for  $\varphi_n^{\leq 0}$  is similar. It remains to prove the last assertion. We have

$$\begin{aligned} \varphi_n^{\geq 0}(\mathbf{k}'_\lambda) &= \prod_{i=-n}^{n-1} \begin{bmatrix} \mathbf{k}_i; 0 \\ t_i \end{bmatrix} \cdot \begin{bmatrix} v^r \mathbf{k}_{-n}^{-1} \mathbf{k}_{-n+1}^{-1} \cdots \mathbf{k}_{n-1}^{-1}; 0 \\ \lambda_n \end{bmatrix} \\ &= \sum_{\mu \in \Lambda(\infty, r)} \left( \prod_{i=-n}^{n-1} \begin{bmatrix} \mu_i \\ \lambda_i \end{bmatrix} \begin{bmatrix} r - \sum_{i=-n}^{n-1} \mu_i \\ \lambda_n \end{bmatrix} \right) [\text{diag}(\mu)]. \end{aligned}$$

Since  $\lambda \in \Lambda([-n, n], r)$  we have  $\sum_{i=-n}^{n-1} (\mu_i - \lambda_i) + (r - \sum_{i=-n}^{n-1} \mu_i - \lambda_n) = r - \sum_{i=-n}^n \lambda_i = 0$ . Hence, for  $\mu \in \Lambda(\infty, r)$  we have

$$\begin{aligned} \prod_{i=-n}^{n-1} \begin{bmatrix} \mu_i \\ \lambda_i \end{bmatrix} \begin{bmatrix} r - \sum_{i=-n}^{n-1} \mu_i \\ \lambda_n \end{bmatrix} \neq 0 &\iff \mu_i \geq \lambda_i \text{ for } -n \leq i \leq n-1 \text{ and } r - \sum_{i=-n}^{n-1} \mu_i \geq \lambda_n \\ &\iff \mu_i = \lambda_i \text{ for } -n \leq i \leq n-1. \end{aligned}$$

Hence,  $\prod_{i=-n}^{n-1} \begin{bmatrix} \mu_i \\ \lambda_i \end{bmatrix} \begin{bmatrix} r - \sum_{i=-n}^{n-1} \mu_i \\ \lambda_n \end{bmatrix} = 1$ , and (3.7.1) follows.  $\square$

**Remark 3.8.** The algebra monomorphisms  $\varphi_n^{\geq 0}$  and  $\varphi_n^{\leq 0}$  can't be glued as an algebra homomorphism from  $\mathbf{V}([-n, n], r)$  to  $\mathbf{V}(\infty, r)$  since the last relation in [14, 4.1] does not hold in  $\mathbf{V}(\infty, r)$ .

#### 4. DRINFELD–JIMBO TYPE PRESENTATION FOR $\mathbf{V}(\infty, r)$

We first record the handy result [12, (6.6.2)] for later use. See (1.1.1) and (3.4.1) for relevant notations.

**Lemma 4.1.** For  $A \in \Xi^\pm([-n, n])$ , let

$$\Lambda_{n,r,A} = \{\lambda \in \Lambda([-n, n], r) \mid \lambda \geq \sigma(A)\} \quad \text{and} \quad \Lambda'_{n,r,A} = \{\mathbf{j} \in \mathbb{N}^{[-n, n-1]} \mid \sigma(\mathbf{j}) + \sigma(A) \leq r\}.$$

Then  $|\Lambda_{n,r,A}| = |\Lambda'_{n,r,A}|$  and  $\det \left( v^{\sum_{i=-n}^{n-1} \lambda_i j_i} \right)_{\lambda \in \Lambda_{n,r,A}, \mathbf{j} \in \Lambda'_{n,r,A}} \neq 0$ .

We are now ready to describe a presentation for  $\mathbf{V}(\infty, r)$ .

**Definition 4.2.** Let  $\tilde{\mathbf{V}}(\infty, r)$  be the associative algebra over  $\mathbb{Q}(v)$  generated by the elements

$$\mathbf{e}_i, \mathbf{f}_i, \mathbf{k}_i \quad (i \in \mathbb{Z}),$$

subject to the relations:

- (a)  $\mathbf{k}_i \mathbf{k}_j = \mathbf{k}_j \mathbf{k}_i$ ;
- (b)  $\prod_{i \in \mathbb{Z}} [\mathbf{k}_i; t_i]^1 = 0$  for all  $\mathbf{t} \in \mathbb{N}^\infty$  with  $\sigma(\mathbf{t}) = r + 1$ ;
- (c)  $\mathbf{e}_i \mathbf{e}_j = \mathbf{e}_j \mathbf{e}_i, \mathbf{f}_i \mathbf{f}_j = \mathbf{f}_j \mathbf{f}_i$  ( $|i - j| > 1$ );
- (d)  $\mathbf{e}_i^2 \mathbf{e}_j - (v + v^{-1}) \mathbf{e}_i \mathbf{e}_j \mathbf{e}_i + \mathbf{e}_j \mathbf{e}_i^2 = 0$  ( $|i - j| = 1$ );
- (e)  $\mathbf{f}_i^2 \mathbf{f}_j - (v + v^{-1}) \mathbf{f}_i \mathbf{f}_j \mathbf{f}_i + \mathbf{f}_j \mathbf{f}_i^2 = 0$  ( $|i - j| = 1$ );
- (f)  $\mathbf{k}_i \mathbf{e}_j = v^{\delta_{i,j} - \delta_{i,j+1}} \mathbf{e}_j \mathbf{k}_i, \quad \mathbf{k}_i \mathbf{f}_j = v^{-\delta_{i,j} + \delta_{i,j+1}} \mathbf{f}_j \mathbf{k}_i$ ;
- (g)  $\mathbf{e}_i \mathbf{f}_j - \mathbf{f}_j \mathbf{e}_i = \delta_{i,j} \frac{\tilde{\mathbf{k}}_i - \tilde{\mathbf{k}}_i^{-1}}{v - v^{-1}}$ , where  $\tilde{\mathbf{k}}_i = \mathbf{k}_i \mathbf{k}_{i+1}^{-1}, i \in \mathbb{Z}$ .

Comparing 4.2 with 2.1, we see that there is an algebra epimorphism  $\mathbf{U}(\infty) \twoheadrightarrow \tilde{\mathbf{V}}(\infty, r)$  satisfying  $E_i \mapsto \mathbf{e}_i$ ,  $F_i \mapsto \mathbf{f}_i$  and  $K_j \mapsto \mathbf{k}_j$ . In particular, for  $A \in \Xi^\pm(\infty)$ , let  $\mathbf{e}^{(A^+)}$ ,  $\mathbf{f}^{(A^-)}$ ,  $\mathbf{k}^{\mathbf{j}}$ , etc., be the images of  $E^{(A^+)}$ ,  $F^{(A^-)}$ ,  $K^{\mathbf{j}}$ , etc., under this homomorphism.

On the other hand, by 3.3(2) and 3.4(1), there is an algebra epimorphism

$$(4.2.1) \quad \pi : \tilde{\mathbf{V}}(\infty, r) \longrightarrow \mathbf{V}(\infty, r),$$

defined by  $\mathbf{e}_i \mapsto \mathbf{e}_i$ ,  $\mathbf{f}_i \mapsto \mathbf{f}_i$ ,  $\mathbf{k}_i \mapsto \mathbf{k}_i$ . We shall prove that  $\pi$  is an isomorphism by displaying a spanning set for  $\tilde{\mathbf{V}}(\infty, r)$  whose image under  $\pi$  is a linearly independent set in  $\mathbf{V}(\infty, r)$ .

We first have the following counterpart of 3.7 by comparing the above defining relations with the relations for Borel subalgebras.

**Lemma 4.3.** *For any given  $n \geq 1$ , there are algebra monomorphisms*

$$\begin{aligned} \psi_n^{\geq 0} : \mathbf{V}^{\geq 0}([-n, n], r) &\rightarrow \tilde{\mathbf{V}}(\infty, r) & \text{satisfying} & \quad \mathbf{e}'_i \mapsto \mathbf{e}_i, \mathbf{k}'_i \mapsto \mathbf{k}_i \quad (-n \leq i \leq n-1) \\ \psi_n^{\leq 0} : \mathbf{V}^{\leq 0}([-n, n], r) &\rightarrow \tilde{\mathbf{V}}(\infty, r) & \text{satisfying} & \quad \mathbf{f}'_i \mapsto \mathbf{f}_i, \mathbf{k}'_i \mapsto \mathbf{k}_i \quad (-n \leq i \leq n-1). \end{aligned}$$

*Proof.* The injectivity follows from 3.7 since  $\varphi_n^{\geq 0}$  is the composition of  $\psi_n^{\geq 0}$  and  $\pi$ .  $\square$

This result shows that the subalgebra  $\mathbf{B}_n^+$  (resp.  $\mathbf{B}_n^-$ ) of  $\tilde{\mathbf{V}}(\infty, r)$  generated by  $\mathbf{e}_i, \mathbf{k}_i$  (resp.  $\mathbf{f}_i, \mathbf{k}_i$ ),  $-n \leq i \leq n-1$ , is a (finite dimensional) Borel subalgebra investigated in [14, §8]. Let  $\mathbf{B}_n^0 = \mathbf{B}_n^+ \cap \mathbf{B}_n^-$ . Then  $\mathbf{B}_n^0$  is the subalgebra generated by  $\mathbf{k}_i$ ,  $-n \leq i \leq n-1$ . We summarize the properties of these algebras; cf. [7, 8.2–3] and [14, 4.7–10].

For any  $\lambda \in \Lambda([-n, n], r)$ , let  $\mathbf{h}_{\lambda, n} := \psi_n^{\geq 0}(\mathbf{k}'_\lambda) = \psi_n^{\leq 0}(\mathbf{k}'_\lambda)$ .

**Corollary 4.4.** *Let  $n \geq 1$ .*

- (1) *Each of the following sets forms a basis for  $\mathbf{B}_n^0$ :*
  - (a)  $\{\mathbf{h}_{\lambda, n} \mid \lambda \in \Lambda([-n, n], r)\}$ ;
  - (b)  $\{\mathbf{k}^{\mathbf{j}} \mid \mathbf{j} \in \mathbb{N}^{[-n, n-1]}, \sigma(\mathbf{j}) \leq r\}$ .
- (2) *For all  $-n \leq i \leq n-1$  and  $\lambda \in \Lambda([-n, n], r)$ ,  $\mathbf{k}_i \mathbf{h}_{\lambda, n} = v^{\lambda_i} \mathbf{h}_{\lambda, n}$ .*
- (3)  $1 = \sum_{\lambda \in \Lambda([-n, n], r)} \mathbf{h}_{\lambda, n}$ .
- (4) *The elements  $\mathbf{h}_{\lambda, n}$  satisfy the commuting relations described in 3.5 with  $\mathbf{k}_\lambda$  replaced by  $\mathbf{h}_{\lambda, n}$  and  $\mathbf{e}^{(A^+)}$ ,  $\mathbf{f}^{(A^-)}$  by  $\mathbf{e}^{(A^+)}$  and  $\mathbf{f}^{(A^-)}$  for all  $A \in \Xi^\pm([-n, n])$  and  $\lambda \in \Lambda([-n, n], r)$ .*

The elements  $\mathbf{h}_{\lambda, n}$  play an important role in a construction of a spanning set for  $\tilde{\mathbf{V}}(\infty, r)$ . Let

$$\mathcal{M}'_n = \{\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} \mid A \in \Xi^\pm([-n, n-1]), \lambda \in \Lambda([-n, n], r), \lambda \geq \sigma(A)\},$$

where  $\lambda \geq \sigma(A)$  means  $\lambda_i \geq \sigma_i(A) \forall i$ , and let  $\mathcal{M}'_\infty = \cup_{n \geq 1} \mathcal{M}'_n$ . Recall from 4.1 and (1.1.2) the set  $\Lambda_{n, r, A}$  and the function  $\deg$ .

**Lemma 4.5.** *For any  $n \geq 1$ ,  $A \in \Xi^\pm([-n, n-1])$ ,  $\lambda \in \Lambda([-n, n], r)$ , if  $\lambda \notin \Lambda_{n, r, A}$ , then there exist  $f_{B, \mu}^{A, \lambda} \in \mathcal{Z}$  such that*

$$(4.5.1) \quad \mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} = \sum_{\substack{B \in \Xi^\pm([-n, n-1]) \\ \mu \in \Lambda_{n, r, A}, \deg(B) < \deg(A)}} f_{B, \mu}^{A, \lambda} \mathbf{e}^{(B^+)} \mathbf{h}_{\mu, n} \mathbf{f}^{(B^-)}.$$

Hence,  $\mathcal{M}'_\infty$  is a spanning set for  $\tilde{\mathbf{V}}(\infty, r)$ .

*Proof.* We first observe, by 4.4(2)&(3), that all  $\mathbf{k}_i$  and  $\left[ \begin{smallmatrix} \mathbf{k}_i \\ t \end{smallmatrix} \right]$  are  $\mathcal{Z}$ -linear combinations of  $\mathbf{h}_{\lambda, n}$  whenever  $-n \leq i \leq n-1$ . So (4.5.1) implies the assertion that  $\mathcal{M}'_\infty$  spans  $\tilde{\mathbf{V}}(\infty, r)$ . We now prove (4.5.1) by applying induction on  $\deg(A)$ ; see (1.1.2).

Suppose  $\lambda \in \Lambda([-n, n], r)$  and  $\lambda_i < \sigma_i(A)$  where  $i$  is minimal. Then  $\deg(A) \geq 1$ . If  $\deg(A) = 1$ , then  $A^+ = E_{i-1, i}$  and  $A^- = 0$  or  $A^+ = 0$  and  $A^- = E_{i, i-1}$ . By 3.5, we have  $\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} = 0$ . Assume now  $\deg(A) \geq 2$  and (4.5.1) is true for all  $A'$  with  $\deg(A') < \deg(A)$ . Let  $A_i = (a_{k, l}^{(i)})$  be the submatrix of  $A$  such that  $a_{k, l}^{(i)} = a_{k, l}$  if  $k, l \leq i$  and  $a_{k, l}^{(i)} = 0$  otherwise. Since  $A \in \Xi^\pm([-n, n-1])$ , we can write  $\mathbf{e}^{(A^+)} = \mathbf{m}_1 \mathbf{e}^{(A_i^+)}$  and  $\mathbf{f}^{(A^-)} = \mathbf{f}^{(A_i^-)} \mathbf{m}'_1$ , where  $\mathbf{m}_1$  is the monomial of  $\mathbf{e}_j^{(a)}$  ( $-n \leq j < n-1$ ,  $a \geq 0$ ) and  $\mathbf{m}'_1$  is the monomial of  $\mathbf{f}_j^{(a)}$  ( $-n \leq j < n-1$ ,  $a \geq 0$ ). Then

$$\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} = \mathbf{m}_1 \mathbf{e}^{(A_i^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A_i^-)} \mathbf{m}'_1.$$

Since  $\lambda_j \geq \sigma_j(A_i^+)$  for all  $j \leq i$ , 4.4(4) implies

$$\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} = \mathbf{m}_1 \mathbf{h}_{\lambda^*, n} \mathbf{e}^{(A_i^+)} \mathbf{f}^{(A_i^-)} \mathbf{m}'_1,$$

where  $\lambda^* \in \Lambda([-n, n], r)$  with  $\lambda_i^* = \lambda - co(A_i^+) - ro(A_i^+) = \lambda_i - \sum_{j < i} a_{ji} = \lambda_i - \sigma_i(A^+) \geq 0$ . On the other hand, the commutator formula in [25, 4.1(a)] (see the remarks right after [14, 2.3]) implies

$$\mathbf{e}^{(A_i^+)} \mathbf{f}^{(A_i^-)} = \mathbf{f}^{(A_i^-)} \mathbf{e}^{(A_i^+)} + f,$$

where  $f$  is a  $\mathcal{Z}$ -linear combination of monomials  $\mathbf{m}_j^e \mathbf{h}_j \mathbf{m}_j^f$  with  $\deg(\mathbf{m}_j^e \mathbf{m}_j^f) < \deg(A_i)$ . Here,  $\mathbf{m}_j^e$  is a product of some  $\mathbf{e}_s^{(a)}$ ,  $\mathbf{m}_j^f$  is a product of some  $\mathbf{f}_s^{(b)}$ , and  $\mathbf{h}_j$  is a product of some  $\left[ \begin{smallmatrix} \tilde{\mathbf{k}}_{t; c} \\ m \end{smallmatrix} \right]$  where<sup>2</sup>  $-n \leq s, t \leq n-2$  and  $a, b, c, m \in \mathbb{Z}$ . Thus,  $\deg(\mathbf{m}_1 \mathbf{m}_j^e \mathbf{m}_j^f \mathbf{m}'_1) < \deg(A)$ . Now  $\lambda_i < \sigma_i(A)$  implies  $\lambda_i^* < \sigma_i(A^-) = \sigma_i(A_i^-)$ . Hence,  $\mathbf{h}_{\lambda^*, n} \mathbf{f}^{(A_i^-)} = 0$  by 4.4(4). Since  $\mathbf{h}_j \in \mathbf{B}_n^0$  is a  $\mathcal{Z}$ -linear combination of  $\mathbf{h}_{\mu, n}$  by 4.4(1-3), it follows from 4.4(4) that  $\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)}$  is a  $\mathcal{Z}$ -linear combination of  $\mathbf{m}_1 \mathbf{m}_j^e \mathbf{h}_{\mu, n} \mathbf{m}_j^f \mathbf{m}'_1$  with  $\deg(\mathbf{m}_1 \mathbf{m}_j^e \mathbf{m}_j^f \mathbf{m}'_1) < \deg(A)$  and  $\mu \in \Lambda([-n, n], r)$ . Now, since  $\tilde{\mathbf{V}}(\infty, r)$  is a homomorphic image of  $\mathbf{U}(\infty)$ , 2.2 implies that each  $\mathbf{m}_1 \mathbf{m}_j^e$  (resp.,  $\mathbf{m}_j^f \mathbf{m}'_1$ ) is a  $\mathcal{Z}$ -linear combination of  $\mathbf{e}^{(B)}$ ,  $B \in \Xi^+([-n, n-1])$  (resp.,  $\mathbf{f}^{(B')}$ ,  $B' \in \Xi^-([-n, n-1])$ ) with  $\deg(B) = \deg(\mathbf{m}_1 \mathbf{m}_j^e)$  (resp.,  $\deg(B') = \deg(\mathbf{m}_j^f \mathbf{m}'_1)$ ). Thus, each  $\mathbf{m}_1 \mathbf{m}_j^e \mathbf{h}_{\mu, n} \mathbf{m}_j^f \mathbf{m}'_1$  is a  $\mathcal{Z}$ -linear combination of  $\mathbf{e}^{(A')^+} \mathbf{h}_{\mu, n} \mathbf{f}^{(A')^-}$  with  $A' \in \Xi^\pm([-n, n-1])$  and  $\deg(A') < \deg(A)$ . Thus, our assertion follows from induction.  $\square$

Though each  $\mathcal{M}'_n$  is linearly independent as part of the integral basis for a  $q$ -Schur algebra (cf. (4.6.1) and 4.8 below), the set  $\mathcal{M}'_\infty$  is unfortunately not linearly independent. In fact, we can see that  $\text{span}(\mathcal{M}'_n)$  is a subspace of  $\text{span}(\mathcal{M}'_{n+1})$ , but  $\mathcal{M}'_n$  is not a subset of  $\mathcal{M}'_{n+1}$ . However, we shall use  $\mathcal{M}'_\infty$  to construct several bases below.

Let

$$\tilde{\mathcal{N}}_\infty := \{ \mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)} \mid A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{N}^\infty, \sigma(\mathbf{j}) + \sigma(A) \leq r \} \subseteq \tilde{\mathbf{V}}(\infty, r)$$

and, for  $n \geq 1$ , let

$$\tilde{\mathcal{N}}_n := \{ \mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)} \mid A \in \Xi^\pm([-n, n]), \mathbf{j} \in \Lambda'_{n, r, A} \},$$

where  $\Lambda'_{n, r, A}$  is defined in 4.1. Then  $\tilde{\mathcal{N}}_n \subseteq \tilde{\mathcal{N}}_{n+1}$  and  $\tilde{\mathcal{N}}_\infty = \bigcup_{n \geq 1} \tilde{\mathcal{N}}_n$ . Similarly, we can define  $\mathcal{N}_\infty$  and  $\mathcal{N}_n$  in the algebra  $\mathbf{V}(\infty, r)$  so that  $\mathcal{N}_\infty = \pi(\tilde{\mathcal{N}}_\infty)$ .

**Corollary 4.6.** *The set  $\tilde{\mathcal{N}}_\infty$  spans  $\tilde{\mathbf{V}}(\infty, r)$ .*

<sup>2</sup>Our assumption on  $A$  guarantees that  $\mathbf{e}_{n-1}$  and  $\mathbf{f}_{n-1}$  do not appear in  $\mathbf{e}^{(A^+)}$  and  $\mathbf{f}^{(A^-)}$ . Thus, only the  $\tilde{\mathbf{k}}_t = \mathbf{k}_t \mathbf{k}_{t+1}^{-1}$  with  $-n \leq t \leq n-2$  occur in the commutator formula; cf. 4.2(g).

*Proof.* By 4.5, it is enough to prove that, for any  $n \geq 1$ , the set  $\mathcal{M}'_n$  lies in the span of  $\tilde{\mathcal{N}}_n$ . In other words, we need to prove that, for any  $A \in \Xi^\pm([-n, n-1])$ , and  $\lambda \in \Lambda([-n, n], r)$  with  $\lambda \geq \sigma(A)$ , the element  $\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} \in \text{span} \tilde{\mathcal{N}}_n$ . We proceed by induction on  $\text{deg}(A)$ . If  $\text{deg}(A) = 0$ , then by 4.4(1) we have  $\mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} = \mathbf{h}_{\lambda, n} \in \text{span} \tilde{\mathcal{N}}_n$  for  $\lambda \in \Lambda([-n, n], r)$ . Assume now that  $\text{deg}(A) \geq 1$ . By 4.4(1), we may write, for any  $\mathbf{j} \in \mathbb{N}^{[-n, n]}$  with  $j_n = 0$ ,

$$(4.6.1) \quad \mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)} = \sum_{\mu \in \Lambda_{n, r, A}} v^{\mu \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{h}_{\mu, n} \mathbf{f}^{(A^-)} + \sum_{\nu \notin \Lambda_{n, r, A}} v^{\nu \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{h}_{\nu, n} \mathbf{f}^{(A^-)}.$$

By (4.5.1) and induction, the second sum is in the span of  $\tilde{\mathcal{N}}_n$ . If  $\mathbf{j}$  runs over the set  $\Lambda'_{n, r, A}$ , then the determinant  $\det(v^{\mu \cdot \mathbf{j}})_{\mu, \mathbf{j}} \neq 0$  (see 4.1). Hence,  $\mathbf{e}^{(A^+)} \mathbf{h}_{\mu, n} \mathbf{f}^{(A^-)} \in \text{span} \tilde{\mathcal{N}}_n$  for all  $\mu \in \Lambda_{n, r, A}$ .  $\square$

**Theorem 4.7.** *The algebra epimorphism  $\pi : \tilde{\mathbf{V}}(\infty, r) \rightarrow \mathbf{V}(\infty, r)$  defined in (4.2.1) is an isomorphism. In particular, if a coefficient  $f_{B, \mu}^{A, \lambda}$  given in (4.5.1) is nonzero, then  $B \prec A$ .*

*Proof.* Since  $\tilde{\mathbf{V}}(\infty, r)$  is spanned by  $\tilde{\mathcal{N}}_\infty$  by 4.6, it is enough to prove that its image  $\pi(\tilde{\mathcal{N}}_\infty) = \mathcal{N}_\infty$  is linearly independent. Fix  $n \geq 1$ . By 3.4 and (3.6.2), for  $A \in \Xi^\pm([-n, n])$  and  $\mathbf{j} \in \mathbb{N}^{[-n, n-1]}$ , there exist  $f_{B, A} \in \mathbb{Q}(v)$  such that

$$\begin{aligned} \mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)} &= \sum_{\substack{\lambda \in \Lambda(\infty, r) \\ \forall j, \lambda_j \geq \sigma_j(A)}} v^{\lambda \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)} + \sum_{\substack{\mu \in \Lambda(\infty, r) \\ \exists j, \mu_j < \sigma_j(A)}} v^{\mu \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{k}_\mu \mathbf{f}^{(A^-)} \\ &= \sum_{\substack{\lambda \in \Lambda([-n, n], r) \\ \forall i, \lambda_i \geq \sigma_i(A)}} v^{\lambda \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)} + \sum_{\substack{\lambda \notin \Lambda([-n, n], r) \\ \forall i, \lambda_i \geq \sigma_i(A)}} v^{\lambda \cdot \mathbf{j}} \mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)} + \sum_{\substack{B \in \Xi(\infty, r) \\ B \prec A}} f_{B, A} \mathbf{m}^{(B)}. \end{aligned}$$

Hence, by (3.6.1) and 4.1 the set  $\mathcal{N}$  is linearly independent. Since  $\mathcal{N}_\infty = \bigcup_{n \geq 1} \mathcal{N}_n$  and  $\mathcal{N}_n \subseteq \mathcal{N}_{n+1}$ , it follows that the set  $\mathcal{N}_\infty$  is linearly independent.

The last assertion follows from 3.6.  $\square$

With the above result we will identify  $\tilde{\mathbf{V}}(\infty, r)$  with  $\mathbf{V}(\infty, r)$ . In particular we will identify  $\mathbf{h}_{\lambda, n}$  with  $\mathbf{h}_{\lambda, n}$ ,  $\mathbf{e}_i$  with  $\mathbf{e}_i$  etc. For  $A \in \Xi^\pm(\infty)$  and  $\mathbf{j} \in \mathbb{N}^\infty$ , let

$$\mathbf{n}^{(A, \mathbf{j})} := \mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)}.$$

**Corollary 4.8.** *The set  $\mathcal{N}_\infty = \{\mathbf{n}^{(A, \mathbf{j})} \mid A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{N}^\infty, \sigma(\mathbf{j}) + \sigma(A) \leq r\}$  forms a basis for  $\mathbf{V}(\infty, r)$ . Moreover, if  $\mathbf{j} \in \mathbb{N}^{[-n, n-1]}$  and  $\mathbf{j} \notin \Lambda'_{n, r, A}$ , then we have, for some  $p_{B, \mathbf{j}'} \in \mathbb{Q}(v)$ ,*

$$\mathbf{e}^{(A^+)} \mathbf{k}^{\mathbf{j}} \mathbf{f}^{(A^-)} = \sum_{\substack{B \in \Xi^\pm([-n, n-1]) \\ B \prec A, \mathbf{j}' \in \Lambda'_{n, r, A}}} p_{B, \mathbf{j}'} \mathbf{n}^{(B, \mathbf{j}')}.$$

*Proof.* Using (4.6.1) and 4.5, the last assertion can be proved as the proof of [12, 6.7].  $\square$

The basis  $\mathcal{N}_\infty$  is the counterpart of the basis [12, 6.6(1)] for  $q$ -Schur algebras. Since the formula similar to [1, 5.5(c)] holds in  $\mathbf{V}(\infty, r)$ , by 4.8 we get the following result which is the counterpart of the basis given in [12, 6.6(2)].

**Corollary 4.9.** *The set  $\mathcal{B}_\infty = \{A(\mathbf{j}, r) \mid A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{N}^\infty, \sigma(\mathbf{j}) + \sigma(A) \leq r\}$  forms a basis for  $\mathbf{V}(\infty, r)$ .*

We end this section with another application of the spanning set  $\mathcal{M}'_\infty$  by displaying an integral monomial basis for the  $\mathcal{Z}$ -form of  $\mathbf{V}(\infty, r)$ ; cf. [14, 5.4].

Let  $V(\infty, r) = \xi_r(U(\infty))$ . We further put  $V^+(\infty, r) = \xi_r(U^+(\infty))$ ,  $V^-(\infty, r) = \xi_r(U^-(\infty))$  and  $V^0(\infty, r) = \xi_r(U^0(\infty))$ .

**Lemma 4.10.** (1) *The set  $\{\mathbf{e}^{(A)} \mid A \in \Xi^+(\infty), \sigma(A) \leq r\}$  (resp.,  $\{\mathbf{f}^{(A)} \mid A \in \Xi^-(\infty), \sigma(A) \leq r\}$ ) forms a  $\mathcal{Z}$ -basis of  $V^+(\infty, r)$  (resp.,  $V^-(\infty, r)$ ).*

(2) *The set  $\mathcal{M}_\infty^0 := \{\mathbf{k}_\lambda \mid \lambda \in \mathbb{N}^\infty, \sigma(\lambda) \leq r\}$  forms a  $\mathcal{Z}$ -basis of  $V^0(\infty, r)$ .*

*Proof.* Let  $A \in \Xi^+(\infty)$ . If  $\sigma(A) = \sum_i \sigma_i(A) > r$ , then  $\mathbf{e}^{(A)} = \sum_{\lambda \in \Lambda(\infty, r)} \mathbf{e}^{(A)} \mathbf{k}_\lambda = 0$  by 3.4 and 3.5. Now (1) follows from 2.2 and (3.6.1). For  $n \geq 1$  we let  $\mathcal{M}_n^0 := \left\{ \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mathbf{k}_i; 0 \\ \mu_i \end{bmatrix} \mid \mu \in \Lambda([-n, n], r) \right\}$ . We fix  $n \geq 1$ . For  $\lambda, \mu \in \Lambda([-n, n], r)$  we write  $\lambda \leq_n \mu$  if and only if  $\lambda_i \leq \mu_i$  for  $-n \leq i \leq n-1$ . If  $\lambda_i < \mu_i$  for some  $-n \leq i \leq n-1$  then we write  $\lambda <_n \mu$ . For  $\lambda \in \Lambda([-n, n], r)$  we have by 4.4(2)&(3)

$$(4.10.1) \quad \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mathbf{k}_i; 0 \\ \lambda_i \end{bmatrix} = \mathbf{h}_{\lambda, n} + \sum_{\mu \in \Lambda([-n, n], r), \lambda <_n \mu} \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mu_i \\ \lambda_i \end{bmatrix} \mathbf{h}_{\mu, n}.$$

Hence, the set  $\mathcal{M}_n^0$  is linearly independent and we have  $\text{span}_{\mathcal{Z}} \mathcal{M}_n^0 = \text{span}_{\mathcal{Z}} \{\mathbf{h}_{\lambda, n} \mid \lambda \in \Lambda([-n, n], r)\}$ . Since  $\mathcal{M}_n^0 \subseteq \mathcal{M}_{n+1}^0$  and  $\mathcal{M}_\infty^0 = \bigcup_{n \geq 1} \mathcal{M}_n^0$ , the set  $\mathcal{M}_\infty^0$  is linearly independent. Since  $V^0(\infty, r) = \text{span}_{\mathcal{Z}} \{\mathbf{h}_{\lambda, n} \mid \lambda \in \Lambda([-n, n], r), n \geq 1\}$ , we have  $V^0(\infty, r) = \text{span}_{\mathcal{Z}} \mathcal{M}_\infty^0$ , proving (2).  $\square$

The proof above shows that every  $\mathbf{h}_{\lambda, n} \in V(\infty, r)$ . Hence, the set  $\mathcal{M}'_\infty$  is a subset of the  $\mathcal{Z}$ -algebra  $V(\infty, r)$ . Thus, 4.5 implies immediately the following.

**Corollary 4.11.** *The set  $\mathcal{M}'_\infty$  is a spanning set of the  $\mathcal{Z}$ -algebra  $V(\infty, r)$ .*

**Proposition 4.12.** *The set*

$$\mathcal{M}_\infty := \{\mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)} \mid A \in \Xi^\pm(\infty), \lambda \in \mathbb{N}^\infty, \lambda \geq \sigma(A), \sigma(\lambda) \leq r\}$$

*forms a  $\mathcal{Z}$ -basis for  $V(\infty, r)$ .*

*Proof.* For any  $n \geq 1$ , let

$$\mathcal{M}_n := \left\{ \mathbf{e}^{(A^+)} \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mathbf{k}_i; 0 \\ \lambda_i \end{bmatrix} \mathbf{f}^{(A^-)} \mid A \in \Xi^\pm([-n, n-1]), \lambda \in \Lambda_{n, r, A} \right\}.$$

Then,  $\mathcal{M}_n \subseteq \mathcal{M}_{n+1}$ ,  $|\mathcal{M}_n| = |\mathcal{M}'_n|$  and  $\mathcal{M}_\infty = \bigcup_{n \geq 1} \mathcal{M}_n$ . Since, for  $A \in \Xi^\pm([-n, n-1])$  and  $\lambda \in \Lambda([-n, n], r)$  with  $\lambda \geq \sigma(A)$ , (4.10.1) implies

$$(4.12.1) \quad \mathbf{e}^{(A^+)} \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mathbf{k}_i; 0 \\ \lambda_i \end{bmatrix} \mathbf{f}^{(A^-)} = \mathbf{e}^{(A^+)} \mathbf{h}_{\lambda, n} \mathbf{f}^{(A^-)} + \sum_{\substack{\mu \in \Lambda([-n, n], r) \\ \lambda <_n \mu}} \prod_{-n \leq i \leq n-1} \begin{bmatrix} \mu_i \\ \lambda_i \end{bmatrix} \mathbf{e}^{(A^+)} \mathbf{h}_{\mu, n} \mathbf{f}^{(A^-)},$$

and  $\mu >_n \lambda$  implies  $\mu \geq \sigma(A)$  (as  $\sigma_i(A) = 0$  for  $i \notin [-n, n-1]$ ), it follows that all  $\mathbf{e}^{(A^+)} \mathbf{h}_{\mu, n} \mathbf{f}^{(A^-)} \in \mathcal{M}'_n$  and  $\text{span}_{\mathcal{Z}} \mathcal{M}_n = \text{span}_{\mathcal{Z}} \mathcal{M}'_n$ . Consequently,  $\text{span}_{\mathcal{Z}} \mathcal{M}_\infty = \text{span}_{\mathcal{Z}} \mathcal{M}'_\infty = V(\infty, r)$  by 4.11. On the other hand, the linear independence of  $\mathcal{M}'_n$  implies that  $\mathcal{M}_n$  is also linearly independent. Hence,  $\mathcal{M}_\infty$  is linearly independent.  $\square$

**Corollary 4.13.** *Let  $A \in \Xi^\pm(\infty)$  and  $\lambda \in \mathbb{N}^\infty$  with  $\sigma(\lambda) \leq r$ . If  $\lambda_i < \sigma_i(A)$  for some  $i \in \mathbb{Z}$  then*

$$\mathbf{e}^{(A^+)} \mathbf{k}_\lambda \mathbf{f}^{(A^-)} = \sum_{\substack{B \leq A \\ \mu \geq \sigma(A), \sigma(\mu) \leq r}} g_{B, \mu} \mathbf{e}^{(B^+)} \mathbf{k}_\mu \mathbf{f}^{(B^-)} \quad (g_{B, \mu} \in \mathcal{Z}, B \in \Xi^\pm(\infty), \mu \in \mathbb{N}^\infty)$$

*Proof.* The assertion follows from (4.10.1), (4.5.1) and (4.12.1).  $\square$

5. INFINITE  $q$ -SCHUR ALGEBRAS

In this section, we will establish the isomorphism between  $\mathbf{U}(\infty, r)$  and  $\mathbf{V}(\infty, r)$  and discuss the relationship between  $\mathbf{U}(\infty, r)$  and the infinite  $q$ -Schur algebra  $\mathbf{S}(\infty, r)$ . We shall mainly work over the field  $\mathbb{Q}(v)$  though results like 5.1 and 5.2 below continue to hold over  $\mathcal{Z}$ .

As in §2, let  $\mathfrak{S}_r$  be the symmetric groups on  $r$  letters. For  $\lambda \in \Lambda(\infty, r)$ , let  $\mathfrak{S}_\lambda$  be the Young subgroup of  $\mathfrak{S}_r$ , and let  $\mathfrak{D}_\lambda$  be the set of distinguished right  $\mathfrak{S}_\lambda$ -coset representatives. Then,  $\mathfrak{D}_{\lambda\mu} = \mathfrak{D}_\lambda \cap \mathfrak{D}_\mu^{-1}$  is the set of distinguished double  $(\mathfrak{S}_\lambda, \mathfrak{S}_\mu)$ -coset representatives.

Let  $\mathcal{H}$  be the associated Hecke algebras over  $\mathbb{Q}(v)$  with basis  $\{T_w := v^{\ell(w)}\mathcal{T}_w\}_{w \in \mathfrak{S}_r}$ , and let

$$x_\lambda = \sum_{w \in \mathfrak{S}_\lambda} T_w.$$

Recall from §2 the  $\mathcal{H}$ -module  $\Omega_\eta^{\otimes r}$ . Clearly, there is an  $\mathcal{H}$ -module embedding  $i_n : \Omega_{[-n, n]}^{\otimes r} \rightarrow \Omega_{[-n-1, n+1]}^{\otimes r}$ , and  $\{(\Omega_{[-n, n]}^{\otimes r}, i_n)\}_{n \geq 1}$  forms a direct system.

**Lemma 5.1.** *The  $\mathcal{H}$ -module  $\Omega_\infty^{\otimes r}$  is the direct limit of  $\mathcal{H}$ -modules  $\{(\Omega_{[-n, n]}^{\otimes r}, i_n)\}_{n \geq 1}$ . Thus,  $\mathcal{H}$ -module  $\Omega_\infty^{\otimes r}$  is isomorphic to the  $\mathcal{H}$ -module  $\bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H}$ .*

*Proof.* By [16, 5.1], we know there is a  $\mathcal{H}$ -module isomorphism  $\alpha_n : \Omega_{[-n, n]}^{\otimes r} \rightarrow \bigoplus_{\lambda \in \Lambda([-n, n], r)} x_\lambda \mathcal{H}$  and the following diagram are commutative.

$$\begin{array}{ccc} \Omega_{[-n, n]}^{\otimes r} & \xrightarrow{\alpha_n} & \bigoplus_{\lambda \in \Lambda([-n, n], r)} x_\lambda \mathcal{H} \\ i_n \downarrow & & \downarrow i'_n \\ \Omega_{[-n-1, n+1]}^{\otimes r} & \xrightarrow{\alpha_{n+1}} & \bigoplus_{\lambda \in \Lambda([-n-1, n+1], r)} x_\lambda \mathcal{H}, \end{array}$$

where  $i_n$  and  $i'_n$  are natural injections. Since  $\Omega_\infty^{\otimes r}$  is the direct limit of  $\{\Omega_{[-n, n]}^{\otimes r}\}_{n \geq 1}$  and  $\bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H}$  is the direct limit of  $\{\bigoplus_{\lambda \in \Lambda([-n, n], r)} x_\lambda \mathcal{H}\}_{n \geq 1}$ , the result follows.  $\square$

**Proposition 5.2.** *We have the following algebra isomorphisms*

$$\begin{aligned} \mathcal{K}(\infty, r) &\cong \bigoplus_{\mu \in \Lambda(\infty, r)} \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H}) \\ \mathcal{S}(\infty, r) &\cong \prod_{\mu \in \Lambda(\infty, r)} \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H}). \end{aligned}$$

*In particular, the algebra  $\mathcal{K}(\infty, r)$  can be regarded as a subalgebra (without the identity) of  $\mathcal{S}(\infty, r)$ .*

*Proof.* It is known (see, e.g., [11]) that, for any  $n \geq 0$ , the  $q$ -Schur algebras  $\mathcal{K}([-n, n], r)$  is isomorphic to  $\bigoplus_{\lambda, \mu \in \Lambda([-n, n], r)} \text{Hom}_{\mathcal{H}}(x_\lambda \mathcal{H}, x_\mu \mathcal{H})$ . Since  $\mathcal{K}(\infty, r)$  is the direct limit of  $\{\mathcal{K}([-n, n], r)\}_{n \geq 1}$ , the first assertion follows. By 5.1, we have

$$\mathcal{S}(\infty, r) \cong \prod_{\mu \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, \bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H}).$$

Now, one checks easily that, for each  $\mu \in \Lambda(\infty, r)$ , there is a natural isomorphism

$$\bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H}) \cong \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, \bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H}),$$

which induces the required algebra isomorphism.  $\square$

The isomorphism  $\theta$  can be made explicit. For  $\lambda \in \Lambda(\infty, r)$ , let  $Y(\lambda)$  be the Young diagram of  $\lambda$  which is a collection of boxes, arranged in left justified rows with  $\lambda_i$  boxes in row  $i$  for all  $i \in \mathbb{Z}$ , and let  $t^\lambda$  be the  $\lambda$ -tableau in which the numbers  $1, 2, \dots, r$  appear in order from left to right down successive (non-empty) rows of  $Y(\lambda)$ . Let  $R_i^\lambda$  ( $i \in \mathbb{Z}$ ) be the set of entries in the  $i$ -th row of  $t^\lambda$ . Clearly,  $R_i^\lambda \neq \emptyset$  if and only if  $\lambda_i > 0$ . Let

$$(5.2.1) \quad \mathfrak{D}(\infty, r) = \{(\lambda, d, \mu) \mid \lambda, \mu \in \Lambda(\infty, r), d \in \mathfrak{D}_{\lambda\mu}\}.$$

By [18, (1.3.10)], every element  $(\lambda, d, \mu) \in \mathfrak{D}(\infty, r)$  defines a matrix  $A = (a_{i,j})_{i,j \in \mathbb{Z}} \in \Xi(\infty, r)$  such that  $a_{i,j} = |R_i^\lambda \cap dR_j^\mu|$ . This defines a bijective map

$$j : \mathfrak{D}(\infty, r) \longrightarrow \Xi(\infty, r).$$

Define, for any  $\lambda, \mu \in \Lambda(\infty, r)$  and  $w \in \mathfrak{D}_{\lambda\mu}$ , a map

$$(5.2.2) \quad \phi_{\lambda\mu}^w : \bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H} \rightarrow \bigoplus_{\lambda \in \Lambda(\infty, r)} x_\lambda \mathcal{H}$$

by setting  $\phi_{\lambda\mu}^w(x_\nu h) = \delta_{\mu,\nu} \sum_{x \in \mathfrak{S}_{\lambda w \mathfrak{S}_\mu}} T_x h$ . Then, the set  $\{\phi_{\lambda\mu}^w \mid (\lambda, d, \mu) \in \mathfrak{D}(\infty, r)\}$  forms a basis for  $\bigoplus_{\mu \in \Lambda(\infty, r)} \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H})$ . By [11, 1.4], the isomorphism  $\theta$  is induced by  $j$ . In other words, if  $j(\lambda, w, \mu) = A$ , then  $\theta(e_A) = \phi_{\lambda\mu}^w$ . In particular, by 3.4, we have  $\theta(\mathbf{k}_\lambda) = \phi_{\lambda\lambda}^1$ .

We now identify  $\mathcal{S}(\infty, r)$  as  $\prod_{\mu \in \Lambda(\infty, r)} \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H})$  under  $\tilde{\theta}$ . Thus, the elements of  $\mathcal{S}(\infty, r)$  have the form  $\left( \sum_{\lambda \in \Lambda(\infty, r)} f_{\lambda,\mu} \right)_{\mu \in \Lambda(\infty, r)}$ , where  $f_{\lambda,\mu} \in \text{Hom}_{\mathcal{H}}(x_\mu \mathcal{H}, x_\lambda \mathcal{H})$  for all  $\lambda, \mu \in \Lambda(\infty, r)$ , which represents the map given by

$$\left( \sum_{\lambda \in \Lambda(\infty, r)} f_{\lambda,\mu} \right)_{\mu \in \Lambda(\infty, r)} \left( \sum_{\nu \in \Lambda(\infty, r)} a_\nu \right) = \sum_{\lambda, \mu \in \Lambda(\infty, r)} f_{\lambda,\mu}(a_\mu),$$

where  $\sum_\nu a_\nu \in \bigoplus_\lambda x_\lambda \mathcal{H}$ . This identification allows us to regard the infinite  $q$ -Schur algebra  $\mathcal{S}(\infty, r)$  as the  $\dagger$ -completion of  $\mathcal{K}(\infty, r)$  (see 1.1).

**Proposition 5.3.** *Let  $\widehat{\mathcal{K}}^\dagger(\infty, r)$  be the completion algebra of  $\mathcal{K}(\infty, r)$  constructed as in 1.1. Then there is an algebra isomorphism*

$$\varsigma_r : \widehat{\mathcal{K}}^\dagger(\infty, r) \xrightarrow{\sim} \mathcal{S}(\infty, r), \quad \sum_{A \in \Xi(\infty, r)} \beta_A [A] \longmapsto \left( \sum_{\substack{\text{co}(A) = \mu \\ A \in \Xi(\infty, r)}} \beta_A [A] \right)_{\mu \in \Lambda(\infty, r)}.$$

*Proof.* Clearly,  $\varsigma_r$  is induced by the natural monomorphism  $\varsigma_r : \mathcal{K}(\infty, r) \rightarrow \mathcal{S}(\infty, r)$  given in 5.2.  $\square$

By 1.1, there is another completion algebra  $\widehat{\mathcal{K}}(\infty, r)$  which is a subalgebra of  $\widehat{\mathcal{K}}^\dagger(\infty, r)$  and which contains  $\mathbf{V}(\infty, r)$  as a subalgebras. Thus, restriction gives an algebra monomorphism  $\varsigma_r : \mathbf{V}(\infty, r) \rightarrow \mathcal{S}(\infty, r)$ .

We are now ready to establish the isomorphism  $\mathbf{U}(\infty, r) \cong \mathbf{V}(\infty, r)$ .

There is a similar basis  $\{\phi_{\lambda\mu}^d \mid \lambda, \mu \in \Lambda([-n, n], r), d \in \mathfrak{D}_{\lambda\mu}\}$  for  $\mathcal{S}([-n, n], r)$  by [6], and  $[A] = v^{-d_A} \phi_{\lambda\mu}^d$ , where  $A = j(\lambda, d, \mu)$ , if we identify  $\mathbf{U}([-n, n], r)$  with  $\mathcal{S}([-n, n], r)$  (cf. [10, A.1]).

**Theorem 5.4.** *The following diagram is commutative*

$$\begin{array}{ccc} \mathbf{V}(\infty) & \xrightarrow{\xi_r} & \mathbf{V}(\infty, r) \\ \sim \downarrow & & \downarrow \varsigma_r \\ \mathbf{U}(\infty) & \xrightarrow[\varsigma_r]{} & \mathcal{S}(\infty, r) \end{array}$$

Hence, we have an isomorphism  $\mathbf{U}(\infty, r) \cong \mathbf{V}(\infty, r)$ . Moreover, the map  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathcal{S}(\infty, r)$  is not surjective for any  $r \geq 1$ .

*Proof.* Recall from 3.3 that we may identify  $\mathbf{U}(\infty)$  with  $\mathbf{V}(\infty)$ . Fix some  $i \in \mathbb{Z}$ . For  $\omega \in \Omega_\infty^{\otimes r}$ , we choose  $n \geq 1$  such that  $-n \leq i < n$  and  $\omega \in \Omega_{[-n, n]}^{\otimes r}$ . Then  $E_i \cdot \omega = E_{i, i+1}(\mathbf{0}, r)_{[-n, n]} \cdot \omega$  by a result for  $q$ -Schur algebras. It is clear that we have  $E_{i, i+1}(\mathbf{0}, r)_{[-n, n]} \cdot \omega = E_{i, i+1}(\mathbf{0}, r)_\infty \cdot \omega$ . Hence,  $E_i \cdot \omega = E_{i, i+1}(\mathbf{0}, r)_\infty \cdot \omega$  for any  $\omega \in \Omega_\infty^{\otimes r}$ . Similarly, we have  $F_i \cdot \omega = E_{i+1, i}(\mathbf{0}, r)_\infty \cdot \omega$  and  $K_i \cdot \omega = 0(e_i, r)_\infty \cdot \omega$  for any  $\omega \in \Omega_\infty^{\otimes r}$ . The first assertion follows.

To see the last statement, it suffices to prove that the injective map  $\zeta_r : \widehat{\mathcal{K}}(\infty, r) \rightarrow \mathcal{S}(\infty, r)$  is not surjective. Fix a  $\lambda \in \Lambda(\infty, r)$ . For any  $\mu \in \Lambda(\infty, r)$ , construct a matrix  $A_\mu \in \Xi(\infty, r)$  such that  $ro(A_\mu) = \lambda$  and  $co(A_\mu) = \mu$ . Clearly, the element  $([A_\mu])_{\mu \in \Lambda(\infty, r)}$  belongs to  $\mathcal{S}(\infty, r)$ , but not in  $\widehat{\mathcal{K}}(\infty, r)$ .  $\square$

**Remark 5.5.** Recall from [20] that, in the finite case, the  $(\mathbf{U}(n), \mathcal{H})$ -bimodule structure on  $\Omega_n^{\otimes r}$  induces two algebra epimorphisms

$$\mathbf{U}(n) \twoheadrightarrow \text{End}_{\mathcal{H}}(\Omega_n^{\otimes r}), \quad \mathcal{H} \twoheadrightarrow \text{End}_{\mathbf{U}(n)}(\Omega_n^{\otimes r})$$

(see [15] for the roots-of-unity case). This is the so-called *Schur–Weyl duality*. The theorem above shows that this duality is no longer true in the infinite case. However, the second epimorphism continues to hold; see 8.3 below.

With the above result, we will identify  $\mathbf{U}(\infty, r)$  with  $\mathbf{V}(\infty, r)$ . Thus, the algebra homomorphism  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathcal{S}(\infty, r)$  sends  $E_i, K_i, F_i$  to  $e_i, k_i, f_i$ , respectively. Now, by applying  $\zeta_r$  to the graded components in (2.1.1), 3.5 implies immediately the following.

**Corollary 5.6.** *Let  $\lambda \in \Lambda(\infty, r)$  and  $\nu', \nu'' \in \mathbb{Z}\Pi(\infty)$ . If  $t' \in \mathbf{U}(\infty)_{\nu'}$  and  $t'' \in \mathbf{U}(\infty)_{\nu''}$ , then*

$$\zeta_r(t')[\text{diag}(\lambda)] = \begin{cases} [\text{diag}(\lambda + \nu')] \zeta_r(t') & \text{if } \lambda + \nu' \in \Lambda(\infty, r); \\ 0 & \text{otherwise,} \end{cases}$$

and

$$[\text{diag}(\lambda)] \zeta_r(t'') = \begin{cases} \zeta_r(t'')[\text{diag}(\lambda - \nu'')] & \text{if } \lambda - \nu'' \in \Lambda(\infty, r); \\ 0 & \text{otherwise.} \end{cases}$$

## 6. MODIFIED QUANTUM $\mathfrak{gl}_\infty$ AND RELATED ALGEBRAS

In this section, we identify  $\mathbf{U}(\infty)$  with  $\mathbf{V}(\infty)$  and  $\mathbf{U}(\infty, r)$  with  $\mathbf{V}(\infty, r)$ . Thus,  $E_h = E_{h, h+1}(\mathbf{0})$  and  $F_h = E_{h+1, h}(\mathbf{0})$  for all  $h \in \mathbb{Z}$ .

The interpretation of the infinite  $q$ -Schur algebra  $\mathcal{S}(\infty, r)$  as the  $\dagger$ -completion  $\widehat{\mathcal{K}}^\dagger(\infty, r)$  of  $\mathcal{K}(\infty, r)$  suggests us to introduce the  $\dagger$ -completion  $\widehat{\mathcal{K}}^\dagger(\infty)$  of the algebra  $\mathcal{K}(\infty)$  which contains  $\widehat{\mathcal{K}}(\infty)$ , and hence  $\mathbf{U}(\infty)$ , as a subalgebra. Thus, it is natural to expect that the map  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathcal{S}(\infty, r)$  given in (2.2.1) may be extended to an epimorphism from  $\widehat{\mathcal{K}}^\dagger(\infty)$  to  $\mathcal{S}(\infty, r)$ . In this section, we will establish this epimorphism through an investigation of the epimorphism  $\zeta_r : \mathcal{K}(\infty) \rightarrow \mathcal{K}(\infty, r)$  induced by  $\zeta_r$  via the modified quantum group  $\widehat{\mathbf{U}}(\infty)$  of  $\mathbf{U}(\infty)$  (see [28, 23.1]).

Recall from 3.3 that  $\mathbf{U}(\infty)$  has a basis  $\{A(\mathbf{j})\}_{A, \mathbf{j}}$ . Thus, for any  $\lambda, \mu \in \mathbb{Z}^\infty$ , there is a linear map from  $\mathbf{U}(\infty)$  to  $\mathcal{K}(\infty)$  sending  $u$  to  $[\text{diag}(\lambda)]u[\text{diag}(\mu)]$ . Let

$$\lambda \mathbf{K}_\mu = \sum_{\mathbf{j} \in \mathbb{Z}^\infty} (K^{\mathbf{j}} - v^{\lambda \cdot \mathbf{j}}) \mathbf{U}(\infty) + \sum_{\mathbf{j} \in \mathbb{Z}^\infty} \mathbf{U}(\infty) (K^{\mathbf{j}} - v^{\mu \cdot \mathbf{j}}) \quad \text{and} \quad \lambda \overline{\mathbf{U}(\infty)}_\mu := \mathbf{U}(\infty) / \lambda \mathbf{K}_\mu.$$



Since  $[\text{diag}(\lambda)]_\lambda \mathbf{K}_\mu [\text{diag}(\mu)] = 0$ , this map induces a linear map  ${}_\lambda \overline{\mathbf{U}(\infty)}_\mu \rightarrow \mathcal{K}(\infty)$  sending  $\pi_{\lambda\mu}(u)$  to  $[\text{diag}(\lambda)]u[\text{diag}(\mu)]$ , where  $\pi_{\lambda\mu} : \mathbf{U}(\infty) \rightarrow {}_\lambda \overline{\mathbf{U}(\infty)}_\mu$  is the canonical projection. Thus, we obtain a linear map

$$f : \dot{\mathbf{U}}(\infty) := \bigoplus_{\lambda, \mu \in \mathbb{Z}^\infty} {}_\lambda \overline{\mathbf{U}(\infty)}_\mu \longrightarrow \mathcal{K}(\infty).$$

We will introduce a multiplication in  $\dot{\mathbf{U}}(\infty)$  and prove that  $f$  is an algebra isomorphism. We need some preparation.

**Lemma 6.1.** *Let  $A \in \Xi^\pm(\infty)$  and  $\lambda, \mu \in \mathbb{Z}^\infty$ . If  $\lambda - \mu \neq \text{ro}(A) - \text{co}(A)$ , then we have  $\pi_{\lambda\mu}(A(\mathbf{j})) = 0$  for any  $\mathbf{j} \in \mathbb{Z}^\infty$ .*

*Proof.* Since  $\mu - \text{co}(A) \neq \lambda - \text{ro}(A)$ , there exist  $\mathbf{j}' \in \mathbb{Z}^\infty$  such that  $\mathbf{j}' \cdot (\mu - \text{co}(A)) \neq \mathbf{j}' \cdot (\lambda - \text{ro}(A))$ . Since  $v^{-\mathbf{j}' \cdot \text{ro}(A)} 0(\mathbf{j}') A(\mathbf{j}) = A(\mathbf{j} + \mathbf{j}') = v^{-\mathbf{j}' \cdot \text{co}(A)} A(\mathbf{j}) 0(\mathbf{j}')$  we have for any  $\mathbf{j} \in \mathbb{Z}^\infty$

$$(v^{\mathbf{j}' \cdot (\mu - \text{co}(A))} - v^{\mathbf{j}' \cdot (\lambda - \text{ro}(A))}) A(\mathbf{j}) = v^{-\mathbf{j}' \cdot \text{ro}(A)} (K^{\mathbf{j}'} - v^{\lambda \cdot \mathbf{j}'}) A(\mathbf{j}) - v^{-\mathbf{j}' \cdot \text{co}(A)} A(\mathbf{j}) (K^{\mathbf{j}'} - v^{\mu \cdot \mathbf{j}'}).$$

The assertion follows.  $\square$

Recall the algebra grading of  $\mathbf{U}(\infty)$  given in (2.1.1).

**Lemma 6.2.** *For  $\lambda, \mu \in \mathbb{Z}^\infty$  and  $\nu \in \mathbb{Z}\Pi(\infty)$ . If  $\nu \neq \lambda - \mu$  then we have  $\pi_{\lambda\mu}(\mathbf{U}(\infty)_\nu) = 0$ . Hence,  ${}_\lambda \overline{\mathbf{U}(\infty)}_\mu = \pi_{\lambda\mu}(\mathbf{U}(\infty)_{\lambda - \mu})$ .*

*Proof.* Let  $t \in \mathbf{U}(\infty)_\nu$ . Since  $K^{\mathbf{j}t} = v^{\nu \cdot \mathbf{j}t} K^{\mathbf{j}}$  for  $\mathbf{j} \in \mathbb{Z}^\infty$ , we have  $v^{\lambda \cdot \mathbf{j}} \pi_{\lambda, \mu}(t) = \pi_{\lambda\mu}(K^{\mathbf{j}t}) = \pi_{\lambda\mu}(v^{\nu \cdot \mathbf{j}t} K^{\mathbf{j}}) = v^{\mu \cdot \mathbf{j} + \nu \cdot \mathbf{j}} \pi_{\lambda\mu}(t)$  for all  $\mathbf{j} \in \mathbb{Z}^\infty$ . Since  $\nu \neq \lambda - \mu$  there exist  $\mathbf{j} \in \mathbb{Z}^\infty$  such that  $\nu \cdot \mathbf{j} \neq (\lambda - \mu) \cdot \mathbf{j}$ . Hence,  $\pi_{\lambda\mu}(t) = 0$ .  $\square$

For any  $\lambda', \mu', \lambda'', \mu'' \in \mathbb{Z}^\infty$  with  $\lambda' - \mu', \lambda'' - \mu'' \in \mathbb{Z}\Pi(\infty)$  and any  $t \in \mathbf{U}(\infty)_{\lambda' - \mu'}$ ,  $s \in \mathbf{U}(\infty)_{\lambda'' - \mu''}$ , define the product in  $\dot{\mathbf{U}}(\infty)$

$$\pi_{\lambda'\mu'}(t) \pi_{\lambda''\mu''}(s) = \begin{cases} \pi_{\lambda'\mu''}(ts), & \text{if } \mu' = \lambda'' \\ 0 & \text{otherwise.} \end{cases}$$

Using 6.2 one can easily check the above product defines an associative  $\mathbb{Q}(v)$ -algebra structure on  $\dot{\mathbf{U}}(\infty)$ .

**Theorem 6.3.** *The linear map  $f : \dot{\mathbf{U}}(\infty) \rightarrow \mathcal{K}(\infty)$  sending  $\pi_{\lambda\mu}(u)$  to  $[\text{diag}(\lambda)]u[\text{diag}(\mu)]$  for all  $u \in \mathbf{U}(\infty)$  and  $\lambda, \mu \in \mathbb{Z}^\infty$ , is an algebra isomorphism.*

*Proof.* Since, for any  $A \in \tilde{\Xi}^\pm(\infty)$ ,  $\mathbf{j} \in \mathbb{Z}^\infty$  and  $\lambda, \mu \in \mathbb{Z}^\infty$  with  $\lambda - \mu = \text{ro}(A) - \text{co}(A)$ , we have

$$(6.3.1) \quad [\text{diag}(\lambda)]A(\mathbf{j})[\text{diag}(\mu)] = v^{(\lambda - \text{ro}(A)) \cdot \mathbf{j}} [A + \text{diag}(\lambda - \text{ro}(A))] = v^{(\mu - \text{co}(A)) \cdot \mathbf{j}} [A + \text{diag}(\mu - \text{co}(A))],$$

it follows that  $[A] = [\text{diag}(\text{ro}(A))]A^\pm(\mathbf{0})[\text{diag}(\text{co}(A))]$  for all  $A \in \tilde{\Xi}(\infty)$ , and so

$$\mathcal{K}(\infty) = \bigoplus_{\lambda, \mu \in \mathbb{Z}^\infty} [\text{diag}(\lambda)]\mathbf{U}(\infty)[\text{diag}(\mu)].$$

So  $f$  is surjective. Suppose now  $\sum_{\lambda, \mu} \pi_{\lambda\mu}(u) \in \ker(f)$ , where  $u = \sum_{\substack{A \in \tilde{\Xi}^\pm(\infty) \\ \mathbf{j} \in \mathbb{Z}^\infty}} \beta_{A, \mathbf{j}} A(\mathbf{j})$  with  $\beta_{A, \mathbf{j}} \in \mathbb{Q}(v)$ . Then,  $[\text{diag}(\lambda)]u[\text{diag}(\mu)] = 0$  for all  $\lambda, \mu$ . Let

$$u_{\lambda\mu} = \sum_{\substack{A \in \tilde{\Xi}^\pm(\infty), \mathbf{j} \in \mathbb{Z}^\infty \\ \lambda - \mu = \text{ro}(A) - \text{co}(A)}} \beta_{A, \mathbf{j}} A(\mathbf{j}).$$

By 6.1 we have  $\pi_{\lambda\mu}(u) = \pi_{\lambda\mu}(u_{\lambda\mu})$ . On the other hand, (6.3.1) implies

$$0 = [\text{diag}(\lambda)]u_{\lambda\mu}[\text{diag}(\mu)] = \sum_{\substack{A \in \Xi^\pm(\infty) \\ \lambda - \mu = \text{ro}(A) - \text{co}(A)}} \left( \sum_{\mathbf{j} \in \mathbb{Z}^\infty} v^{(\lambda - \text{ro}(A)) \cdot \mathbf{j}} \beta_{A, \mathbf{j}} \right) [A + \text{diag}(\lambda) - \text{ro}(A)].$$

Hence,  $\sum_{\mathbf{j} \in \mathbb{Z}^\infty} v^{(\lambda - \text{ro}(A)) \cdot \mathbf{j}} \beta_{A, \mathbf{j}} = 0$  for any  $A \in \Xi^\pm(\infty)$  with  $\lambda - \mu = \text{ro}(A) - \text{co}(A)$ . Since  $A(\mathbf{j}) = v^{-\mathbf{j} \cdot \text{ro}(A)} 0(\mathbf{j}) A(\mathbf{0})$  we have

$$\begin{aligned} u_{\lambda\mu} &= \sum_{\substack{A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{Z}^\infty \\ \lambda - \mu = \text{ro}(A) - \text{co}(A)}} \beta_{A, \mathbf{j}} A(\mathbf{j}) - \sum_{\substack{A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{Z}^\infty \\ \lambda - \mu = \text{ro}(A) - \text{co}(A)}} v^{(\lambda - \text{ro}(A)) \cdot \mathbf{j}} \beta_{A, \mathbf{j}} A(\mathbf{0}) \\ &= \sum_{\substack{A \in \Xi^\pm(\infty), \mathbf{j} \in \mathbb{Z}^\infty \\ \lambda - \mu = \text{ro}(A) - \text{co}(A)}} (K^{\mathbf{j}} - v^{\lambda \cdot \mathbf{j}}) v^{-\mathbf{j} \cdot \text{ro}(A)} \beta_{A, \mathbf{j}} A(\mathbf{0}) \in {}_\lambda \mathbf{K}_\mu. \end{aligned}$$

Hence,  $\pi_{\lambda\mu}(u) = \pi_{\lambda\mu}(u_{\lambda\mu}) = 0$  for all  $\lambda, \mu$ . So  $f$  is injective.

We now prove  $f$  is an algebra homomorphism. Let  $u_1, u_2 \in \mathbf{U}(\infty)$  and  $\lambda', \mu', \lambda'', \mu'' \in \mathbb{Z}^\infty$ . If  $\mu' \neq \lambda''$  or  $\lambda' - \mu' \notin \mathbb{Z}\Pi(\infty)$  or  $\lambda'' - \mu'' \notin \mathbb{Z}\Pi(\infty)$  then by definition and 6.2

$$f(\pi_{\lambda'\mu'}(u_1)\pi_{\lambda''\mu''}(u_2)) = 0 = f(\pi_{\lambda'\mu'}(u_1))f(\pi_{\lambda''\mu''}(u_2)).$$

It remains to prove the case when  $\mu' = \lambda''$ ,  $\lambda' - \mu' \in \mathbb{Z}\Pi(\infty)$  and  $\lambda'' - \mu'' \in \mathbb{Z}\Pi(\infty)$ . By 6.2, we may assume  $u_1 \in \mathbf{U}(\infty)_{\lambda' - \mu'}$  and  $u_2 \in \mathbf{U}(\infty)_{\lambda'' - \mu''}$ . Observe that, for  $u = E^{(A^+)} K^{\mathbf{j}} F^{(A^-)} \in \mathbf{U}(\infty)_{\lambda - \mu}$ , where  $\lambda, \mu \in \mathbb{Z}^\infty$  with  $\lambda - \mu \in \mathbb{Z}\Pi(\infty)$ , since  $\lambda - \mu = \sum_{i \leq h < j} (a_{ij} - a_{ji}) \alpha_h$ , it follows from 3.2 that

$$[\text{diag}(\lambda)]u[\text{diag}(\mu)] = [\text{diag}(\lambda)][\text{diag}(\mu + \sum_{i \leq k < j} (a_{ij} - a_{ji}) \alpha_k)]u = [\text{diag}(\lambda)]u.$$

Hence, by 2.2,  $[\text{diag}(\lambda)]u'[\text{diag}(\mu)] = [\text{diag}(\lambda)]u'$  for any  $u' \in \mathbf{U}(\infty)_{\lambda - \mu}$ . Thus, if  $\mu' = \lambda''$ , then

$$\begin{aligned} f(\pi_{\lambda', \mu'}(u_1)\pi_{\lambda'', \mu''}(u_2)) &= f(\pi_{\lambda', \mu''}(u_1 u_2)) = [\text{diag}(\lambda')]u_1 u_2 [\text{diag}(\mu'')] \\ &= ([\text{diag}(\lambda')]u_1 [\text{diag}(\mu')])u_2 [\text{diag}(\mu'')] \\ &= f(\pi_{\lambda', \mu'}(u_1))f(\pi_{\lambda'', \mu''}(u_2)). \end{aligned}$$

as required.  $\square$

With the above result, we shall identify  $\dot{\mathbf{U}}(\infty)$  with  $\mathbf{K}(\infty)$ . In particular, we shall identify  $\pi_{\lambda\mu}(u)$  with  $[\text{diag}(\lambda)]u[\text{diag}(\mu)]$  for  $u \in \mathbf{U}(\infty)$  and  $\lambda, \mu \in \mathbb{Z}^\infty$ .

Since both algebras  $\mathbf{K}(\infty) = \dot{\mathbf{U}}(\infty)$  and  $\mathbf{V}(\infty) = \mathbf{U}(\infty)$  are subalgebras of  $\widehat{\mathbf{K}}(\infty)$ , the definition of  $\mathbf{V}(\infty)$  implies that the algebra  $\mathbf{K}(\infty)$  is a  $\mathbf{U}(\infty)$ -bimodule with an action induced by multiplication. We now have the following interpretation of this bimodule structure.

**Corollary 6.4.** *The algebra  $\mathbf{U}(\infty)$  acts on  $\dot{\mathbf{U}}(\infty)$  by the following rules:*

$$t' \pi_{\lambda\mu}(s) t'' = \pi_{\lambda + \nu', \mu - \nu''}(t' s t'')$$

for all  $t' \in \mathbf{U}(\infty)(\nu')$ ,  $t'' \in \mathbf{U}(\infty)(\nu'')$  and  $s \in \mathbf{U}(\infty)$ , where  $\lambda, \mu \in \mathbb{Z}^\infty$  and  $\nu', \nu'' \in \mathbb{Z}\Pi(\infty)$ .

*Proof.* By 3.2 and 2.2 we have  $t[\text{diag}(\lambda)] = [\text{diag}(\lambda + \nu)]t$  and  $[\text{diag}(\lambda)]t = t[\text{diag}(\lambda - \nu)]$  for  $t \in \mathbf{U}(\infty)_\nu$ . Hence, we have  $t' \pi_{\lambda\mu}(s) t'' = t' [\text{diag}(\lambda)] s [\text{diag}(\mu)] t'' = [\text{diag}(\lambda + \nu')] t' s t'' [\text{diag}(\mu + \nu'')] = \pi_{\lambda + \nu', \mu - \nu''}(t' s t'')$ .  $\square$

Similar to [29, 3.4], we define a map  $\zeta_r$  from  $\dot{\mathbf{U}}(\infty)$  to  $\mathbf{U}(\infty, r)$  as follows.

**Theorem 6.5.** *The map  $\dot{\zeta}_r : \dot{\mathbf{U}}(\infty) \rightarrow \mathbf{U}(\infty, r)$  defined by*

$$\dot{\zeta}_r(\pi_{\lambda\mu}(u)) = \begin{cases} [\text{diag}(\lambda)]\zeta_r(u)[\text{diag}(\mu)] & \text{if } \lambda, \mu \in \Lambda(\infty, r); \\ 0 & \text{otherwise} \end{cases}$$

for  $u \in \mathbf{U}(\infty)$  and  $\lambda, \mu \in \mathbb{Z}^\infty$  is an algebra homomorphism.

*Proof.* We first prove that  $\dot{\zeta}_r$  is well-defined. Assume  $\pi_{\lambda\mu}(u) = 0$  where  $u \in \mathbf{U}(\infty)$  and  $\lambda, \mu \in \Lambda(\infty, r)$ . By the definition of  ${}_\lambda\mathbf{U}(\infty)_\mu$  we can write

$$u = \sum_{\mathbf{j} \in \mathbb{Z}^\infty} (K^{\mathbf{j}} - v^{\lambda \cdot \mathbf{j}})u'_j + \sum_{\mathbf{j} \in \mathbb{Z}^\infty} u''_j(K^{\mathbf{j}} - v^{\mu \cdot \mathbf{j}})$$

where  $u'_j, u''_j \in \mathbf{U}(\infty)$ . By 3.4 we have  $[\text{diag}(\lambda)](\mathbf{k}^{\mathbf{j}} - v^{\lambda \cdot \mathbf{j}}) = 0 = (\mathbf{k}^{\mathbf{j}} - v^{\mu \cdot \mathbf{j}})[\text{diag}(\mu)]$  for  $\mathbf{j} \in \mathbb{Z}^\infty$ . It follows that  $[\text{diag}(\lambda)]\zeta_r(u)[\text{diag}(\mu)] = 0$ . Hence,  $\dot{\zeta}_r$  is well defined.

For convenience, we let  $[\text{diag}(\lambda)] = 0 \in \mathbf{U}(\infty, r)$  for  $\lambda \notin \Lambda(\infty, r)$ . Let  $t \in \mathbf{U}(\infty)(\lambda' - \mu')$  and  $s \in \mathbf{U}(\infty)(\lambda'' - \mu'')$  where  $\lambda', \mu', \lambda'', \mu'' \in \mathbb{Z}^\infty$  such that  $\lambda' - \mu', \lambda'' - \mu'' \in \mathbb{Z}\Pi(\infty)$ . If  $\mu' \neq \lambda''$  then by the definition of  $\dot{\zeta}_r$  we have  $\dot{\zeta}_r(\pi_{\lambda'\mu'}(t)\pi_{\lambda''\mu''}(s)) = 0 = \dot{\zeta}_r(\pi_{\lambda'\mu'}(t))\dot{\zeta}_r(\pi_{\lambda''\mu''}(s))$ . Now we assume  $\mu' = \lambda''$ . Then by 5.6 we have  $\dot{\zeta}_r(\pi_{\lambda'\mu'}(t))\dot{\zeta}_r(\pi_{\mu'\mu''}(s)) = [\text{diag}(\lambda')]\zeta_r(t)[\text{diag}(\mu')]\zeta_r(s)[\text{diag}(\mu'')] = [\text{diag}(\lambda')]\zeta_r(t)\zeta_r(s)[\text{diag}(\mu'')] = [\text{diag}(\lambda')]\zeta_r(ts)[\text{diag}(\mu'')] = \dot{\zeta}_r(\pi_{\lambda'\mu''}(ts)) = \dot{\zeta}_r(\pi_{\lambda'\mu'}(t)\pi_{\mu'\mu''}(s))$ . The result follows.  $\square$

**Proposition 6.6.** *The map  $\dot{\zeta}_r$  satisfies the following property:*

$$\dot{\zeta}_r(u_1 u_2 u_3) = \zeta_r(u_1)\dot{\zeta}_r(u_2)\zeta_r(u_3)$$

where  $u_1, u_3 \in \mathbf{U}(\infty)$  and  $u_2 \in \dot{\mathbf{U}}(\infty)$ .

*Proof.* We assume  $u' \in \mathbf{U}(\infty)(\nu')$ ,  $u'' \in \mathbf{U}(\infty)(\nu'')$  and  $u \in \mathbf{U}(\infty)(\lambda - \mu)$  where  $\nu', \nu'', \lambda - \mu \in \mathbb{Z}\Pi(\infty)$  and  $\lambda, \mu \in \mathbb{Z}^\infty$ . By 6.4 we have

$$\dot{\zeta}_r(u'\pi_{\lambda\mu}(u)u'') = \dot{\zeta}_r(\pi_{\lambda+\nu', \mu-\nu''}(u'uu'')) = \begin{cases} \zeta_r(u')\zeta_r(u)\zeta_r(u'')[\text{diag}(\mu - \nu'')] & \text{if } \mu - \nu'' \in \Lambda(\infty, r); \\ 0 & \text{otherwise.} \end{cases}$$

Hence, by 5.6 we have  $\zeta_r(u')\dot{\zeta}_r(\pi_{\lambda\mu}(u))\zeta_r(u'') = \zeta_r(u')\zeta_r(u)[\text{diag}(\mu)]\zeta_r(u'') = \dot{\zeta}_r(u'\pi_{\lambda\mu}(u)u'')$ .  $\square$

**Proposition 6.7.** *Let  $A \in \tilde{\Xi}(\infty)$ . Then we have*

$$\dot{\zeta}_r([A]) = \begin{cases} [A] & \text{if } A \in \Xi(\infty, r); \\ 0 & \text{otherwise.} \end{cases}$$

In particular we have  $\dot{\zeta}_r(\mathcal{K}(\infty)) = \dot{\zeta}_r(\dot{\mathbf{U}}(\infty)) = \mathcal{K}(\infty, r)$ .

*Proof.* Let  $\lambda = ro(A)$  and  $\mu = co(A)$ . If either  $\lambda \notin \Lambda(\infty, r)$  or  $\mu \notin \Lambda(\infty, r)$ , then we have  $\dot{\zeta}_r([A]) = \dot{\zeta}_r([\text{diag}(\lambda)]A^\pm(\mathbf{0})[\text{diag}(\mu)]) = 0$ . Now we assume  $\lambda, \mu \in \Lambda(\infty, r)$ . Then we have

$$\begin{aligned} \dot{\zeta}_r([A]) &= \dot{\zeta}_r([\text{diag}(\lambda)]A^\pm(\mathbf{0})[\text{diag}(\mu)]) \\ &= [\text{diag}(\lambda)]\zeta_r(A^\pm(\mathbf{0}))[\text{diag}(\mu)] \\ &= [\text{diag}(\lambda)]A^\pm(\mathbf{0}, r)[\text{diag}(\mu)] \\ &= \begin{cases} [A] & \text{if } A \in \Xi(\infty, r); \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

The result follows.  $\square$

**Remark 6.8.** The natural linear map from  $\mathcal{K}(\infty, r)$  to  $\mathcal{K}(\infty)$  by sending  $[A]$  to  $[A]$  for  $A \in \Xi(\infty, r)$  is not an algebra homomorphism. For example, in the algebra  $\mathcal{K}(\infty, r)$  we have  $[E_{1,2}] \cdot [E_{2,1}] = [E_{1,1}]$ . However we have  $[E_{1,2}] \cdot [E_{2,1}] = [E_{1,1}] + [A]$ , where  $A = \begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix}$  in the algebra  $\mathcal{K}(\infty)$ .

By 1.1 we may construct the completion algebra  $\widehat{\mathcal{K}}^\dagger(\infty)$  of  $\mathcal{K}(\infty)$  such that  $\widehat{\mathcal{K}}(\infty)$  becomes an subalgebra of  $\widehat{\mathcal{K}}^\dagger(\infty)$ .

**Theorem 6.9.** *There is an algebra epimorphism  $\tilde{\zeta}_r$  from  $\widehat{\mathcal{K}}^\dagger(\infty)$  to  $\mathcal{S}(\infty, r)$  defined by sending  $\sum_{A \in \tilde{\Xi}(\infty)} \beta_A [A]$  to  $\sum_{A \in \Xi(\infty, r)} \beta_A [A]$ . Moreover we have  $\tilde{\zeta}_r(\widehat{\mathcal{K}}(\infty)) = \widehat{\mathcal{K}}(\infty, r)$  and  $\tilde{\zeta}_r|_{\mathbf{U}(\infty)} = \zeta_r$ .*

*Proof.* By 6.7 there is an algebra homomorphism  $\tilde{\zeta}_r$  from  $\widehat{\mathcal{K}}^\dagger(\infty)$  to  $\mathcal{S}(\infty, r)$  by sending  $\sum_{A \in \tilde{\Xi}(\infty)} \beta_A [A]$  to  $\sum_{A \in \tilde{\Xi}(\infty)} \beta_A \zeta_r([A]) = \sum_{A \in \Xi(\infty, r)} \beta_A [A]$ . It is clear we have  $\tilde{\zeta}_r(\widehat{\mathcal{K}}(\infty)) = \widehat{\mathcal{K}}(\infty, r)$ . By 5.2 the map  $\tilde{\zeta}_r$  is surjective. By 3.3 and 6.7 we have  $\zeta_r(A(\mathbf{j})) = A(\mathbf{j}, r) = \tilde{\zeta}_r(A(\mathbf{j}))$  for  $A \in \Xi^\pm(\infty)$  and  $\mathbf{j} \in \mathbb{Z}^\infty$ . Hence, we have  $\tilde{\zeta}_r|_{\mathbf{U}(\infty)} = \zeta_r$ .  $\square$

**Remark 6.10.** By [1, 3.10] there is an antiautomorphism  $\tau_r$  on  $\mathcal{K}(\infty, r)$  defined by  $\tau_r([A]) = [A^T]$  for  $A \in \Xi(\infty, r)$ . This induces, by 3.1, an antiautomorphism  $\tau$  on  $\mathcal{K}(\infty)$  defined by  $\tau([A]) = [A^T]$  for  $A \in \Xi(\infty)$ . Hence, we have  ${}^\dagger\widehat{\mathcal{K}}(\infty) \cong (\widehat{\mathcal{K}}^\dagger(\infty))^{\text{op}}$  and  ${}^\dagger\widehat{\mathcal{K}}(\infty, r) \cong (\widehat{\mathcal{K}}^\dagger(\infty, r))^{\text{op}}$ .

## 7. HIGHEST WEIGHT REPRESENTATIONS OF $\mathbf{U}(\infty)$

In the rest of the paper, we will discuss the representation theory of  $\mathbf{U}(\infty)$ . There are two types of representations for  $\mathbf{U}(\infty)$ . The first type consists of highest weight representations, while the second type consists of ‘‘polynomial representations’’ arising from representations of infinite  $q$ -Schur algebras. In the quantum  $\mathfrak{gl}_n$  case, it is well-known that polynomial representations more or less determine all highest weight representations. However, we shall see in Theorem 9.3 that polynomial representations and highest weight representations of  $\mathbf{U}(\infty)$  are almost mutually exclusive. In this section, we first discuss the highest weight representations of  $\mathbf{U}(\infty)$ , following the approach used in [28].

For a consecutive segment  $\eta$  of  $\mathbb{Z}$ , let

$$(7.0.1) \quad \begin{aligned} X(\eta) &= \{\lambda = (\lambda_i)_{i \in \eta} \mid \lambda_i \in \eta\}, \quad \text{and} \\ X^+(\eta) &= \{\lambda \in X(\eta) \mid \lambda_i \geq \lambda_{i+1}, \forall i \in \eta^{-1}\} \end{aligned}$$

be the sets of weights and dominant weights, respectively. Thus, both  $\mathbb{Z}^\infty$  and  $\mathbb{N}^\infty$  are subsets of  $X(\infty) := X(\mathbb{Z})$ . Recall from §2 that, for  $i \in \mathbb{Z}$ ,  $\mathbf{e}_i = (\dots, 0, 1, 0 \dots) \in \mathbb{Z}^\infty$  and  $\alpha_i = \mathbf{e}_i - \mathbf{e}_{i+1} \in \mathbb{Z}^\infty$ . Let  $\Pi(\infty) = \{\alpha_i \mid i \in \mathbb{Z}\}$ . We introduce a partial ordering  $\leq_{\text{wt}}$  on  $X(\infty)$  by setting  $\mu \leq_{\text{wt}} \lambda$  if  $\lambda - \mu \in \mathbb{Z}^+ \Pi(\infty)$ . For  $\alpha = \sum_{i \in \mathbb{Z}} k_i \alpha_i \in \mathbb{Z} \Pi(\infty)$  the number  $\text{ht}(\alpha) := \sum_{i \in \mathbb{Z}} k_i$  is called the height of  $\alpha$ .

We first discuss the ‘standard’ representation theory of  $\mathbf{U}(\infty)$  which covers the category  $\mathcal{C}$  of weight modules, the category<sup>3</sup>  $\mathcal{C}^{\text{int}}$  of integrable weight modules, the category  $\mathcal{C}^{\text{hi}}$ , and the category  $\mathcal{O}$ . All these categories are full subcategories of the category  $\mathbf{U}(\infty)\text{-Mod}$  of  $\mathbf{U}(\infty)$ -modules.

Let  $M$  be a  $\mathbf{U}(\infty)$ -module. For  $\lambda \in X(\infty)$ , let  $M_\lambda = \{x \in M \mid K_i x = v^{\lambda_i} x \text{ for } i \in \mathbb{Z}\}$ . If  $M_\lambda \neq 0$ , then  $\lambda$  is called a weight of  $M$ ,  $M_\lambda$  is called a weight space and a nonzero vector in  $M_\lambda$  is called a weight vector of  $M$ . Let  $\text{wt}(M) = \{\lambda \in X(\infty) \mid M_\lambda \neq 0\}$  denote the set of weights of  $M$ . It is clear we have the following lemma.

**Lemma 7.1.** *Let  $M$  be a  $\mathbf{U}(\infty)$ -module. Then  $E_i M_\lambda \subseteq M_{\lambda + \alpha_i}$  and  $F_i M_\lambda \subseteq M_{\lambda - \alpha_i}$  for  $i \in \mathbb{Z}$ .*

<sup>3</sup>This category is denoted as  $\mathcal{C}'$  in [28].

A  $\mathbf{U}(\infty)$ -module  $M$  is called a *weight module*, if  $M = \bigoplus_{\lambda \in X(\infty)} M_\lambda$ . Let  $\mathcal{C}$  denote the full subcategory of  $\mathbf{U}(\infty)\text{-Mod}$  consisting of all weight modules. It is clear every submodule of a weight module and every quotient of a weight module are weight modules. Hence,  $\mathcal{C}$  is an abelian category.

An object  $M \in \mathcal{C}$  is said to be integrable if, for any  $x \in M$  and any  $i \in \mathbb{Z}$ , there exist  $n_0 \geq 1$  such that  $E_i^n x = F_i^n x = 0$  for any  $n \geq n_0$ . Let  $\mathcal{C}^{int}$  be the full subcategory of  $\mathcal{C}$  whose objects are integrable  $\mathbf{U}(\infty)$ -modules.

Let  $\mathcal{C}^{hi}$  be the full subcategory of  $\mathcal{C}$  whose objects are the  $\mathbf{U}(\infty)$ -module  $M$  with the following property: for any  $x \in M$  there exist  $n_0 \geq 1$  such that  $u.x = 0$  where  $u$  is a monomial in the  $E_i$ 's and  $\deg(u) \geq n_0$ .

For  $\lambda \in X(\infty)$  set  $(-\infty, \lambda] = \{\mu \in X(\infty) \mid \mu \leq_{\text{wt}} \lambda\}$ . The category  $\mathcal{O}$  is defined as follows. Its objects are  $\mathbf{U}(\infty)$ -modules  $M$  which are weight modules with finite dimensional weight spaces and such that there exists a finite number of elements  $\lambda^{(1)}, \dots, \lambda^{(s)} \in X(\infty)$  such that  $\text{wt}(M) \subseteq \bigcup_{i=1}^s (-\infty, \lambda^{(i)}]$ .

**Proposition 7.2.** *The category  $\mathcal{O}$  is a full subcategory of  $\mathcal{C}^{hi}$ .*

*Proof.* Let  $M$  be a object of  $\mathcal{O}$ . Then there exist  $\lambda^{(i)} (1 \leq i \leq s)$  such that  $\text{wt}(M) \subseteq \bigcup_{i=1}^s (-\infty, \lambda^{(i)}]$ . Let  $x_\mu \neq 0 \in M_\mu$ . Let  $n_\mu = \max\{\text{ht}(\lambda^{(i)} - \mu) \mid \lambda^{(i)} \geq_{\text{wt}} \mu, 1 \leq i \leq s\}$ . We claim that if  $k \geq n_\mu + 1$ , then for any  $i_1, \dots, i_k \in \mathbb{Z}$  we have  $\mu + \alpha_{i_1} + \alpha_{i_2} + \dots + \alpha_{i_k} \notin \text{wt}(M)$ . Indeed, if there exists  $1 \leq j \leq s$  such that  $\mu + \alpha_{i_1} + \alpha_{i_2} + \dots + \alpha_{i_k} \leq \lambda^{(j)}$ , then  $\text{ht}(\lambda^{(j)} - \mu) \geq k \geq n_\mu + 1$ , contrary to the definition of  $n_\mu$ . Thus, for any  $k \geq n_\mu + 1$  and any  $i_1, \dots, i_k \in \mathbb{Z}$ , we have  $E_{i_1} \dots E_{i_k} x_\mu \in M_{\mu + \alpha_{i_1} + \dots + \alpha_{i_k}} = 0$ . Hence,  $M$  is a object of  $\mathcal{C}^{hi}$ .  $\square$

A  $\mathbf{U}(\infty)$ -module  $M$  is called a *highest weight module* with highest weight  $\lambda \in X(\infty)$  if there exists a nonzero vector  $x_0 \in M_\lambda$  such that  $E_i x_0 = 0$ ,  $K_i x_0 = v^{\lambda_i} x_0$  for all  $i \in \mathbb{Z}$  and  $\mathbf{U}(\infty)x_0 = M$ . The vector  $x_0$  is called a highest weight vector. By a standard argument (see, e.g., [25]) we have the following lemma.

**Lemma 7.3.** *Let  $M$  be a highest weight  $\mathbf{U}(\infty)$ -module with highest weight  $\lambda$ . Then  $M = \bigoplus_{\mu \leq \lambda} M_\mu$  and  $\dim M_\lambda = 1$ . Moreover  $M$  contains a unique maximal submodule.*

For  $\lambda \in X(\infty)$ , let

$$M(\lambda) = \mathbf{U}(\infty) / \left( \sum_{i \in \mathbb{Z}} \mathbf{U}(\infty) E_i + \sum_{i \in \mathbb{Z}} \mathbf{U}(\infty) (K_i - v^{\lambda_i}) \right) = \mathbf{U}^-(\infty) 1_\lambda,$$

where  $1_\lambda$  is the image of 1 in  $M(\lambda)$ . The module  $M(\lambda)$  is called a *Verma module*. By the above lemma, we know that  $M(\lambda)$  has a unique irreducible quotient module  $L(\lambda)$ . It is clear, for  $\lambda \in X(\infty)$ , the modules  $M(\lambda)$  and  $L(\lambda)$  are all in the category  $\mathcal{C}^{hi}$  and the category  $\mathcal{O}$ .

The next result classifies the irreducible modules in the categories  $\mathcal{C}^{hi}$  and  $\mathcal{O}$ .

**Theorem 7.4.** (1) *The map  $\lambda \rightarrow L(\lambda)$  defines a bijection between  $X(\infty)$  and the set of isomorphism classes of irreducible  $\mathbf{U}(\infty)$ -modules in the category  $\mathcal{C}^{hi}$ .*

(2) *The map  $\lambda \rightarrow L(\lambda)$  defines a bijection between  $X(\infty)$  and the set of isomorphism classes of irreducible  $\mathbf{U}(\infty)$ -modules in the category  $\mathcal{O}$ .*

*Proof.* Let  $M$  be a irreducible  $\mathbf{U}(\infty)$ -module in the category  $\mathcal{C}^{hi}$ . Choose a nonzero vector  $x_\mu \in M_\mu$  for some  $\mu \in \text{wt}(M)$ . Let

$$n_0 = \max\{\deg(u) \mid \text{there exists some monomial } u \text{ in the } E_i\text{'s such that } u.x_\mu \neq 0\}.$$

Since  $M \in \mathcal{C}^{hi}$ ,  $n_0 < \infty$ . Let  $u$  be the monomial in the  $E_i$ 's such that  $\deg(u) \geq n_0$  and  $u.x_\mu \neq 0$ . Then  $u.x_\mu$  is a highest weight vector of  $M$ . Let  $\lambda \in X(\infty)$  be the weight of  $u.x_\mu$ . Then there exists

a surjective homomorphism from  $M(\lambda)$  to  $M$ . Since  $L(\lambda)$  is a unique irreducible quotient module of  $M(\lambda)$  we have  $M \cong L(\lambda)$ , proving (1). The statement (2) follows from (1) and 7.2.  $\square$

**Corollary 7.5.** *An irreducible module  $L(\lambda)$  in  $\mathcal{C}^{hi}$  is integrable if and only if  $\lambda \in X^+(\infty)$ .*

*Proof.* If  $\lambda \in X^+(\infty)$ , then the module  $L(\lambda)$  is a homomorphic image of the module  $M(\lambda)/I(\lambda)$ , where  $I(\lambda) = \sum_{i \in \mathbb{Z}} \mathbf{U}^-(\infty) F_i^{(\lambda_i - \lambda_{i+1} + 1)} 1_\lambda$ . Note that  $I(\lambda)$  is a proper submodule of  $M(\lambda)$ . By [28, 3.5.3],  $M(\lambda)/I(\lambda)$  is integrable. Hence,  $L(\lambda)$  is integrable. Conversely, let  $L(\lambda)$  be a irreducible module in the category  $\mathcal{C}^{hi}$ , and assume that  $L(\lambda)$  is integrable. Then, there exists  $x_0 \in M_\lambda$ ,  $x_0 \neq 0$  and  $E_i x_0 = 0$  for all  $i \in \mathbb{Z}$ . By [28, (3.5.8)],  $\lambda \in X^+(\infty)$ .  $\square$

Thus, using 7.2, we have the following classification theorem.

**Theorem 7.6.** (1) *The map  $\lambda \rightarrow L(\lambda)$  defines a bijection between  $X^+(\infty)$  and the set of isomorphism classes of irreducible  $\mathbf{U}(\infty)$ -modules in the category  $\mathcal{C}^{int} \cap \mathcal{C}^{hi}$ .*

(2) *The map  $\lambda \rightarrow L(\lambda)$  defines a bijection between  $X^+(\infty)$  and the set of isomorphism classes of irreducible  $\mathbf{U}(\infty)$ -modules in the category  $\mathcal{C}^{int} \cap \mathcal{O}$ .*

The integrable module  $L(\lambda)$  has actually structure similar to finite dimensional irreducible  $\mathbf{U}([-n, n])$ -modules. Recall from [19, 5.15] that, if  $\mu \in X^+([-n, n])$ , then the irreducible  $\mathbf{U}([-n, n])$ -module  $L(\mu) \cong \mathbf{U}([-n, n])/I_n(\mu)$  where

$$I_n(\mu) = \left( \sum_{-n \leq i < n} (\mathbf{U}([-n, n]) E_i + \mathbf{U}([-n, n]) F_i^{\mu_i - \mu_{i+1} + 1}) + \sum_{-n \leq j \leq n} \mathbf{U}([-n, n]) (K_j - v^{\mu_j}) \right).$$

We want to prove that a similar isomorphism holds for  $\mathbf{U}(\infty)$ .

For any given  $\lambda \in X^+(\infty)$ , let  $\widetilde{L(\lambda)} = M(\lambda)/I(\lambda)$ , and let  $\lambda_{[-n, n]} = (\lambda_i)_{-n \leq i \leq n} \in X^+([-n, n])$ . Let  $\pi_{n, \lambda} : \mathbf{U}([-n, n]) \rightarrow L(\lambda_{[-n, n]})$  be the map sending  $u$  to  $u \bar{1}_\lambda$ , where  $\bar{1}_\lambda$  is the image of  $1_\lambda$ . Since  $I_n(\lambda_{[-n, n]}) \subset I_{n+1}(\lambda_{[-n-1, n+1]})$ , there is a natural  $\mathbf{U}([-n, n])$ -module homomorphism  $\iota_{n, \lambda} : L(\lambda_{[-n, n]}) \rightarrow L(\lambda_{[-n-1, n+1]})$  by sending  $\pi_{n, \lambda}(u)$  to  $\pi_{n+1, \lambda}(u)$  for all  $u \in \mathbf{U}([-n, n])$ , which is compatible with the inclusion  $\mathbf{U}([-n, n]) \hookrightarrow \mathbf{U}([-n-1, n+1])$ . Thus, we obtain a direct system  $\{L(\lambda_{[-n, n]})\}_{n \geq 1}$  whose direct limit  $\lim_{\rightarrow n} L(\lambda_{[-n, n]})$  is naturally a  $\mathbf{U}(\infty)$ -module. Further, since  $\iota_{n, \lambda}(\pi_{n, \lambda}(1)) = \pi_{n+1, \lambda}(1) \neq 0$ , the map  $\iota_{n, \lambda}$  is injective. So we may identify  $L(\lambda_{[-n, n]})$  as a  $\mathbf{U}([-n, n])$ -submodule of  $L(\lambda_{[-n-1, n+1]})$ . Hence,  $\lim_{\rightarrow n} L(\lambda_{[-n, n]}) = \bigcup_{n \geq 1} L(\lambda_{[-n, n]})$ .

**Theorem 7.7.** *We have, for  $\lambda \in X^+(\infty)$ ,*

$$L(\lambda) \cong \lim_{\rightarrow n} L(\lambda_{[-n, n]}) \cong \widetilde{L(\lambda)}.$$

*Proof.* Let  $W$  be a  $\mathbf{U}(\infty)$ -submodule of  $\lim_{\rightarrow n} L(\lambda_{[-n, n]})$ . Since  $L(\lambda_{[-n, n]})$  is irreducible  $\mathbf{U}([-n, n])$ -module for  $n \geq 1$  we have  $W \cap L(\lambda_{[-n, n]}) = 0$  or  $W \cap L(\lambda_{[-n, n]}) = L(\lambda_{[-n, n]})$ . If there exist  $n_0 \geq 1$  such that  $W \cap L(\lambda_{[-n, n]}) \neq 0$  then  $W \cap L(\lambda_{[-n, n]}) \neq 0$  for  $n \geq n_0$ . Hence,  $W \cap L(\lambda_{[-n, n]}) = L(\lambda_{[-n, n]})$  for  $n \geq n_0$ . It follows that

$$W = \bigcup_{n \geq 1} (W \cap L(\lambda_{[-n, n]})) = \bigcup_{n \geq n_0} (W \cap L(\lambda_{[-n, n]})) = \bigcup_{n \geq n_0} L(\lambda_{[-n, n]}) = \lim_{\rightarrow n} L(\lambda_{[-n, n]}).$$

On the other hand, if  $W \cap L(\lambda_{[-n, n]}) = 0$  for  $n \geq 1$  then  $W = 0$ . Hence,  $\lim_{\rightarrow n} L(\lambda_{[-n, n]})$  is an irreducible  $\mathbf{U}(\infty)$ -module. Let  $x_0 := \pi_{1, \lambda}(1) \in \lim_{\rightarrow n} L(\lambda_{[-n, n]})$ . Then  $x_0 = \pi_{n, \lambda}(1)$  for any  $n \geq 1$ . Hence, for  $n \geq 1$  and  $1 \leq i < n$ ,  $1 \leq j \leq n$  we have  $E_i x_0 = \pi_{n, \lambda}(E_i) = 0$  and

$K_j x_0 = \pi_{n,\lambda}(K_j) = v^{\lambda_j} x_0$ . Hence,  $\lim_{\xrightarrow{n}} L(\lambda_{[-n,n]})$  is a irreducible highest weight module with highest weight  $\lambda$ . It follows that  $L(\lambda) \cong \lim_{\xrightarrow{n}} L(\lambda_{[-n,n]})$ . For any  $n \geq 1$  there is a natural  $\mathbf{U}([-n,n])$ -module homomorphism  $f_n$  from  $L(\lambda_{[-n,n]})$  to  $\widetilde{L(\lambda)}$  by sending  $\pi_{n,\lambda}(u)$  to  $\bar{u}$  for  $u \in \mathbf{U}([-n,n])$ . The maps  $f_n$  ( $n \geq 1$ ) induce a surjective  $\mathbf{U}(\infty)$ -homomorphism  $f$  from  $\lim_{\xrightarrow{n}} L(\lambda_{[-n,n]})$  to  $\widetilde{L(\lambda)}$ . Since  $\lim_{\xrightarrow{n}} L(\lambda_{[-n,n]})$  is irreducible  $\mathbf{U}(\infty)$ -module and  $\text{Im}(f) = \widetilde{L(\lambda)} \neq 0$ ,  $f$  has to be an isomorphism.  $\square$

**Remark 7.8.** By [13, 4.4] and [4, §6] we can use the PBW basis  $\{A(\mathbf{0}) \mid A \in \Xi^-(\infty)\}$  of  $\mathbf{U}^-(\infty)$  to define the canonical basis  $\{\mathbf{c}_A \mid A \in \Xi^-(\infty)\}$  of  $\mathbf{U}^-(\infty)$ . For  $\lambda \in X^+(\infty)$ , let  $x_\lambda$  be the highest weight vector of  $L(\lambda)$ . By [27, 8.10] and 7.7, one can easily show that the set  $\{\mathbf{c}_A x_\lambda \mid A \in \Xi^-(\infty)\} \setminus \{0\}$  forms a  $\mathbb{Q}(v)$  basis of  $L(\lambda)$ .

Finally, it is interesting to point out that there are not many *finite dimensional* weight  $\mathbf{U}(\infty)$ -modules. For  $n \geq 0$ , let  $J(n)$  be the two sided ideal of  $\mathbf{U}(\infty)$  generated by  $E_i, F_i, i \in (-\infty, -n) \cup [n, \infty)$ .

**Lemma 7.9.** (1) For  $n \geq 0$ , we have  $J(n) = J(0)$ .

(2) Let  $M$  be a finite dimensional  $\mathbf{U}(\infty)$ -module in the category  $\mathcal{C}$ . Then  $E_i M = F_i M = 0$  for all  $i \in \mathbb{Z}$  and  $\text{wt}(M) \subseteq \{\lambda \in X(\infty) \mid \lambda = k\mathbf{1}\}$ , where  $\mathbf{1} = (\dots, 1, 1, \dots, 1, \dots) \in X(\infty)$ .

*Proof.* Since  $E_n, F_n \in J(n)$ , by 2.1(e) we have  $\widetilde{K}_n - \widetilde{K}_n^{-1} \in J(n)$ . Hence,  $K_n^2 - K_{n+1}^2 \in J(n)$ . By 2.1(b) we have  $K_n^2 E_{n-1} = v^{-2} E_{n-1} K_n^2$  and  $K_{n+1}^2 E_{n-1} = E_{n-1} K_{n+1}^2$ . Hence,  $v^{-2} E_{n-1} K_n^2 - E_{n-1} K_{n+1}^2 = (K_n^2 - K_{n+1}^2) E_{n-1} \in J(n)$ . It follows that  $(v^{-2} - 1) E_{n-1} K_n^2 = (v^{-2} E_{n-1} K_n^2 - E_{n-1} K_{n+1}^2) - E_{n-1} (K_n^2 - K_{n+1}^2) \in J(n)$ . Hence,  $E_{n-1} \in J(n)$ . It is similar we can show  $F_{n-1}, E_{-n}, F_{-n} \in J(n)$ . Hence, we have  $J(n) = J(n-1)$  for  $n \geq 1$ , and (1) follows.

Since  $M$  is finite dimensional weight module, we see that  $\text{wt}(M)$  is a finite set. Assume  $\text{wt}(M) = \{\lambda^{(i)} \mid 1 \leq i \leq s\}$ . By 7.1 for  $1 \leq i \leq s$  there exist  $n_i$  such that  $E_j M_{\lambda^{(i)}} = F_j M_{\lambda^{(i)}} = 0$  for  $j \notin [-n_i, n_i]$ . Let  $n_0 = \max\{n_i \mid 1 \leq i \leq s\}$ . then  $E_i M = F_i M = 0$  for  $i \notin [-n_0, n_0]$ . Hence, by (1) we have  $E_i M = F_i M = 0$  for all  $i \in \mathbb{Z}$ . Let  $\lambda \in \text{wt}(M)$  and  $x_0$  be a nonzero vector in  $M_\lambda$ . By 2.1(e) we have  $\widetilde{K}_i^2 x_0 = v^{2\lambda_i - 2\lambda_{i+1}} x_0 = 0$ . Hence,  $\lambda_i = \lambda_{i+1}$  for all  $i \in \mathbb{Z}$ , proving (2).  $\square$

Let  $\mathcal{C}^{fd}$  be the category of finite dimensional weight  $\mathbf{U}(\infty)$ -module. Now by the above result we have the following classification of finite dimensional  $\mathbf{U}(\infty)$ -modules.

**Theorem 7.10.** The modules  $L(k\mathbf{1})$  ( $k \in \mathbb{Z}$ ) are all non-isomorphic finite dimensional irreducible  $\mathbf{U}(\infty)$ -module in the category  $\mathcal{C}^{fd}$  and  $\dim(L(k\mathbf{1})) = 1$ . Moreover every finite dimensional  $\mathbf{U}(\infty)$ -module in the category  $\mathcal{C}$  is complete reducible.

## 8. REPRESENTATIONS OF $\mathcal{S}(\infty, r)$

In this section, we investigate the weight modules for  $\mathcal{S}(\infty, r)$  and classify the irreducible ones.

Recall from §2 and §5 the basis  $\{T_w\}_{w \in \mathfrak{S}_r}$  for the Hecke algebra  $\mathcal{H}$  and the basis  $\{\phi_{\lambda\mu}^w\}$  for the algebra  $\mathcal{K}(\infty, r)$  given in (5.2.2). Recall also the various notations for the idempotents

$$\mathbf{k}_\lambda = [\text{diag}(\lambda)] = \phi_{\lambda\lambda}^1, \quad \lambda \in \Lambda(\infty, r).$$

Let

$$(8.0.1) \quad \varpi = (\varpi_i)_{i \in \mathbb{Z}} \in \Lambda(\infty, r), \quad \text{where } \varpi_i = \begin{cases} 1, & \text{if } 1 \leq i \leq r; \\ 0, & \text{otherwise.} \end{cases}$$

We first observe the following which is clear from the definition of the  $\dagger$ -completion  $\widehat{\mathcal{K}}^\dagger(\infty, r)$  and the identification  $\mathcal{S}(\infty, r) = \widehat{\mathcal{K}}^\dagger(\infty, r)$  (see 5.3).

**Lemma 8.1.** (1) *Let  $\lambda, \mu \in \Lambda(\infty, r)$ . Then  $\mathcal{S}(\infty, r)\mathbf{k}_\lambda = \mathbf{U}(\infty, r)\mathbf{k}_\lambda = \mathcal{K}(\infty, r)\mathbf{k}_\lambda$ , and  $\mathbf{k}_\lambda\mathcal{S}(\infty, r)\mathbf{k}_\mu = \text{Hom}_{\mathcal{H}}(x_\mu\mathcal{H}, x_\lambda\mathcal{H})$ .*

(2) *The Hecke algebra  $\mathcal{H}$  is isomorphic to  $\mathbf{k}_\varpi\mathcal{S}(\infty, r)\mathbf{k}_\varpi$  by sending  $T_u$  to  $\phi_{\varpi, \varpi}^u$  ( $u \in \mathfrak{S}_r$ ).*

(3) *Let  $\eta$  be a consecutive segment of  $\mathbb{Z}$ . Then, whenever  $|\eta| \geq r$ ,  $\mathcal{H}$  is isomorphic to a (centralizer) subalgebra  $e\mathcal{S}(\eta, r)e$  of  $\mathcal{S}(\eta, r)$  for some idempotent  $e$ .*

An  $\mathcal{S}(\infty, r)$ -module  $M$  is called a *weight module* if  $M = \bigoplus_{\lambda \in \Lambda(\infty, r)} \mathbf{k}_\lambda M$ . Clearly, 8.1(1) implies that  $\mathcal{K}(\infty, r)$  is a weight  $\mathcal{S}(\infty, r)$ -module. Let  $\mathcal{S}(\infty, r)\text{-mod}$  be the category of weight  $\mathcal{S}(\infty, r)$ -modules. Since all  $\mathbf{k}_\lambda \in \mathcal{K}(\infty, r)$ , we can define similarly the categories  $\mathcal{K}(\infty, r)\text{-mod}$  and  $\mathbf{U}(\infty, r)\text{-mod}$ .

**Proposition 8.2.** (1) *Up to category isomorphism, the categories  $\mathcal{K}(\infty, r)\text{-mod}$ ,  $\mathbf{U}(\infty, r)\text{-mod}$  and  $\mathcal{S}(\infty, r)\text{-mod}$  are all the same.*

(2) *The category  $\mathcal{S}(\infty, r)\text{-mod}$  is a full subcategory of  $\mathcal{C}$ .*

*Proof.* The statement (1) follows easily from part (1) of the lemma above. We now prove (2). Using the homomorphism  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathcal{S}(\infty, r)$ , every  $\mathcal{S}(\infty, r)$ -module  $M$  is a  $\mathbf{U}(\infty)$ -module. It suffices to prove that  $\mathbf{k}_\lambda M = M_\lambda$  for all  $\lambda \in \Lambda(\infty, r)$ . If  $x \in M_\lambda$ , then

$$\mathbf{k}_\lambda x = \prod_{i \in \mathbb{Z}} \left( \begin{bmatrix} \mathbf{k}_i; 0 \\ \lambda_i \end{bmatrix} x \right) = \prod_{i \in \mathbb{Z}} \begin{bmatrix} \lambda_i \\ \lambda_i \end{bmatrix} x = x.$$

Hence,  $M_\lambda \subseteq \mathbf{k}_\lambda M$ . Conversely, the inclusion  $\mathbf{k}_\lambda M_\lambda \subseteq M_\lambda$  follows from 3.4(3).  $\square$

Let  $\mathbf{\Omega}(\infty, r) := \mathcal{S}(\infty, r)\mathbf{k}_\varpi$ . If we identify  $\mathcal{H}$  with the subalgebra  $\mathbf{k}_\varpi\mathcal{S}(\infty, r)\mathbf{k}_\varpi$  of  $\mathcal{S}(\infty, r)$  via the isomorphism given in 8.1(2), then  $\mathbf{\Omega}(\infty, r) = \bigoplus_{\mu \in \Lambda(\infty, r)} \phi_{\mu, \varpi}^1 \mathcal{H}$ . Hence,  $\mathbf{\Omega}(\infty, r)$  is an  $(\mathcal{S}(\infty, r), \mathcal{H})$ -bimodule.

**Proposition 8.3.** *The evaluation map*

$$\text{Ev} : \mathbf{\Omega}(\infty, r) \longrightarrow \mathbf{\Omega}_\infty^{\otimes r}, \quad f \mapsto f(T_1)$$

*defines an  $(\mathcal{S}(\infty, r), \mathcal{H})$ -bimodule isomorphism. Moreover, we have*

$$\mathcal{H} \cong \text{End}_{\mathcal{S}(\infty, r)}(\mathbf{\Omega}_\infty^{\otimes r}) \cong \text{End}_{\mathbf{U}(\infty)}(\mathbf{\Omega}_\infty^{\otimes r}).$$

*Proof.* Since the set  $\{\phi_{\mu, \varpi}^d \mid \mu \in \Lambda(\infty, r), d \in \mathfrak{D}_\mu\}$  forms a basis for  $\mathbf{\Omega}(\infty, r)$ , and the set  $\{\phi_{\mu, \varpi}^d(T_1) \mid \mu \in \Lambda(\infty, r), d \in \mathfrak{D}_\mu\}$  forms a basis for  $\mathbf{\Omega}_\infty^{\otimes r}$ . So the assertion follows easily.  $\square$

For  $\lambda \in \Lambda(\infty, r)$ , let  $\lambda^t = (\lambda_i^t)_{i \in \mathbb{Z}} \in \Lambda(\infty, r)$ , where  $\lambda_i^t = \#\{j \in \mathbb{Z} \mid \lambda_j \geq i\}$  for  $i \geq 1$  and  $\lambda_i^t = 0$  for  $i \leq 0$ . Let  $w_\lambda$  be the unique element in  $\mathfrak{D}_{\lambda, \lambda^t}$  such that  $w_\lambda^{-1} \mathfrak{S}_{w_\lambda} \cap \mathfrak{S}_{\lambda^t} = \{1\}$ , and let  $z_\lambda = \phi_{\lambda, \varpi}^1 T_{w_\lambda} y_{\lambda^t}$ , where  $y_{\lambda^t} = \sum_{w \in \mathfrak{S}_{\lambda^t}} (-v^2)^{-l(w)} T_w$ .

For any  $\mu \in \Lambda([-n, n], r)$ , let

$$(8.3.1) \quad \mathcal{S}^\mu = z_\mu \mathcal{H}, \quad \text{and} \quad W(\infty, \mu) = \mathcal{S}(\infty, r) z_\mu$$

be the *Specht module* of  $\mathcal{H}$  and the *Weyl module* of  $\mathcal{S}(\infty, r)$ , respectively. Note that, if  $\mu \in \Lambda([-n, n], r)$  and  $n \geq r$ , then  $\varpi \in \Lambda([-n, n], r)$ . So

$$W([-n, n], \mu) := \mathcal{S}([-n, n], r) z_\mu$$

is well-defined. This is a Weyl module of  $\mathcal{S}([-n, n], r)$ . Since  $\mathcal{S}([-n, n], r)$  is a finite dimensional semisimple algebra, by [6, 4.6],  $W([-n, n], \mu)$  is an irreducible  $\mathcal{S}([-n, n], r)$ -module. Note that



$z_\mu = \mathbf{k}_\mu z_\mu$ . So 8.1 implies  $W(\infty, \mu) = \mathcal{S}(\infty, r) \mathbf{k}_\mu z_\mu = \mathcal{K}(\infty, r) \mathbf{k}_\mu z_\mu = \mathcal{K}(\infty, r) z_\mu$ . Hence, for any  $\mu \in \Lambda(\infty, r)$ , we obtain

$$(8.3.2) \quad W(\infty, \mu) = \bigcup_{n \geq r} W([-n, n], \mu) \cong \varinjlim_n W([-n, n], \mu).$$

Let  $\Lambda^+(r)$  be the set of all partitions of  $r$ . We will regard  $\Lambda^+(r)$  as the subset

$$\{\lambda \in \Lambda(\infty, r) \mid \lambda_i \geq \lambda_{i+1} \forall i \geq 1, \lambda_i = 0 \forall i \leq 0\}$$

of  $\Lambda(\infty, r)$ . Define a map from  $\Lambda(\infty, r)$  to  $\Lambda^+(r)$  by sending  $\lambda$  to  $\lambda^+$  where  $\lambda^+$  is the unique element in  $\Lambda^+(r)$  obtained by reordering the parts of  $\lambda$ .

**Proposition 8.4.** (1) *For  $\mu \in \Lambda(\infty, r)$  we have*

$$W(\infty, \mu) \cong \text{Hom}_{\mathcal{H}}(S^\mu, \Omega_\infty^{\otimes r}) \cong \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}(S^\mu, x_\lambda \mathcal{H}).$$

(2) *Let  $\mu \in \Lambda(\infty, r)$ . Then the module  $W(\infty, \mu)$  is an irreducible  $\mathcal{S}(\infty, r)$ -module. Moreover, we have  $W(\infty, \mu) \cong W(\infty, \mu^+)$ .*

*Proof.* The bimodule isomorphism  $\text{Ev}$  given in 8.3 induces  $\mathcal{S}(\infty, r)$ -module isomorphism

$$\text{Hom}_{\mathcal{H}}(S^\mu, \Omega_\infty^{\otimes r}) \cong \text{Hom}_{\mathcal{H}}(S^\mu, \Omega(\infty, r)) \cong \bigoplus_{\lambda \in \Lambda(\infty, r)} \text{Hom}(S^\mu, \phi_{\lambda\varpi}^1 \mathcal{H}).$$

Now 5.1 implies

$$\text{Hom}_{\mathcal{H}}(S^\mu, \Omega_\infty^{\otimes r}) \cong \varinjlim_n \text{Hom}_{\mathcal{H}}(S^\mu, \Omega_{[-n, n]}^{\otimes r}).$$

Choose  $m_0$  such that  $\mu \in \Lambda([-m_0, m_0], r)$  and let  $r_0 = \max\{r, m_0\}$ . Then, for any  $n \geq r_0$ , we have, by [6, 8.1, 8.7], isomorphisms  $\text{Hom}_{\mathcal{H}}(S^\mu, \Omega_{[-n, n]}^{\otimes r}) \cong W([-n, n], \mu)$  which are compatible with the direct systems. Now, the first isomorphism in (1) is obtained by taking direct limits.

The assertions in (2) follow easily from (8.3.2) (cf. [6, 3.9]).  $\square$

For  $\lambda, \mu \in \Lambda(\infty, r)$  we write  $\mu \leq_{\text{wt}}^+ \lambda$  if  $\mu^+ \leq_{\text{wt}} \lambda^+$ . Note that  $\mu^+ \leq_{\text{wt}} \lambda^+$  is equivalent to the dominance order  $\mu^+ \trianglelefteq \lambda^+$  on partitions.

We also recall the notation of Young tableaux. Suppose  $\lambda \in \Lambda(\infty, r)$  and  $\mu \in \Lambda^+(r)$ . A  $\mu$ -tableau of type  $\lambda$  is a  $\mu$ -tableau with (possibly) repeated entries, where for each  $i$ , the number of entries  $i$  is equal to  $\lambda_i$ . We denote the set of  $\mu$ -tableaux of type  $\lambda$  by  $\mathcal{T}(\mu, \lambda)$ . For  $T \in \mathcal{T}(\mu, \lambda)$ , we say that  $T$  is row-standard (resp., strictly row-standard) if the numbers are weakly increasing (resp., increasing) along each row of  $T$ . The column-standard and strictly column-standard tableaux can be defined similarly. A tableau  $T$  is semistandard if it is row-standard and strictly column-standard. Let  $\mathcal{T}_0(\mu, \lambda)$  denote the set of semistandard  $\mu$ -tableaux of type  $\lambda$ .

**Proposition 8.5.** *Let  $\lambda \in \Lambda(\infty, r)$  and  $\mu \in \Lambda^+(r)$ . We have  $W(\infty, \mu)_\lambda \cong \text{Hom}_{\mathcal{H}}(S^\mu, x_\lambda \mathcal{H})$ . Hence,  $\dim W(\infty, \mu)_\lambda = \#\mathcal{T}_0(\mu, \lambda)$  ([6, 8.7]). In particular,  $\dim W(\infty, \mu)_\mu = 1$ , and  $W(\infty, \mu)_\lambda \neq 0$  implies  $\lambda \leq_{\text{wt}}^+ \mu$ .*

*Proof.* Let  $f \in \text{Hom}_{\mathcal{H}}(S^\mu, x_\lambda \mathcal{H})$ . Assume  $f(z_\mu) = x_\lambda h_\lambda$  where  $h_\lambda \in \mathcal{H}$ . Then we have

$$\begin{aligned} (\mathbf{k}_i f)(z_\lambda h) &= \mathbf{k}_i f(z_\lambda h) = \mathbf{k}_i(x_\lambda h_\lambda h) = \left( \sum_{\nu \in \Lambda(\infty, r)} v^{\nu_i} \phi_{\nu\nu}^1 \right) (x_\lambda h_\lambda h) \\ &= \sum_{\nu \in \Lambda(\infty, r)} v^{\nu_i} \phi_{\nu\nu}^1(x_\lambda h_\lambda h) = v^{\lambda_i} x_\lambda h_\lambda h = v^{\lambda_i} f(z_\lambda h), \end{aligned}$$

where  $i \in \mathbb{Z}$  and  $h \in \mathcal{H}$ . Hence,  $\mathbf{k}_i f = v^{\lambda_i} f$  for  $i \in \mathbb{Z}$ . This means  $f \in W(\infty, \mu)_\lambda$ . By 8.4(1), we obtain  $(W(\infty, \mu))_\lambda = \text{Hom}_{\mathcal{H}}(S^\mu, x_\lambda \mathcal{H})$ .  $\square$

Though the double centralizer property in the Schur-Weyl duality is no longer true in the infinite case, the following decomposition for the tensor space  $\Omega_\infty^{\otimes r}$  continue to hold.

**Theorem 8.6.** *We have the following  $(\mathcal{S}(\infty, r), \mathcal{H})$ -bimodule isomorphism*

$$\Omega_\infty^{\otimes r} \cong \bigoplus_{\mu \in \Lambda^+(r)} W(\infty, \mu) \otimes S^\mu.$$

Hence, as a (left)  $\mathcal{S}(\infty, r)$ -module,  $\Omega_\infty^{\otimes r}$  is completely reducible.

*Proof.* It is well-known that, for all  $n \geq r$ , there are  $(\mathcal{S}([-n, n], r), \mathcal{H})$ -bimodule isomorphisms

$$\Omega_{[-n, n]}^{\otimes r} \cong \bigoplus_{\mu \in \Lambda^+(r)} W([-n, n], \mu) \otimes S^\mu.$$

The require isomorphism is obtained by taking direct limits; cf. 5.1 and (8.3.2).  $\square$

**Corollary 8.7.** *For each  $\lambda \in \Lambda(\infty, r)$ , the  $\mathcal{S}(\infty, r)$ -module  $\mathcal{S}(\infty, r)\mathbf{k}_\lambda$  is completely reducible.*

*Proof.* Consider the  $\mathcal{S}(\infty, r)$ -module homomorphism

$$f : \mathcal{S}(\infty, r)\mathbf{k}_\lambda \longrightarrow \mathcal{S}(\infty, r)\mathbf{k}_\varpi \cong \Omega_\infty^{\otimes r}, \quad uk_\lambda \mapsto uk_\lambda \phi_{\lambda\varpi}^1 \quad \forall u \in \mathcal{S}(\infty, r).$$

Since  $f$  is induced from the surjective map  $\phi_{\lambda\varpi}^1 : \mathcal{H} \rightarrow x_\lambda \mathcal{H}$ , it follows that  $f$  is injective. Hence,  $\mathcal{S}(\infty, r)\mathbf{k}_\lambda$  is isomorphic to a submodule of  $\Omega_\infty^{\otimes r}$ . Hence, the assertion follows from 8.6.  $\square$

We are now ready to classify irreducible objects in the category  $\mathcal{S}(\infty, r)\text{-mod}$ . We first observe the following.

**Theorem 8.8.** (1) *The map  $\mu \rightarrow W(\infty, \mu)$  defines a bijection between  $\Lambda^+(r)$  and the set of isomorphism classes of irreducible  $\mathcal{S}(\infty, r)$ -modules in the category  $\mathcal{S}(\infty, r)\text{-mod}$ .*

(2) *Any module in  $\mathcal{S}(\infty, r)\text{-mod}$  is completely reducible.*

*Proof.* By 8.4 and 8.5, the modules  $W(\infty, \mu)$  ( $\mu \in \Lambda^+(r)$ ) are irreducible objects in  $\mathcal{S}(\infty, r)\text{-mod}$ . Let  $M$  be an irreducible  $\mathcal{S}(\infty, r)$ -module in  $\mathcal{S}(\infty, r)\text{-mod}$ . Let  $\mu \in \text{wt}(M)$  and  $x_0$  is a nonzero vector in  $M_\mu$ . Then we have a surjective homomorphism  $f$  from  $\mathcal{S}(\infty, r)$  to  $M$  by sending  $u$  to  $ux_0$  for  $u \in \mathcal{S}(\infty, r)$ . By the proof of 8.2(2), we have  $x_0 = \mathbf{k}_\mu x_0$ , and so  $f(\mathcal{S}(\infty, r)\mathbf{k}_\mu) = \mathcal{S}(\infty, r)f(\mathbf{k}_\mu) = \mathcal{S}(\infty, r)\mathbf{k}_\mu x_0 = \mathcal{S}(\infty, r)x_0 = M$ . Hence,  $M$  is a quotient module of  $\mathcal{S}(\infty, r)\mathbf{k}_\mu$ . By 8.7,  $M$  is isomorphic to an irreducible component of  $\mathcal{S}(\infty, r)\mathbf{k}_\mu$ , and hence, to an irreducible component of  $\Omega_\infty^{\otimes r}$ . Now, 8.6 implies that  $M$  is isomorphic to  $W(\infty, \mu)$  for some  $\mu \in \Lambda^+(r)$ , proving (1).

Let  $N$  be an arbitrary module in  $\mathcal{S}(\infty, r)\text{-mod}$ . Let  $0 \neq x \in N$ . Since  $N \in \mathcal{S}(\infty, r)\text{-mod}$ , we have  $\mathcal{S}(\infty, r)x = \mathcal{K}(\infty, r)x$  is a quotient module of  $\mathcal{K}(\infty, r)$ . But, by 8.1,

$$\mathcal{K}(\infty, r) = \bigoplus_{\lambda \in \Lambda(\infty, r)} \mathcal{K}(\infty, r)\mathbf{k}_\lambda = \bigoplus_{\lambda \in \Lambda(\infty, r)} \mathcal{S}(\infty, r)\mathbf{k}_\lambda.$$

Hence, 8.7 implies that the  $\mathcal{S}(\infty, r)$ -module  $\mathcal{K}(\infty, r)$  is completely reducible, and so  $\mathcal{S}(\infty, r)x$  is completely reducible. Consequently,  $N$  is completely reducible.  $\square$

**Remark 8.9.** Following the construction in [10], we can use the PBW type basis

$$\{[A] = v^{-d_A} \phi_{\lambda\mu}^d \mid A = j(\lambda, d, \mu), (\lambda, d, \mu) \in \mathfrak{D}(\infty, r)\}$$

for  $\mathcal{K}(\infty, r)$  to define the canonical basis  $\{\theta_{\lambda\nu}^d \mid (\lambda, d, \mu) \in \mathfrak{D}(\infty, r)\}$  of  $\mathcal{K}(\infty, r)$ . By [9, 5.3] one can easily show that, for each  $\mu \in \Lambda^+(r)$ , the set  $\{\theta_{\lambda\nu}^d z_\mu \mid (\lambda, d, \nu) \in \mathfrak{D}(\infty, r)\} \setminus \{0\}$  forms a  $\mathbb{Q}(v)$ -basis for  $W(\infty, \mu)$ . This basis is called the *canonical basis* of  $W(\infty, \mu)$ .

## 9. POLYNOMIAL REPRESENTATIONS OF $\mathbf{U}(\infty)$

We are now ready to classify all irreducible polynomial representations of  $\mathbf{U}(\infty)$ . Let  $\mathcal{C}^{pol}$  be the full subcategory of  $\mathcal{C}$  consisting of weight  $\mathbf{U}(\infty)$ -modules  $M$  such that  $\text{wt}(M) \subseteq \mathbb{N}^\infty$ . We call the objects in  $\mathcal{C}^{pol}$  *polynomial representations*. Let  $r$  be a positive integer and let  $\mathcal{C}_r$  be the category of  $\mathbf{U}(\infty, r)$ -modules which are weight  $\mathbf{U}(\infty)$ -modules via the homomorphism  $\zeta_r : \mathbf{U}(\infty) \rightarrow \mathbf{U}(\infty, r)$ .

**Proposition 9.1.** (1) *The category  $\mathcal{C}_r$  is a full subcategory of  $\mathcal{C}^{pol} \cap \mathcal{C}^{int}$ .*

(2) *For each  $r \in \mathbb{N}$ , there is a surjective homomorphism from  $\mathbf{U}(\infty, r+1)$  to  $\mathbf{U}(\infty, r)$  by sending the generators  $\mathbf{e}_i, \mathbf{f}_i, \mathbf{k}_i$  for  $\mathbf{U}(\infty, r+1)$  to the generators  $\mathbf{e}_i, \mathbf{f}_i, \mathbf{k}_i$  for  $\mathbf{U}(\infty, r)$ , respectively. Hence,  $\mathcal{C}_r$  is a full subcategory of  $\mathcal{C}_{r+1}$ .*

*Proof.* Let  $M$  be a  $\mathbf{U}(\infty, r)$ -module in the category  $\mathcal{C}_r$ . As a  $\mathbf{U}(\infty)$ -module,  $E_i, K_i, F_i$  act on  $M$  as the action of  $\mathbf{e}_i, \mathbf{k}_i, \mathbf{f}_i$ , respectively. But, by 3.5, we have  $\mathbf{e}_i^n = \mathbf{f}_i^n = 0$  for all  $n \geq r+1$ . Hence,  $M$  is an integrable  $\mathbf{U}(\infty)$ -module.

We now prove  $\text{wt}(M) \subseteq \mathbb{Z}^\infty$ . Suppose this is not the case. Then there exists  $\lambda \in \text{wt}(M)$  and  $\lambda \notin \mathbb{Z}^\infty$ . Thus,  $\lambda$  has an infinite support. Choose  $r+1$  integers  $i_1 < i_2 < \dots < i_{r+1}$  such that  $\lambda_{i_1} \neq 0, \dots, \lambda_{i_{r+1}} \neq 0$  and  $\mathbf{t} \in \Lambda(\infty, r)$  such that  $t_i = 1$  for  $i \in \{i_1, \dots, i_{r+1}\}$  and  $t_i = 0$  for all other  $i$ . Let  $u_\lambda$  be a nonzero vector in  $M_\lambda$ . Then

$$0 \neq (v^{\lambda_{i_1}} - 1) \cdots (v^{\lambda_{i_{r+1}}} - 1) u_\lambda = \prod_{i \in \mathbb{Z}} [v^{\lambda_i}; t_i]^! u_\lambda = \prod_{i \in \mathbb{Z}} [K_i; t_i]^! u_\lambda = \prod_{i \in \mathbb{Z}} [\mathbf{k}_i; t_i]^! u_\lambda \stackrel{4.2}{=} 0 u_\lambda = 0,$$

a contradiction. Hence,  $\text{wt}(M) \subseteq \mathbb{Z}^\infty$ . On the other hand, by 4.2 again, we have  $[\mathbf{k}_i; r+1]^! u_\lambda = (v^{\lambda_i} - 1) \cdots (v^{\lambda_i} - v^r) u_\lambda = 0$  for  $i \in \mathbb{Z}$ , forcing  $0 \leq \lambda_i \leq r$  for any  $i \in \mathbb{Z}$ . Hence,  $\lambda \in \mathbb{N}^\infty$ , proving (1).

Let  $J_r$  be the ideal of  $\mathbf{U}(\infty)$  generated by  $\prod_{i \in \mathbb{Z}} [K_i; t_i]^!$ , where  $\mathbf{t} = (t_i)_{i \in \mathbb{Z}} \in \Lambda(\infty, r+1)$ . By 4.7 we have  $\mathbf{U}(\infty, r)$  is isomorphic to  $\mathbf{U}(\infty)/J_r$ . Let  $\mathbf{t} \in \Lambda(\infty, r+2)$ . Choose  $i_0 \in \mathbb{Z}$  such that  $t_{i_0} > 0$ . Then  $\prod_{i \in \mathbb{Z}} [K_i; t_i]^! = (K_{i_0} - v^{t_{i_0}-1}) (\prod_{i \neq i_0} [K_i; t_i]^!) [K_{i_0}; t_{i_0} - 1]^! \in J_r$ . Hence,  $J_{r+1} \subseteq J_r$ , and (2) follows.  $\square$

**Lemma 9.2.** *Let  $M$  be a irreducible  $\mathbf{U}(\infty)$ -module in the category  $\mathcal{C}^{pol}$ . Then  $M \in \mathbf{U}(\infty, r)\text{-mod}$  for some  $r \geq 0$ .*

*Proof.* Let  $\lambda \in \text{wt}(M)$  and  $u_\lambda$  be a nonzero vector in  $M_\lambda$ . Then  $M = \mathbf{U}(\infty)u_\lambda$  since  $M$  is irreducible. Hence, by 7.1, we know  $\sigma(\mu) = \sigma(\lambda)$  for any  $\mu \in \text{wt}(M)$ . Let  $r = \sigma(\lambda)$ . Since  $\text{wt}(M) \subseteq \mathbb{N}^\infty$  we have  $\text{wt}(M) \subseteq \Lambda(\infty, r)$ . Thus, for any  $\mathbf{t} \in \Lambda(\infty, r+1)$ , there exist  $i_0 \in \mathbb{Z}$  such that  $t_{i_0} > \lambda_{i_0} \geq 0$ . Hence,  $[v^{\lambda_{i_0}}; t_{i_0}]^! = 0$ , and  $\prod_{i \in \mathbb{Z}} [\mathbf{k}_i; t_i]^! u_\lambda = \prod_{i \in \mathbb{Z}} [v^{\lambda_i}; t_i]^! u_\lambda = 0$ . This shows that  $M$  is a  $\mathbf{U}(\infty, r)$ -module (by 4.7). Since  $M \in \mathcal{C}$ , it follows that  $M \in \mathbf{U}(\infty, r)\text{-mod}$ .  $\square$

**Theorem 9.3.** (1) *For  $r \in \mathbb{N}$  the modules  $W(\infty, \mu)$  ( $\mu \in \Lambda^+(r'), r' \leq r$ ) are all non-isomorphic irreducible  $\mathbf{U}(\infty, r)$ -modules in the category  $\mathcal{C}_r$ .*

(2) *The modules  $W(\infty, \mu)$  ( $\mu \in \Lambda^+(r), r \in \mathbb{N}$ ) are all non-isomorphic irreducible  $\mathbf{U}(\infty)$ -modules in the category  $\mathcal{C}^{pol}$ . Moreover, there is only one irreducible  $\mathbf{U}(\infty)$ -module, the trivial module  $L(0)$ , in the category  $\mathcal{C}^{pol} \cap \mathcal{C}^{hi}$ .*

*Proof.* By 8.4(2) and 9.1(2) we know  $W(\infty, \mu)$ , where  $\mu \in \Lambda^+(r')$  with  $r' \leq r$ , are irreducible  $\mathbf{U}(\infty, r)$ -modules in  $\mathcal{C}_r$ . On the other hand, let  $M$  be a irreducible  $\mathbf{U}(\infty, r)$ -module in the category  $\mathcal{C}_r$ . By 9.1(1) and 9.2 we have  $M \in \mathbf{U}(\infty, r')\text{-mod}$  for some  $r' \geq 0$ . We claim that  $r' \leq r$ . Indeed,

suppose the contrary  $r' > r$ . Then, for any  $\lambda \in \text{wt}(M) \subseteq \Lambda(\infty, r')$ , there exists  $\mu \in \Lambda(\infty, r+1)$  such that  $\mu_i \leq \lambda_i$  for all  $i \in \mathbb{Z}$ . Thus, for any nonzero vector  $x_0$  in  $M_\lambda$ ,

$$\prod_{i \in \mathbb{Z}} [K_i; \mu_i]^! x_0 = \prod_{i \in \mathbb{Z}} [k_i; \mu_i]^! x_0 = \prod_{i \in \mathbb{Z}} [v^{\lambda_i}; \mu_i]^! x_0 \neq 0$$

for  $i \in \mathbb{Z}$ . However, since  $M$  is a  $\mathbf{U}(\infty, r)$ -module, the presentation of  $\mathbf{U}(\infty, r)$  implies  $\prod_{i \in \mathbb{Z}} [K_i; \mu_i]^! x_0 = 0$ , a contradiction. Hence,  $r' \leq r$ . Now 8.8 implies that there exists  $\mu \in \Lambda^+(r')$  such that  $M \cong W(\infty, \mu)$ , proving (1).

The first assertion in (2) follows from 8.8 and 9.2. By 9.1,  $W(\infty, \mu)$  is integrable. Since  $\text{wt}(W(\infty, \mu)) \subseteq \Lambda(\infty, r)$ , by the classification of irreducible integral modules in  $\mathcal{C}^{hi}$  given in 7.6, we conclude  $W(\infty, \mu) \notin \mathcal{C}^{hi}$  for any  $r \geq 1$  and  $\mu \in \Lambda(\infty, r)$ , proving the last assertion in (2).  $\square$

**Remark 9.4.** (1) Classification of irreducible integrable modules over a Kac-Moody algebra is an open problem (see [23, Ex. 10.23]). However, irreducible integrable  $\mathbf{U}(\infty, r)$ -modules can be classified for all  $r \in \mathbb{N}$ . In fact, by the above result, the modules  $W(\infty, \mu)$  ( $\mu \in \Lambda^+(r')$ ,  $r' \leq r$ ) are all irreducible integrable  $\mathbf{U}(\infty, r)$ -modules for any  $r \in \mathbb{N}$ .

It is clear that  $\mathcal{S}(\infty, r) \cong \prod_{\mu \in \Lambda(\infty, r)} \mathcal{S}(\infty, r)_{\mathbf{k}_\mu}$  as a  $\mathcal{S}(\infty, r)$ -module. Hence,  $\mathcal{S}(\infty, r) \notin \mathcal{C}$  as a  $\mathbf{U}(\infty)$ -module. Since  $\mathcal{S}(\infty, r)_{\mathbf{k}_\mu}$  is a direct sum of finitely many irreducible  $\mathcal{S}(\infty, r)$ -modules in  $\mathcal{S}(\infty, r)$ -**mod**,  $\mathcal{S}(\infty, r)$  is the direct product of irreducible  $\mathcal{S}(\infty, r)$ -modules in  $\mathcal{S}(\infty, r)$ -**mod**.

(2) It is interesting to make a comparison between the finite and infinite cases for quantum  $\mathfrak{gl}_\eta$ . In the finite case, if we only consider finite dimensional representations, then the categories  $\mathcal{C}^{hi}$  and  $\mathcal{C}^{int}$  are the same as the category  $\mathcal{C}^{fd}$ , and contain  $\mathcal{S}(n, r)$ -**mod** as a full subcategory of polynomial representations of degree  $r$  which more or less determine  $\mathcal{C}^{hi}$  and  $\mathcal{C}^{int}$ . They are all completely reducible and the irreducible modules in these categories are all indexed by dominant weights. In the infinite case, the situation is completely different. For example, the categories  $\mathcal{C}^{hi} \cap \mathcal{C}^{int}$ ,  $\mathcal{S}(\infty, r)$ -**mod** and  $\mathcal{C}^{fd}$  become three different categories. The complete reducibility continues to hold in the last two categories, but seems not true in the first category. The irreducible modules in  $\mathcal{C}^{hi} \cap \mathcal{C}^{int}$  are also indexed by dominant weights. Moreover, there are only a few irreducible objects in  $\mathcal{C}^{fd}$ , and there is only one irreducible object in the category  $\mathcal{C}^{pol} \cap \mathcal{C}^{hi}$ .

## REFERENCES

- [1] A.A. Beilinson, G. Lusztig and R. MacPherson, *A geometric setting for the quantum deformation of  $GL_n$* , Duke Math.J. **61** (1990), 655-677.
- [2] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra  $\mathfrak{gl}(m|n)$* , J. Amer. Math. Soc. **16** (2003), 185-231.
- [3] M. Date, M. Jimbo, M. Kashiwara and T. Miwa, *Transformation groups for soliton equations*, Publ. RIMS, Kyoto Univ. **18** (1982), 1077-1110.
- [4] B. Deng and J. Du, *Bases of quantized enveloping algebras*, Pacific J. Math. **220** (2005), 33-48.
- [5] B. Deng, J. Du, B. Parshall and J. Wang, *Finite Dimensional Algebras and Quantum Groups*, Mathematical Surveys and Monographs Volume 150, Amer. Math. Soc., Providence 2008.
- [6] R. Dipper and G. James,  *$q$ -Tensor spaces and  $q$ -Weyl modules*, Trans. Amer. Math. Soc. **327** (1991), 251-282.
- [7] S. Doty and A. Giaquinto, *Presenting Schur algebras*, International Mathematics Research Notices IMRN, **36** (2002), 1907-1944.
- [8] S. R. Doty and R. M. Green, *Presenting affine  $q$ -Schur algebras*, Math. Z. **256** (2007), 311-345.
- [9] J. Du, *Canonical bases for irreducible representations of quantum  $GL_n$* , Bull. London Math. Soc. **24** (1992), 325-334.
- [10] J. Du, *Kazhdan-Lusztig bases and isomorphism theorems for  $q$ -Schur algebras*, Contemp. Math. **139** (1992), 121-140.
- [11] J. Du, *A note on the quantized Weyl reciprocity at roots of unity*, Alg. Colloq. **2**(1995), 363-372.
- [12] J. Du, Q. Fu and J.-P. Wang, *Infinitesimal quantum  $\mathfrak{gl}_n$  and little  $q$ -Schur algebras*, J. Algebra **287** (2005), 199-233.

- [13] J. Du and B. Parshall, *Linear quivers and the geometric setting of quantum  $GL_n$* , Indag. Math. **13** (2002), 459-481.
- [14] J. Du and B. Parshall, *Monomial bases for  $q$ -Schur algebras*, Trans. Amer. Math. Soc. **355** (2003), 1593-1620.
- [15] J. Du, B. Parshall and L. Scott, *Quantum Weyl reciprocity and tilting modules*, Comm. Math. Phys. **195** (1998), 321-352.
- [16] J. Du, B. Parshall, and Jian-pan Wang, *Two-parameter quantum linear groups and the hyperbolic invariance of  $q$ -Schur algebras*, J. London Math. Soc. **44** (1991), 420-436.
- [17] T. Hayashi,  *$Q$ -analogues of Clifford and Weyl algebras—spinor and oscillator representations of quantum enveloping algebras*, Comm. Math. Phys. **127** (1990), 129-144.
- [18] G. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Encyclopedia of Math. and its Appl. no. 6, Addison-Wesley, 1981 London.
- [19] J. C. Jantzen, *Lectures on quantum groups*, Graduate Studies in Mathematics, **6**. American Mathematical Society, Providence, RI, 1996.
- [20] M. Jimbo, *A  $q$ -analogue of  $U(\mathfrak{gl}(N+1))$ , Hecke algebra, and the Yang-Baxter equation*, Lett. Math. Phys. **11** (1986), no. 3, 247-252.
- [21] V. G. Kac, and D. H. Peterson, *Spin and wedge representations of infinite-dimensional Lie algebras and groups*, Proc. Nat. Acad. Sci. U.S.A. **78** (1981), 3308-3312.
- [22] V. G. Kac and A. K. Raina, *Bombay lectures on highest weight representations of infinite-dimensional Lie algebras* Advanced Series in Mathematical Physics, 2. World Scientific Publishing Co., Inc., Teaneck, NJ, 1987.
- [23] V. G. Kac, *Infinite-dimensional Lie algebras*, Third edition. Cambridge University Press, Cambridge, 1990.
- [24] S. Levendorskii and Y. Soibelman, *Quantum group  $A_\infty$* , Comm. Math. Phys. **140**(1991), 399-414.
- [25] G. Lusztig, *Modular representations and quantum groups*, Comtemp. Math. **82** (1989), 59-77.
- [26] G. Lusztig, *Finite-dimensional Hopf algebras arising from quantized universal enveloping algebra* J. Amer. Math. Soc. **3** (1990), 257-296.
- [27] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447-498.
- [28] G. Lusztig, *Introduction to quantum groups*, Progress in Math. **110**, Birkhäuser, 1993.
- [29] G. Lusztig, *Transfer maps for quantum affine  $\mathfrak{sl}_n$* , In Representations and quantizations (Shanghai, 1998), China High. Educ. Press, Beijing (2000), 341-356.
- [30] K. McGerty, *Generalized  $q$ -Schur algebras and quantum Frobenius*, Adv. Math. **214** (2007), 116-131.
- [31] K. Misra and T. Miwa, *Crystal base for the basic representation of  $U_q(\widehat{\mathfrak{sl}}(n))$* , Comm. Math. Phys. **134** (1990), 79-88.
- [32] T. D. Palev, *Highest weight irreducible unitary representations of Lie algebras of infinite matrices. I. The algebra  $\mathfrak{gl}(\infty)$* , J. Math. Phys. **31** (1990), 579-586.
- [33] T. D. Palev, *Highest weight irreducible unitarizable representations of Lie algebras of infinite matrices The algebra  $A_\infty$* , J. Math. Phys. **31** (1990), 1078-1084.
- [34] T. D. Palev and N. I. Stoilova, *Highest weight representations of the quantum algebra  $U_h(\mathfrak{gl}_\infty)$* , J. Phys. A **30** (1997), L699-L705.
- [35] T. D. Palev and N. I. Stoilova, *Highest weight irreducible representations of the quantum algebra  $U_h(A_\infty)$* , J. Math. Phys. **39** (1998), 5832-5849.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY 2052, AUSTRALIA.

Home page: <http://web.maths.unsw.edu.au/~jied>

E-mail address: [j.du@unsw.edu.au](mailto:j.du@unsw.edu.au)

DEPARTMENT OF MATHEMATICS, TONGJI UNIVERSITY, SHANGHAI, 200092, CHINA.

E-mail address: [q.fu@hotmail.com](mailto:q.fu@hotmail.com)