

Modules over Fedosov deformation quantization

S. A. Pol'shin*

Institute for Theoretical Physics

NSC Kharkov Institute of Physics and Technology

Akademicheskaya St. 1, 61108 Kharkov, Ukraine

Abstract

For a given polarization L of an arbitrary symplectic manifold M we construct a Fedosov-type star-product $*_L$ on M such that for an arbitrary leaf φ of L a set $C^\infty(\varphi)[[h]]$ has a natural structure of left module over the star-product algebra $(C^\infty(M)[[h]], *_L)$. With certain additional assumptions on M , $*_L$ becomes a so-called star-product with separation of variables.

MSC 2000: 53D55 53D50 17B55

Key words: Fedosov quantization, polarization, contracting homotopy, adapted star-product, bi-Lagrangian manifold.

1 Introduction

In [9] B.V.Fedosov gave a simple construction of deformation quantization of an arbitrary symplectic manifold (see also [10]). Later J.Donin [6] and D.Farkas [8] show the algebraic nature of Fedosov construction. The problem of constructing modules over Fedosov deformation quantization which generalize the states of usual quantum mechanics is of great interest. To this end, a notion of adapted star-products was recently introduced [2].

Definition 1. A star-product $*$ on M is called adapted to a lagrangian (or, more generally, coisotropic) submanifold $\varphi \subset M$ if the vanishing ideal of φ in the commutative algebra $C^\infty(M)$ is a left ideal in the deformed algebra $\mathcal{A} = (C^\infty(M)[[h]], *)$. Then obviously $C^\infty(\varphi)[[h]]$ has a structure of left \mathcal{A} -module.

Using the algebraic framework of [6, 8], in the present letter we construct a Fedosov-type star-product $*_L$ adapted to all the leaves of given polarization L of a symplectic manifold M . If M is a so-called bi-Lagrangian manifold, $*_L$ becomes a star-product with separation of variables in the sense of Karabegov.

Plan of the present paper is the following. In Sec. 2 we consider the Koszul complex for Weyl algebra, in Sec. 3 we define various ideals associated with L , in Sec. 4 we define Fedosov complex and prove the main result.

The author is grateful to L.L.Vaksman for constant support during the work.

2 Koszul complex and Weyl algebra

Let M be a symplectic manifold, $\dim M = 2\nu$, $A = C^\infty(M)$ an algebra of smooth functions on M with pointwise multiplication, and $E = \Gamma TM$ a set of all smooth vector fields on M

*E-mail: itl593@online.kharkov.ua

with the natural structure of an unitary A -module. By $T(E)$ and $S(E)$ denote the tensor and symmetric algebra of E respectively, and let $\wedge E^*$ be an algebra of smooth differential forms on M . Let $\omega \in \wedge^2 E^*$ be a symplectic form on M and let $u : E \rightarrow \wedge^1 E^*$ be the mapping $u(x)y = \omega(x, y)$, $x, y \in E$. Let

$$a = x_1 \otimes \dots \otimes x_m \otimes y_1 \wedge \dots \wedge y_n \in T^m(E) \otimes \wedge^n E^*.$$

Define the Koszul differential of bidegree $(-1, 1)$ on $T(E) \otimes \wedge E^*$ as

$$\delta a = \sum_i x_1 \otimes \dots \otimes \hat{x}_i \otimes \dots \otimes x_m \otimes u(x_i) \wedge y_1 \dots \wedge y_n.$$

Let λ be an independent variable (physically $\lambda = -i\hbar$). In the sequel we will write A, E etc. instead of $A[\lambda], A[[\lambda]], E[\lambda], E[[\lambda]]$ etc. Let \mathcal{I}_W be a two-sided ideal in $T(E)$ generated by relations $x \otimes y - y \otimes x - \lambda\omega(x, y) = 0$. The factor-algebra $W(E) = T(E)/\mathcal{I}_W$ is called the Weyl algebra of E and let \circ be the multiplication in $W(E)$.

A ν -dimensional distribution $L \subset E$ is called a polarization if it is (a) lagrangian, i.e. $\omega(x, y) = 0$ for all $x, y \in L$ and (b) involutive, i.e. $[x, y] \in L$ for all $x, y \in L$, where $[\cdot, \cdot]$ is the commutator of vector fields on M . It is well known [7] that always we can choose a distribution L' transversal to L , i.e. $L \oplus L' = E$. Let $\alpha, \alpha_1, \dots = 1, \dots, \nu$ and $\beta, \beta_1, \dots = \nu + 1, \dots, 2\nu$. Suppose E has a 2ν -dimensional basis $\{e_i | i = 1, \dots, 2\nu\}$ as A -module. Choose it in such a way that $\{e_\alpha | \alpha = 1, \dots, \nu\}$ and $\{e_\beta | \beta = \nu + 1, \dots, 2\nu\}$ are the bases in L and L' respectively. Let $i_1, \dots, i_p = 1, \dots, 2\nu$ and let $I = (i_1, \dots, i_p)$ be an arbitrary sequence of indices. We write $e_I = e_{i_1} \otimes \dots \otimes e_{i_p}$ and we say that the sequence I is nonincreasing if $i_1 \geq i_2 \geq \dots \geq i_p$. We consider $\{\emptyset\}$ as a nonincreasing sequence and $e_{\{\emptyset\}} = 1$. We say that a sequence I is of α -length n if it contains n elements less or equal ν . Let Υ^n be a set of all nonincreasing sequences of α -length n and $\Upsilon_n = \bigcup_{p=n}^{\infty} \Upsilon^p$. Then a variant of Poincare-Birkhoff-Witt theorem holds [4, 15].

Theorem 1 (Poincare-Birkhoff-Witt). *Let $\tilde{S}(E)$ be an A -submodule of $T(E)$ generated by elements $\{e_I | I \in \Upsilon_0\}$. Then*

(a) *The restrictions $\mu_S|_{\tilde{S}(E)}$ and $\mu_W|_{\tilde{S}(E)}$ of the canonical homomorphisms $\mu_S : T(E) \rightarrow S(E)$ and $\mu_W : T(E) \rightarrow W(E)$ are A -module isomorphisms.*

(b) *$\{\mu_S(e_I) | I \in \Upsilon_0\}$ and $\{\mu_W(e_I) | I \in \Upsilon_0\}$ are A -bases of $S(E)$ and $W(E)$ respectively.*

(c) *$T(E) = \tilde{S}(E) \oplus \mathcal{I}_W$.*

Let ι_m ($m = 1, 2$) be a natural embedding of m th direct summand in the rhs of Theorem 1 (c) into $T(E)$, so $\mu_{S,W}|_{\tilde{S}(E)} = \mu_{S,W}\iota_1$. Then from Theorem 1 (c) it follows that a short exact sequence of A -modules

$$0 \longrightarrow \mathcal{I}_W \xrightarrow{\iota_2} T(E) \xrightarrow{\mu_W} W(E) \longrightarrow 0$$

splits, then we have another short exact sequence of A -modules

$$0 \longrightarrow \mathcal{I}_W \otimes \wedge E^* \xrightarrow{\iota_2 \otimes \text{id}} T(E) \otimes \wedge E^* \xrightarrow{\mu_W \otimes \text{id}} W(E) \otimes \wedge E^* \longrightarrow 0 \quad (1)$$

and $\iota_1 \otimes \text{id}$ is a natural embedding of $\tilde{S}(E) \otimes \wedge E^*$ into $T(E) \otimes \wedge E^*$.

It is easily seen that δ preserves $\mathcal{I}_W \otimes \wedge E^*$, so it induces a well-defined differential on $W(E) \otimes \wedge E^*$ due to (1). It is well known that u is an isomorphism due to nondegeneracy of ω . So we can define the so-called contracting homotopy of bidegree $(1, -1)$ on $S(E) \otimes \wedge E^*$ which to an element

$$a = x_1 \odot \dots \odot x_m \otimes y_1 \wedge \dots \wedge y_n \in S^m(E) \otimes \wedge^n E^*,$$

where \odot is the multiplication in $S(E)$, assigns the element

$$\delta^{-1}a = \frac{1}{m+n} \sum_i (-1)^{i-1} u^{-1}(y_i) \odot x_1 \odot \dots \odot x_m \otimes y_1 \wedge \dots \wedge \hat{y}_i \wedge \dots \wedge y_n$$

at $m + n > 0$ and $\delta^{-1}a = 0$ at $m = n = 0$.

Let $a = \sum_{m,n \geq 0} a_{mn}$, where $a_{mn} \in S^m(E) \otimes \wedge^n E^*$ and $\tau : a \mapsto a_{00}$ is the projection onto a component of bidegree $(0,0)$. Carry δ onto $S(E) \otimes \wedge E^*$ using the canonical homomorphism $\mu_S \otimes \text{id}$. Then it is well known that the following equality

$$\delta\delta^{-1} + \delta^{-1}\delta + \tau = \text{Id} \quad (2)$$

holds. Carry the grading of $S(E)$ onto $W(E)$ using the isomorphism $S(E) \cong W(E)$, then $W^1(E) \cong E$ and we will identify them. It is easily seen that δ preserves $\tilde{S}(E) \otimes \wedge E^*$, then it commutes with A -module isomorphism $\mu_W \iota_1 (\mu_S \iota_1)^{-1} \otimes \text{id}$ due to the remark after eq. (1). So each arrow of the following commutative diagram of A -modules commutes with δ .

$$\begin{array}{ccc}
& T(E) \otimes \wedge E^* & \\
\mu_S \otimes \text{id} \swarrow & \uparrow \iota_1 \otimes \text{id} & \searrow \mu_W \otimes \text{id} \\
& \tilde{S}(E) \otimes \wedge E^* & \\
\mu_S \iota_1 \otimes \text{id} \swarrow & & \searrow \mu_W \iota_1 \otimes \text{id} \\
S(E) \otimes \wedge E^* & \xrightarrow{\mu_W \iota_1 (\mu_S \iota_1)^{-1} \otimes \text{id}} & W(E) \otimes \wedge E^*.
\end{array}$$

Carry the contracting homotopy δ^{-1} and the projection τ from $S(E) \otimes \wedge E^*$ onto $W(E) \otimes \wedge E^*$ via the A -module isomorphism $\mu_W \iota_1 (\mu_S \iota_1)^{-1} \otimes \text{id}$, then the equality (2) remains true. Let $\delta W^\bullet = (W(E) \otimes \wedge^n E^*, \delta)$, then from (2) it follows that

$$H^0(\delta W^\bullet) = A, \quad H^n(\delta W^\bullet) = 0 \text{ at } n > 0. \quad (3)$$

3 The ideals

Let \mathcal{I}_\wedge be an ideal in $\wedge E^*$ those elements annihilate the polarization L , i.e. $\mathcal{I}_\wedge = \sum_{n=1}^{\infty} \mathcal{I}_\wedge^n$, where

$$\mathcal{I}_\wedge^n = \{\alpha \in \wedge^n E^* \mid \alpha(x_1, \dots, x_n) = 0 \ \forall x_1, \dots, x_n \in L\}.$$

It is well known that \mathcal{I}_\wedge is locally generated by ν independent 1-forms which are the basis of \mathcal{I}_\wedge^1 . From the other hand, L is lagrangian, so from the dimensional reasons we obtain $u(L) = \mathcal{I}_\wedge^1$, so

$$\mathcal{I}_\wedge = (u(L)). \quad (4)$$

Let \mathcal{I}_L be a left ideal in $W(E)$ generated by elements of L and $\mathcal{I} = \mathcal{I}_L \otimes \wedge E^* + W(E) \otimes \mathcal{I}_\wedge$ a left ideal in $W(E) \otimes \wedge E^*$. Then from (4) it follows that

$$\delta(\mathcal{I}) \subset \mathcal{I}. \quad (5)$$

Lemma 1. *An ideal $\mathcal{I}_L^{(S)}$ in $S(E)$ generated by elements of L is an image of \mathcal{I}_L under the isomorphism $\mu_S \iota_1 (\mu_W \iota_1)^{-1} : W(E) \xrightarrow{\cong} S(E)$.*

Proof. Since L is lagrangian, we have $e_{\alpha_1} \circ e_{\alpha_2} = e_{\alpha_2} \circ e_{\alpha_1}$. Then from Theorem 1 (b) it follows that \mathcal{I}_L is contained in $\text{span}\{\mu_W(e_I) \mid I \in \Upsilon_1\}$. From the other hand, \mathcal{I}_L contains all the $\mu_W(e_I)$, $I \in \Upsilon_1$, so it contains $\text{span}\{\mu_W(e_I) \mid I \in \Upsilon_1\}$ too and we have $\mathcal{I}_L = \mu_W \iota_1 (\tilde{S}_1(E))$, where $\tilde{S}_i(E) = \text{span}\{e_I \mid I \in \Upsilon_i\}$, $i = 0, 1, 2, \dots$ is a decreasing filtration on $\tilde{S}(E)$. Analogously $\mathcal{I}_L^{(S)} = \mu_S \iota_1 (\tilde{S}_1(E))$, which proves the lemma. \square

From (4) it is easily seen that δ^{-1} preserves the submodule $\mathcal{I}_L^{(S)} \otimes \wedge E^* + S(E) \otimes \mathcal{I}_\wedge$ of $S(E) \otimes \wedge E^*$, then using Lemma 1 we obtain

$$\delta^{-1}(\mathcal{I}) \subset \mathcal{I}. \quad (6)$$

Remark. The choice of $\tilde{S}(E)$ in Theorem 1 is crucial for our construction of contracting homotopy of δW^\bullet . The usual choice of submodule $S'(E)$ of symmetric tensors in $T(E)$ instead of $\tilde{S}(E)$ yields another contracting homotopy of δW^\bullet which does not preserve \mathcal{I} .

Let ∇ be an exterior derivative on $\wedge E^*$ which to an element $\alpha \in \wedge^{n-1} E^*$ assigns the element

$$\begin{aligned} (\nabla \alpha)(x_1, \dots, x_n) &= \sum_{1 \leq i < j \leq n} (-1)^{i+j} \alpha([x_i, x_j], x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_n) \\ &\quad + \sum_{1 \leq i \leq n} (-1)^{i-1} x_i \alpha(x_1, \dots, \hat{x}_i, \dots, x_n). \end{aligned} \quad (7)$$

Let $\nabla_x y \in E$, $x, y \in E$ be a connection on M , then we can extend ∇_x to $T(E)$ by the Leibniz rule. It is well known that we can always choose ∇ to be symplectic, i.e. $\nabla_z \omega(x, y) = \omega(\nabla_z x, y) + \omega(x, \nabla_z y)$ for all $x, y, z \in E$. It is well known that such ∇_x preserve \mathcal{I}_W for all $x \in E$, so it induces a well-defined derivation on $W(E)$. Now consider ∇ as a map $W(E) \rightarrow W(E) \otimes \wedge^1 E^*$ such that $(\nabla a)(x) = \nabla_x a$. Then it is well known that ∇ may be extended to a derivation of bidegree $(0, 1)$ of the whole algebra $W(E) \otimes \wedge E^*$ whose restriction to $\wedge E^*$ coincides with (7)

Lemma 2. *Let $a \in W(E) \otimes \wedge^n E^*$ and $a(x_1, \dots, x_n) \in V$ for all $x_1, \dots, x_n \in L$, where V is a submodule of $W(E)$. Then $a \in V \otimes \wedge E^* + W(E) \otimes \mathcal{I}_\wedge$.*

Proof. Let $\{\tilde{e}^i | i = 1, \dots, 2\nu\}$ be a local basis of E^* dual to $\{e_i\}$, i.e. $\tilde{e}^i(e_j) = \delta_j^i$. Then from the well-known theorem of basic algebra it follows that an arbitrary $a \in W(E) \otimes \wedge^n E^*$ may be represented in the form $a = \sum_{i_1 < \dots < i_n} a_{i_1 \dots i_n} \otimes \tilde{e}^{i_1} \wedge \dots \wedge \tilde{e}^{i_n}$, where $a_{i_1 \dots i_n} \in W(E)$ for all i_1, \dots, i_n .

It is easily seen that $\{\tilde{e}^\beta | \beta = \nu + 1, \dots, 2\nu\}$ locally generate \mathcal{I}_\wedge , so

$$a = \sum_{\alpha_1 < \dots < \alpha_n} a_{\alpha_1 \dots \alpha_n} \otimes \tilde{e}^{\alpha_1} \wedge \dots \wedge \tilde{e}^{\alpha_n} + W(E) \otimes \mathcal{I}_\wedge^n.$$

From the other hand, $a_{\alpha_1 \dots \alpha_n} = a(e_{\alpha_1}, \dots, e_{\alpha_n}) \in V$ due to the lemma's conditions. So $a \in V \otimes \wedge E^* + W(E) \otimes \mathcal{I}_\wedge$. \square

Suppose the integral submanifolds of L are totally geodesic, i.e.

$$\nabla_x y \in L, \quad x, y \in L. \quad (8)$$

For a given L , a symplectic connection which obeys (8) always exists ([14], Lemma 5.6). Then $\nabla_x \mathcal{I}_L \subset \mathcal{I}_L \forall x \in L$, so using Lemma 2 we obtain $\nabla \mathcal{I}_L \subset \mathcal{I}$. From the other hand, the involutivity of L together with (7) yield $\nabla \mathcal{I}_\wedge \subset \mathcal{I}_\wedge$ (Frobenius theorem), so we finally obtain

$$\nabla \mathcal{I} \subset \mathcal{I}. \quad (9)$$

Let $\tilde{R} \in E \otimes E^* \otimes \wedge^2 E^*$ be the curvature tensor of ∇ , i.e.

$$\tilde{R}(x, y)z = \nabla_x \nabla_y z - \nabla_y \nabla_x z - \nabla_{[x, y]} z, \quad x, y, z \in E.$$

Then from (8) using the involutivity of L it follows that $\tilde{R}(x, y)z \in L \forall x, y, z \in L$. Then Lemma 2 yields $\tilde{R}(x, y) \in \mathcal{I} \forall x, y \in L$. From the other hand, $\tilde{R}(x, y) \in W^1(E) \otimes \wedge^1 E^*$, so an

element $R \in W^2(E) \otimes \wedge^2 E^*$ there exists such that $R(x, y) = \delta^{-1}(\tilde{R}(x, y)) \quad \forall x, y \in E$. Then using (6) and Lemma 2 we obtain

$$R \in \mathcal{I}. \quad (10)$$

Let $R^i_{jkl} = \tilde{e}^i(\tilde{R}(e_k, e_l)e_j)$ be the components of curvature tensor. Denote $R^{ij}_{kl} = \omega^{jp}R^i_{pkl}$, where ω^{ij} is the matrix inverse to $\omega(e_i, e_j)$. Then it is well known [11] that $R^{ij}_{kl} = R^{ji}_{kl}$, so we have

$$R = R^{ij}_{kl} e_i \circ e_j \otimes \tilde{e}^k \wedge \tilde{e}^l$$

which coincides with the notations of [9, 10].

Let $T \in W^1(E) \otimes \wedge^2 E^*$, $T(x, y) = \nabla_x y - \nabla_y x - [x, y]$ be the torsion of ∇ . Then analogously to (10) we obtain

$$T \in \mathcal{I}. \quad (11)$$

Suppose \wp is a leaf of the distribution L , $\Phi = \{f \in A \mid f|_{\wp} = 0\}$ is the vanishing ideal of \wp in A , and $\mathcal{I}_{\Phi} = \Phi W(E) \otimes \wedge E^*$, $\mathcal{I}_{\text{fin}} = \mathcal{I} + \mathcal{I}_{\Phi}$ are homogeneous ideals in $W(E) \otimes \wedge E^*$. Then due to (5),(6) we can define the subcomplex $\delta \mathcal{I}_{\text{fin}}^{\bullet} = (\mathcal{I}_{\text{fin}}, \delta)$ with the same contracting homotopy δ^{-1} . Note that $\tau(\mathcal{I}_{\text{fin}}) = \Phi$, then using (2) we obtain

$$H^0(\delta \mathcal{I}_{\text{fin}}^{\bullet}) = \Phi, \quad H^i(\delta \mathcal{I}_{\text{fin}}^{\bullet}) = 0, \quad i = 1, 2, \dots \quad (12)$$

It is easily seen that vector fields of L preserve Φ , i.e. $(\nabla f)(x) \in \Phi \quad \forall f \in \Phi, x \in L$. Then from Lemma 2 we obtain $\nabla \Phi \in \mathcal{I}_{\Phi} + \mathcal{I}_{\wedge}^1$, so we finally obtain

$$\nabla \mathcal{I}_{\Phi} \subset \mathcal{I}_{\text{fin}}. \quad (13)$$

4 Fedosov complex and star-product

Let $W^{(i)}(E)$ be the grading in $W(E)$ which coincides with $W^i(E)$ except for the $\lambda \in W^{(2)}(E)$, and let $W_{(i)}(E)$ be the decreasing filtration generated by $W^{(i)}(E)$. Suppose $\widehat{W}(E)$, $\widehat{\mathcal{I}}$ are completions of $W(E)$, \mathcal{I} with respect to this filtration, then $\widehat{\mathcal{I}}$ is a left ideal in $\widehat{W}(E) \otimes \wedge E^*$. Let A_i , $i = 0, 1, 2, \dots$ be an (λ) -adic filtration in A , then $\tau(W_{(i)}(E)) \subset A_{\{i/2\}}$. Then $W_{(2i)}(E) \subset \tau^{-1}(A_i)$, so τ is continuous in the topologies generated by $W_{(i)}(E)$ and A_i and thus can be extended to a mapping $\widehat{W}(E) \rightarrow \widehat{A}$. Since δ, δ^{-1} and ∇ have fixed bidegrees with respect to bigrading $W^{(i)}(E) \otimes \wedge^n E^*$, we can extend them to derivations of $\widehat{W}(E) \otimes \wedge E^*$ in such a way that Eq. (2) remains true and they preserve $\widehat{\mathcal{I}}$. So we will write $A, W(E)$ etc. instead of $\widehat{A}, \widehat{W}(E)$ etc.

Let

$$r_0 = \delta^{-1}T, \quad r_{n+1} = \delta^{-1} \left(R + \nabla r_n + \frac{1}{\lambda} r_n^2 \right), \quad n = 0, 1, 2, \dots$$

Then it is well known that the sequence $\{r_n\}$ has a limit $r \in W_{(2)}(E) \otimes \wedge^1 E^*$. Then we can define well-known Fedosov complex $DW^{\bullet} = (W(E) \otimes \wedge^n E^*, D)$ with the differential ([9, 10], see also [6, 13] for the case of nonzero torsion)

$$D = \delta + \nabla - \frac{i}{\lambda} [[r, \cdot]],$$

where $[[\cdot, \cdot]]$ is the commutator in $W(E) \otimes \wedge E^*$.

Let F be an Abelian group which is complete with respect to its decreasing filtration F_i , $i = 0, 1, 2, \dots$, $\cup F_i = F$, $\cap F_i = \emptyset$. Let $\deg a = \max\{i : a \in F_i\}$ for $a \in F$.

Lemma 3 ([6]). *Let $\varphi : F \rightarrow F$ be a set-theoretic map such that $\deg(\varphi(a) - \varphi(b)) > \deg(a - b)$ for all $a, b \in F$. Then the map $Id + \varphi$ is invertible.*

Let $Q = Id + \delta^{-1}(D - \delta)$, then it is well known that $\delta Q = QD$ and from Lemma 3 it follows that Q is invertible, so it is a chain equivalence and we obtain

$$H^n(Q) : H^n(DW^\bullet) \cong H^n(\delta W^\bullet), \quad n = 0, 1, 2, \dots$$

Using (6),(9),(10),(11) and taking into account that \mathcal{I} is a left ideal in $W(E) \otimes \wedge E^*$ we have $r_n \in \mathcal{I}$ for all n , so $r \in \mathcal{I}$. Using (5),(6),(9),(13) we see that $D\mathcal{I}_{\text{fin}} \subset \mathcal{I}_{\text{fin}}$ and $Q\mathcal{I}_{\text{fin}} \subset \mathcal{I}_{\text{fin}}$, so we can define the subcomplex $D\mathcal{I}_{\text{fin}}^\bullet = (\mathcal{I}_{\text{fin}}, D)$ and using Lemma 3 we obtain

$$H^n(Q) : H^n(D\mathcal{I}_{\text{fin}}^\bullet) \cong H^n(\delta\mathcal{I}_{\text{fin}}^\bullet), \quad n = 0, 1, 2, \dots$$

Then due to (3),(12) we have the following diagram

$$\begin{array}{ccc} A = H^0(\delta W^\bullet) & \xrightarrow[\cong]{Q} & H^0(DW^\bullet) \\ \uparrow \cup & & \uparrow \cup \\ \Phi = H^0(\delta\mathcal{I}_{\text{fin}}^\bullet) & \xrightarrow[\cong]{Q} & H^0(D\mathcal{I}_{\text{fin}}^\bullet). \end{array} \quad (14)$$

Then we can define the Fedosov-type star-product $A \times A \ni (f, g) \mapsto f *_L g \in A$ on M carrying the multiplication from $H^0(DW^\bullet)$ onto $H^0(\delta W^\bullet)$:

$$f *_L g = Q(Q^{-1}f \circ Q^{-1}g). \quad (15)$$

Then from (14) we see that Φ is a left ideal in $\mathcal{A} = (A, *_L)$ since $H^0(D\mathcal{I}_{\text{fin}}^\bullet)$ is a left ideal in $H^0(DW^\bullet)$. Since the choice of φ does not affects $*_L$, we have proved the following theorem.

Theorem 2. *Let M be a symplectic manifold and let $L \subset TM$ be a polarization on M . Suppose ΓTM has a $\dim M$ -dimensional basis as a $C^\infty(M)$ -module. Then there exists a star-product on M adapted to all the leaves of L .*

This extends the results of Xu [14] who constructed commutative subalgebra of \mathcal{A} associated with a lagrangian submanifold.

Corollary 1. *$A/\Phi \cong C^\infty(\varphi)$ has a natural structure of left \mathcal{A} -module.*

Suppose L' is a polarization, then M is a *bi-Lagrangian manifold* [7] (called the Fedosov manifold of Wick type in [5]) and we can choose a connection on M such that $\nabla_z x \in L$ and $\nabla_z y \in L'$, $x \in L$, $y \in L'$, $z \in E$. Let φ' be a leaf of L' and $\Phi' = \{f \in A \mid f|_{\varphi'} = 0\}$. Then analogously to Theorem 2 we see that Φ' is a right ideal in \mathcal{A} . It is well known that for any two transversal involutive distributions L, L' we can choose a local coordinate system $\{x_i\}$ on M such that $\{\partial/\partial x_\alpha \mid \alpha = 1, \dots, \nu\}$ and $\{\partial/\partial x_\beta \mid \beta = \nu + 1, \dots, 2\nu\}$ are local bases in L and L' respectively ([1], Ch.1, Problem 30; see also [16], Sec.4.9 for the case when L, L' are polarizations). Write the star-product (15) as a bidifferential operator on M :

$$f *_L g = \sum_{r,s} \sum_{\substack{i_1, \dots, i_r \\ j_1, \dots, j_s}} \Lambda^{i_1 \dots i_r | j_1 \dots j_s} \frac{\partial^r f}{\partial x^{i_1} \dots \partial x^{i_r}} \frac{\partial^s g}{\partial x^{j_1} \dots \partial x^{j_s}}.$$

Since Φ is a left ideal in \mathcal{A} , we see that $\Lambda^{i_1 \dots i_r | \beta_1 \dots \beta_s} \in \Phi$ and analogously $\Lambda^{\alpha_1 \dots \alpha_r | j_1 \dots j_s} \in \Phi'$ since Φ' is a right ideal. But φ, φ' are arbitrary, so $*_L$ is a star-product with separation of variables in the sense of Karabegov [12] (called a star-product of Wick type in [3]):

$$f *_L g = \sum_{r,s} \sum_{\substack{\beta_1 \dots \beta_r \\ \alpha_1 \dots \alpha_s}} \Lambda^{\beta_1 \dots \beta_r | \alpha_1 \dots \alpha_s} \frac{\partial^r f}{\partial x^{\beta_1} \dots \partial x^{\beta_r}} \frac{\partial^s g}{\partial x^{\alpha_1} \dots \partial x^{\alpha_s}}$$

and we have the following theorem.

Theorem 3. *Let M be a bi-Lagrangian manifold and suppose ΓTM has a $\dim M$ -dimensional basis as a $C^\infty(M)$ -module. Then there exists a star-product with separation of variables on M .*

This extends the results of Bordemann and Waldman [3] who constructed a Fedosov star-product of Wick type on arbitrary Kähler manifold.

References

1. Bishop R.L. and Crittenden R.J.: Geometry of manifolds. N.Y. and London: Academic Press, 1964
2. Bordemann M., Ginot G., Halbout G., Herbig H.-C., Waldmann S.: Formalité G_∞ adaptée et star-représentations sur des sous-variétés coisotropes, arxiv:math.QA/0504276
3. Bordemann, M., Waldmann, S.: A Fedosov Star Product of Wick Type for Kähler Manifolds, Lett. Math. Phys. 41 243–253 (1997)
4. Bourbaki N.: Groupes et algèbres de Lie, ch.1. Algèbres de Lie. P.:Hermann, 1971
5. Dolgushev V.A., Lyakhovich S.L. and Sharapov A.A.: Wick type deformation quantization of Fedosov manifolds, Nucl. Phys. B 606 647-672 (2001)
6. Donin J.: On the quantization of Poisson brackets, Adv.Math. 127 73-93 (1997)
7. Etayo Gordejuela F. and Santamaria R.: The canonical connection on a bi-lagrangian manifold, J.Phys.A: Math.Gen. 34 981-987 (2001)
8. Farkas D.: A ring-theorist's description of Fedosov quantization, Lett. Math. Phys. 51 161-177 (2000)
9. Fedosov B.V.: A simple geometrical construction of deformation quantization, J.Diff.Geom. 40 213-238 (1994)
10. Fedosov B.V.: Deformation quantization and index theory, Akademie, B., 1996
11. Gelfand I., Retakh V. and Shubin M.: Fedosov manifolds, Adv.Math.136 104-140 (1998)
12. Karabegov A.V.: Deformation quantization with separation of variables on a Kähler manifolds, Commun. Math. Phys. 180 745-755 (1996)
13. Karabegov A.V. and Schlichenmaier M.: Almost Kähler deformation quantization, Lett. Math. Phys. 57 135-148 (2001)
14. Xu P.: Fedosov *-products and quantum momentum maps, Commun. Math. Phys. 197 167-197 (1998)
15. Sridharan R.: Filtered algebras and representations of Lie algebras, Trans. Amer. Math. Soc. 100 530-550 (1961)
16. Woodhouse N.M.J.: Geometric quantization, Clarendon Press, Oxford, 1991