Generalized Relativistic Chapman-Enskog Solution of the Boltzmann Equation

A. L. García-Perciante¹, A. Sandoval-Villalbazo², L. S. García-Colín³

¹Depto. de Matemáticas Aplicadas y Sistemas, Universidad Autónoma Metropolitana-Cuajimalpa, Artificios #40, México DF 01120, México.

²Depto. de Física y Matemáticas, Universidad Iberoamericana,

Prolongación Paseo de la Reforma 880, México D. F. 01210, México

³Depto. de Física, Universidad Autónoma Metropolitana-Iztapalapa,

Av. Purísima y Michoacán S/N, México D. F. 09340,

México. Also at El Colegio Nacional, Luis González Obregón 23, Centro Histórico, México D. F. 06020, México.

(Dated: September 28, 2018)

Abstract

The Chapman-Enskog method of solution of the relativistic Boltzmann equation is generalized in order to admit a time-derivative term associated to a thermodynamic force in its first order solution. Both existence and uniqueness of such a solution are proved based on the standard theory of integral equations. The mathematical implications of the generalization here introduced are thoroughly discussed regarding the nature of heat as chaotic energy transfer in the context of relativity theory.

I. INTRODUCTION

The standard treatment of the Boltzmann equation (BE) provides microscopic support to the transport equations commonly used in science and engineering problems [1][2]. It is well known that whereas the local equilibrium solution to the homogeneous BE leads to the Euler equations of hydrodynamics, its first order correction in the Knudsen parameter yields the Navier-Stokes-Fourier transport equations of fluid mechanics for a dilute gas. In the standard non-relativistic method of establishing the first order correction to the local equilibrium function, all partial time derivatives are replaced with spatial gradients using the Euler equations. Both existence and uniqueness of the solutions built in this scheme have been shown using the theory of integral equations [3].

On the other hand, in relativity theory time is simply a new coordinate $ct = x^4$. According to this fact, the direct application of the conventional method used to obtain the first order correction to the local equilibrium function is shown here to lead to a first order correction to the distribution function in terms of both spatial and time derivatives of the local variables namely, the generalized four-component thermodynamic forces. This is done following the tenets of relativistic linear irreversible thermodynamics as shown in Ref. [4]. We wish to emphasize that in this work the Meixner version of non-equilibrium thermodynamics is developed. However, in most standard works on relativistic kinetic theory [5, 6, 7, 8] these ideas are ignored. The consequence of the absence of time components in the structure of the thermodynamic forces is on the one hand, the obtention of transport equations which apparently violate causality, and on the other hand the tensorial structure of the heat flux in relativity, which is still a subject of debate [9]. The main motivation of this work is precisely to critically analyze these fine points.

The mathematical procedure here used to show that the solution obtained, considering four component thermodynamic forces, exists and is unique is based on the standard theory of integral equations. One requires orthogonality between the inhomogeneous part of the equation and the collisional invariants. This relation implies the validity of Euler equations. Nevertheless the use of these equations is not required for the construction of the solution to first order in the gradients. These results have strong implications on the structure of the four component heat flux as will be discussed in this work. Finally, the usual subsidiary conditions are invoked in order to adjust integration constants. This has, as will be discussed in this work, strong implications on the structure of the heat flux tensor.

The paper is divided as follows. In Section 2 the relativistic BE and some elements of the Chapman-Enskog method of solution are briefly reviewed. The general form of the first order in the gradients solution to the relativistic BE is proposed in Section 3 and both its existence and uniqueness are proved in Section 4. Finally, Section 5 includes a discussion of the implications of the results and some concluding remarks.

II. THE CHAPMAN-ENSKOG EXPANSION

The starting point of the generalized formalism is the special relativistic Boltzmann equation for a simple system in the absence of external forces [5, 6]:

$$v^{\alpha}f_{,\alpha} = J\left(f,\,f'\right)\,,\tag{1}$$

which describes the evolution of the single particle distribution function $f = f(x^{\nu}, v^{\nu} | n(x^{\nu}, t), u(x^{\nu}, t), \varepsilon(x^{\nu}, t))$ namely, the molecular number density in phase space. In Eq. (1)

$$f_{,\alpha} = \begin{bmatrix} \frac{\partial f}{\partial x^k} \\ \frac{1}{c} \frac{\partial f}{\partial t} \end{bmatrix}, \qquad (2)$$

and the index α , as well as all greek indices in the rest of this work, runs from 1 to 4 while the latin ones run up to 3. The local variables, number density n, hydrodynamic velocity u^k and internal energy ε per unit of mass are defined in terms of f as follows

$$n = \int f\gamma dv^* \,, \tag{3}$$

$$nu^k = \int f\gamma v^k dv^* \,, \tag{4}$$

$$n\varepsilon = mc \int f\gamma v^4 dv^* \,. \tag{5}$$

Here v^{μ} is the molecular velocity four-vector which we denote in terms of the three velocity \vec{w} as

$$v^{\alpha} = \begin{bmatrix} \gamma \vec{w} \\ \gamma c \end{bmatrix} , \tag{6}$$

where $\gamma = (1 - w^2/c^2)^{-1/2}$ is the usual relativistic factor. Also, the differential velocity element can be written as [7]

$$dv^* = \gamma^5 \frac{cd^3w}{v^4} = 4\pi c^3 \sqrt{\gamma^2 - 1} d\gamma \,, \tag{7}$$

and, as usual, c is the speed of light. Equation (1) implies that changes in the distribution function are due to collisions, represented on the right side of Boltzmann's equation through the so called collision kernel J(f, f') which is given by

$$J(f, f') = \int \int (f' f'_1 - f f_1) \sigma(\Omega) g d\Omega dv_1^*.$$
(8)

In Eq. (8) primes denote quantities after a binary collision between particles with velocity v and v_1 takes place; $\sigma(\Omega) d\Omega$ is the differential cross section element and g is the relative velocity.

To solve Eq. (1) following what is now known in the literature as the Chapman-Enskog method [1], originally due to Hilbert, the distribution function is expanded in a power series of the Knudsen's parameter ϵ around the local equilibrium distribution function $f^{(0)}$

$$f = f^{(0)} \left(1 + \epsilon \phi^{(1)} + \epsilon^2 \phi^{(2)} + \cdots \right) \,. \tag{9}$$

The expansion parameter ϵ is a measure of the relative magnitude of the gradients of the local variables within a mean free path and the characteristic size of the system. The local equilibrium distribution function, solution to the homogeneous relativistic BE J(f, f') = 0, is the well known Jüttner function. For particles of rest mass m, relativistic parameter $z = \frac{kT}{mc^2}$ and in the non-degenerate case [5, 6]:

$$f^{(0)} = \frac{n}{4\pi c^3 z \mathcal{K}_2\left(\frac{1}{z}\right)} e^{-\frac{\gamma}{z}} \,. \tag{10}$$

As usual, k is the Boltzmann constant, n and T are the local density and temperature, respectively, and \mathcal{K}_2 represents the modified Bessel function of the second kind.

Substitution of Eq. (9) in Eq. (1) and considering only linear deviations in ϵ from local equilibrium yields

$$v^{\alpha} f_{,\alpha}^{(0)} = J \left(f^{(0)} \left(1 + \epsilon \phi \right), \, f'^{(0)} \left(1 + \epsilon \phi' \right) \right) \,, \tag{11}$$

where $\epsilon \phi^{(1)} \equiv \phi$ and $f^{(0)}$, as pointed above, is the solution to the homogeneous equation namely,

$$J(f^{(0)}, f^{(0)}) = 0,$$

or

$$f^{(0)}f'^{(0)} = f^{(0)}f_1^{(0)}.$$
(12)

Introducing the relation given by Eq. (12) in the right hand side of Eq. (11) leads to

$$J(f, f') = f^{(0)} \int \int [\phi'_1 + \phi' - \phi_1 - \phi] f_1^{(0)} \sigma(\Omega) g d\Omega dv_1^*.$$
(13)

Thus, this procedure yields a linear inhomogeneous integral equation for ϕ , namely

$$v^{\alpha} f^{(0)}_{,\alpha} = f^{(0)} \mathbb{C} (\phi) ,$$
 (14)

where

$$\mathbb{C}(\phi) = \int \int \left\{ \phi_1' + \phi' - \phi_1 - \phi \right\} f_1^{(0)} \sigma(\Omega) g d\Omega dv_1^*, \qquad (15)$$

is the linearized Boltzmann collision kernel. The following section is devoted to the obtention of the solution to Eq. (14).

III. SOLUTION TO FIRST ORDER IN THE GRADIENTS

Following the theory of integral equations as presented in Ref. [3], the general solution to Eq. (14) may be written as the sum of a solution to the inhomogeneous equation ϕ_P and an arbitrary linear combination of the solutions of the homogeneous equation ϕ_H :

$$\phi = \phi_H + \phi_P \,. \tag{16}$$

The homogeneous equation has five solutions, the collisional invariants namely,

$$\mathbb{C}\begin{pmatrix} mv^{\mu} \\ m\gamma v^{4} \end{pmatrix} = 0.$$
(17)

Notice that, for $\mu = 4$ Eq. (17) corresponds to the particle number conservation. A solution given by Eq. (16) exists if these invariants in ϕ_H are orthogonal to the left hand side of Eq. (14) namely

$$\int \left(\begin{array}{c} mv^{\mu} \\ m\gamma v^4 \end{array} \right) v^{\alpha} f^{(0)}_{,\alpha} dv^* = 0.$$
(18)

The relations in Eq. (18) correspond precisely to the relativistic Euler equations and thus the orthogonality conditions are satisfied. However we recall that there is an infinite number of solutions since ϕ_H is an arbitrary linear combination of collisional invariants. The unicity is achieved when the arbitrary constants are determined with the use of the subsidiary conditions as we shall argue below.

In order to find a particular solution to the inhomogeneous equation, we start by calculating the left hand side of Eq. (14). In the comoving frame and assuming that no external forces are present

$$v^{\alpha}f^{(0)}_{,\alpha} = v^{\alpha}\left(\frac{\partial f^{(0)}}{\partial n}n_{,\alpha} + \frac{\partial f^{(0)}}{\partial T}T_{,\alpha}\right).$$
(19)

Since $\alpha = 1, 2, 3, 4$ each gradient on the right side of Eq. (19) has four components. It is precisely at this point where our formalism deviates from the standard one [1, 5, 6]. In conventional relativistic treatments, one separates time and space derivatives using a projection operator and introduces Euler's equations in order to eliminate the time derivatives of the local variables. However, special relativity is a four dimensional formalism in which time corresponds to a coordinate. Because of this feature we here propose that the four-gradients are to be considered with the same status as the generalized thermodynamic forces [10]. *This is why we do not resort to the Euler equations to eliminate time derivatives*. Also, in most standard relativistic treatments, the temperature gradient is written in terms of the pressure gradient in order to introduce, through the momentum balance equation, an acceleration term so that an Eckart-type constitutive equation is sustained [5]. However, in the Meixner-type scheme here considered, such a substitution needs not to be carried out since the thermodynamic forces are $T_{,\nu}$ and $n_{,\nu}$ exclusively.

In order to proceed, we first notice that since the calculations will be performed in the comoving frame, $\frac{\partial u^{\alpha}}{\partial t} = 0$ and $p_{,k} = 0$. Thus, using the equation of state for an ideal gas one can write for the space components of the gradients that

$$\frac{n_{,k}}{n} = -\frac{T_{,k}}{T} \,. \tag{20}$$

For the time component of the density gradient, in the comoving frame, one has

$$n_{,4} = 0$$
, (21)

since the continuity equation is satisfied to any order in ϵ . Introducing Eqs. (20) and (21) in Eq. (19) and calculating explicitly the derivatives of the Jüttner function, the integral equation for ϕ_P can be written as

$$\frac{v^k T_{,k}}{zT} f^{(0)} \left[\gamma - \mathcal{G} \left(\frac{1}{z} \right) \right] + \frac{v^4 T_{,4}}{T} f^{(0)} \left[\frac{1}{z} \left(\gamma - \mathcal{G} \left(\frac{1}{z} \right) \right) + 1 \right] = f^{(0)} \mathbb{C} \left(\phi_P \right) , \qquad (22)$$

where $\mathcal{G} \left(\frac{1}{z} \right) = \mathcal{K}_3 \left(\frac{1}{z} \right) / \mathcal{K}_2 \left(\frac{1}{z} \right).$

Since the thermodynamic forces present on the left hand side of Eq. (22) are independent, we seek a solution as a linear combination of such forces:

$$\phi_P = \mathcal{A}^{\nu} \frac{T_{,\nu}}{T} \,, \tag{23}$$

which is a direct generalization of the structure proposed in the non-relativistic case. The coefficients \mathcal{A}^{ν} , which are in general functions of v^{α} , n and T, are solutions to

$$\frac{v^k}{z} \left[\gamma - \mathcal{G}\left(\frac{1}{z}\right) \right] = \mathbb{C}\left(\mathcal{A}^k\right) \,, \tag{24}$$

for the first three components and,

$$v^{4}f^{(0)}\left[\frac{1}{z}\left(\gamma - \mathcal{G}\left(\frac{1}{z}\right)\right) + 1\right] = f^{(0)}\mathbb{C}\left(\mathcal{A}^{4}\right), \qquad (25)$$

for the fourth component. It now remains to show that one may obtain a unique solution to Eq. (14) by using the subsidiary conditions. This task will be performed in the following section.

IV. SUBSIDIARY CONDITIONS AND UNIQUENESS OF THE SOLUTION

According to the previous discussion, the general solution to the relativistic BE, Eq. (1), to first order in the Knudsen parameter, has the general form

$$\phi = \mathcal{A}^{\nu} \frac{T_{,\nu}}{T} + \alpha + \tilde{\alpha}_{\nu} v^{\nu} , \qquad (26)$$

where the first term corresponds to the particular solution, Eq. (23), while the remaining ones, as mentioned above, are an arbitrary linear combination of the solutions to the homogeneous equation, namely the collisional invariants.

The components of \mathcal{A}^{ν} are given by the solution to Eqs. (24) and (25). However, in order to uniquely determine the complete solution ϕ , the constants α and $\tilde{\alpha}_{\mu}$ must be determined. This is usually achieved by arbitrarily assuming that the local macroscopic variables are determined only from the local equilibrium state. That is,

$$\int \phi f^{(0)} \left(\begin{array}{c} m v^{\mu} \\ m \gamma v^4 \end{array} \right) dv^* = 0 \,, \tag{27}$$

which are the so-called subsidiary conditions [1, 2]. Since spatial and temporal terms in ϕ_P have opposite parities (in v), it is convenient at this point to explicitly separate such terms by proposing that

$$\mathcal{A}^{\nu} = \begin{bmatrix} a'(\gamma) v^k \\ a''(\gamma) v^4 \end{bmatrix}.$$
 (28)

Thus, Eq. (27) can be written as

$$\int \left(a'(\gamma) \, v^k T_{,k} + a''(\gamma) \, v^4 T_{,4} + \alpha + \tilde{\alpha}_{\nu} v^{\nu} \right) \psi f^{(0)} dv^* = 0 \,, \tag{29}$$

where ψ are the collisional invariants v^{μ} and γv^4 . For $\psi = v^4$, the odd parity in $a'(\gamma) v^k T_{,k}$ and $\tilde{\alpha}_k v^k$ yields zero values for the respective integrals and thus

$$\int \left(a''(\gamma) v^4 T_{,4} + \alpha + \tilde{\alpha}_4 v^4 \right) \gamma f^{(0)} dv^* = 0.$$
(30)

In a similar way, for $\psi = v^{\ell}$ the only non-vanishing terms due to parity are $a'(\gamma) v^k T_{k}$ and $\tilde{\alpha}_k v^k$, whence

$$\int \left(a'(\gamma) v^k T_{,k} + \tilde{\alpha}_k v^k\right) v^\ell f^{(0)} dv^* = 0, \qquad (31)$$

which is non-zero only for $\ell = k$. In this case

$$\int \left(a'(\gamma) v^k T_{,k} + \tilde{\alpha}_k v^k\right) v^\ell f^{(0)} dv^* = \frac{1}{3} \int \left(a'(\gamma) T_{,k} + \tilde{\alpha}_k\right) \gamma^2 w^2 f^{(0)} dv^* \tag{32}$$

Thus, the three subsidiary conditions for $\psi = v^{\ell}$ may be written as

$$\int (a'(\gamma) T_{,k} + \tilde{\alpha}_k) \gamma^2 w^2 f^{(0)} dv^* = 0.$$
(33)

while the subsidiary condition corresponding to the fifth collisional invariant, $\psi = \gamma v^4$, is

$$\int \left(a''(\gamma) v^4 T_{,4} + \alpha + \tilde{\alpha}_4 v^4\right) \gamma v^4 f^{(0)} dv^* = 0.$$
(34)

From Eq. (33), it is straightforward to obtain the relation

$$\tilde{\alpha}_{k} = -T_{,k} \frac{\int a'(\gamma) \,\gamma^{2} w^{2} f^{(0)} dv^{*}}{\int \gamma^{2} w^{2} f^{(0)} dv^{*}}, \qquad (35)$$

which implies $\tilde{\alpha}_k \propto T_{,k}$. Thus, one can redefine the coefficients appearing in Eq. (29) as follows

$$a'(\gamma) v^k T_{,k} + \tilde{\alpha}_k v^k \longrightarrow a'(\gamma) v^k T_{,k} , \qquad (36)$$

This implies that the subsidiary condition given in Eq. (31) now reads

$$\int a'(\gamma) \gamma^2 w^2 f^{(0)} dv^* = 0.$$
(37)

On the other hand, from Eqs. (30) and (34) an *inhomogeneous* system of equations for α and $\tilde{\alpha}_4$ is obtained:

$$g_{11}\alpha + g_{12}\tilde{\alpha}_4 = s_1 T_{,4} \,, \tag{38}$$

$$g_{21}\alpha + g_{22}\tilde{\alpha}_4 = s_2 T_{,4} \,, \tag{39}$$

where the coefficients are

$$g_{11} = \int \gamma f^{(0)} dv^* \qquad g_{12} = g_{21} = \int \gamma^2 c f^{(0)} dv^* \qquad g_{22} = \int c^2 \gamma^3 f^{(0)} dv^* \,, \tag{40}$$

and the coefficients in the inhomogeneous terms are given by

$$s_1 = -\int a''(\gamma) \,\gamma^2 c f^{(0)} dv^* \qquad s_2 = -\int a''(\gamma) \,c^2 \gamma^3 f^{(0)} dv^* \,, \tag{41}$$

The solution to this system of equations is

$$\alpha = T_{,4} \frac{s_1 g_{22} - s_2 g_{12}}{g_{11} g_{22} - g_{12} g_{21}}, \tag{42}$$

$$\tilde{\alpha}_4 = T_{,4} \frac{s_2 g_{11} - s_1 g_{21}}{g_{11} g_{22} - g_{12} g_{21}} \,. \tag{43}$$

Since, from Eqs. (42) and (43), both α and $\tilde{\alpha}_4$ are proportional to $T_{,4}$ one can redefine

$$a''(\gamma) v^4 T_{,4} + \alpha + \tilde{\alpha}_4 v^4 \longrightarrow a''(\gamma) v^4 T_{,4} , \qquad (44)$$

and therefore the two corresponding subsidiary conditions are now given by

$$\int a''(\gamma) \,\gamma^2 f^{(0)} dv^* = 0\,, \tag{45}$$

$$\int a''(\gamma) \gamma^3 f^{(0)} dv^* = 0, \qquad (46)$$

Putting these results together, the unique solution to the relativistic BE to first order in the gradients according to the Chapman-Enskog expansion has the general structure

$$f = f^{(0)} \left(1 + \mathcal{A}^{\nu} \frac{T_{,\nu}}{T} \right) , \qquad (47)$$

where \mathcal{A}^{ν} is defined in Eq. (28) and the coefficients $a'(\gamma)$ and $a''(\gamma)$ are given by the solution of the integral equations

$$\frac{v^{k}}{z}f^{(0)}\left(\gamma - \mathcal{G}\left(\frac{1}{z}\right)\right) = \mathbb{C}\left(a'\left(\gamma\right)v^{k}\right), \qquad (48)$$

$$v^{4}f^{(0)}\left[\frac{1}{z}\left(\gamma - \mathcal{G}\left(\frac{1}{z}\right)\right) + 1\right] = \mathbb{C}\left(a''\left(\gamma\right)v^{4}\right), \qquad (49)$$

subject to the subsidiary conditions given in Eqs. (37), (45), and (46).

Notice that, in the non-relativistic case as well as in the relativistic treatments given in the literature [5, 6], the absence of a fourth component of \mathcal{A}^{ν} leads to a homogeneous system instead of Eqs. (38) and (39) and is thus responsible for the appearance of the trivial solutions $\alpha = \tilde{\alpha}_4 = 0$. The consequences of this substantial difference are discussed in the next section.

V. DISCUSSION AND CONCLUDING REMARKS

By considering the four-dimensional gradients of local variables as thermodynamic forces within the Chapman-Enskog method of solution to the BE, a generalized first correction to the equilibrium distribution function has been found in the context of Meixner's relativistic irreversible thermodynamics. In the comoving frame, this procedure leads to a first order in ϵ solution with the general structure given by Eqs. (47) and (28). The proportionality coefficients, functions of the local variables and γ , can be obtained by an appropriate decomposition in orthogonal functions which simplifies the integral equations (48) and (49). Thus, the solution here obtained is suitable for the calculation of relativistic transport coefficients consistent with the phenomenological formalism proposed in Ref. [4]. Such a procedure is beyond the scope of this work and will be addressed elsewhere.

However one important point should be discussed here, namely the definition of heat flux in the context of relativity. We recall that in the non-relativistic theory this quantity is related to the transport of chaotic kinetic energy in agreement with the interpretation proposed by Clausius in 1857 [11]. The question here is: can this definition be extended within a relativistic framework? This, in principle, may be achieved by resorting to the relativistic version of the Maxwell-Enskog transport equation. Its explicit form was derived seven years ago [12] by two of us. This equation, in the notation here introduced reads

$$\left(n\left\langle\psi\frac{v^{\alpha}}{\gamma}\right\rangle\right)_{;\alpha} = 0\tag{50}$$

where $\psi = mv^{\mu}$, $m\gamma v^4$ is a collisional invariant and the bracket represents the statistical average defined, for any function $g(\gamma)$, as

$$\langle g \rangle = \frac{1}{n} \int f g \gamma dv^* \tag{51}$$

For $g = mv^4$ the heat transport equation is obtained. In particular, the heat flux will be given by

$$J^{\nu}_{[Q]} = mc^2 \int v^{\nu} f \gamma dv^*$$

This is a rather striking result because the fourth component of this vector turns out to be precisely the energy density as defined in Eq. (5). This implies that, because of the subsidiary conditions, such component will be identically equal to zero for any perturbation $\phi^{(n)}$ with $n \ge 1$. Thus we recover the usual form which reduces to the equation obtained by other authors resorting to the use of a spatial projector [5, 6, 8, 13].

Indeed to clarify this statement, we recall that Eckart proposed the following stress-energy tensor [8] for a simple, non-viscous fluid

$$T_{\mu\nu} = \frac{n\varepsilon}{c^2} u_{\mu} u_{\nu} + p h_{\mu\nu} + \frac{1}{c^2} u_{\mu} J_{[Q]\nu} + \frac{1}{c^2} u_{\nu} J_{[Q]\mu}$$
(52)

where $h_{\mu\nu} = g_{\mu\nu} - u_{\mu}u_{\nu}/c^2$ is the projector introduced by Eckart in order to eliminate dissipation in the time-axis. This work thus provides further support for this assumption. However, $T_{\mu\nu}$ as in Eq. (52) is in principle coupled to the Einstein tensor $G_{\mu\nu}$ by means of the field equation:

$$G_{\mu\nu} = \mathcal{K}T_{\mu\nu} \tag{53}$$

where \mathcal{K} is the coupling constant. The point here is how heat would affect the gravitational field if it is placed in the stress energy tensor. Indeed, heat is identified with molecular motion and its capability of transforming itself to mechanical energy by any process is restricted. On the other hand, the introduction of $J_{[Q]\nu}$ in Eq. (53) seems to allow a transformation of heat to work, via the gravitational field, with unitary efficiency. To the authors' knowledge, direct attempts to obtain solutions to the system (53) in the presence of heat and to analyze the dynamical consequences of the solutions are practically absent in the literature.

An alternative way of introducing heat in relativistic irreversible thermodynamics consists in a generalization of Meixner's approach [4]. In it, heat is not included in the stress energy tensor but introduced directly in the total energy flux. The construction of the entropy production, consistent with Clausius' idea of uncompensated heat, and consequent enforcement of the second law of thermodynamics in this context leads to the proposal of the constitutive equation

$$J^{\mu}_{[Q]} = -\kappa T^{,\mu} \tag{54}$$

where κ is the thermal conductivity. Equation (54) coincides with the constitutive equation in Eckart's formalism in the comoving frame where the acceleration term vanishes.

It is important to emphasize at this point that for the purpose of this discussion, the key difference between both formalism is the role of the heat flux. On the one hand it is included in the stress energy tensor in Eckart's formalism, as pointed out above. On the other hand, it is conceived only as energy in transit in the Meixner-like formalism and is introduced separately in the total energy flux. Clearly the latter conception does not predict any effects on the geometry of space-time due to heat since $J^{\mu}_{[Q]}$ is not included in T^{μ}_{ν} and thus the Einstein field equation remains unchanged. An experiment that could clarify this difference has already been suggested in Ref. [9].

On the other hand, the validity of Eq. (54) should be sustained by solving the relativistic Boltzmann equation. The calculations performed in the previous sections show that this is not the case. Indeed, the heat flux has no fourth component which is in agreement with other theories but for entirely different reasons.

We believe that this poses a rather intriguing question. Is it possible or not to think

of heat as a chaotic molecular motion in a relativistic framework? The correct answer to this question is crucial when considering both the structure of transport equations and the consistency of any relativistic kinetic theory with the second law of thermodynamics [4]. These questions merit further study.

- Chapman S. and Cowling T. G. 1970 Third Edition The Mathematical Theory of Non-Uniform Gases (Cambridge: Univ. Press)
- [2] Ferziger J. H. and Kaper H. G. 1972 Mathematical Theory of Transport Processes in Gases (Amsterdam: North Holland)
- [3] Courant C. and Hilbert D. 1989 Methods of Mathematical Physics (New York: J. Wiley)
- [4] García-Colín L. S. and Sandoval-Villalbazo A. 2006 Relativistic non-equilibrum thermodynamics revisited J. Non Equilib. Thermodyn. 31 11-22
- [5] Cercignani C. and Medeiros Kremer G. 2002 The relativistic Boltzmann equation: theory and applications (Berlin: Birkhäuser)
- [6] de Groot S. R., van Leeuwen W. A. and van Weert Ch. G. 1980 Relativistic Kinetic Theory (Amsterdam: North Holland)
- [7] Liboff R. L. and Liboff R. C. 2003 Kinetic theory: classical, quantum and relativistic descriptions (Berlin: Springer)
- [8] Eckart C. 1940 The thermodynamics of irreversible processes III: the relativistic theory of a simple fluid *Phys. Rev.* 58 919-924
- [9] Sandoval-Villalbazo A. and García-Colín L. S. 2006 Light scattering test regarding the relativistic nature of heat Proc. of the XXVIII Spanish Relativity Meeting, Eds. L. Mornas, J. Diaz-Alonso, AIP Press 603-606.
- [10] de Groot S. R. and Mazur P. 1984 Non equilibrium thermodynamics (Mineola: Dover)
- [11] Brush S. 1976 The kind of motion we call heat (Amsterdam: North Holland)
- [12] Sandoval-Villalbazo A. and García-Colín L. S. 2000 The relativistic kinetic formalism revisited Physica A 278 428-439
- [13] Israel W. 1963 Relativistic kinetic theory of a simple gas J. Math. Phys 4 1163-1181