G-STABLE PIECES AND LUSZTIG'S DIMENSION ESTIMATES

XUHUA HE

ABSTRACT. We use G-stable pieces to construct some equidimensional varieties and as a consequence, obtain Lusztig's dimension estimates [L2, section 4]. This is a generalization of [HL].

In Lemma 1.1 and Proposition 1.2, we assume that G is arbitrary connected algebraic group and \tilde{G} is an algebraic group with identity component G.

Lemma 1.1. Let $g \in \tilde{G}$. Define $i : \tilde{G} \to \tilde{G}$ by $i(h) = ghg^{-1}h^{-1}$. For any closed subgroup A of Z_G with $gAg^{-1} = A$, set $H_A = \{h \in G; i(h) \in A\}$. Then

(1) H_A is an algebraic group and $i|_{H_A} : H_A \to A$ is a morphism of algebraic groups.

(2) $i(A)^0 = i(H_A)^0$. (3) $\dim(H_A) = \dim(Z_G(g)) + \dim(A) - \dim(Z_A(g))$.

If $h, h' \in H_A$, then

$$\begin{split} i(hh') &= ghh'g^{-1}(h')^{-1}h^{-1} = (ghg^{-1}h^{-1})h(gh'g^{-1}(h')^{-1})h^{-1} \\ &= i(h)hi(h')h^{-1} = i(h)i(h') \in A \end{split}$$

and $hh' \in H_A$. If $h \in H_A$, then $i(h^{-1}) = h^{-1}i(h)^{-1}h = i(h)^{-1} \in A$ and $h^{-1} \in H_A$. Part (1) is proved.

Now $i(A)^0$ is a connected subgroup of $i(H_A)$. Define $\delta : A \to A$ by $\delta(z) = gzg^{-1}$. Then

$$\dim(i(A)) = \dim(A) - \dim(A^{\delta}).$$

Define $\sigma : A \to A$ by $\sigma(z) = \delta^{m-1}(z)\delta^{m-2}(z)\cdots z$, where *m* is the order of the automorphism δ . Then σ is a group homomorphism and

$$i(H_A) \subset \{z \in Z; \sigma(z) = 1\}$$

Notice that $\sigma(A^{\delta}) = \{t^m; t \in A^{\delta}\}$ is of dimension dim (A^{δ}) . Thus

$$\dim(i(H_A)) \leq \dim(A) - \dim(\sigma(A)) \leq \dim(A) - \dim(\sigma(A^{\delta}))$$
$$= \dim(A) - \dim(A^{\delta}).$$

Therefore, $\dim(i(A)) = \dim(i(H_A)) = \dim(A) - \dim(A^{\delta})$. Part (2) is proved.

The author is partially supported by NSF grant DMS-0700589.

Since the kernal of $i|_{H_A}$ is $Z_G(g)$,

$$\dim(H_A) = \dim(Z_G(g)) + \dim(i(H_A))$$
$$= \dim(Z_G(g)) + \dim(A) - \dim(A^{\delta})$$

Part (3) is proved.

Proposition 1.2. Let D, D' be connected group of \tilde{G} . Define $\delta : Z_G \to Z_G$ by $\delta(z) = gzg^{-1}$ for any $g \in D$ and $\delta' : Z_G \to Z_G$ by $\delta'(z) = g'z(g')^{-1}$ for any $g' \in D'$. Let c be a G-conjugacy class in D and Z be a closed subgroup of Z_G with $\delta(Z) = Z$. Set

$$X = \{(g,g'); g \in cZ, g' \in D', gg'g^{-1}(g')^{-1} \in Z\}.$$

If $X \neq \emptyset$, then DD' = D'D and $\delta\delta'(z) = \delta'\delta(z)$ for $z \in Z_G$. Moreover, X is of pure dimension $\dim(G) + \dim(Z) - \dim(\frac{\delta'(Z)}{Z \cap \delta'(Z)})^{\delta}$.

It is easy to see that if $X \neq \emptyset$, then DD' = D'D. Thus for $g \in D$ and $g' \in D'$, $g^{-1}(g')^{-1}gg' \in G$ and $\delta^{-1}(\delta')^{-1}\delta\delta'(z) = z$ for all $z \in Z_G$.

Consider the projection map $X \to D$ defined by $(g, g') \mapsto g$. Let g be in the image and X_g be the fiber over g. Fix $g' \in X_g$. Set $i: \tilde{G} \to \tilde{G}$ by $i(h) = ghg^{-1}h^{-1}$. Then i(hg') = i(h)i(g') for $h \in G$. Hence $X_g = H_Z g'$. Let $z \in Z$. Then for $h \in G$,

$$(zg)(hg')(zg)^{-1}(hg')^{-1} = zi(h)i(g')\delta'(z)^{-1}.$$

If $zi(h)i(g')\delta'(z)^{-1} \in Z$, then $i(h) \in Z\delta'(Z)$. Hence $h \in H_{Z\delta'(Z)}$ and $\delta'(z) \in i(H_{Z\delta'(Z)})Z$. On the other hand, if $z \in Z$ with $\delta'(z) \in i(H_{Z\delta'(Z)})Z$, then there exists $h \in G$ with $zi(h)i(g')\delta'(z)^{-1} \in Z$. Therefore, for $z \in Z$, $X_{zg} \neq \emptyset$ if and only if $z \in Z'$, where $Z' = \{z \in Z; \delta'(z) \in i(H_{Z\delta'(Z)})Z\}$. It is easy to see that Z' is an algebraic group.

By part (2) of the previous lemma, $i(H_{Z\delta'(Z)})^0 \subset i(Z\delta'(Z))$. Hence $\delta'(Z')^0 = (\delta'(Z) \cap i(Z\delta'(Z))Z)^0 = (\delta'(Z) \cap i(\delta'(Z))Z)^0$. Notice that $i(Z) \subset \delta(Z)Z = Z$ and $i\delta'(Z) \subset \delta\delta'(Z)\delta'(Z) = \delta'(Z)$. Now *i* induces a group morphism $\overline{i} : \delta'(Z)/(Z \cap \delta'(Z)) \to \delta'(Z)/(Z \cap \delta'(Z))$. We have that $z \in \delta'(Z) \cap i(\delta'(Z))Z$ if and only if $z(Z \cap \delta'(Z))$ is contained in the image of \overline{i} . Hence $\dim(Z') = \dim(\frac{\delta'(Z)}{Z \cap \delta'(Z)}) - \dim(\frac{\delta'(Z)}{Z \cap \delta'(Z)})^{\delta} + \dim(Z \cap \delta'(Z))$ $\delta'(Z)) = \dim(Z) - \dim(\frac{\delta'(Z)}{Z \cap \delta'(Z)})^{\delta}$.

Set $Y = \{(h, z); z \in Z', h \in X_{zg}\}$. Then we have the projection map $Y \to Z'$ and each fiber is isomorphic to H_Z . Hence Y is of pure dimension dim $(Z') + \dim(H_Z)$.

Consider the morphism $G \times Y \to X$ defined by

$$(g_1, h, z) \mapsto (g_1 h g_1^{-1}, z g_1 g g_1^{-1}).$$

Then this morphism is surjective and the fiber over (h, zg) is

$$\{(g_1, g_1^{-1}hg_1, i(g_1)z); g_1 \in H_Z\},\$$

which is of pure dimension $\dim(H_Z)$. Therefore X is of pure dimension $\dim(G) + \dim(Z')$.

 $\mathbf{2}$

1.3. From now on, we assume that G be a simply-connected, semisimple algebraic group over an algebraically closed field **k**. We fix a Borel subgroup B of G and a maximal torus $T \subset B$. Let I be the set of simple roots determined by B and T.

For any $J \subset I$, let P_J be the standard parabolic subgroup corresponding to J and \mathcal{P}_J be the set of parabolic subgroups that are G-conjugate to P_J . We simply write \mathcal{P}_{\emptyset} as \mathcal{B} . Let L_J be the Levi subgroup of P_J that contains T.

For any parabolic subgroup P, let U_P be the unipotent radical of Pand H_P be the inverse image of the connected center of P/U_P under the projection map $\pi_P : P \to P/U_P$. We simply write U for U_B .

For $J \subset I$, we denote by W_J the standard parabolic subgroup of W generated by J and by W^J (resp. JW) the set of minimal coset representatives in W/W_J (resp. $W_J \setminus W$). For $J, K \subset I$, we simply write $W^J \cap {}^KW$ as ${}^KW^J$.

For $P \in \mathcal{P}_J$ and $Q \in \mathcal{P}_K$, we write pos(P,Q) = w if $w \in {}^JW^K$ and there exists $g \in G$ such that $P = gP_Jg^{-1}$, $Q = g\dot{w}P_K\dot{w}^{-1}g^{-1}$, where \dot{w} is a representative of w in N(T).

For $g \in G$ and $H \subset G$, we write ${}^{g}H$ for gHg^{-1} .

For any algebraic group H, let H^0 be its identity component.

1.4. Let σ be a diagram automorphism of G, i.e., an automorphism of G that stabilizes B and T and the order of σ as an automorphism of G coincides with the order of the associated permutation on I. We use the same symbol σ for the associated automorphism on W and associated permutation on I. Set $\tilde{G} = G \rtimes \langle \sigma \rangle$, where $\langle \sigma \rangle$ is the finite subgroup of G generated by σ . We simply write the element $(g, \sigma^n) \in \tilde{G}$ as $g\sigma^n$. For each element $g \in \tilde{G}$, we write g_s for its semisimple part and g_u its unipotent part.

Let $D = (G, \sigma)$ be a connected component of G. We have the following result.

Proposition 1.5. Let $g \in D$. Then g is G^0 -conjugate to an element of the form $t\sigma u$, where $t \in (T^{\sigma})^0$ and u is a unipotent element in $Z^0_G(t\sigma_s)$.

By [St, Lemma 7.3], after conjugate by an element in G^0 , we may assume that $g \in B\sigma \subset B \ltimes < \sigma >$. Then $g_s \in B\sigma_s$ and $g_u \in B\sigma_u$. Then after conjugate by an element in B, we may assume that $g_s = t_1\sigma_s$ and $g_u = t_2\sigma_u u$ for $t_1, t_2 \in T$ and $u \in U$. By [L2, 1.2], after conjugate by an element in T, we may assume furthermore that $t_2 \in (T^{\sigma_u})^0$. Consider the group morphism $B \ltimes < \sigma > \to T \ltimes < \sigma >$. Since g_u is unipotent, then $t_2\sigma_u$ is also unipotent. Notice that t_2 commutes with σ_u . Then t_2 is unipotent and $t_2 = 1$.

Since σ is a diagram automorphism, σ_s and σ_u are also diagram automorphisms. In particular, $\rho^{\vee}(t) \in T^{\sigma_{\sigma}} \cap T^{\sigma_u}$ for all $t \in \mathbf{k}^{\times}$. Hence $\operatorname{Ad}(\rho^{\vee}(t))g_u \in Z_G(g_s)$. Since σ_u is contained in the closure

XUHUA HE

of $\{\operatorname{Ad}(\rho^{\vee}(t))g_u)\}$ and $Z_G(g_s)$ is closed, we have that $\sigma_u \in Z_G(g_s)$. We also have that $u \in Z_G^0(g_s)$.

Now σ_u commutes with $t_1\sigma_s$. Hence $t_1 \in T^{\sigma_u}$ By [Bo, 9.6], (T^{σ_u}) is connected. Notice that σ_s is an automorphism on T^{σ_u} . Then by [L2, 1.2], after conjugate by T^{σ_u} , $t_1 \in ((T^{\sigma_u})^{\sigma_s})^0 \subset (T^{\sigma})^0$.

1.6. Let D//G be set of closed *G*-conjugacy classes in *D*. By geometric invariant theory, D//G has a natural structure of affine variety and there is a well-defined morphism St : $D \to D//G$ which maps the element $g \in D$ to the unique closed *G*-conjugacy class in *D* that is contained in the closure of the *G*-conjugacy class of *g*. If σ is trivial, then St is just the Steinberg morphism of *G*. Thus for arbitrary σ , we call St the Steinberg morphism of *D* and the fibers the Steinberg fibers of *D*.

By the previous proposition, any element $g \in D$ is of the form $t\sigma u$, where $t \in (T^{\sigma})^0$ and u is a unipotent element in $Z^0(t\sigma_s)$. Moreover, $t\sigma_s$ is contained in the closure of the *G*-conjugacy class of g. Hence $\operatorname{St}(g) = \operatorname{St}(t\sigma_s)$. Notice that $t\sigma$ is quasi-semisimple in the sense of [St, Sect.9], i.e. the automorphism of *G* obtained by conjugation by $t\sigma_s$ will fix a Borel subgroup and a maximal torus thereof. As a consequence, the *G*-conjugacy class of $t\sigma_s$ in *D* is closed [Sp, II.1.15(f)]. We conclude that any Steinberg fiber of *D* is of the form

$$\bigcup_{g \in G \ u \text{ is unipotent in } Z_G(t\sigma_s)^0} \bigcup_{g \in G \ u \text{ is unipotent in } Z_G(t\sigma_s)^0} g(t\sigma u) g^{-1}$$

for some $t \in (T^{\sigma})^0$. It is known that $Z_G(t\sigma_s)^0$ is reductive and the set of unipotent elements in a reductive group is an irreducible variety. Thus

(a) each Steinberg fiber is irreducible.

Moreover, there are only finitely many unipotent conjugacy classes in a reductive group [L1]. Therefore

(b) each Steinberg fiber contains finitely many G-conjugacy classes.

Lemma 1.7. Let **a** be a Steinberg fiber in D and $J \subset I$ with $\sigma(J) = J$. Then there exists finitely many L_J -conjugacy classes c_1, \dots, c_m in $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$ such that $\{g; g \in N_{\tilde{G}}(P_J) \cap D, St(g) = \mathbf{a}\} = \sqcup_i c_i U_{P_J}$.

Let $l \in N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J)$ and $u \in U_{P_J}$. Then it is easy to see that l is contained in the closure of $\{tlut^{-1}; t \in (Z(L_J)^{\sigma})^0\}$. Hence St(l) = St(lu). In other words,

$$\{g; g \in N_{\tilde{G}}(P_J) \cap D, St(g) = \mathbf{a}\}$$

= $\{lu; l \in N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D, u \in U_{P_J}, St(l) = \mathbf{a}\}.$

By [L2, Proposition 1.14], any quasi-semisimple element in D (resp. $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$) is G-conjugate (resp. L_J -conjugate) to $T_1\sigma$, where $T_1 = (T^{\sigma})^0$. Notice that $\{t \in T_1; St(t\sigma) = \mathbf{a}\}$ is a finite

set. Hence there are only finitely many quasi-semisimple L_J -conjugacy classes in $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$ that are contained in $St^{-1}(\mathbf{a})$. One can see that a L_J -conjugacy class c is contained in $St^{-1}(\mathbf{a})$ if and only if the unique L_J -conjugacy class c' that is contained in c is also contained in $St^{-1}(\mathbf{a})$. Then the lemma follows from Lemma 1.7 (b).

1.8. For $J \subset I$, set

$$Z_J = \{ (P, P', gU_P); P, P' \in \mathfrak{P}_J, g \in G, P' = {}^gP \}, Z'_J = \{ (P, P', gH_P); P, P' \in \mathfrak{P}_J, g \in G, P' = {}^gP \}$$

with the $G \times G$ -action defined by

$$(g_1, g_2) \cdot (P, P', gU_P) = ({}^{g_2}P, {}^{g_1}P', g_1gU_Pg_2^{-1}), (g_1, g_2) \cdot (P, P', gH_P) = ({}^{g_2}P, {}^{g_1}P', g_1gH_Pg_2^{-1}).$$

Set $h_J = (P_J, P_J, U_{P_J}) \in Z_J$ and $h'_J = (P_J, P_J, H_{P_J}) \in Z'_J$. By [L3, section 3], [H1, section 1] and the remark of [H2, Corollary 5.4], we have partitions

(a)
$$Z_J = \sqcup_{w \in {}^J W} Z_{J;w}$$
 and $Z'_J = \sqcup_{w \in {}^J W} Z'_{J;w}$,

where $Z_{J;w} = G_{\Delta} \cdot (BwB, B)h_J$ and $Z'_{J;w} = G_{\Delta} \cdot (BwB, B)h'_J$. The subvarieties $Z_{J;w}$ (resp. $Z'_{J;w}$) are called *G*-stable pieces of Z_J (resp. Z'_J).

Fix $w \in {}^{J}W$. Let K = I(J, id; w). Then by [L3, 3.14],

(b) there is a canonical bijection between the G_{Δ} -orbits on $Z_{J,w}$ and the L_K -conjugacy classes on wL_K .

By [L3, section 3], we have G-equivariant morphisms $pr : Z_{J;w} \to G/P_K$ and $pr' : Z'_{J;w} \to G/P_K$, where G acts diagonally on $Z_{J;w}$ and $Z'_{J;w}$ and acts in the natural way on G/P_K . Moreover, by [H1, Proposition 1.10],

(c)
$$pr(z) = P_K$$
 if and only if $z = (pw, 1) \cdot h_J$ for some $p \in P_K$,
 $pr'(z) = P_K$ if and only if $z = (pw, 1) \cdot h'_J$ for some $p \in P_K$.

Also we have that the closure of any G-stable piece is a union of G-stable pieces.

(d) $\overline{Z_{J;w}} = \bigsqcup_{w' \in J, w' \leq J, idw} Z_{J,w'}$. See [H1, Proposition 4.6] and [H2, Proposition 5.8].

1.9. If $\sigma(J) = J$, then the action of $G \times G$ on Z_J and Z'_J can be extended in a natural way to an action of $\tilde{G} \times \tilde{G}$.

Now set

$$\Lambda_{J,D} = \{ (z,g) \in Z_J \times D; (g,g) \cdot z = z \}, \Lambda'_{J,D} = \{ (z,g) \in Z'_J \times D; (g,g) \cdot z = z \}.$$

For $w \in W^J$, set $\Lambda_{J,D;w} = \{(z,g) \in \Lambda_{J,D}; z \in Z_{J;w}\}$ and $\Lambda'_{J,D;w} = \{(z,g) \in \Lambda'_{J,D}; z \in Z'_{J;w}\}.$

1.10. Set $P = N_{\tilde{G}}P$ for any parabolic subgroup P of G. Define the action of P_J on $G \times \tilde{P}_J$ by $p \cdot (g, p') = (gp^{-1}, pp'p^{-1})$. Let $G \times_{P_J} \tilde{P}_J$ be the quotient space. Then we may identify $G \times_{P_J} \tilde{P}_J$ with $\{(P,g); P \in \mathcal{P}_J, g \in \tilde{P}\}$ via $(g, p) \mapsto ({}^gP_J, gpg^{-1})$.

Let c be a subvariety of $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$ that is stable under the conjugation action of L_J . Then cU_{P_J} and cH_{P_J} are stable under the conjugation action of P_J . So we may define $G \times_{P_J} cU_{P_J} \subset G \times_{P_J} cH_{P_J} \subset$ $G \times_{P_J} \tilde{P}_J$.

Now set

$$\Lambda_{J,c} = \{ (P, P', gU_P, h) \in \Lambda_{J,D}; (P, h), (P', h) \in G \times_{P_J} cU_{P_J} \}, \\ \Lambda'_{J,c} = \{ (P, P', gH_P, h) \in \Lambda'_{J,D}; (P, h), (P', h) \in G \times_{P_J} cH_{P_J} \}.$$

For $w \in W^J$, set $\Lambda_{J,c;w} = \Lambda_{J,D;w} \cap \Lambda_{J,c}$ and $\Lambda'_{J,c;w} = \Lambda'_{J,D;w} \cap \Lambda'_{J,c}$.

Lemma 1.11. Let $w \in {}^{J}W$ and K = I(J, id; w). Let c be a L_{K} conjugacy class in $N_{\tilde{G}}(P_K) \cap N_{\tilde{G}}(L_K) \cap D$. Set $X_{J,c;w} = \{(z,g) \in \Lambda_{J,c;w}; pr(z) = P_K\}$ and $X'_{J,c;w} = \{(z,g) \in \Lambda'_{J,c;w}; pr'(z) = P_K\}$. Then
(1) If $X_{J,c;w} \neq \emptyset$, then $X_{J,c;w}$ is of pure dimension dim (P_K) .
(2) If $X'_{J,c;w} \neq \emptyset$, then $X'_{J,c;w}$ is of pure dimension

$$\dim(P_K) - \dim(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J) \cap \operatorname{Ad}(w)Z^0(L_J)})^{\sigma}.$$

Remark. (1) If $X_{J,c;w} \neq \emptyset$ or $X'_{J,c;w} \neq \emptyset$, then as we will see in the proof, $\sigma \operatorname{Ad}(w)(z) = \operatorname{Ad}(w)\sigma(z)$ for $z \in Z^0(L_J)$. So we have that $\sigma Z^0(L_J) = Z^0(L_J)$ and $\sigma \operatorname{Ad}(w)Z^0(L_J) = \operatorname{Ad}(w)Z^0(L_J)$. Therefore $(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J)\cap\operatorname{Ad}(w)Z^0(L_J)})^{\sigma}$ is defined.

(2) If w(J) = J, then $\operatorname{Ad}(w)Z^0(L_J) = Z^0(L_J)$ and

$$\dim(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J)\cap\operatorname{Ad}(w)Z^0(L_J)})^{\sigma} = 0$$

If $w(J) \neq J$, then there exists $j \in J$ such that w(j) is not spanned by the simple roots in J. Set $z_t = \prod_{j \notin J} \omega_j(t)$ for $t \in \mathbf{k}^*$. Then $z_t \notin \operatorname{Ad}(w)^{-1}Z^0(L_J)$ for all $t \in \mathbf{k}^* - \{1\}$. Moreover, $\sigma \operatorname{Ad}(w)(z_t) =$ $\operatorname{Ad}(w)\sigma(z_t) = \operatorname{Ad}(w)(z_t)$. Hence $\operatorname{Ad}(w)z_t \in (\operatorname{Ad}(w)Z^0(L_J))^{\sigma}$. So $\dim(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J)\cap\operatorname{Ad}(w)Z^0(L_J)})^{\sigma} = 0$ if and only if w(J) = J.

We only prove part (1) here. Part (2) can be proved in a similar way. Let $\bar{p}: P_K \to L_K$ be the projection map. By (c), $pr(z) = P_K$ if and only if $z = (pw, 1) \cdot h_J$ for some $p \in P_K$. Moreover, the morphism $f: \{z \in Z_{J;w}; pr(z) = P_K\} \to L_K$ defined by $(pw, 1) \cdot h_{J,D} \mapsto \bar{p}(p)$ is well-defined. Now consider the morphism $X_{J,c;w} \to L_K \times c$ which sends (z,g) to $(f(z), \bar{p}(g))$. We see that the image is contained in $\{(l, l') \in L_K \times c; lwl' = l'lw\}$, which is of pure dimension dim (L_K) .

Let $l \in L_K$ and $l' \in c$ with lwl' = l'lw. Then the fiber Y over (l, l') is $\{(z, ul'); z \in (U_{P_K} lw, 1) \cdot h_J, u \in U_{P_K}, (ul', ul') \cdot z = z\}$. Define the action

of U_{P_K} on Y by $u_1 \cdot (z, ul') = ((u_1, u_1) \cdot z, u_1 ul' u_1^{-1})$. Then the projection map $Y \to (U_{P_K} lw, 1) \cdot h_J = (U_{P_K})_{\Delta}(lw, 1) \cdot h_J$ is U_{P_K} -equivariant for the diagonal U_{P_K} -action on $(U_{P_K} lw, 1) \cdot h_J$. Since U_{P_K} acts transitively on $(U_{P_K} lw, 1) \cdot h_J$, the projection map is a locally trivially fibration with fibers isomorphic to $\{u \in U_{P_K}; (ulw, u) \cdot h_J = (lw, 1) \cdot h_J\}$. In particular, Y is of pure dimension $\dim(U_{P_K})$. Therefore $X_{J,c;w}$ is of pure dimension $\dim(L_K) + \dim(U_{P_K}) = \dim(P_K)$. \Box

Proposition 1.12. Let **a** be a Steinberg fiber of $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$. (1) If $\Lambda_{J,\mathbf{a};w} \neq \emptyset$, then $\Lambda_{J,\mathbf{a};w}$ is of pure dimension dim(G). (2) If $\Lambda'_{J,\mathbf{a};w} \neq \emptyset$, then $\Lambda'_{J,\mathbf{a};w}$ is of pure dimension

$$\dim(G) - \dim(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J) \cap \operatorname{Ad}(w)Z^0(L_J)})^{\sigma}.$$

We only prove part (1) here. Part (2) can be proved in a similar way. Let K = I(J, id; w). For $(z, g) \in X_{J,\mathbf{a};w}$, we have that ${}^{g}P_{K} = g \cdot pr(z) = pr((g, g) \cdot z) = pr(z) = P_{K}$. Hence $g \in P_{K}$. By lemma 1.7, $X_{J,\mathbf{a};w} = \bigsqcup_{i} X_{J,c_{i};w}$ for finitely many L_{K} -conjugacy classes c_{i} . By lemma 1.11, $X_{J,\mathbf{a};w}$ is of pure dimension dim (P_{K}) .

Let $\pi : \Lambda_{J,a;w} \to Z_{J;w}$ be the projection map. It is easy to see that π is *G*-equivariant for the diagonal *G*-action. Thus $pr \circ \pi : \Lambda_{J,\mathbf{a};w} \to G/P_K$ is also *G*-equivariant. Since *G* acts transitively on G/P_K , $pr \circ \pi$ is a locally trivial fibration with fibers isomorphism to $X_{J,\mathbf{a};w}$. Thus $\Lambda_{J,w;\mathbf{a}}$ is of pure dimension dim(*G*).

Lemma 1.13. Let c be a L_J -conjugacy class in $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D$. Set

$$\mathcal{Z}_{J,c} = \{ (P, P', g); (P, g), (P', g) \in G \times_{P_J} cU_{P_J} \}, \\ \mathcal{Z}'_{J,c} = \{ (P, P', g); (P, g), (P', g) \in G \times_{P_J} cH_{P_J} \}.$$

(1) Define the map $\Lambda_{J,c} \to \mathcal{Z}_{J,c}$ by $(P, P', kU_P, g) \mapsto (P, P', g)$. If $\mathcal{Z}_{J,c} \neq \emptyset$, then the map is surjective and each fiber is of pure dimension $\dim(L_J) - \dim(c)$.

(2) Define the map $\Lambda'_{J,c} \to \mathcal{Z}'_{J,c}$ by $(P, P', kH_P, g) \mapsto (P, P', g)$. If $\mathcal{Z}'_{J,c} \neq \emptyset$, then the map is surjective and each fiber is of pure dimension $\dim(L_J) - \dim(c) - \dim(Z(L_J)^{\sigma})$.

We only prove part (1) here. Part (2) can be proved in the same way.

Let $(P, P', g) \in \mathcal{Z}_{J,c}$. Then there exists $k \in G$ such that $P' = {}^{k}P$. By definition, $kgk^{-1}U_{P'}$ and $gU_{P'}$ are P'-conjugate. Therefore, there exists $l \in P'$, such that $lkgk^{-1}l^{-1} \in gU_{P'}$. In other words, $(g,g) \cdot (P,P', lkU_P) = (P,P', lkU_P)$. So the map is surjective.

Assume that $(P, P', kU_P, g), (P, P', k'U_P, g) \in \Lambda_{J,c}$. Then $k^{-1}k' \in P$ and $(k^{-1}k')gU_P(k^{-1}k')^{-1} = gU_P$. Thus the fibers of the map $\Lambda_{J,c} \rightarrow \mathcal{Z}_{J,c}$ are isomorphic to $\{(lU_P; l \in P; lgU_P l^{-1} = gU_P\}$. Notice that $(P,g) \in G \times_{P_J} cU_{P_J}$. is of dimension $\dim(L_J) - \dim(c)$. \Box

XUHUA HE

Now combining Proposition 1.12 and Lemma 1.13, we have the following result.

Corollary 1.14. Let c be a L_J -conjugacy class in $N_{\tilde{G}}(P_J) \cap N_{\tilde{G}}(L_J) \cap D'$. Then

(1) $\dim(\mathcal{Z}_{J,c}) \leq \dim(G) - \dim(L_J) + \dim(c).$

(2) $\dim(\mathcal{Z}'_{J,c}) \leq \dim(G) - \dim(L_J) + \dim(c) + \dim(Z(L_J)^{\sigma})$. More precisely, define $\mathcal{Z}'_{J,c;w} = \{(P, P', g) \in \mathcal{Z}'_{J,c}; (P, P') \in G_{\Delta} \cdot (P_J, {}^wP_{J'})\}$ for $w \in {}^JW^{J'}$. Then

$$\dim(\mathcal{Z}'_{J,c;w}) \leq \dim(G) - \dim(L_J) + \dim(c) + \dim(Z(L_J)^{\sigma}) - \dim(\frac{\operatorname{Ad}(w)Z^0(L_J)}{Z^0(L_J) \cap \operatorname{Ad}(w)Z^0(L_J)})^{\sigma}.$$

Remark. Part (1) was first proved in [L2, Proposition 4.2 (d)]. By the remark of Lemma 1.11, part (2) is a stronger version of [L2, Proposition 4.2(c)].

References

- [Bo] A. Borel, Groupes linéaires algébriques, Annals of Math. (2) 64 (1956), 20–82.
- [H1] X. He, The G-stable pieces of the wonderful compactification, Trans. Amer. Math. Soc. 359 (2007), 3005-3024.
- [H2] X. He, Minimal length elements in some double cosets of Coxeter groups, Adv. Math. (2007), 215 (2007), 469-503.
- [HL] X. He and G. Lusztig, Singular supports for character sheaves on a group compactification, GAFA, to appear, math.RT/0511050.
- [L1] G. Lusztig, On the finiteness of the number of unipotent classes, Invent. Math. 34 (1976), 201–213.
- [L2] G. Lusztig, Character sheaves on disconnected groups, I, Represent. Theory, 7 (2003), 374–403.
- [L3] G. Lusztig, Parabolic character sheaves, I, II, Mosc. Math. J. 4 (2004), no. 1, 153–179; no. 4, 869–896.
- [Sp] N. Spaltenstein, Classes unipotentes et sous-groupes de Borel, Lecture Notes in Math., 946, Springer, Berlin, 1982.
- [St] R. Steinberg, Endomorphisms of linear algebraic groups, Memoirs of Amer. Math. Soc., vol. 80, 1968.