A crystal theoretic method for finding rigged configurations from paths

Reiho Sakamoto

Department of Physics, Graduate School of Science, University of Tokyo, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan reiho@spin.phys.s.u-tokyo.ac.jp

Abstract

The Kerov–Kirillov–Reshetikhin (KKR) bijection gives one to one correspondences between the set of highest paths and the set of rigged configurations. In this paper, we give a crystal theoretic reformulation of the KKR map from the paths to rigged configurations, using the combinatorial R and energy functions. This formalism provides tool for analysis of the periodic box-ball systems.

1 Introduction

The Kerov-Kirillov-Reshetikhin (KKR) bijection [1, 2, 3] gives the combinatorial one to one correspondences between the set of highest weight elements of tensor products of crystals [4, 5] (which we call highest paths) and the set of combinatorial objects called the rigged configurations. This bijection was originally introduced as essential tool to derive new expression (called fermionic formulae) of the celebrated Kostka–Foulkas polynomials [6]. Background of this expression is the Bethe ansatz for the Heisenberg spin chain [7] and, in this context, the rigged configurations form index set of the eigenvalues and eigenvectors of the Hamiltonian. To date the fermionic formulae have been extended to wider class of representations, and proved in several cases (see, e.g., [8, 9, 10, 11, 12] for the current status of the study).

Recently, the KKR bijection itself becomes active subject of the study. Fundamental observation [13] is that the KKR bijection is inverse scattering transform of the box-ball systems, which is the prototypical example of the ultradiscrete soliton systems introduced by Takahashi–Satsuma [14, 15]. In this context, the rigged configurations are regarded as action and angle variables for the box-ball systems. This observation leads to derivation [16, 17] of general solutions for the box-ball systems for the first time.

Therefore it is natural to ask what the representation theoretic origin of the KKR bijection is. Partial answer was given in the previous paper [16], and it is substantially used in the derivation in [17]. However, the formalism in [16] works only for the map from the rigged configuration to paths, and extension of the formalism to the inverse direction seems to have essential obstructions. Up to now, crystal interpretation for the map from the paths to rigged configurations (which we denote by ϕ) remains open. Closely related problem is what the representation theoretic origin of the mysterious combinatorial algorithm of the original definition of ϕ is. In fact, the formalism in [16] gives alternative representation theoretic map for ϕ^{-1} while it does not give meanings to the combinatorial procedures like vacancy numbers or singular rows. We remark that in Section 2.7 of [13], there is decomposition of the \mathfrak{sl}_n type ϕ into successive computation of \mathfrak{sl}_2 type ϕ . However it finally uses combinatorial version of ϕ , hence it is not a complete crystal interpretation of ϕ .

One of the main aims of the present article is to give a crystal interpretation for the \mathfrak{sl}_2 type ϕ by clarifying the representation theoretic origin of the original combinatorial procedure of ϕ (see Theorem 3.3). In our formalism, the combinatorial procedure of ϕ is identified with differences of energy functions called local energy distribution and indeed we can read off all information about the rigged configurations from them. In terms of the box-ball systems, the local energy distributions clarify which letters of given path correspond to which soliton even if they are in multiply scattering state.

Another aim of the paper is to provide a tool for analysis of the periodic box-ball systems [18, 19]. In our \mathfrak{sl}_2 case formalism, there is remarkably nice property (see Proposition 4.4). Namely, the solitons appeared in the local energy distribution are always separated from each other. This leads to alternative version of our procedure as given in Theorem 3.6. Importance of this reformulation is that when we apply the formalism to find the action and angle variables of the periodic box-ball systems [20], we do not need to cut paths and treat them as non-periodic paths. This improves the inverse scattering formalism of [20] and theta function formulae of [21, 22] in a sense that we treat paths genuinely as periodic. As a byproduct, we give intuitive picture of the basic operators which are the key to define angle variables in [20] (see Remark 3.7). We remark that there is another approach to the initial value problem of the periodic box-ball systems [23]. Although their combinatorial method and our representation theoretic method are largely different, it will be important to clarify the relationship between these two approaches.

The present paper is organized as follows. In Section 2, we review the combinatorial R and energy functions following [24]. In Section 3, we formulate our main results (Theorem 3.3 and Theorem 3.6). In Section 4, we give proof of these theorems. Section 5 is summary. In Appendix A, we recall the KKR bijection and collect necessary facts, and in Appendix B, we collect necessary facts about the time evolution operators T_l .

2 Combinatorial *R* and energy functions

In this section, we introduce necessary facts from the crystal bases theory, namely, the combinatorial R and energy functions. Let B_k be the crystal of k-fold symmetric powers of the vector (or natural) representation of $U_q(\mathfrak{sl}_2)$. As the set, it is

$$B_k = \{ (x_1, x_2) \in \mathbb{Z}_{>0}^2 \, | \, x_1 + x_2 = k \}.$$
(1)

We usually identify elements of B_k as the semi-standard Young tableaux

$$(x_1, x_2) = \underbrace{\left[\underbrace{x_1 \ x_2}_{1 \dots 1 \ 2 \dots 2} \right]}_{(2)},$$
 (2)

i.e., the number of letters i contained in a tableau is x_i .

For two crystals B_k and B_l of $U_q(\mathfrak{sl}_2)$, one can define the tensor product $B_k \otimes B_l = \{b \otimes b' \mid b \in B_k, b' \in B_l\}$. Then we have a unique isomorphism $R : B_k \otimes B_l \xrightarrow{\sim} B_l \otimes B_k$, i.e. a unique map which commutes with actions of the Kashiwara operators \tilde{e}_i , \tilde{f}_i . We call this map combinatorial R and usually write the map R simply by \simeq .

In calculation of the combinatorial R, it is convenient to use the diagrammatic technique due to Nakayashiki–Yamada (Rule 3.11 of [24]). Consider the two elements $x = (x_1, x_2) \in B_k$ and $y = (y_1, y_2) \in B_l$. Then we draw the following diagram to express the tensor product $x \otimes y$.



The combinatorial R matrix and energy function H for $x \otimes y \in B_k \otimes B_l$ (with $k \geq l$) are calculated by the following rule.

- 1. Pick any dot, say \bullet_a , in the right column and connect it with a dot \bullet'_a in the left column by a line. The partner \bullet'_a is chosen from the dots whose positions are higher than that of \bullet_a . If there is no such a dot, we return to the bottom, and the partner \bullet'_a is chosen from the dots in the lower row. In the former case, we call such a pair "unwinding," and, in the latter case, we call it "winding."
- 2. Repeat procedure (1) for the remaining unconnected dots (l-1) times.

- 3. Action of the combinatorial R matrix is obtained by moving all unpaired dots in the left column to the right horizontally. We do not touch the paired dots during this move.
- 4. The energy function H is given by the number of unwinding pairs.

The number of winding (or unwinding) pairs is sometimes called the winding (or unwinding, respectively) number of tensor product. It is known that the resulting combinatorial R matrix and the energy functions are not affected by the order of making pairs ([24], Propositions 3.15 and 3.17). In the above description, we only consider the case $k \ge l$. The other case $k \le l$ can be done by reversing the above procedure, noticing the fact $R^2 = id$. For more properties, including that the above definition indeed satisfies the axiom, see [24].

Example 2.1 Corresponding to the tensor product $1122 \otimes 122$, we draw the diagram given in the left hand side of:



By moving one unpaired dot to the right, we obtain

$$\boxed{1122} \otimes \boxed{122} \simeq \boxed{112} \otimes \boxed{1222}. \tag{3}$$

Since we have two unwinding pair, the energy function is $H(1122 \otimes 122) = 2$.

Consider the affinization of the crystal B. As the set, it is

$$\operatorname{Aff}(B) = \{ b[d] \mid b \in B, \, d \in \mathbb{Z} \}.$$

$$\tag{4}$$

Integers d of b[d] are called modes. For the tensor product $b_1[d_1] \otimes b_2[d_2] \in \text{Aff}(B_k) \otimes \text{Aff}(B_l)$, we can lift the above definition of the combinatorial R as follows:

$$b_1[d_1] \otimes b_2[d_2] \stackrel{R}{\simeq} b'_2[d_2 - H(b_1 \otimes b_2)] \otimes b'_1[d_1 + H(b_1 \otimes b_2)], \tag{5}$$

where $b_1 \otimes b_2 \simeq b'_2 \otimes b'_1$ is the combinatorial R defined in the above.

Remark 2.2 Piecewise linear formula to obtain the combinatorial R and the energy function is also available [25]. This is suitable for computer implementation. For the affine combinatorial $R: x[d] \otimes y[e] \simeq \tilde{y}[e - H(x \otimes y)] \otimes \tilde{x}[d + H(x \otimes y)]$, we have

$$\tilde{x}_{i} = x_{i} + Q_{i}(x, y) - Q_{i-1}(x, y), \qquad \tilde{y}_{i} = y_{i} + Q_{i-1}(x, y) - Q_{i}(x, y),
H(x \otimes y) = Q_{0}(x, y),
Q_{i}(x, y) = \min(x_{i+1}, y_{i}),$$
(6)

where we have expressed $x = (x_1, x_2)$, $y = (y_1, y_2)$, $\tilde{x} = (\tilde{x}_1, \tilde{x}_2)$ and $\tilde{y} = (\tilde{y}_1, \tilde{y}_2)$. All indices *i* should be considered as $i \in \mathbb{Z}/2\mathbb{Z}$.

3 Local energy distribution and the KKR bijection

In this section, we reformulate the combinatorial procedure ϕ in terms of the energy functions of crystal base theory. See Appendix A for explanation of ϕ . In order to do this, it is convenient to express actions of the combinatorial R by vertex type diagrams. First, we express the isomorphism of the combinatorial R matrix

$$a \otimes b_1 \simeq b_1' \otimes a' \tag{7}$$

and the corresponding value of the energy function $e_1 := H(a \otimes b_1)$ by the following vertex diagram:



If we apply combinatorial R successively as

$$a \otimes b_1 \otimes b_2 \simeq b_1' \otimes a' \otimes b_2 \simeq b_1' \otimes b_2' \otimes a'', \tag{8}$$

with the energy function $e_2 := H(a' \otimes b_2)$, then we express this by joining two vertices as follows:



Definition 3.1 For a given path $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L$, we define local energy $E_{l,j}$ by $E_{l,j} := H(u_l^{(j-1)} \otimes b_j)$. Here, in the diagrammatic expression, $u_l^{(j-1)}$ are defined as follows (see also (29) with convention $u_l^{(0)} := u_l$).

Here, we denote $T_l(b) = b'_1 \otimes b'_2 \otimes \cdots \otimes b'_L$. We define $E_{0,j} = 0$ for all $1 \leq j \leq L$. We also use the following notation:

$$E_l := \sum_{j=1}^{L} E_{l,j}.$$
 (9)

In other words, $u_l[0] \otimes b \stackrel{R}{\simeq} T_l(b) \otimes u_l^{(L)}[E_l]$, where we have omitted modes for b and $T_l(b)$.

For a given path $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L$ $(b_i \in B_{\lambda_i})$, we create a path $b' = b \otimes \boxed{1}^{\otimes \Lambda}$, where $\Lambda > \lambda_1 + \cdots + \lambda_L$. Then we always have $u_l^{(L+\Lambda)} = u_l$ for arbitrary l (Proposition B.1 (1)). In such a circumstance, it is known that the sum E_l is conserved quantities of the box-ball system; $E_l(T_k(b')) = E_l(b')$. The proof is based on successive application of the Yang-Baxter equation (see Theorem 3.2 of [26] and section 3.4 of [25]). However, for our purpose, we need more detailed information such as $E_{l,j}$.

Lemma 3.2 For a given path $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L$, we have $E_{l,j} - E_{l-1,j} = 0$ or 1, for all l > 0 and for all $1 \le j \le L$.

Proof. First we give a proof when l = 1, i.e., we show $E_{1,j} - E_{0,j} = 0$ or 1. In this case, we have $u_l, u_l^{(i)} \in B_1$ and $E_{0,j} = 0$. Since $H(x \otimes y) = 0$ or 1 for all $x \in B_1$ and all $y \in B_k$, the proof follows.

Now we consider possible values for $E_{l,j} - E_{l-1,j}$. In order to do this, we show that the difference between tableaux representations of $u_l^{(j)}$ and $u_{l-1}^{(j)}$ is only one letter. More precisely, we show that if $u_{l-1}^{(j)} = (x_1, x_2)$, then $u_l^{(j)} = (x_1 + 1, x_2)$ or $u_l^{(j)} = (x_1, x_2 + 1)$. We show this claim by induction on j. For j = 0 case, it is true because $u_{l-1}^{(0)} = u_{l-1} = (l-1, 0)$ and $u_l^{(0)} = u_l = (l, 0)$, by definition. Suppose that the above claim holds for all j < kfor some k. In order to compare $u_{l-1}^{(k)}$ and $u_l^{(k)}$, consider the isomorphism $u_{l-1}^{(k-1)} \otimes b_k \simeq$ $b'_{l-1,k} \otimes u_{l-1}^{(k)}$ and $u_l^{(k-1)} \otimes b_k \simeq b'_{l,k} \otimes u_l^{(k)}$. By assumption, the difference between $u_{l-1}^{(k-1)}$ and $u_l^{(k-1)}$ is one letter. Recall that in calculating the combinatorial R, order of making pairs can be chosen arbitrary. Therefore, in $u_l^{(k-1)} \otimes b_k$, first we can make all pairs that appear in $u_{l-1}^{(k-1)} \otimes b_k$, and next we make remaining one pair. This means the difference of number of unwinding pairs, i.e., $E_{l,k} - E_{l-1,k}$ is 0 or 1. To make the induction proceeds, note that this fact means the difference between $u_{l-1}^{(k)}$ and $u_l^{(k)}$ is also one letter.

The following theorem gives crystal theoretic reformulation of the KKR map ϕ . See Appendix A for explanation of the unrestricted rigged configurations.

Theorem 3.3 Let $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots \otimes B_{\lambda_L}$ be an arbitrary path. b can be highest weight or non-highest weight. Set $N = E_1(b)$. We determine the pair of numbers $(\mu_1, r_1), (\mu_2, r_2), \cdots, (\mu_N, r_N)$ by the following procedure from Step 1 to Step 4. Then the resulting $(\lambda, (\mu, r))$ coincides with the (unrestricted) rigged configuration $\phi(b)$.

- 1. Draw a table containing $(E_{l,j} E_{l-1,j} = 0, 1)$ at the position (l, j), i.e., at the l th row and the j th column. We call this table local energy distribution.
- 2. Starting from the rightmost 1 in the l = 1 st row, pick one 1 from each successive row. The one in the (l + 1) th row must be weakly right of the one selected in the lth row. If there is no such 1 in the (l + 1) th row, the position of the lastly picked 1 is called (μ_1, j_1) . Change all selected 1 into 0.
- 3. Repeat Step 2 for (N-1) times to further determine $(\mu_2, j_2), \dots, (\mu_N, j_N)$ thereby making all 1 into 0.
- 4. Determine r_1, \cdots, r_N by

$$r_k = \sum_{i=1}^{j_k - 1} \min(\mu_k, \lambda_i) + E_{\mu_k, j_k} - 2\sum_{i=1}^{j_k} E_{\mu_k, i}.$$
 (10)

Proof of Theorem 3.3 will be given in the next section. As we will see in Proposition 4.4, the groups obtained in the above theorem have no crossing with each other. Therefore, when we search 1 of the (l+1) th row in the above Step 2, we have at most one candidate, i.e., we can uniquely determine such 1.

Example 3.4 For example of Theorem 3.3, we consider the following path

$$b = \boxed{1111} \otimes \boxed{11} \otimes \boxed{2} \otimes \boxed{1122} \otimes \boxed{1222} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{22} \tag{11}$$

Corresponding to Step 1, the local energy distribution is given by the following table (j stands for column coordinate of the table).

	1111	11	2	1122	1222	1	2	22
$E_{1,j} - E_{0,j}$	0	0	1	0	1	0	1	0
$E_{2,j} - E_{1,j}$	0	0	0	1	0	0	0	1
$E_{3,j} - E_{2,j}$	0	0	0	1	0	0	0	0
$E_{4,j} - E_{3,j}$	0	0	0	0	1	0	0	0
$E_{5,j} - E_{4,j}$	0	0	0	0	1	0	0	0
$E_{6,j} - E_{5,j}$	0	0	0	0	0	0	0	1
$E_{7,j} - E_{6,j}$	0	0	0	0	0	0	0	0

Following Step 2 and Step 3, letters 1 contained in the above table are found to be classified into 3 groups, as indicated in the following table.

	1111	11	2	1122	1222	1	2	22
$E_{1,j} - E_{0,j}$			3		2^{*}		1	
$E_{2,j} - E_{1,j}$				3				1^*
$E_{3,j} - E_{2,j}$				3				
$E_{4,j} - E_{3,j}$					3			
$E_{5,j} - E_{4,j}$					3			
$E_{6,j} - E_{5,j}$								3^*
$E_{7,j} - E_{6,j}$								

From the above table, we see that the cardinalities of groups 1, 2 and 3 are 2, 1 and 6, respectively. Also, in the above table, positions of (μ_1, j_1) , (μ_2, j_2) and (μ_3, j_3) are indicated by asterisks. Their explicit locations are $(\mu_1, j_1) = (2, 8)$, $(\mu_2, j_2) = (1, 5)$ and $(\mu_3, j_3) = (6, 8)$.

Now we evaluate riggings r_i according to equation (10).

$$r_{1} = \sum_{i=1}^{8-1} \min(2, \lambda_{i}) + E_{2,8} - 2 \sum_{i=1}^{8} E_{2,i}$$

$$= (2+2+1+2+2+1+1) + 1 - 2(0+0+1+1+1+0+1+1)$$

$$= 2,$$

$$r_{2} = \sum_{i=1}^{5-1} \min(1, \lambda_{i}) + E_{1,5} - 2 \sum_{i=1}^{5} E_{1,i}$$

$$= (1+1+1+1) + 1 - 2(0+0+1+0+1)$$

$$= 1,$$

$$r_{3} = \sum_{i=1}^{8-1} \min(6, \lambda_{i}) + E_{6,8} - 2 \sum_{i=1}^{8} E_{6,i}$$

$$= (4+2+1+4+4+1+1) + 2 - 2(0+0+1+2+3+0+1+2)$$

$$= 1.$$

Therefore we obtain $(\mu_1, r_1) = (2, 2)$, $(\mu_2, r_2) = (1, 1)$ and $(\mu_3, r_3) = (6, 1)$, which coincide with the following computation according to the original definition of ϕ . In the following, we use Young diagrammatic expression for the rigged configurations, and we put riggings and vacancy numbers on the right and on the left of the corresponding rows of the configuration, respectively.



In the above diagrams, newly added boxes are indicated by circles " \circ ". The reader should compare the local energy distribution and the above box adding procedure. Then one will observe that the local energy distribution and box addition on μ part have close relationships. This relation will be established in Lemma 4.2. In other words, the original combinatorial procedure for ϕ is embedded into rather automatic applications of the combinatorial R and energy functions.

Example 3.5 By using Theorem 3.3, we can easily grasp the large scale structure of combinatorial procedures of the KKR bijection from calculations of the combinatorial R and energy functions. In order to show the typical example, consider the following long path (length 40).



Then, the local energy distribution takes the following form.



In the above table, letters 1 in the local energy distribution are represented by "•", and letters 0 are suppressed. By doing Step 2 and Step 3, we obtain classifications of letters 1. In the above table, letters 1 belonging to the same group are joined by thick lines. We see there are 15 groups whose cardinalities are 3, 7, 1, 4, 1, 22, 1, 6, 2, 3, 4, 2, 6, 2, 2 from left to right, respectively.

By using equation (10), we obtain the unrestricted rigged configuration as follows: $(\mu_1, r_1) = (2, 13), (\mu_2, r_2) = (2, 13), (\mu_3, r_3) = (6, 15), (\mu_4, r_4) = (2, 12), (\mu_5, r_5) = (4, 14),$ $(\mu_6, r_6) = (3, 12), (\mu_7, r_7) = (2, 11), (\mu_8, r_8) = (6, 5), (\mu_9, r_9) = (1, 3), (\mu_{10}, r_{10}) =$ $(22, -17), (\mu_{11}, r_{11}) = (1, 1), (\mu_{12}, r_{12}) = (4, 1), (\mu_{13}, r_{13}) = (1, 1), (\mu_{14}, r_{14}) = (7, -3)$ and $(\mu_{15}, r_{15}) = (3, -2)$. The vacancy numbers for each row is $p_{22} = -15, p_7 = 15,$ $p_6 = 19, p_4 = 21, p_3 = 18, p_2 = 14$ and $p_1 = 10$. Note that since the path in this example is not highest weight, the resulting unrestricted rigged configuration has negative values of the riggings and vacancy numbers.

We have alternative form of Theorem 3.3.

Theorem 3.6 In the above Theorem 3.3, Step 2 can be replaced by the following procedure (Step 2'). The resulting groups are the same as those obtained in Theorem 3.3 up to reordering in subscripts.

2'. Pick one of the lowest 1 of the local energy distribution arbitrary, and denote it by (μ_1, j_1) . Starting from (μ_1, j_1) , choose one 1 from each row successively as follows. Assume that we have chosen 1 at (l, k_l) . Then $(l-1, k_{l-1})$ is the rightmost 1 among the part of row (l-1, 1), (l-1, 2), \cdots , $(l-1, k_l)$. Change all selected 1 into 0. \Box

Note that comparing both Theorem 3.3 and Theorem 3.6, the resulting $(\lambda, (\mu, r))$ can be different in ordering of μ . However, this difference has no role in the KKR theory. By the same reason, an ambiguity in choosing the lowest 1 in Step 2' brings no important difference. Proof of Theorem 3.6 will be given in the next section.

The formalism in Theorem 3.6 is suitable for analysis of the periodic box-ball system. In particular, consider the case when there are more than one longest group in the local energy distribution. Choose any successive longest groups and apply the above Step 2' to these two groups. Then due to non-crossing property of groups (Proposition 4.4), we can concentrate on the region between these two groups and determine all groups between them ignoring other part of the path.

Remark 3.7 Let us remark how the above formalism works for analysis of the periodic box-ball systems. We concentrate on the path b of the form $B_1^{\otimes L}$, where number of [2] is equal to or less than that of [1]. We define $v_l \in B_l$ by the relation $u_l \otimes b \simeq T_l(b) \otimes v_l$ with $T_l(b) \in B_1^{\otimes L}$. Then we have $v_l \otimes b \simeq \overline{T_l}(b) \otimes v_l$ with $\overline{T_l}(b) \in B_1^{\otimes L}$ (Proposition 2.1 of [20]). $\overline{T_l}$'s are the time evolution operator of the periodic box-ball systems, and $\overline{T_1}$ is simply the cyclic shift operator.

Consider the path $b^{\otimes N} = b \otimes \cdots \otimes b$. Then, from the property of v_l , we have $T_l(b^{\otimes N}) = T_l(b) \otimes \overline{T}_l(b) \otimes \cdots \otimes \overline{T}_l(b)$, i.e., we can embed the periodic box-ball system into the usual linear system with operator T_l . Let us consider the local energy distribution for $b^{\otimes N}$. From the property $v_l \otimes b \simeq \overline{T}_l(b) \otimes v_l$, we see that under the right (N-1) copies of b in $b^{\otimes N}$, we have (N-1) copies of the same pattern of the local energy distribution.

Look at the local energy distribution below the rightmost b. In view of Theorem 3.6 and the comments following it, convenient way to find the structure of it is as follows. Instead of using u_l , we put v_l on the left of the path, and draw the local energy distribution. In step 2' of Theorem 3.6, we choose the rightmost 1 from (l-1)th row. If there is not such 1, we return to the right end of the (l-1)th row, and find such 1.

In such periodic extension of the local energy distribution, we can always find a boundary of successive two columns where none of groups crossing the boundary. By applying \bar{T}_1 , we can move such boundary to the left end of the path. We assume that b has already such property. In our case, we can always do such procedure, since by appropriate choice of d, we can always make $\bar{T}_1^d(b)$ highest weight (such d is not unique). Then this $\bar{T}_1^d(b)$ meet the condition (see Lemma C.1 of [20] and Lemma 4.2).

To summarize, by applying appropriate cyclic shifts, we can always make $b^{\otimes N}$ whose local energy distribution is N times repetition of the pattern for single b. On this property, we can apply the arguments of [21, 22] (combined with the tau function of [17]) to get the tau function in terms of the ultradiscrete Riemann theta function. More systematic treatment is given in [27]. Finally, we remark one thing without giving details (see section 3.3 of [27]). Recall that there is ambiguity in the choice of the cyclic shifts in the last paragraph. Let $\bar{T}_1^d(b)$ and $\bar{T}_1^{d'}(b)$ are the two such possible choices (we assume d' = 0 for the sake of simplicity). Consider the local energy distribution for b. If the left d columns contain a group of cardinality l (or, in other words, if the difference between b and $\bar{T}_1^d(b)$ is a soliton of length l), then the riggings corresponding to b and $\bar{T}_1^d(b)$ differ by the operator σ_l called the slide (see section 3.2 of [20] for definition of slides). The slides σ_l are closely related to the period matrix of the tau functions of [21, 22] (see section 4 of [20]).

4 Proof of Theorem 3.3 and Theorem 3.6

For the proof of Theorem 3.3, we prepare some lemmas.

Lemma 4.1 Let $(\lambda, (\mu, r))$ be the (unrestricted) rigged configuration corresponding to the path $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L$. Then we have (see (22) for definition of $Q_l^{(1)}$)

$$E_l = Q_l^{(1)}.$$
 (12)

Proof. We consider the path $b' := b \otimes \boxed{1}^{\otimes \Lambda}$, where $\Lambda \gg |\lambda|$. If b' is not highest, apply Lemma A.2 and we can use the same argument which is given below. Since E_l is conserved quantity on b', we have $E_l(T_{\infty}^{t_0}(b')) = E_l(b')$. We take t_0 large enough with the condition $\Lambda \geq t_0|\lambda|$ (the last inequality serves to assure that both $T_{\infty}^{t_0}(b')$ and b' contain the same number of letters 2). As we will see in the following, $T_{\infty}^{t_0}(b')$ has simplified structure, so that we can evaluate $E_l(T_{\infty}^{t_0}(b'))$ explicitly.

Now we use Proposition B.1. Since the actions of T_{∞} cause linear evolution of riggings, we can assume the (unrestricted) rigged configuration corresponding to $T_{\infty}^{t_0}(b')$ as $(\lambda \cup (1^{\Lambda}), (\mu, \bar{r}))$. By the assumption $t_0 \gg 1$, these \bar{r} have simple property. Recall that in (31), if we apply T_{∞} for one time, the rigging r_i corresponding to the row μ_i becomes $r_i + \mu_i$. Therefore the riggings \bar{r}_i and \bar{r}_j corresponding to the rows μ_i and μ_j satisfy $\bar{r}_i \gg \bar{r}_j$ if $\mu_i > \mu_j$.

Using these observations, we determine the shape of $T_{\infty}^{t_0}(b')$ from $(\lambda \cup (1^{\Lambda}), (\mu, \bar{r}))$. By the assumption $t_0 \gg 1$, all letters 2 in $T_{\infty}^{t_0}(b')$ are contained in $B_1^{\otimes \Lambda}$ part of the path. Therefore, corresponding to the row μ_i , there is a soliton of the form $2^{\otimes \mu_i}$. For example, in the following path,

$$\cdots \otimes \underline{1} \otimes \underline{2} \otimes \underbrace{1} \otimes \underbrace{1} \otimes \underbrace{1} \otimes \cdots \otimes \underline{1} \otimes \underline{1} \otimes \underline{2} \otimes \underline{2} \otimes \underline{1} \otimes \underline{1} \otimes \underline{2} \otimes \underline{2} \otimes \underline{1} \otimes \cdots$$
(13)

there are one soliton of length 1 and two solitons of length 2. Since the riggings satisfy $\bar{r}_i \gg \bar{r}_j$ if $\mu_i > \mu_j$, the shorter solitons are located on the far left of longer solitons (see the above example).

Assume there are solitons of the same length such as $2^{\otimes \mu_2} \otimes 1^{\otimes \sigma} \otimes 2^{\otimes \mu_1}$ ($\mu_1 = \mu_2$). Then we show $\sigma \geq \mu_1 = \mu_2$. Let the riggings corresponding to the rows μ_1 and μ_2 be r_1 and r_2 , respectively. In order to minimize σ , we choose $r_1 = r_2$. Now we consider ϕ^{-1} on rows μ_1 and μ_2 . Since we are assuming $t_0 \gg 1$, we do not need to consider the rows whose widths are different from μ_1 . From $r_1 = r_2$, rows μ_1 and μ_2 become simultaneously singular, and we can choose one of them arbitrary. We remove μ_1 first. While removing boxes from μ_1 one by one, the shortened row μ_1 is always made singular, and the rows whose lengths are shorter than μ_1 are not singular. Therefore we can remove entire row μ_1 successively. After removing row μ_1 , $Q_{\mu_2}^{(0)}$ decrease by μ_1 (note that the shape of the removed part of the quantum space is (1^{μ_1})), and $Q_{\mu_2}^{(1)}$ is also decrease by μ_1 (because of the removal of μ_1). Since the vacancy number is defined by $Q_{\mu_2}^{(0)} - 2Q_{\mu_2}^{(1)}$, the vacancy number for the row μ_2 increase by μ_1 compared to the one calculated before removing μ_1 . Therefore, in order to make the row μ_2 singular again, we have to remove extra μ_1 boxes from the quantum space, without removing boxes of μ part. Hence we have $\sigma \geq \mu_1$, as requested.

Now we are ready to evaluate $E_l(T^{t_0}_{\infty}(b'))$. From definition of the combinatorial R, we have

$$\begin{bmatrix}
l \\
11\cdots 1 \\
\otimes 2
\end{bmatrix}^{\otimes m} \otimes 1^{\otimes M} \\
\simeq 1^{\otimes \min(l,m)} \otimes 2^{\otimes \max(m-l,0)} \otimes \underbrace{1^{\otimes max(l-m,0)\min(l,m)}}_{11\cdots 1 22\cdots 2} \otimes 1^{\otimes M} \tag{14}$$

As we have seen, if there is a soliton of length m, there is always interval longer than m, i.e., it has the form $\dots \otimes \boxed{2}^{\otimes m} \otimes \boxed{1}^{\otimes M} \otimes \dots$ with $m \leq M$. This makes $\boxed{11 \dots 122 \dots 2}$ of (14) into the form $\boxed{11 \dots 1} = u_l$ when it comes to the left of the next (or right) soliton. Therefore, in order to evaluate $E_l(T^{t_0}_{\infty}(b'))$, we only have to consider the situation like (14). Noticing the fact that the energy function, i.e., unwinding number gains its value from the unwinding pair $\boxed{1 \dots \otimes 2}$ appearing in the each tensor product (more precisely, E_l gains min(l, m) corresponding to the procedure (14)), the proof of lemma finishes.

Combining the property of the combinatorial R with Lemma 4.1, the relationship between the local energy distribution and the KKR bijection can be clarified as follows.

Lemma 4.2 For the given path $b = b_1 \otimes \cdots \otimes b_i \otimes \cdots \otimes b_L$, draw local energy distribution. Within the *i* th column, denote the locations of 1 as $(j_1, i), (j_2, i), \cdots, (j_k, i)$ $(j_1 < j_2 < \cdots < j_k)$. Consider the calculation of $\phi(b) = (\lambda, (\mu, r))$. During the whole process of $\phi(b)$, when we create part of the (unrestricted) rigged configuration from b_i of b, we add boxes to columns j_1, j_2, \cdots, j_k of μ in this order.

Proof. Denoting $b_i = (x_1, x_2)$, let us define $b_{i,s} = (0, s)$ for $s \le x_2$. Consider the path $b_s = b_1 \otimes b_2 \otimes \cdots \otimes b_{i,s}$, and draw local energy distribution for this b_s . In the local energy distribution, from the 1 st column to i - 1 th column are identical to the ones in the case for the original b. On the other hand, from i + 1 th column to L th column, local energies are all 0. These are obvious from the construction of b_s .

Now consider the *i* th column of the local energy distribution for b_s . Then we show that the *i* th column is obtained from the one corresponding to the original *b* by making *s* letters 1 from the top as it is, and letting all other letters be 0. This follows from the property of the combinatorial *R*, that is, the order of making pairs of dots do not affect the resulting image of the combinatorial *R*. In fact, in calculating $E_{l,i}$, we have $u_l^{(i-1)} \otimes b_{i,s}$. Compare this with $u_l^{(i-1)} \otimes b_i$. Then we can make pair of dots in $u_l^{(i-1)} \otimes b_i$ such that first we join unwinding pairs, and then we join winding pairs. Note that all letters 1 contained in b_i here cannot contribute as unwinding pairs. Therefore, we see that when we consider $u_l^{(i-1)} \otimes b_{i,s}$, $E_{l,i}$ $(l = 1, 2, \cdots)$ are the same with $u_l^{(i-1)} \otimes b_i$ case up to the first *s* unwinding pairs, and we do not have the rest of the unwinding pairs. This verifies the assertion for *i* th column of the local energy distribution for b_s .

Compare the local energy distribution for $b_1 \otimes \cdots \otimes b_{s-1}$ and $b_1 \otimes \cdots \otimes b_s$. Then, from the above observation, there is extra one 1 at (j_s, i) . Now we apply the relation $E_l = Q_l^{(1)}$ (Lemma 4.1) to both $b_1 \otimes \cdots \otimes b_{s-1}$ and $b_1 \otimes \cdots \otimes b_s$. Then we see that the letter 1 at (j_s, i) corresponds to addition of one box at column j_s of μ of $\phi(b_{s-1})$. Since μ part of $\phi(b)$ is obtained by adding boxes to μ recursively as $\phi(b_1), \phi(b_2), \cdots, \phi(b_{i,1}), \phi(b_{i,2}), \cdots$, this gives the proof of lemma.

Lemma 4.3 Let $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots \otimes B_{\lambda_L}$ be an arbitrary path. b can be highest weight or non-highest weight. Set $N = E_1(b)$. We determine the numbers $\mu_1, \mu_2, \cdots, \mu_N$ by the following procedure from Step 1 to Step 3.

- 1. Draw a table containing $(E_{l,j} E_{l-1,j} = 0, 1)$ at the position (l, j), i.e., at the *l* th row and the *j* th column. We call this table local energy distribution.
- 2. Starting from the rightmost 1 in the l = 1 st row, pick the nearest 1 from each successive row. The one in the (l + 1) th row must be weakly right of the one selected in the l th row. If there is no such 1 in the (l + 1) th row, the position of the lastly picked 1 is called (μ_1, j_1) . Change all selected 1 into 0.
- 3. Repeat Step 2 for (N-1) times to further determine $(\mu_2, j_2), \dots, (\mu_N, j_N)$ thereby making all 1 into 0. Then μ coincides with μ of the (unrestricted) rigged configuration $\phi(b) = (\lambda, (\mu, r)).$

Proof. We first interpret Step 2 in terms of the original combinatorial procedure ϕ . In Step 2, we choose the rightmost 1 of the first row of local energy distribution. From Lemma 4.2, this 1 corresponds to the leftmost box of the lastly created row of μ . Suppose we choose letters 1 up to l th row according to Step 2. Next we choose 1 in l + 1 th row, whose position is weakly right of 1 in l th row. Since the lastly created row grows by adding boxes one by one during the procedure ϕ , this means that these two 1 at l th and l + 1 th rows of the local energy distribution belong to the same row (lastly created row) of μ . Note that if there are more than one row with the same length l, we can always add a box to the lastly created row, since it has maximal riggings among the rows with the same length. This follows from the fact that after adding a box at l th column of lastly created row of μ , the row is made to be singular, i.e., the row is assigned the maximal possible rigging for the row with length l. To summarize, Step 2 ensures us to identify all 1 in the local energy distribution which correspond to the lastly created row of μ .

In Step 3, we do the same procedure for the rest of 1 in the local energy distribution. Since we omit all letters 1 which are already identified with some rows of μ , we can always use Step 2 to determine the next row. Therefore, Step 3 ensures us to identify all 1 in the local energy distribution with the rows of μ .

For the proof of Theorem 3.3, we show that the groups of letters 1 obtained in Lemma 4.3 have simplified structure.

Proposition 4.4 The groups of letters 1 obtained in Step 2 and Step 3 of Lemma 4.3 have no crossing with each other.

Proof. The proof is divided into 6 steps. In Step 1, we analyze the geometric property of the crossing of groups. In Step 2, we make the assumptions about crossing. In Step 3, we analyze the behavior of corigging (= vacancy number - rigging) under the operation of ϕ . Then we introduce a convenient graphical method to analyze the coriggings by using " \circ " and " \bullet ". Here " \bullet " represents the letters 1 contained in the local energy distribution and " \circ " represents the change of the quantum space induced by letters 1 contained in the path. In Step 4, we derive the relations from the assumption made in Step 2. In Step 5, we consider how to minimize the number of " \circ " for given pattern of " \bullet " of the local energy distribution. Finally, in Step 6, we show that combination of the relations derived in Step 4 and arguments in Step 5 lead to the contradiction, hence completes the proof.

Step 1. Consider the path $b = b_1 \otimes b_2 \otimes \cdots \otimes b_L$, and calculate the local energy distribution corresponding to $b_{[k]} = b_1 \otimes b_2 \otimes \cdots \otimes b_k$ $(k \leq L)$. Recall that in Lemma 4.2, we have shown that patterns of the local energy distribution represent the combinatorial procedures of the KKR bijection ϕ . Suppose that there are two groups of letters 1 whose cardinalities are m_1 and m_2 , respectively, below $b_{[k]}$. We name these two groups as M_1 and M_2 , respectively. Here we take the top end of the group M_1 is located to the left of that of the group M_2 .

From geometric property of crossing, we show that, by appropriate choice of k, we can assume $m_1 < m_2$ without crossing beneath $b_{[k]}$. Assume that there is a crossing between lth row and (l+1)th row, whereas there is no crossing above it. Denote the elements of M_1 and M_2 at lth row by (l, m_{1-}) and (l, m_{2-}) where $m_{1-} < m_{2-}$, and the elements of m_1 and m_2 at (l+1)th row by $(l+1, m_{1+})$ and $(l+1, m_{2+})$ where $m_{2+} < m_{1+}$, respectively. From the procedure given in Step 2 of Lemma 4.3, we have $m_{2-} \le m_{2+}$. Now we choose k such that $m_{2+} \le k < m_{1+}$ is satisfied. Since k satisfies $m_{1-} < m_{2-} \le m_{2+} \le k$, cardinality of the group M_1 beneath $b_{[k]}$ is l, whereas that of the group M_2 is equal to or greater than l+1, which gives the claim.

Step 2. We keep the notation like m_{i+} etc., as before, therefore we have $m_1 < m_2$ beneath $b_{[m_{1+}-1]}$. Again, we are assuming that the crossing of M_1 and M_2 occurs beneath $b_{[m_{1+}]}$, and also that there is no crossing beneath $b_{[m_{1+}-1]}$. We denote the cardinalities of the groups M_1 and M_2 under $b_{[m_{1+}-1]}$ by m_1 and m_2 , respectively. At the end of the proof, we will show that the existence of the crossing leads to contradiction. We consider the crossing of two groups, since this is the fundamental situation. The general case involving more than two crossings follows from this fundamental case. The situation here is depicted in the following diagram.



Here letters 1 in the local energy distribution are represented by "•" and all letters 0 are suppressed. Note that we have introduced the domain D(m) on the right of the bottom point of M_2 , occupying from the first row to the *m*th row. Since we are considering the

crossing caused by two groups, we can assume that groups contained in $D(m_2)$ are, in fact, contained in $D(m_1-1)$ (in the above diagram, it is indicated by the gray rectangle). If we can put "•" indicated by "?" in the above diagram, then the crossing of the groups M_1 and M_2 occurs.

In the following, we first treat the case that M_2 and other groups on the right of M_2 are well separated. This means the other groups on the right of M_2 are located on the right of the bottom point of M_2 , and between the top and the bottom point of M_2 , there is no "o" on the right of M_2 (see Step 3 for meaning of "o"). This assumption is only for the sake of simplicity, and the general case will be mentioned at the end of the proof.

Step 3. We summarize the basic properties of the vacancy numbers (or, at the same time, that of the coriggings). Recall the definition of the vacancy numbers $p_j = Q_j^{(0)} - 2Q_j^{(1)}$ corresponding to the pair (λ, μ) . Consider the box adding procedure of ϕ . If we add boxes to $\lambda|_{\leq j}$ and $\mu|_{\leq j}$ simultaneously, then the vacancy number p_j decrease by 1. On the contrary, if we add a box to $\lambda|_{\leq j}$ and do not add box to $\mu|_{\leq j}$, then the vacancy number increase by 1. Note that If we do not add box to both $\lambda|_{\leq j}$ and $\mu|_{\leq j}$, then the vacancy number do not change. Recall also that the procedure ϕ only refers to corigging.

In order to analyze the above change of coriggings induced by box adding procedure of ϕ , it is convenient to supplement the local energy distribution with the information of change of the quantum space corresponding to letters 1 contained in the path. In the local energy distribution, we replace letters 1 by "•", and suppress all letters 0. We join "•" belonging to the same group by thick lines. Then, corresponding to the letters 1 contained in b_s , we put "o" on the right of "•" corresponding to the letters 2 contained in b_s . The row coordinates of "o" are taken to be the same as the column coordinate of the added box of the quantum space induced by the corresponding letters 1. Here we give examples of such diagram for $22 \otimes 12 \otimes 1 \otimes 22$ and $22 \otimes 112 \otimes 22$, respectively.



In this diagram, the KKR map ϕ proceeds from the left to right, and within the same column (distinguishing columns of "•" and "o"), it proceeds from the top to bottom. In the above examples, the left group containing three "•" became singular (i.e., corigging = 0) after the bottom "•" is added. Then the two "o" increase the corigging by 2, thereby the right group containing two "•" stays independently from the left group. This kind of analysis of change of the coriggings is a prototype of the arguments given in Step 5 and Step 6. Note that along each group of "•", the notion of left and right of the group is well defined. Let us remark the convenient method to determine locations of "o". Given an element $1 \cdots 12 \cdots 2$, we reverse the orderings of numbers as $2 \cdots 21 \cdots 1$. Choose the specific letter 1, and denote by p the number of letters 1 and 2 on the left of it. Then, corresponding to the chosen 1, we put "o" on the (p+1)th row on the local energy distribution.

Step 4. Assume that we are going to add the box corresponding to $(m_1 + 1, m_{1+})$ of the local energy distribution. In order to add a box corresponding to $(m_1 + 1, m_{1+})$, or in other words, in order to make crossing, the row of μ corresponding to M_2 cannot be singular when we add the box corresponding to $(m_1 + 1, m_{1+})$. This follows from the assumption $m_1 < m_2$ and the fact that we add a box to the longest possible singular row in the procedure ϕ . Also, the row of μ corresponding to M_1 have to be singular in order to add a box corresponding to $(m_1 + 1, m_{1+})$. We consider the implications of these two conditions.

Recall that the row of μ corresponding to the group M_2 is singular when the bottom "•" is added to the end of M_2 . On the other hand, we have to make M_2 non-singular as we have seen in the above. This means

(number of "o" within
$$D(m_2)$$
) > (number of "•" within $D(m_2)$). (15)

On the other hand, in order to make M_1 singular, we have

(number of "o" within
$$D(m_1)$$
) \leq (number of "•" within $D(m_1)$). (16)

Note that all "•" are contained in $D(m_1 - 1)$. From these two restrictions, we see there are at least one "o" in $D(m_2) \setminus D(m_1)$.

Step 5. If we are given the pattern of "•", there remains various possibilities about pattern of "o". Now we are going to consider the patterns of "o" that minimize the number of "o". To say the result at first, we see that we need "o" as much as "•", therefore in order to meet the condition (16), we have to minimize the number of "o".

Suppose there are two groups in the local energy distribution, the group s_1 on the left, and s_2 on the right. Let the top "•" be located at columns k_1 and k_2 . Then, in order to make s_1 and s_2 as separated groups, we need $\min(s_1, s_2)$ "o" within the region between (or surrounded by) s_1 and column k_2 . To make the situation transparent, we consider the concrete path 22222222 \otimes 111111 \otimes 2 \otimes 1122 \otimes 22 \otimes 22 \otimes 111111 \otimes 2222222. Then the corresponding diagram is as follows.



We see there are 4 groups, labeled by M_2 , s_1 , s_2 , s_3 from left to right. These groups are indicated by thick lines. We can analyze the situation as follows (as for the method for analysis of change of the coriggings, see latter part of Step 3):

1. In order to make M₂ and s₁ separated, we need 6 "○" between M₂ and s₁. Precise meaning of "between M₂ and s₁" etc., is given after the example. In this example, they are supplied by 6 "○" coming from the left <u>111111</u>. This situation is indicated by thin lines, which join the corresponding "○" and "●". Of course, there is ambiguity in the way of joining "○" and "●", however this ambiguity brings no important effect, hence we neglected. For example, we can join top 5 "○" coming from the left <u>111111</u> and the bottom "○" coming from <u>1122</u> with 6 "●" of s₁. In such a case, the bottom "○" coming from the left <u>111111</u> should be connected with the bottom "●" of s₃.

- 2. In order to make s_1 and s_2 separated, we need 1 " \circ " between s_1 and s_2 . In this example, it is supplied by 1 " \circ " coming from 1122.
- 3. In order to make s_2 and s_3 separated, we need 1 "o" between s_2 and s_3 . In this example, it is supplied by the top "o" coming from the right 111111.
- 4. In order to make s_1 and s_3 separated, we need 5 " \circ " between s_1 and s_3 . In this example, it is supplied by the bottom 5 " \circ " coming from the right 111111.
- 5. In order to make M_2 and s_3 separated, we need 1 " \circ " between M_2 and s_3 . In this example, it is supplied by 1 " \circ " coming from 1122.

From this example, we can infer the general case. Assume there are groups s_1, \dots, s_n (from left to right) on the right of M_2 . We define the region between s_i and s_j (i < j) as the region surrounded by s_i , the first row and the row containing the bottom point of s_i , s_j and the column containing the bottom point of s_j (except s_i , s_j and the column containing the bottom point of s_j , see the following diagram).



We choose ordered subsequence s_{j_1}, \dots, s_{j_p} of groups s_1, \dots, s_n such that it is the longest subsequence which satisfies $s_{j_1} > \dots > s_{j_p} > s_{n-1}$. Between s_{n-1} and the s_n , we need at least min (s_n, s_{n-1}) "o" in order to make s_{n-1} and s_n separate. If $s_n > s_{n-1}$, then we compare s_n and s_{j_p} . Then we need at least min $(s_n - s_{n-1}, s_{j_p} - s_{n-1})$ "o" between s_{j_p} and the s_n in order to make s_{j_p} and s_n separate. We continue this process and, in conclusion, we need "o" as much as "•", compared within the right of M_2 .

Step 6. Based on the ground of the arguments given in Step 5, we derive the contradiction against the statement "there are at least one " \circ " in $D(m_2) \setminus D(m_1)$ ".

Again assume there are groups s_1, \dots, s_n on the right of M_2 . Denote the column coordinate of the top "•" of s_j by k_j and that of the bottom of s_j by k'_j . In order to meet the condition (16), we choose the pattern of "o" which minimizes the number of "o". From observations made in Step 5, we have at most s_n "o" between s_{n-1} and the s_n . If all "o" are located between the k_{n-1} th column and k_n th column, then the number of "o" (i.e. s_n) is too short to make "o" appear in $D(m_2) \setminus D(m_1)$. Let us analyze the case when some of s_n "o" appear between the k_n th column and k'_n th column. To be specific, take some k between k_n and k'_n , and write the row of the lowest "•" of the column k belonging to the group s_n by s. In order to make the upper s "•" of s_n separated from s_{n-1} , it needs at least s "o" on the left of the kth column. See the following schematic diagram.



Thus, we have at most $s_n - s$ " \circ " below the *s*th row of the *k*th column. If we attach a column of $s_n - s$ " \circ " to the *k*th row, it has to begin from the s + 1th row. Therefore, it is also too short to make " \circ " appear in $D(m_2) \setminus D(m_1)$. Similarly, between s_{n-2} and the s_{n-1} , we have at most $\max(s_n, s_{n-1})$, again too short to make " \circ " appear in $D(m_2) \setminus D(m_1)$. Continuing in this way, we see that no " \circ " appear in $D(m_2) \setminus D(m_1)$, which gives contradiction.

As we have claimed at the end of Step 2, so far we are dealing only with the case that M_2 and the other groups on the right of it are well separated. However, we can treat the general case by similar arguments. First, by applying the same argument of Step 5, we can show that we need " \circ " as much as " \bullet " within the region on the right of the group M_2 . Then, by applying the same argument in Step 6, all " \circ " on the right of M_2 are included in the first $m_1 - 1$ rows of the local energy distribution. Therefore, in order to " \circ " appear in $D(m_2) \setminus D(m_1)$, we have to add at least one " \circ " within the first m_1 rows. This makes M_1 non-singular, hence the crossing does not occur in this case.

Hence we complete the proof of proposition.

Proof of Theorem 3.3. From Proposition 4.4, we can remove the procedure to find "the nearest 1" from Step 2 of Lemma 4.3. This gives the proof of Step 1 to Step 3 of Theorem 3.3.

Finally, we clarify the meaning of Step 4. In Step 3 of Lemma 4.3, we determined (μ_k, j_k) , which corresponds to the rightmost box of row μ_k of μ . Since the row μ_k is not lengthened in calculation of $\phi(b)$ after b_{j_k} , the rigging of row μ_k is equal to the vacancy number at the time when we add the rightmost box to μ_k . At this moment, the quantum space takes the form

$$(\lambda_1, \lambda_2, \cdots, \lambda_{j_k-1}, E_{\mu_k, j_k}). \tag{17}$$

The meaning of the last E_{μ_k,j_k} is as follows. E_{μ_k,j_k} counts all letters 1 contained in the first μ_k rows of j_k th column of local energy distribution. This means that, from Lemma 4.2, we added E_{μ_k,j_k} boxes to μ before the rightmost box of the row μ_k is added (while considering b_{j_k}). In the procedure ϕ , we use letters 2 of b_{j_k} first, and then use the rest of letters 1 of b_{j_k} . Since letters 2 in b_{j_k} means simultaneous addition of box to the quantum space and μ , we can conclude that the quantum space has the row with length $E_{\mu_k,j_k}(\leq \mu_k)$. From this shape of the quantum space, we have

$$Q_{\mu_k}^{(0)} = \sum_{i=1}^{j_k - 1} \min(\mu_k, \lambda_i) + E_{\mu_k, j_k}$$
(18)

From Lemma 4.1 applying to the path $b_1 \otimes b_2 \otimes \cdots \otimes b_{j_k}$, we deduce the following:

$$Q_{\mu_k}^{(1)} = \sum_{i=1}^{j_k} E_{\mu_k,i}.$$
(19)

Hence we obtain the formula in Step 4, and complete the proof of Theorem 3.3. $\hfill \Box$

Proof of Theorem 3.6. This follows immediately from non-crossing property of Proposition 4.4. In Theorem 3.3, we determine groups from right to left. More precisely, after determining one group, all letters 1 belonging to the group are made to be 0, and we determine the rightmost group again. However, if we start from the bottom point of the longest group, the elimination made in the right has no effect, and we can determine the same group by virtue of non-crossing property. By doing these procedure from longer groups to shorter groups, we obtain the same groups as in the case Theorem 3.3.

5 Summary

In this paper, we consider crystal interpretation of the KKR map ϕ from paths to rigged configurations. In Section 3, we introduce table called the local energy distribution. The entries of the table are differences of the energy functions, and we show in Theorem 3.3 that this table have complete information about ϕ so that we can read off the rigged configuration from it. As we see in Lemma 4.2, this table can be viewed as giving crystal interpretation of the combinatorial procedures appearing in the original definition of ϕ .

As we see in Proposition 4.4, our formalism has simple property. This enables us to reformulate Theorem 3.3 as described in Theorem 3.6. The latter formalism is particularly important when we consider inverse scattering formalism for the periodic box-ball systems. As we see in Remark 3.7, advantage of our formalism, compared with the formalism given in [20], is that we can treat states of the periodic box-ball system directly without sending them to linear semi-infinite systems.

A Kerov–Kirillov–Reshetikhin bijection

In this section, we prepare notations and basic properties corresponding to the Kerov– Kirillov–Reshetikhin (KKR) bijection. As for the definitions of the rigged configurations as well as combinatorial procedure of the bijection, we refer to Section 2 of [16] (ϕ there should be read as ϕ^{-1} here) or Appendix A to [20], and we only prepare necessary notations.

Assume we have given a highest weight path b:

$$b = b_1 \otimes b_2 \otimes \cdots \otimes b_L \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots \otimes B_{\lambda_L}.$$
 (20)

Then we have one to one correspondence ϕ between b and the rigged configuration

$$\phi: b \longrightarrow \mathrm{RC} = \left((\lambda_i)_{i=1}^L, \, (\mu_i, r_i)_{i=1}^N \right). \tag{21}$$

Here $(\mu_i)_i \in \mathbb{Z}_{\geq 0}^N$ is called configuration and we depict $(\lambda_i)_i$ and $(\mu_i)_i$ by Young diagrammatic expression whose rows are given by λ_i and μ_i , respectively. Integers r_i are called riggings and we associate them with the corresponding row μ_i . In the KKR bijection,

orderings within integer sequences $(\lambda_i)_i$ or $(\mu_i, r_i)_i$ does not make any differences. On this rigged configurations, we use the symbols $Q_j^{(a)}$ (a = 0, 1) defined by

$$Q_j^{(0)} := \sum_{k=1}^{L} \min(j, \lambda_k), \qquad Q_j^{(1)} := \sum_{k=1}^{N} \min(j, \mu_k).$$
(22)

The vacancy number p_j for length j rows of μ is then defined by

$$p_j := Q_j^{(0)} - 2Q_j^{(1)}.$$
(23)

If row μ_i has property $p_{\mu_i} = r_i$, then the row is called singular. For the highest weight paths, the corresponding rigged configurations are known to satisfy $0 \le r_i \le p_{\mu_i}$. The quantity $p_{\mu_i} - r_i$ is sometimes called corigging.

One of the most important properties of ϕ or ϕ^{-1} is that if we consider isomorphic paths $b \simeq b'$, then the corresponding rigged configuration is the same (Lemma 8.5 of [3]). We express this property in terms of the map ϕ^{-1} as follows.

Theorem A.1 Take successive two rows from the quantum space λ of the rigged configuration arbitrary, and denote them by λ_a and λ_b . When we remove λ_a at first and next λ_b by the KKR map ϕ^{-1} , then we obtain two tableaux, which we denote by a_1 and b_1 , respectively. Next, on the contrary, we first remove λ_b and second λ_a (keeping the order of other removal invariant) and we get b_2 and a_2 . Then we have

$$b_1 \otimes a_1 \simeq a_2 \otimes b_2, \tag{24}$$

under the isomorphism of \mathfrak{sl}_2 combinatorial R matrix.

We remark that there is an extension of ϕ which covers all non-highest weight elements as well. Let b be an arbitrary element of arbitrary tensor products of crystals; $b \in B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_L}$. In particular, b can be non-highest weight element. Then we can apply the same combinatorial procedure for ϕ and obtain $\phi(b)$ as extension of the rigged configurations. Following [28, 29], we call such $\phi(b)$ unrestricted rigged configuration. Let us denote $\phi(b) = (\lambda, (\mu, r))$. Then, from definition of ϕ , we see that $|\lambda|$ represents the number of all letters 1 and 2 contained in the path b, whereas $|\mu|$ represents the number of letters 2 contained in b. Note, in particular, that $|\lambda| \geq |\mu|$ holds for unrestricted rigged configurations. These unrestricted rigged configurations contain the rigged configurations as the special case.

Let b be a non-highest weight element as above. Consider the following modification of b:

$$b' := \boxed{1}^{\otimes \Lambda} \otimes b, \tag{25}$$

where Λ is an integer satisfying $\Lambda \geq \lambda_1 + \cdots + \lambda_L$. Then b' is highest weight. Under these notations, we have the following:

Lemma A.2 Let the unrestricted rigged configuration corresponding to b be

$$((\lambda_i)_{i=1}^L, (\mu_j, r_j)_{j=1}^N).$$
 (26)

Then the rigged configuration corresponding to the highest path b' is given by

$$((\lambda_i)_{i=1}^L \cup (1^\Lambda), (\mu_j, r_j + \Lambda)_{j=1}^N).$$
 (27)

Proof. Let the vacancy number of row μ_j of the pair (λ, μ) of (26) be p_{μ_j} . Then the vacancy number of row μ_j of (27) is equal to $p_{\mu_j} + \Lambda$, because of the addition of (1^{Λ}) on λ . Now we apply ϕ^{-1} on (27). From $\lambda \cup (1^{\Lambda})$ of the quantum space, we remove λ first, and next remove (1^{Λ}) . Recall that in the combinatorial procedure of ϕ^{-1} , we only refer to corigging, and it does not refer to actual values of the riggings. Therefore, when we remove λ from the quantum space of (27), we obtain b as the corresponding part of the image. Then, remaining rigged configuration has the quantum space (1^{Λ}) without μ part. On this rigged configuration, the map ϕ^{-1} becomes trivial and obtain b' as the image corresponding to (27).

B Operators T_l

In this section, we introduce the operators T_l which are defined by the combinatorial R. T_l 's serve as the time evolution operators of the box-ball systems [25]. Let u_l be a highest weight element of B_l , i.e., in a tableau representation, it is $u_l = \underbrace{11\cdots 1}^l$. We consider the path

$$b = b_1 \otimes b_2 \otimes \dots \otimes b_L \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \dots \otimes B_{\lambda_L}.$$
(28)

Then its time evolution $T_l(b)$ $(l \in \mathbb{Z}_{>0})$ is defined by successively sending u_l to the right of b under the isomorphism of the combinatorial R as follows:

$$u_{l} \otimes b = u_{l} \otimes b_{1} \otimes b_{2} \otimes \cdots \otimes b_{L}$$

$$\stackrel{R}{\simeq} b_{1}' \otimes u_{l}^{(1)} \otimes b_{2} \otimes \cdots \otimes b_{L}$$

$$\stackrel{R}{\simeq} b_{1}' \otimes b_{2}' \otimes u_{l}^{(2)} \otimes \cdots \otimes b_{L}$$

$$\stackrel{R}{\simeq} \cdots \cdots$$

$$\stackrel{R}{\simeq} b_{1}' \otimes b_{2}' \otimes \cdots \otimes b_{L}' \otimes u_{l}^{(L)}$$

$$=: T_{l}(b) \otimes u_{l}^{(L)}.$$
(29)

According to Proposition 2.6 of [13], operators T_l on highest paths can be linearized by the KKR bijection. Since, in the main text, we use the similar property for general case including non-highest paths, we include here a proof for generalized version.

Proposition B.1 (1) Consider the path *b* of the form (28). Here *b* can be non-highest weight element. Define $b' = b \otimes \boxed{1}^{\otimes \Lambda}$, where the integer Λ satisfies $\Lambda > \lambda_1 + \lambda_2 + \cdots + \lambda_L$. Then, we have $u_l \otimes b' \simeq T_l(b') \otimes u_l$.

(2) Denote the (unrestricted) rigged configuration corresponding to b' as

$$b' \xrightarrow{\text{KKR}} (\lambda \cup (1^{\Lambda}), (\mu_j, r_j)_{j=1}^N).$$
 (30)

Then, corresponding to $T_l(b')$, we have

$$T_l(b') \xrightarrow{\text{KKR}} \left(\lambda \cup (1^\Lambda), (\mu_j, r_j + \min(\mu_j, l))_{j=1}^N \right).$$
(31)

Proof. (1) Consider the elements $u_l^{(i)}$ defined in (29). In our case, $u_l^{(L)}$ contains letters 2 for at most $\lambda_1 + \lambda_2 + \cdots + \lambda_L$ times. Then, by calculating combinatorial R along (29) with $u_l^{(L)}$ and $\boxed{1}^{\otimes \Lambda}$, we see that $u_l^{(L+\Lambda)} = u_l$.

(2) Consider the following rigged configuration:

$$(\lambda \cup (1^{\Lambda}) \cup (l), (\mu_j, r_j + \min(\mu_j, l))_{j=1}^{N}),$$
(32)

i.e., we added a row with width l to the quantum space. Compare the coriggings of (32) and (30). Recall that the vacancy number is defined by $Q_{\mu_j}^{(0)} - 2Q_{\mu_j}^{(1)}$. As for $Q_{\mu_j}^{(1)}$, both (32) and (30) give the same value, since we have μ in the both second terms. On the contrary, $Q_{\mu_j}^{(0)}$ for (32) is greater than the one for (30) by $\min(\mu_j, l)$, since we have the extra row of width l in the quantum space of (32). In (32), riggings are increased by value $\min(\mu_j, l)$, therefore we conclude that the coriggings for both (32) and (30) coincide.

Now we apply ϕ^{-1} on (32) in two different ways. First, we remove $\lambda \cup (1^{\Lambda})$ from the quantum space of (32) (order of removal is the same as ϕ^{-1} on (30) to obtain b'). Since the coriggings for both μ_j of (32) and (30) coincide, we obtain b' as the corresponding part of the image. Then we are left with the rigged configuration $(l, (\emptyset, \emptyset))$, which yields u_l . Therefore we obtain $u_l \otimes b'$ as the image.

Next, we apply ϕ^{-1} on (32) in different way. In this case, we remove the row l of the quantum space as the first step. Note that in the (unrestricted) rigged configuration $(\lambda, (\mu, r))$ corresponding to the path b, all riggings r_j are smaller than the corresponding vacancy numbers. By definition of Λ , we have $\Lambda > \min(\mu_j, l)$ for all j (recall that from definition of the unrestricted rigged configuration, we always have $\lambda_1 + \cdots + \lambda_L \ge \mu_1 + \cdots + \mu_N$). As the result, if we remove the row l from the quantum space of (32), rows μ_j do not become singular even if the riggings are increased as $r_j + \min(\mu_j, l)$, since the vacancy numbers are also increased by Λ by the addition of (1^{Λ}) . Thus, we obtain u_l as the corresponding part of the image. Then we are left with $(\lambda \cup (1^{\Lambda}), (\mu_j, r_j + \min(\mu_j, l))_{j=1}^N)$, whose corresponding path we denote by $\tilde{b'}$. In conclusion, we obtain $\tilde{b'} \otimes u_l$ as the image.

In the above two calculation of ϕ^{-1} , the only difference is the order of removing rows of the quantum space of (32). Therefore we can apply Theorem A.1 to get the isomorphism

$$u_l \otimes b' \stackrel{R}{\simeq} \tilde{b'} \otimes u_l. \tag{33}$$

If b' is non-highest weight, we apply Lemma A.2 and $u_l \otimes u_m \simeq u_m \otimes u_l$, then we can use the same argument. From (1), this means $\tilde{b'} = T_l(b')$, which completes a proof.

Acknowledgements: The author would like to thank Atsuo Kuniba and Taichiro Takagi for collaboration in an early stage of the present study and valuable comments on this manuscript. He is a research fellow of the Japan Society for the Promotion of Science.

References

- S. V. Kerov, A. N. Kirillov and N. Yu. Reshetikhin, Combinatorics, the Bethe ansatz and representations of the symmetric group. J. Soviet Math. 41 (1988) 916–924.
- [2] A. N. Kirillov and N. Yu. Reshetikhin, The Bethe ansatz and the combinatorics of Young tableaux. J. Soviet Math. 41 (1988) 925–955.
- [3] A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood– Richardson tableaux and rigged configurations, Selecta Math. (N.S.) 8 (2002) 67–135, arXiv:math.CO/9901037.

- [4] M. Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras, Duke Math. J. 63 (1991) 465–516.
- [5] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Affine crystals and vertex models, Int. J. Mod. Phys. A7 Suppl. 1A (1992) 449–484.
- [6] I. Macdonald, Symmetric functions and Hall polynomials, 2nd edition, Oxford Univ. Press, New York (1995).
- [7] H. A. Bethe, Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, Z. Physik 71 (1931) 205–231.
- [8] M. Okado, X = M conjecture, MSJ Memoirs **17** (2007) 43–73.
- [9] A. Schilling, X = M Theorem: Fermionic formulas and rigged configurations under review, MSJ Memoirs **17** (2007) 75–104, arXiv:math.QA/0512161.
- [10] G. Lusztig, Fermionic form and Betti numbers, arXiv:math.QA/0005010.
- [11] H. Nakajima, t-analogs of q-characters of quantum affine algebras of type E_6 , E_7 , E_8 , arXiv:math/0606637.
- [12] M. Okado and A. Schilling, Existence of Kirillov-Reshetikhin crystals for nonexceptional types, Representation Theory 12 (2008) 186–207, arXiv:0706.2224.
- [13] A. Kuniba, M. Okado, R. Sakamoto, T. Takagi and Y. Yamada, Crystal interpretation of Kerov-Kirillov-Reshetikhin bijection, Nucl. Phys. B740 (2006) 299–327, arXiv:math.QA/0601630.
- [14] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Japan, 59 (1990) 3514–3519.
- [15] D. Takahashi, On some soliton systems defined by using boxes and balls, Proceedings of the International Symposium on Nonlinear Theory and Its Applications (NOLTA '93), (1993) 555–558.
- [16] R. Sakamoto, Crystal interpretation of Kerov–Kirillov–Reshetikhin bijection II. Proof for \mathfrak{sl}_n case, J. Algebraic Combinatorics **27** (2008) 55–98, arXiv:math.QA/0601697.
- [17] A. Kuniba, R. Sakamoto and Y. Yamada, Tau functions in combinatorial Bethe ansatz, Nucl. Phys. B786 (2007) 207–266, arXiv:math.QA/0610505.
- [18] F. Yura and T. Tokihiro, On a periodic soliton cellular automaton, J. Phys. A: Math. Gen. 35 (2002) 3787–3801, arXiv:nlin/0112041.
- [19] D. Yoshihara, F. Yura and T. Tokihiro, Fundamental cycle of a periodic box-ball system, J. Phys. A: Math. Gen. 36 (2003) 99–121. arXiv:nlin/0208042.
- [20] A. Kuniba, T. Takagi and A. Takenouchi, Bethe ansatz and inverse scattering transform in a periodic box-ball system, Nucl. Phys. B747 (2006) 354–397, arXiv:math.QA/0602481.

- [21] A. Kuniba and R. Sakamoto, The Bethe ansatz in a periodic box-ball system and the ultradiscrete Riemann theta function, J. Stat. Mech. (2006) P09005, 1–12, arXiv:math.QA/0606208.
- [22] A. Kuniba and R. Sakamoto, Combinatorial Bethe ansatz and ultradiscrete Riemann theta function with rational characteristics, Lett. Math. Phys. 80 (2007) 199–209, arXiv:nlin/0611046.
- [23] J. Mada, M. Idzumi and T. Tokihiro, On the initial value problem of a periodic box-ball system J. Phys. A: Math. Gen. 39 (2006) L617–L623, arXiv:nlin/0608037.
- [24] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Math. (N.S.) 3 (1997) 547–599, arXiv:q-alg/9512027.
- [25] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, The $A_M^{(1)}$ automata related to crystals of symmetric tensors, J. Math. Phys. **42** (2001) 274–308, arXiv:math.QA/9912209.
- [26] K. Fukuda, M. Okado and Y. Yamada, Energy functions in box-ball systems, Int. J. Mod. Phys. A15 (2000) 1379–1392, arXiv:math.QA/9908116.
- [27] A. Kuniba and R. Sakamoto, Combinatorial Bethe ansatz and generalized periodic box-ball system, Rev. Math. Phys. 20 (2008) 1–35, arXiv:0708.3287.
- [28] A. Schilling, Crystal structure on rigged configurations, Int. Math. Res. Notices (2006) Article ID 97376, 1–27, arXiv:math.QA/0508107.
- [29] L. Deka and A. Schilling, New fermionic formula for unrestricted Kostka polynomials, J. Comb. Theor. Ser. A, 113 (2006) 1435–1461, arXiv:math.CO/0509194.