

## Comment on “Limits on the Time Variation of the Electromagnetic Fine-Structure Constant...”

In their Letter [1] (also [2]), Srianand *et al.* analysed optical spectra of heavy-elements in 23 absorbers along background quasar sight-lines, reporting limits on variations in the fine-structure constant,  $\alpha$ :  $\Delta\alpha/\alpha = (-0.06 \pm 0.06) \times 10^{-5}$ . This would contradict previous evidence [e.g. 3, 4] for a smaller  $\alpha$  in the absorption clouds compared to the laboratory:  $\Delta\alpha/\alpha = (-0.57 \pm 0.11) \times 10^{-5}$  [5]. Here we demonstrate basic flaws in the analysis of [1] using the same data and absorption profile fits.

For each absorber,  $\Delta\alpha/\alpha$  is measured using a  $\chi^2$  minimization of a multiple-component Voigt profile fit to the absorption profiles of several transitions. The column densities, Doppler widths and redshifts defining the components are varied iteratively until the decrease in  $\chi^2$  between iterations falls below a specified tolerance,  $\Delta\chi_{\text{tol}}^2$ . In our approach, we simply add  $\Delta\alpha/\alpha$  as an additional free parameter whereas [1] keep it as an external one: for each fixed input value of  $\Delta\alpha/\alpha$  the other, free parameters are varied to minimize  $\chi^2$ . The functional form of  $\chi^2$  implies that, in the vicinity of the best-fitting  $\Delta\alpha/\alpha$ , the ‘ $\chi^2$  curve’ – the value of  $\chi^2$  as a function of  $\Delta\alpha/\alpha$  – should be near parabolic and smooth. That is,  $\Delta\chi_{\text{tol}}^2$  should be  $\ll 1$  to ensure that fluctuations on the  $\chi^2$  curve are also  $\ll 1$ . This is crucial for deriving the  $1\text{-}\sigma$  uncertainty in  $\Delta\alpha/\alpha$  from the width of the  $\chi^2$  curve at  $\chi_{\text{min}}^2 + 1$ .

However, none of Srianand *et al.*’s  $\chi^2$  curves – figure 2 in [1], 14 in [2] – are smooth at the  $\ll 1$  level; many fluctuations exceed unity. Two examples are reproduced in Fig. 1. The fluctuations can only be due to failings in the  $\chi^2$  minimization: even when [2] fit *simulated* spectra (their figure 2) jagged  $\chi^2$  curves result, leading to a strongly non-Gaussian distribution of  $\Delta\alpha/\alpha$  values and a large range of  $1\text{-}\sigma$  uncertainties (their figure 6). Clearly, these basic flaws in the parameter estimation will yield underestimated uncertainties and spurious  $\Delta\alpha/\alpha$  values.

To demonstrate these failings, we apply the *same profile fits* to the *same data* but with a robust  $\chi^2$  minimization. The spectra were kindly provided by B. Aracil who confirmed that the wavelength and flux arrays are identical to those in [1]. For each absorber, the best-fitting profile parameters of [2] were treated as first guesses in our  $\chi^2$  minimization procedure (detailed in [4]). The relationships between the Doppler widths of corresponding velocity components in different transitions were also the same, as were the relevant atomic data. The relative tolerance for halting the  $\chi^2$  minimization was  $\Delta\chi_{\text{tol}}^2/\chi^2 = 2 \times 10^{-7}$ . All absorbers yield smooth  $\chi^2$  curves in new our analysis; Fig. 1 shows two examples.




FIG. 1: Example  $\chi^2$  curves from our minimization (circles) and that of [1] (triangles). Fluctuations in the latter indicate failings in the minimization. Points and error-bars indicate best-fitting values and  $1\text{-}\sigma$  uncertainties; for our curves  $\Delta\alpha/\alpha$  was a free parameter. Note the different vertical scales: left-hand scales for our curves, right-hand scales for [1].

By-products of this analysis are revised values of  $\Delta\alpha/\alpha$  and  $1\text{-}\sigma$  errors. We find 14 of the 23  $\Delta\alpha/\alpha$  values deviate by  $> 0.3 \times 10^{-5}$  from those of [1]. Moreover, the errors are almost always larger, typically by a factor of  $\sim 3$ . The formal weighted mean over the 23 absorbers becomes  $\Delta\alpha/\alpha = (-0.44 \pm 0.16) \times 10^{-5}$  but the scatter in the values is well beyond that expected from the errors. This probably arises from many sources, including overly simplistic profile fits (see [6]). Allowing for additional, unknown random errors by increasing the error-bars to match the scatter (i.e.  $\chi_{\nu}^2 = 1$  about the weighted mean), a more conservative result from the data and fits of [1] is  $\Delta\alpha/\alpha = (-0.64 \pm 0.36) \times 10^{-5}$  – a 6-fold larger uncertainty than quoted by [1]. We conclude that the latter offers no stringent test of previous evidence for varying  $\alpha$ ; this must await a future, extensive statistical approach.

M. T. Murphy,<sup>1,2</sup> J. K. Webb,<sup>3</sup> V. V. Flambaum<sup>3</sup>

<sup>1</sup>Centre for Astrophysics & Supercomputing, Swinburne University of Technology, Victoria 3122, Australia

<sup>2</sup>Institute of Astronomy, University of Cambridge Madingley Road, Cambridge CB3 0HA, UK

<sup>3</sup>School of Physics, University of New South Wales Sydney, NSW 2052, Australia

- 
- [1] R. Srianand *et al.*, Phys. Rev. Lett. **92**, 121302 (2004).
  - [2] H. Chand *et al.*, Astron. Astrophys. **417**, 853 (2004).
  - [3] J.K. Webb *et al.*, Phys. Rev. Lett. **82**, 884 (1999).
  - [4] M.T. Murphy *et al.*, Mon. Not. Roy. Soc. **345**, 609 (2003).
  - [5] M.T. Murphy *et al.*, Lecture Notes Phys. **648**, 131 (2004).
  - [6] M.T. Murphy *et al.*, arXiv:astro-ph/0612407.

## Discussion of Srianand *et al.*'s Reply to our Comment

Our *Comment* (arXiv:0708.3677) sought to demonstrate that the results of Srianand *et al.* [1] (also [2]) were not robust and were based on a measurement technique which failed in a fundamental way. The numbers emerging from the failed algorithm are meaningless, as discussed again below; they cannot even be considered approximately correct. Indeed, when we apply the same measurement technique (without the failure) to the *same data*, using the *same profile fits* as [1, 2], we find very different values of  $\Delta\alpha/\alpha$  and errors which are typically a factor of  $\sim 3$  larger. We present much more detail of that analysis in [6].

In their *Reply* (arXiv:0711.1742), Srianand *et al.* state or argue several points, all of which we dismiss below either because they are demonstrably incorrect or because they rely on a flawed application of simple statistical arguments. In order of importance:

**Point (i):** Despite demonstrating in our *Comment* that the measurement procedure used by [1, 2] failed, their *Reply* argued that indeed their “procedure is robust as shown in” [2]. Much of the discussion and Fig. 1 in our *Comment* – and, indeed, Srianand *et al.*'s own figures (2 in [1], 14 in [2]) – demonstrate the precise opposite, i.e. that large fluctuations on the  $\chi^2$  curves are clearly present. This means that at any given point on a  $\chi^2$  curve (i.e. for a given input value of  $\Delta\alpha/\alpha$ ), the true minimum of  $\chi^2$  cannot have been reached. Therefore, the *true*  $\chi^2$  curve must lie entirely *beneath* that derived by [1, 2] for most absorbers. That is to say, the  $\chi^2$  curves of [1, 2] simply cannot be used to infer any values of  $\Delta\alpha/\alpha$  or their  $1\text{-}\sigma$  errors whatsoever. Nor can values of  $\Delta\alpha/\alpha$  (or uncertainties) one chooses to infer from them even be considered ‘approximately correct’ in any meaningful way. These basic and fundamental aspects of  $\chi^2$  fitting cannot be overemphasized:  **$\chi^2$  curves with large fluctuations provide no meaningful measurement of any kind.**

In their *Reply*, Srianand *et al.* also “point out that fluctuations in  $\chi^2$  curves get indeed smoothed after a large number of iterations but the results from the first and last iterations are found to be very similar”. It must again be strongly emphasized that fluctuations on a  $\chi^2$  curve indicate nothing but the simple fact that  $\chi^2$  has not been reduced to its true minimum value at some, if not all points on the curve. One can not “smooth” these fluctuations in any way, not by averaging many “iterations” together (as may be implied by the above statement), nor by fitting a parabolic or polynomial line through the  $\chi^2$  curve as one would fit a model to noisy data. It is not “noise” at all, but just an indication that the algorithm for reducing  $\chi^2$  has failed. One simply has to identify the coding error, bug or mis-use of the algorithm which is preventing  $\chi^2$  from reaching its minimum.

Srianand *et al.* [1, 2] also argue that their measure-

ment procedure is robust based on measurements using simulated absorption systems and these are referred to again in their *Reply*. However, as we pointed out in our *Comment* (and in more detail in [6]), those simulations actually demonstrate the precise opposite: strong fluctuations even appear in the  $\chi^2$  curves for these simulations (figure 2 in [2]). This leads to spurious  $\Delta\alpha/\alpha$  values: figure 6 in [2] shows the results from 30 realizations of a simulated single-component Mg/Fe II absorber. At least 15  $\Delta\alpha/\alpha$  values deviate by  $\geq 1\sigma$  from the input value; 8 of these deviate by  $\geq 2\sigma$  and 4 by  $\geq 3\sigma$ . There is even a  $\approx 5\text{-}\sigma$  value. The distribution of  $\Delta\alpha/\alpha$  values should be Gaussian in this case but these outliers demonstrate that it obviously is not. The  $\chi^2$  fluctuations also cause the uncertainty estimates from the different realizations to range over a factor of  $\approx 4$  even though all had the same simulated spectral signal-to-noise and input profile fitting parameters. None of these problems arise in our own simulations of either single- or multiple-component systems (see [4] for detailed discussion).

**Point (ii):** We used smaller error spectra in our *Comment* compared to the original analysis of [1, 2], as described in detail in [6]. Although Srianand *et al.* point out this fact in their *Reply*, they do not discuss its import. We do so in [6]. To summarize: The main argument in our *Comment* is that the error bars of [1, 2] are underestimated. The fact that we used *smaller* error arrays and still found much *larger* errors on  $\Delta\alpha/\alpha$  than [1, 2] only emphasizes this argument more. Using somewhat larger error arrays would simply lead to somewhat larger uncertainties on  $\Delta\alpha/\alpha$  and change the actual measured values of  $\Delta\alpha/\alpha$  negligibly. Thus, “Point 2” in the *Reply* actually reinforces our conclusion that the data and fits do not offer a stringent test of previous evidence for a varying  $\alpha$ .

**Point (iii):** The *Reply* argues that many of our revised values of  $\Delta\alpha/\alpha$  “match” the original values of [2] “at  $\leq 1\sigma$  level” and, therefore, that the original results are robust. Three simple points can be made here; the first two (1 & 2) are practical while the third (3) is a more important general one:

1. A matching criterion of “ $\leq 1\sigma$ ” is used for each value of  $\Delta\alpha/\alpha$ , but it isn't clear if “ $\sigma$ ” is the original uncertainty from [1, 2] or our revised value which is typically larger by a factor of  $\sim 3$ . More importantly, we cannot replicate the number of systems which supposedly “match”, i.e. 16. Using the uncertainty values from [1, 2] as “ $\sigma$ ” we find 11 ‘matching’ values of  $\Delta\alpha/\alpha$ . Using our new uncertainties gives 13 ‘matches’. Finally, adding the two uncertainty values in quadrature and treating that as “ $\sigma$ ” gives 14 systems.
2. Even if the old and new  $\Delta\alpha/\alpha$  values were statistically independent, how many values would one expect to “match” given the general distribution of values and uncertainties in both measurement sets?

Whatever that number, even more values would “match” when we consider that the new and old values are in fact correlated. Therefore, at least some of the 16 ‘matches’ the *Reply* cites cannot be argued to bolster a case for the robustness of either the old or new values. Only if one found many more matches than one expects, given the two sets of values and their errors, would there be some case for that.

3. The more important general point is that the usual “1  $\sigma$ ” threshold is *meaningless* here because the new and old  $\Delta\alpha/\alpha$  values are not independent in any respect. When such non-independent data are being compared, it should actually be *very few values* (not 32% as for truly independent values) which would deviate by more than one formal standard deviation. Only if the deviation was very small with respect to the (old or new) 1- $\sigma$  uncertainties could one claim that the faults in the  $\chi^2$  minimization algorithm had negligible importance and that the original results were robust. The large deviations we observe – 14 of the new 23  $\Delta\alpha/\alpha$  values deviate from the old by  $> 0.3 \times 10^{-5}$  – clearly imply that the old results were not robust at all.

**Point (iv):** The *Reply* discusses how two  $\Delta\alpha/\alpha$  values

in our revised set deviate by more than 4  $\sigma$  (presumably with respect to  $\Delta\alpha/\alpha = 0$ ) and Srianand *et al.* choose to remove them from the sample. How can one decide which systems are best to remove based only on the parameter of interest, in this case the value (or significance) of  $\Delta\alpha/\alpha$ ? This is obviously a very biased selection method. In our *Comment*, this is why we chose to increase the error bars on all  $\Delta\alpha/\alpha$  values to match the observed scatter. Although itself not ideal, this is at least an unbiased procedure. Our main point remains whatever procedure one chooses: the constraints obtained from the data and fits of [1, 2] can not be regarded as robust, nor are they a stringent test of previous evidence for varying  $\alpha$ .

**Point (v):** Srianand *et al.*’s *Reply* states that their “procedure takes into account the differences in spectral resolution in different settings ... while this is not the case with VPFIT”. This is simply incorrect, as is easily verified by reading the VPFIT manual at <http://www.ast.cam.ac.uk/~rfc/vpfit.html> [for the current versions (8 & 9), this is mentioned in the first few lines of the introduction]. VPFIT has *always* has this feature. Differences in resolution between different portions of input spectra were taken into account in all previous analyses of Keck/HIRES spectra in [3, 4, 5] using VPFIT.