

On dynamical smash product

A. I. Mudrov

St.-Petersburg Department of Steklov Mathematical Institute,
Fontanka 27, 191023 St.-Petersburg, Russia,

and

Department of Mathematics, University of York, YO10 5DD, UK

e-mail: mudrov@pdmi.ras.ru

Abstract

In the theory of dynamical Yang-Baxter equation, with any Hopf algebra H and a certain H -module and H -comodule algebra L (base algebra) one associates a monoidal category. Given an algebra A in that category, one can construct an associative algebra $A \rtimes L$, which is a generalization of the ordinary smash product when A is an ordinary H -algebra. We study this "dynamical smash product" and its modules induced from one-dimensional representation of the subalgebra L . In particular, we construct an analog of the Galois map $A \otimes_{AH} A \rightarrow A \otimes H^*$.

1 Introduction

This work is motivated by a recently discovered connection between equivariant deformation quantization and the dynamical Yang-Baxter equation (YBE), see [DM1, AL, KMST, EEM]. One goal of this paper is to give a deeper algebraic insight to constructions arising along those lines, in particular, to non-associative algebras participating in quantization, [DM1]. To a certain extent, this work is also about non-commutative "principal bundles". Such a point of view was already suggested in [DM1], and here we develop that idea one step further.

Conventionally, the function algebra of a non-commutative principal fiber bundle is understood as Hopf-Galois extension of the subalgebra of invariants. What is special about our point of view as compared to the standard one is that we do not assume the principal bundle

to be an algebra. The necessity of such a generalization is dictated by the fact that for a great deal of interesting examples the principal bundle cannot be quantized in the class of associative algebras. Rather, we consider functions on a principal bundle as sections of the associated vector bundle whose fiber is the function algebra on the structure group (viewed as the co-regular module). In this sense it can be quantized for important homogeneous spaces of simple groups. Thus we view non-commutative principal bundles as modules over non-commutative quotient spaces subject to a Galois-like condition, which we introduce and study in this paper.

The term "dynamical" takes its origin in the mathematical physics literature, where the corresponding generalization of the YBE appears in connection with integrable conformal field theories. Contrary to the purely algebraic ordinary YBE, its dynamical analog is differential in the classical version and difference-like in the quantum. The theory of dynamical YBE provides a natural environment for deformation quantization of G -spaces alongside with vector bundles over them. In this geometrical interpretation the set of dynamical variables (more precisely, its invariant classical points) appears to be the moduli space of quantized principal bundles. This moduli space admits an action of the "Picard group" of the associated linear bundles.

The ground object of our theory is the so called dynamical base, which is the pair (H, L) of a Hopf algebra H and a certain H -algebra L . Algebraically, L is a braided-commutative algebra in the quasitensor category of modules over the double of H . Our present interest is centered on a dynamical algebra over (H, L) , which is an algebra in a certain monoidal category rather than in the usual sense. In the simplest non-trivial case $L = H$, the dynamical algebras can be described as follows. Consider an associative algebra B containing H as a subalgebra. We endow B with the structure of left H -module under the adjoint action. Suppose there exists an H -submodule $A \subset B$ freely generating B over H . Then A is a dynamical algebra and B is a generalization of the smash product of A and H . A similar description is valid for general L , and we reserve for B the notation $A \rtimes L$.

For every dynamical base (H, L) we define and study a homomorphism onto a base (H', L') with commutative H' and L' , which reduction can be accompanied with a reduction of a dynamical algebra A ; the reduced dynamical algebra is a certain submodule in A . We study the reduced case in detail and classify all reduced dynamical bases. This can serve to a rough classification of dynamical bases and dynamical algebras.

With any H -invariant character χ of the algebra L we relate an associative algebra A_χ^L and a pair of right and left A_χ^L -modules A_χ and ${}_{\bar{\chi}}A$ on the same underlying vector space

A. Here $\tilde{\chi}$ is another character canonically constructed out of χ . These modules play the role of non-commutative left and right "principal bundles". They are constructed as $A \rtimes L$ -modules via right and left induction from the one dimensional representations χ and $\tilde{\chi}$ of the subalgebra $L \subset A \rtimes L$; then A_χ^L is realized as the algebra of $A \rtimes L$ -endomorphisms of A_χ and ${}_{\tilde{\chi}}A$. This is the algebraic meaning of A_χ^L , which appeared in [DM1] as the deformed function algebra on a coadjoint orbit (conjugacy class) of a simple complex group. In the special case of [DM1] the vector space A_χ^L coincides with H -invariants in A .

For every quadruple (H, L, A, χ) we construct a B -bimodule $A \otimes_\chi H^*$ on the tensor product $A \otimes H^*$. Here H^* is the dual Hopf algebra to H . We introduce a B -bimodule map $A_\chi \otimes_{A_\chi^H} {}_{\tilde{\chi}}A \rightarrow A \otimes_\chi H^*$, where $A_\chi^H \subseteq A_\chi^L$ stands for the subalgebra of invariants. This is an analog of the Galois map for dynamical algebras and it turns into the ordinary Galois map when A is an ordinary H -algebra. We call the pair (A, A_χ^H) Galois extension if this map is an isomorphism. Restriction of the dynamical Galois map to $A_\chi^L \otimes_{A_\chi^H} A_\chi^L$ coincides with the usual Galois map for the extension A_χ^L/A_χ^H .

Finally, for every invariant character χ in the left regular representation we construct a Morita context between $A \rtimes H$ and A_χ^H . We construct it also between $A \rtimes L$ and A_χ^L , out of a χ -generator $e \in L$ in the left regular representation of L .

Throughout the paper H is assumed to be a finite dimensional Hopf algebra over a field \mathbb{k} . Some of our results can be generalized to a certain class of infinite dimensional Hopf algebras that are close to quantum groups, but we make this simplifying assumption by technical reasons in order to avoid numerous stipulations.

Acknowledgements This research is supported by the EPSRC grant C511166 and partly supported by the RFBR grant 06-01-00451. The author is grateful to the Max-Planck Institute für Mathematik in Bonn for hospitality.

2 Preliminaries (dynamical base)

Fix a finite dimensional Hopf algebra H over a field \mathbb{k} with the comultiplication Δ , the counit ε and the antipode γ . The antipode is invertible in a finite dimensional Hopf algebra, and we denote its inverse by $\bar{\gamma}$. We shall use the same notation for the structure operations in the dual Hopf algebra H^* ; this will not cause any confusion, as elements of H and H^* will be always explicitly stated.

We denote by $\mathfrak{D}(H)$ the double Hopf algebra of H and we choose its realization in the

form $H \bowtie H_{op}^*$, where op designates the opposite multiplication. In other words, the double is constructed on the tensor product of $H \otimes H_{op}^*$, where the factors are embedded as subalgebras. Given a basis $\{h_i\} \subset H$ and its dual $\{\eta^i\} \subset H^*$, the element $\Theta := \sum_i \eta^i \otimes h_i \in \mathfrak{D}(H)$ is a quasitriangular structure on $\mathfrak{D}(H)$; it is independent on the choice of basis. The Yang-Baxter equation on Θ is equivalent to the commutation relations between elements of H and of H_{op}^* and thus determines the algebra structure of $\mathfrak{D}(H)$.

The symbol \otimes stands for the tensor product over the ground field \mathbb{k} ; the tensor product over other rings will be indicated explicitly. We adopt the following Sweedler-like convention denoting symbolically the coproducts and coactions:

$$\Delta(h) = h^{(1)} \otimes h^{(2)}, \quad \Delta(\eta) = \eta^{(1)} \otimes \eta^{(2)},$$

for $h \in H$ and $\eta \in H^*$. Thus we use different indication for the Sweedler H - and H^* -components. For a left H -comodule V and a right H^* -comodule W we use the symbolic presentation

$$\delta(v) = v^{(1)} \otimes v^{[\infty]}, \quad \delta(w) = w^{[0]} \otimes w^{(1)},$$

where $v \in V$ and $w \in W$. We also suppress summation when labelling tensor factors of tensor objects and denote them simply as $\Theta = \Theta_1 \otimes \Theta_2$ for $\Theta \in \mathfrak{D}(H)^{\otimes 2}$ etc.

We denote by \leftarrow and \rightarrow the right and, respectively, left co-regular actions of H on H^* : Explicitly they are given by

$$h \rightarrow \eta = \eta^{(1)} \langle \eta^{(2)}, h \rangle, \quad \eta \leftarrow h = \langle \eta^{(1)}, h \rangle \eta^{(2)}, \quad (2.1)$$

for $h \in H$ and $\eta \in H^*$. Here $\langle \cdot, \cdot \rangle$ designates the canonical pairing between H^* and H .

Recall that a Yetter-Drinfeld (YD) module V over H is a left H -module and left H -comodule obeying

$$\delta(h \triangleright v) = h^{(1)} v^{(1)} \gamma(h^{(3)}) \otimes h^{(2)} \triangleright v^{[\infty]}$$

for all $v \in V$ and $h \in H$; the symbol \triangleright stands for the action of H on V . Such modules form a quasitensor category, which for finite dimensional H is isomorphic to the category of $\mathfrak{D}(H)$ -modules.

Recall that an H -module algebra B is an algebra in the monoidal category of H -modules. In other words, the multiplication $B \otimes B \rightarrow B$ is supposed to be equivariant. Similarly, an H -comodule algebra is an algebra in the category of H -comodules. That means that the coaction is an algebra homomorphism. An algebra in the YD category is simultaneously a YD module, H -module algebra and H -comodule algebra. In the finite dimensional case under consideration, a YD algebra is a module algebra over $\mathfrak{D}(H)$.

Every YD module V is equipped with an H -equivariant permutation $\tau_{V,W}: V \otimes W \rightarrow W \otimes V$ with every H -module W . The permutation acts by the assignment $\tau_{V,W}(v \otimes w) = v^{(1)} \triangleright w \otimes v^{[\infty]}$ for all $v \in V$ and $w \in W$. A YD-algebra is called braided-commutative if its multiplication is stable under this permutation.

The following definition is the key ingredient of the theory of dynamical Yang-Baxter equation.

Definition 2.1. A *dynamical base* (or simply base) is a pair (H, L) of a Hopf algebra H and a braided commutative YD algebra L .

We call L an H -base algebra or simply a base algebra when H is clear from the context. We always assume L to be unital. Here we write down, for the reader's convenience, the defining axiom of base algebra L :

1. L is a left H -module algebra under the action $\triangleright: H \otimes L \rightarrow L$,
2. L is a left H -comodule algebra under the coaction $\delta: L \rightarrow H \otimes L$.
3. the action and coaction satisfy the YD condition $\delta(h \triangleright \lambda) = h^{(1)} \lambda^{(1)} \gamma(h^{(3)}) \otimes h^{(2)} \triangleright \lambda^{[\infty]}$ for all $h \in H, \lambda \in L$.
4. the braided commutativity condition $\lambda \mu = (\lambda^{(1)} \triangleright \mu) \lambda^{[\infty]}$ for all $\lambda, \mu \in L$ holds true for all $\lambda, \mu \in L$.

The above definition works for arbitrary Hopf algebras. for finite dimensional H a base algebra L is precisely a module algebra over the double $\mathfrak{D}(H)$ satisfying the braided commutativity condition. In terms of the universal R-matrix Θ , the H -coaction can be written as

$$\lambda \mapsto \Theta_2 \otimes \Theta_1 \triangleright \lambda = \lambda^{(1)} \otimes \lambda^{[\infty]}, \quad (2.2)$$

for all $\lambda \in L$. The braided commutativity is then reads $\lambda \mu = (\Theta_2 \triangleright \mu)(\Theta_1 \triangleright \lambda)$, where the H -action \triangleright is extended to the action of the double $\mathfrak{D}(H)$.

We also consider the base algebra L as a right comodule algebra over H^* corresponding to the left H -action by duality. The same applies to every H -module (algebra).

A trivial example of base algebra is given by the ground field \mathbb{k} , considered as the trivial H -module and H -comodule. Below we give two more examples of base algebras of which the first is valid not only for finite dimensional but for arbitrary Hopf algebras.

Example 2.2. Consider H as a left H -module algebra with respect to the adjoint action and a left H -comodule algebra under the comultiplication. Then the conditions 3-4 are satisfied, and (H, H) is a dynamical base.

Example 2.3. Let $\{h_i\} \subset H$ be a basis and $\{\eta^i\} \subset H^*$ be its dual basis. Consider H^* as a left H -module algebra with respect to the action (2.1) and a left H -comodule algebra with respect to the coaction $a \mapsto \sum_{i,j} \gamma(h_i)h_j \otimes \eta^i a \eta^j$; this definition does not depend on the choice of basis. Then conditions 3-4 are satisfied, and (H, H^*) is a dynamical base.

Definition 2.4. A homomorphism $(H_1, L_1) \rightarrow (H_2, L_2)$ of dynamical bases is a pair $\varphi: H_1 \rightarrow H_2, \varpi: L_1 \rightarrow L_2$ of mappings, where φ is a Hopf algebra homomorphism and ϖ an algebra homomorphism respecting the actions and coactions:

$$\varpi(h \triangleright \lambda) = \varphi(h) \triangleright \varpi(\lambda), \quad (\varphi \otimes \varpi)(\delta_1(\lambda)) = \delta_2(\varpi(\lambda)), \quad (2.3)$$

for all $h \in H_1$ and $\lambda \in L_1$.

Example 2.5. Consider H as an H -base algebra, as an Example 2.2. For any group-like element $\alpha \in H^*$ the map $\varpi_\alpha: h \mapsto \alpha \rightharpoonup h = h^{(1)} \langle \alpha, h^{(2)} \rangle$ is an algebra automorphism. It is easy to check that the pair $(\text{id}, \varpi_\alpha)$ an automorphism of dynamical base (H, H) .

Suppose that (H_1, L_1) is a dynamical base, L_2 is a YD algebra over H_2 , and the pair of homomorphisms (φ, ϖ) obeys (2.3). If ϖ is onto, then L_2 is braided commutative, and hence (H_2, L_2) is a dynamical base. Then the pair (φ, ϖ) is a homomorphism of dynamical bases.

3 Invariant characters of base algebra

In the present section we study invariant characters of base algebras, by which we understand unital H -equivariant homomorphisms of L to the ground field regarded as the trivial H -module. We are especially interested in the situation when the corresponding one-dimensional representation is realized as a left ideal in L , e.g. when L is finite dimensional semisimple.

Consider a unital homomorphism $\chi: L \rightarrow \mathbb{k}$ of algebras. We call it invariant character if $\chi(h \triangleright \lambda) = \varepsilon(h)\chi(\lambda)$ for all $h \in H$ and $\lambda \in L$. Note that for $L = H$ from Example 2.2 any character is invariant. On the contrary, the H -base algebra H^* from Example 2.3 has no invariant characters, unless $\dim H = 1$. Denote by \hat{L} the set of invariant characters of L .

The set $\hat{H} \subset H^*$ is a group, with the identity ε and the inverse $\bar{\alpha} = \alpha \circ \gamma = \alpha \circ \bar{\gamma}$; here γ is the antipode in H . The group \hat{H} naturally acts on L on the right by $\lambda \cdot \alpha = (\alpha \otimes \text{id}) \circ \delta(\lambda)$. This action induces a left action on \hat{L} by $(\alpha \cdot \chi)(\lambda) = (\alpha \otimes \chi) \circ \delta(\lambda)$.

Every invariant character defines a homomorphism $\iota_\chi: L \rightarrow H$ through the mapping $\lambda \mapsto \lambda^{(1)}\chi(\lambda^{[\infty]})$ for all $\lambda \in L$. It amounts to a homomorphism of H -bases $(H, L) \rightarrow (H, H)$ identical on H .

For any invariant character $\chi \in \hat{L}$ we call $e \in L$ a left χ -generator if $Le = \chi(L)e$. Let L^χ denote the vector space of left χ -generators. In the special case $L = H$ and $\chi = \varepsilon$, it coincides with the space \int_l of left integrals. By construction, L^χ is a two-sided ideal in L .

Lemma 3.1. *The vector space L^χ is H -invariant.*

Proof. Indeed, for all $h \in H$, $\lambda \in L$, and $e \in L^\chi$:

$$\lambda(h \triangleright e) = (h^{(2)}\bar{\gamma}(h^{(1)}) \triangleright \lambda)(h^{(3)} \triangleright e) = h^{(2)} \triangleright ((\bar{\gamma}(h^{(1)}) \triangleright \lambda)e) = \chi(\lambda)\varepsilon(\bar{\gamma}(h^{(1)})h^{(2)}) \triangleright e.$$

The last expression is equal to $\chi(\lambda)h \triangleright e$. □

We call a character $\chi \in \hat{L}$ projective if the corresponding one dimensional L -module is projective. In other words, if $L^\chi \not\subset \ker \chi$ or, equivalently, if L^χ contains an idempotent. Below we show that $\dim L^\chi = 1$ if the character χ is projective.

Next we introduce an operation on invariant characters through a Drinfeld element in the double of H , which is defined for any finite dimensional quasitriangular Hopf algebra, [Dr2]. Set

$$\vartheta = \Theta_1\gamma^{-2}(\Theta_2), \quad \bar{\vartheta} = \Theta_1\gamma(\Theta_2).$$

It is known from [Dr2] that $\bar{\vartheta} = \vartheta^{-1}$. Conjugation with ϑ implements the squared antipode in $\mathfrak{D}(H)$: $\vartheta h \vartheta^{-1} = \gamma^2(h)$. The element ϑ satisfies the equation

$$\Theta_{21}\Theta_{12}\Delta(\vartheta) = (\vartheta \otimes \vartheta). \tag{3.4}$$

It follows from (3.4) that the action of ϑ on L implements an algebra automorphism $\lambda \mapsto \vartheta \triangleright \lambda$. That is a consequence of braided-commutativity of L , so that Θ -s in the left hand side of (3.4) vanish on L . The automorphism ϑ commutes with the action of $\hat{H} \subset \mathfrak{D}(H)$, as the square antipode is identical on \hat{H} .

For every $\chi \in \hat{L}$ define the adjoint character $\tilde{\chi}$ by the assignment $\tilde{\chi}: \lambda \mapsto \chi(\bar{\vartheta} \triangleright \lambda)$. It is invariant by virtue of $\vartheta h = \gamma^2(h)\vartheta$ for all $h \in H$. Since the action of ϑ commutes with the action of \hat{H} , we have $\alpha \cdot \tilde{\chi} = \widetilde{\alpha \cdot \chi}$ for all $\alpha \in \hat{H}$. We call a character self adjoint if $\tilde{\chi} = \chi$. Next we characterize the mapping $\chi \rightarrow \tilde{\chi}$ in terms of a certain action of L on H^* .

Define a left L -action on H^* by setting $\lambda \xrightarrow{\chi} \eta := \eta \leftarrow (\gamma \circ \iota_\chi)(\lambda)$ for all $\lambda \in L$ and $\eta \in H^*$.

Lemma 3.2. *For all $\lambda \in L$ the element $\lambda^{[0]} \xrightarrow{\chi} \lambda^{(1)} \in H^*$ is the scalar multiple $\tilde{\chi}(\lambda)1$.*

Proof. Consider L as an algebra over the double $\mathfrak{D}(H)$. Evaluating $\lambda^{[0]} \xrightarrow{\chi} \lambda^{(1)}$ at an arbitrary element $h \in H$ we find

$$\langle \lambda^{[0]} \xrightarrow{\chi} \lambda^{(1)}, h \rangle = \chi(\Theta_1 \Theta_{2'} \triangleright \lambda) \langle \Theta_{1'}, \gamma(\Theta_2) h \rangle = \chi(\Theta_1 \gamma(\Theta_2) h \triangleright \lambda) = \chi(\bar{\vartheta} h \triangleright \lambda) = \chi(\bar{\vartheta} \triangleright \lambda) \varepsilon(h).$$

Here we use the dashed subscripts to distinguish between different copies of Θ . The last equality is obtained by pulling $\bar{\vartheta}$ to the right and taking into account $\bar{\vartheta} h \vartheta = \bar{\gamma}^2(h)$. \square

Similarly to left χ -generators one can define χ -generators under the right regular representation of L : those are elements $e \in L$ satisfying $eL = e\chi(L)$.

Proposition 3.3. *For all $\chi \in \hat{L}$, left χ -generators are right $\tilde{\chi}$ -generators and vice versa.*

Proof. Let $e \in L^\chi$ be a left χ -generator. Then, for all $\lambda \in L$,

$$e\lambda = (e^{(1)} \triangleright \lambda) e^{[\infty]} = \langle \lambda^{(1)}, e^{(1)} \rangle \lambda^{[0]} e^{[\infty]} = \langle \lambda^{[0]} \xrightarrow{\chi} \lambda^{(1)}, e^{(1)} \rangle e^{[\infty]} = \tilde{\chi}(\lambda) \varepsilon(e^{(1)}) e^{[\infty]} = \tilde{\chi}(\lambda) e.$$

In the middle equality we have used the identity $e^{(1)} \otimes \mu e^{[\infty]} = (\gamma \circ \iota_\chi)(\mu) e^{(1)} \otimes e^{[\infty]}$ for $\mu = \lambda^{[0]} \in L$. The next equality to the right is due to Lemma 3.2. This calculation proves that e is a right $\tilde{\chi}$ -generator. To check that every right $\tilde{\chi}$ -generator is a left χ -generator, notice that the opposite algebra L_{op} is a base algebra over H_{op} , cf. [DM2]. The subscript op designates the opposite multiplication. The action of H_{op} on L_{op} is given by the assignment $h \otimes \lambda \mapsto \bar{\gamma}(h) \triangleright \lambda$, and the coaction stays the same as the H -coaction on L . This readily implies that every right $\tilde{\chi}$ -generator is a left χ -generator. \square

4 Base reduction

In the present section we study a reduction procedure for dynamical base. Namely, with any base (H, L) such that $\hat{L} \neq \emptyset$ we associate a dynamical base (H', L') with commutative H' and L' and a base homomorphism $(H, L) \xrightarrow{\phi, \varpi} (H', L')$, which is a surjective on the components. Then we classify reduced dynamical bases.

Proposition 4.1. *Suppose an invariant character $\chi \in \hat{L}$ is projective. Then a) the H -module L^χ is trivial and contained in the center of L , b) $\dim L^\chi = 1$, c) χ is self-adjoint.*

Proof. Suppose there exists a generator $e \in L^\chi$ such that $\chi(e) \neq 0$. We can assume that e is an idempotent and hence $\chi(e) = \tilde{\chi}(e) = 1$, due to Proposition 3.3. If $e' \in L^\chi$ is yet another χ -generator, then the equalities

$$\chi(e + e')e = (e + e')e = (e + e')\tilde{\chi}(e)$$

imply $\chi(e')e = e'$. This proves b).

First of all remark that every H -invariant in L are central. Now c) follows from a) and Proposition 3.3. To prove a) recall from Lemma 3.1 that $L^\chi = \mathbb{k}e$ is an H -module. This module is trivial, as the character χ is H -invariant. This proves a). \square

Remark 4.2. Remark that Propositions 3.3 and 4.1 generalize the well known fact that the vector space of left integrals \int_l in a finite dimensional Hopf algebra is one dimensional, [Sw]. On setting $L = H$ an integral $t \in \int_l$ becomes a left ε -generator a right $\tilde{\varepsilon}$ -generator. However $\dim \int_l = 1$ always for $L = H$, contrary to general base algebra case. For example, consider the algebra $L = \text{Span}\{1, e_i\}_{i=1}^k$ with the nil multiplication on $\text{Span}\{e_i\}_{i=1}^k$. Endowed with the structure of trivial $\mathfrak{D}(H)$ -module, L becomes an H -base algebra. The character $\chi(1) = 1$, $\chi(e_i) = 0$ is invariant and $\dim L^\chi = k$, because L^χ in this example coincides with $\ker \chi$.

Consider the subspace \hat{H}^\perp in H annihilated by all characters of H , that is, the intersection $\hat{H}^\perp = \bigcap_{\alpha \in \hat{H}} \ker \alpha$. It is a bi-ideal, and the quotient H/\hat{H}^\perp is a Hopf algebra. The latter is commutative, and can be viewed as the function algebra $\mathbb{k}^{\hat{H}}$ on the group \hat{H} . Put $H' := H/\hat{H}^\perp$ and denote by ϖ the corresponding projection $H \rightarrow H'$.

From now on we suppose that the set of invariant characters of L is not empty. Let \hat{L}^\perp be the intersection $\bigcap_{\chi \in \hat{L}} \ker \chi$, which is an ideal in L . Set $L' := L/\hat{L}^\perp$ and denote by φ the projection $L \rightarrow L'$. As the characters are unital homomorphisms and $\hat{L} \neq \emptyset$ by assumption, one has $L' \neq \{0\}$. Clearly the algebra L' is commutative.

Proposition 4.3. *The pair (H', L') is a dynamical base, and $(H, L) \xrightarrow{\varphi, \varpi} (H', L')$ is a homomorphism of bases.*

Proof. The ideal \hat{L}^\perp is obviously H -invariant. Moreover, $\hat{H}^\perp L \subset L^\perp$. Indeed, for all $h \in \hat{H}^\perp$ and $\lambda \in L$ one has $\chi(h \triangleright \lambda) = \varepsilon(h)\chi(\lambda) = 0$. Hence $L' = L/\hat{L}^\perp$ is a natural module algebra over H' . By construction, $\varpi(h \triangleright \lambda) = \varphi(h)\varpi(\lambda)$, that is, the projection ϖ is φ -equivariant. Under the coaction $(\varphi \otimes \text{id}) \circ \delta$, the ideal \hat{L}^\perp is also a coideal. Thus, L' is a YD-module algebra over H' and hence a base algebra. This proves the assertion. \square

Definition 4.4. A dynamical base (H, L) is called reduced if $\bigcap_{\chi \in \hat{L}} \ker \chi = \{0\}$. The reduced base (H', L') constructed above is called reduction of (H, L) .

If L' is finite dimensional, then it is semisimple. Otherwise, we can select in L' a finite dimensional subalgebra, which is also a base algebra over H' .

Proposition 4.5. *Suppose that the base (H, L) is reduced and $\bigoplus_{\chi \in \hat{L}} L^\chi \neq \{0\}$. Then $\bigoplus_{\chi \in \hat{L}} L^\chi$ is a base algebra over H .*

Proof. It is sufficient to show that $\bigoplus_{\chi} L^\chi$ is a sub-comodule over H' . Suppose $L^\chi \neq 0$ and let $e_\chi \in L^\chi$ be the idempotent generating L^χ , cf. Proposition 4.1. For all $\alpha \in \hat{H}$ the element $\alpha(e_\chi^{(1)})e_\chi^{[\infty]}$ is the idempotent $e_{\bar{\alpha}\chi}$ generating $L^{\bar{\alpha}\chi}$. This readily implies the statement. \square

Note that $\bigoplus_{\chi} L^\chi$ is not only a subalgebra of L but also a quotient algebra. Indeed, $\bigoplus_{\chi} L^\chi$ is the image under the homomorphism $\lambda \mapsto \sum_{\chi \in \hat{L}} e_\chi \lambda$.

Any finite dimensional reduced base algebra may be regarded as a function algebra on the set \hat{L} . The action of the group \hat{H} on \hat{L} stratifies \hat{L} to a disjoint union of orbits. Restriction of L to every orbit is again a base algebra over H ; the whole L splits into a direct sum of these base algebras.

Below we prove a statement, which gives rise to classification of semisimple reduced dynamical bases. We consider a slightly more general situation: a) the group \hat{H}^* acts transitively on a set X , b) L is the function algebra \mathbb{k}^X , c) the characters of \mathbb{k}^X , which are naturally identified with elements of X , are not necessarily invariant.

Theorem 4.6. *Let G be a finite group, $K \subset G$ its subgroup, and $X = G/K$ the set of left cosets. The base algebra structures on \mathbb{k}^X are parameterized by central idempotents $\pi \in \mathbb{k}K$. The base (G, K, π) is reduced if and only if π is the identity, otherwise the set $\widehat{\mathbb{k}^X}$ of invariant characters is empty. Two bases (G, K_i, π_i) are isomorphic if and only if there exists an automorphism $\varphi \in \text{Aut}(G)$ such that $K_2 = \varphi(K_1)$ and $\pi_2 = \varphi(\pi_1)$.*

Proof. The left action of G on X corresponds to the right action of $\mathbb{k}G$ on \mathbb{k}^X , which gives rise to a left \mathbb{k}^G -coaction. Let $e_\psi \in L$ denote the idempotent corresponding to a point $\psi \in G/K$. The commutative algebra \mathbb{k}^G acts on L by $h \triangleright e_\psi = \langle \pi(\psi), h \rangle e_\psi$ for some $\pi(\psi) \in \mathbb{k}G$. The condition $e_\psi e_\chi = \delta_{\psi\chi} e_\chi$, with $\delta_{\psi\chi}$ being the Kronecker symbol, implies that L is a \mathbb{k}^G -module algebra if and only if $\pi(\psi)$ is an idempotent in $\mathbb{k}G$. This idempotent is the group identity if and only if the character ψ is invariant.

Further, the YD condition is equivalent to G -equivariance of the map $\pi: X \rightarrow G$, that is, to the equality $\pi(\alpha \cdot \psi) = \alpha \pi(\psi) \alpha^{-1}$ for all $\alpha \in G$ and $\psi \in X$. Hence $\pi(\psi)$ is determined by its value $\pi(\psi_0)$ where $\psi_0 \in X$ is the origin. The idempotent $\pi(\psi_0)$ commutes with K , the isotropy subgroup of ψ_0 .

The inclusion $\pi(\psi_0) \in \mathbb{k}K$ is equivalent to $\mathfrak{D}(\mathbb{k}^G)$ -commutativity. Indeed, the equation $e_\psi e_\phi = (e_\psi^{(1)} \triangleright e_\phi) e_\psi^{[\infty]}$ turns into $\phi(e_\psi) e_\phi = \langle \pi(\phi), e_\psi^{(1)} \rangle \phi(e_\psi^{[\infty]}) e_\phi$ for all $\phi, \psi \in X$. This is equivalent to $\phi(e_\psi) = (\pi(\phi) \cdot \phi)(e_\psi)$ for all $\phi, \psi \in X$. That is, to $\phi = \pi(\phi) \cdot \phi$ for all $\phi \in X$. This is the case if and only if $\pi(\psi_0) \in K$.

The part concerning the isomorphism is an easy exercise left to the reader. \square

5 The bialgebra H^χ

In the present section we study certain properties of H -modules in the presence of the base algebra L . We shall heavily rely on these technical results in the sequel. As another application, we associate a bialgebra H^χ with every invariant character $\chi \in \hat{L}$. This bialgebra is constructed as a quotient of H and also is used in what follows.

Let V be a left H -module. For every integer k and ordered pair $(\psi, \chi) \in \hat{L} \times \hat{L}$ we define $V[k, \psi, \chi] \subset V$ to be the vector subspace of ψ -generators in V under the representation $L \xrightarrow{\iota_\chi} H \xrightarrow{\gamma^k} H \rightarrow \text{End}(V)$, that is, $v \in V[k, \psi, \chi] \Leftrightarrow (\gamma^k \circ \iota_\chi)(\lambda)v = \psi(\lambda)v$, for all $\lambda \in L$. Note that this representation is left if k is even and right if k is odd. It is easy to check that $V[k, \psi, \chi]$ is an H -submodule in V . For every $\alpha \in \hat{H}$ and all $\chi \in L$ the H -submodule of weight α is obviously contained in $V[k, \chi, \alpha^{(-1)^k} \cdot \chi]$.

Lemma 5.1. *For all $k \in \mathbb{Z}$ one has*

$$V[k, \psi, \chi] = V[k \pm 1, \chi, \psi], \quad V[k, \psi, \chi] = V[k, \tilde{\psi}, \tilde{\chi}].$$

Proof. For all $v \in V[k, \psi, \chi]$ and all $\lambda \in L$ we find

$$\begin{aligned} \gamma^{k \pm 1}(\lambda^{(1)}) \chi(\lambda^{[\infty]}) v &= \gamma^{k \pm 1}(\lambda^{(1)}) \gamma^k(\lambda^{(2)}) \psi(\lambda^{[\infty]}) v = \psi(\lambda) v, \\ \chi(\lambda^{[\infty]}) \gamma^{k \mp 1}(\lambda^{(1)}) v &= \psi(\lambda^{[\infty]}) \gamma^k(\lambda^{(2)}) \gamma^{k \mp 1}(\lambda^{(1)}) v = \psi(\lambda) v, \end{aligned}$$

where we take the upper sign if k is even and the lower sign if k is odd. The left equality in the second line is obtained using the fact that $V[k, \psi, \chi]$ is H -invariant. The above calculation shows that $V[k, \psi, \chi] \subset V[k \pm 1, \chi, \psi]$ for all $k \in \mathbb{Z}$. This implies the first equality of the proposition.

The right equality follows from the left one, because for all λ the operator

$$\tilde{\psi}(\lambda^{(\infty)}) \gamma^k(\lambda^{(1)}) = \psi(\bar{\vartheta} \triangleright \lambda^{(\infty)}) \gamma^k(\lambda^{(1)}) = \psi((\bar{\vartheta} \triangleright \lambda)^{(\infty)}) \gamma^{k+2}((\bar{\vartheta} \triangleright \lambda)^{(1)})$$

is the scalar multiple $\chi(\bar{\vartheta} \triangleright \lambda) = \tilde{\chi}(\lambda)$ on $V[k, \psi, \chi] = V[k+2, \psi, \chi]$. The right equality is obtained using the presentation (2.2) for the coaction, and the standard fact $(\gamma \otimes \gamma)(\Theta) = \Theta$.

The above calculation proves the inclusion $V[k, \tilde{\psi}, \tilde{\chi}] = V[k, \psi, \chi]$. The inverse inclusion is proved similarly. \square

The above proposition implies that the module $V[k, \psi, \chi]$ actually depends on $k \pmod 2$, and $V[k, \chi, \chi]$ is independent on k at all. In what follows, we reserve the notation $V[\psi, \chi] := V[0, \psi, \chi]$ and $V[\chi] := V[\chi, \chi]$.

Remark 5.2. One can equally consider right H -modules and define the corresponding right and left representations of L . The statement analogous to Lemma 5.1 will be true as well. That can be reduced to the considered case by passing to the dynamical base $(H_{op}, L_{op}, \text{cf. the proof of Proposition 3.3.}$

For every pair V, W of left H modules we find that $V[\phi, \psi] \otimes W[\psi, \chi] \subset (V \otimes W)[\phi, \chi]$. Therefore, $V[\phi, \psi] \otimes W[\psi, \chi] \rightarrow U[\phi, \chi]$ under every H -equivariant mapping $V \otimes W \rightarrow U$.

With every invariant character $\chi \in \hat{L}$ we shall associate a bialgebra H^χ . Let \mathcal{M} be the monoidal category of left H -modules. Denote by $\text{Ann}(V, \chi) \subset H$ the annihilator of $V[\chi]$. Consider the intersection $\Xi_\chi := \bigcap_V \text{Ann}(V, \chi)$ taken over all modules from \mathcal{M} . It is an ideal in H , and the inclusion $V[\chi] \otimes W[\chi] \subset (V \otimes W)[\chi]$ implies that it is a bi-ideal: $\Delta(\Xi_\chi) \subset \Xi_\chi \otimes H + H \otimes \Xi_\chi$. For \mathbb{k} regarded as the trivial H -module, we have $\mathbb{k}[\chi] = \mathbb{k}$, hence Ξ_χ is annihilated by the counit. The quotient $H^\chi = H/\Xi_\chi$ is a bialgebra, and it is a Hopf algebra, provided $\gamma(\Xi_\chi) = \Xi_\chi$. By construction, for every V from \mathcal{M} the representation of H in $V[\chi]$ factors through a representation of H^χ .

Recall that $\hat{H}^\chi \subset \hat{H}$ stands for the isotropy subgroup of the point $\chi \in \hat{L}$.

Proposition 5.3. *Characters \widehat{H}^χ of the bialgebra H^χ constitute a subgroup in \hat{H} isomorphic to \hat{H}^χ .*

Proof. As H^χ is a homomorphic image of H , there is an inclusion $\widehat{H}^\chi \subset \hat{H}$. We have to show that $\widehat{H}^\chi = \hat{H}^\chi$. Let \mathbb{k}_α denote the one dimensional representation of H corresponding to $\alpha \in \hat{H}$. If $\alpha \in \hat{H}^\chi$, then $\mathbb{k}_\alpha[\chi] = \mathbb{k}_\alpha$. Hence $\alpha(\Xi_\chi) = \{0\}$ and $\alpha \in \widehat{H}^\chi$; this proves the inclusion $\hat{H}^\chi \subset \widehat{H}^\chi$. Conversely, if $\alpha \in \widehat{H}^\chi$, then $\alpha(\Xi_\chi) = \{0\}$. For every $\lambda \in L$ the element $\iota_\chi(\lambda) - \chi(\lambda)1 \in H$ belongs to Ξ_χ . Hence $0 = \alpha(\iota_\chi(\lambda)) - \chi(\lambda) = \alpha \cdot \chi(\lambda) - \chi(\lambda)$ for all $\lambda \in L$, that is, $\alpha \in \hat{H}^\chi$. \square

Next we give a detailed consideration to a special case, when the bialgebra H^χ becomes a Hopf algebra. Namely, we assume that the character $\chi \in \hat{L}$ is projective. Let $e_\chi \in \hat{L}$ be the corresponding idempotent. The element $t_\chi := \iota_\chi(e_\chi) \in H$ is also an idempotent and it is central in H , because e_χ is invariant.

Lemma 5.4. *For every left H -module V the submodule $V[\chi]$ has the form $V[\chi] = t_\chi V$.*

Proof. Indeed, if $t_\chi v = v$ for some $v \in V$, then

$$\iota_\chi(\lambda)v = \iota_\chi(\lambda)t_\chi v = \iota_\chi(\lambda e_\chi)v = \chi(\lambda)t_\chi v = \chi(\lambda)v,$$

that is, $t_\chi V \subset V[\chi]$. Conversely, if $\iota_\chi(\lambda)v = \chi(\lambda)v$ for all $\lambda \in L$, then

$$t_\chi v = \iota_\chi(e_\chi)v = \chi(e_\chi)v = v.$$

This proves the inverse inclusion $t_\chi V \supset V[\chi]$. □

Regard H as a left and right L -module via the homomorphism $\iota_\chi: L \rightarrow H$. Then $H[\chi]$ is a subring $t_\chi H t_\chi = H t_\chi \subset H$. It is also isomorphic to the quotient of H by the ideal $(1 - t_\chi)H$, which coincides with the ideal Ξ_χ introduced above.

Corollary 5.5. *The idempotent t_χ is fixed under the antipode γ , and $\gamma(H[\chi]) = H[\chi]$.*

Proof. For all $\lambda \in L$ and all $h \in H[\chi]$ we find

$$\gamma(\iota_\chi(\lambda))h = \chi(\lambda)h, \quad h\bar{\gamma}(\iota_\chi(\lambda)) = h\chi(\lambda).$$

The left equality is a direct consequence of Lemma 5.1, the right one holds true due to Remark 5.2. Setting $h = t_\chi$ and $\lambda = e_\chi$ we get $\gamma(t_\chi)t_\chi = t_\chi$ and $t_\chi\bar{\gamma}(t_\chi) = t_\chi$. From this we conclude that $t_\chi = \gamma(t_\chi)$ and hence $\gamma(H[\chi]) = H[\chi]$. □

Denote by H^χ the algebra $H[\chi]$ and consider t_χ as the projector from H to H^χ .

Theorem 5.6. *The comultiplication*

$$H^\chi \hookrightarrow H \xrightarrow{\Delta} H \otimes H \xrightarrow{t_\chi \otimes t_\chi} H^\chi \otimes H^\chi \tag{5.5}$$

along with the counit $\varepsilon|_{H^\chi}$ and the antipode $\gamma|_{H^\chi}$ makes H^χ a Hopf algebra.

Proof. The comultiplication is obviously an algebra homomorphism. Its coassociativity readily follows from the equality

$$\Delta(t_\chi)(t_\chi \otimes t_\chi) = t_\chi \otimes t_\chi.$$

Let us check this equality. We have for the left-hand side:

$$e_\chi^{(1)}t_\chi \otimes \iota_\chi(e_\chi^{[\infty]}e_\chi) = e_\chi^{(1)}t_\chi \otimes \chi(e_\chi^{[\infty]})t_\chi = t_\chi t_\chi \otimes t_\chi = t_\chi \otimes t_\chi,$$

as required.

Further, from $\varepsilon(t_\chi) = \chi(e_\chi) = 1$ we derive

$$(\varepsilon \otimes \text{id})\left(\Delta(h)(t_\chi \otimes t_\chi)\right) = 1 \otimes ht_\chi = 1 \otimes h, \quad (\text{id} \otimes \varepsilon)\left(\Delta(h)(t_\chi \otimes t_\chi)\right) = ht_\chi \otimes 1 = h \otimes 1$$

for all $h \in H^\chi$. Thus the restriction of ε to H^χ is the counit.

To verify the antipode, notice that for all $h \in H^\chi$

$$\gamma(t_\chi)\gamma(h^{(1)})h^{(2)}t_\chi = \varepsilon(h)\gamma(t_\chi)t_\chi = \varepsilon(h)t_\chi, \quad t_\chi h^{(1)}\gamma(h^{(2)})\gamma(t_\chi) = \varepsilon(h)t_\chi\gamma(t_\chi) = \varepsilon(h)t_\chi.$$

Hence $\gamma|_{H^\chi}$ is the antipode indeed, as t_χ is the unit in $H^\chi \subset H$. This completes the proof. \square

Note that for every character $\alpha \in \hat{H}$ the bialgebras H^χ and $H^{\alpha\chi}$ are isomorphic. The isomorphism is included in the commutative diagram

$$\begin{array}{ccc} H & \rightarrow & H \\ \downarrow & & \downarrow \\ H^\chi & \rightarrow & H^{\alpha\chi} \end{array}$$

of bialgebra homomorphisms, where the vertical arrows are projections and the top arrow is given by the assignment $h \mapsto \bar{\alpha}(h^{(1)})h^{(2)}\alpha(h^{(3)})$, $h \in H$.

6 Dynamical algebras

A dynamical or shifted associative algebra A over the base (H, L) is an H -module equipped with an equivariant map (multiplication) $\ast: A \otimes A \rightarrow A \otimes L$ satisfying the (shifted associativity) condition:

$$\begin{array}{ccccccc} A \otimes L \otimes A & \xrightarrow{\tau_A} & A \otimes A \otimes L & \xrightarrow{\ast} & A \otimes L \otimes L & \xrightarrow{m_L} & A \otimes L \\ \ast \otimes \text{id} \uparrow & & & & & & \parallel \\ A \otimes A \otimes A & \xrightarrow{\text{id} \otimes \ast} & A \otimes A \otimes L & \xrightarrow{\ast} & A \otimes L \otimes L & \xrightarrow{m_L} & A \otimes L \end{array} \quad (6.6)$$

Here m_L is the multiplication in L and τ_A is the H -invariant permutation $L \otimes A \rightarrow A \otimes L$ acting by $\lambda \otimes a \mapsto \lambda^{(1)} \triangleright a \otimes \lambda^{[\infty]}$; the obvious identity maps are suppressed.

The dynamical algebra is called unital if there exists an element 1_A such that $1_A \ast a = a \ast 1_A = a \otimes 1_L$ for all $a \in A$. Like an ordinary algebra, a dynamical algebra can be made unital by extension (see the next section).

Example 6.1. Suppose A is a module algebra over a Hopf algebra. Under its multiplication and subsequent embedding $A \simeq A \otimes 1_L \subset A \otimes L$ it becomes a dynamical algebra over an arbitrary base algebra L . We call such dynamical algebras trivial.

Example 6.2. Suppose A is a module algebra over a Hopf algebra U containing H as a Hopf subalgebra. Suppose $F \in U \otimes U \otimes L$ is a dynamical twist, [DM1]. Then the multiplication $a * b := (F_1 \triangleright a)(F_2 \triangleright b) \otimes F_3$ makes A a dynamical algebra.

Further we define a reduction procedure for dynamical algebras. For any dynamical algebra A over the base (H, L) it provides a dynamical algebra A' over the reduced base (H', L') . Consider the H -submodule $A' \subset A$ annihilated by $\hat{H}^\perp \subset H$. It is straightforward to check that the composition $*': A' \otimes A' \rightarrow A \otimes L \rightarrow A \otimes L'$ of the multiplication restricted to A' with the projection $\varpi: L \rightarrow L'$ takes values in $A' \otimes L'$. Indeed, if $h \in \hat{H}^\perp$, then for all $\chi \in \hat{L}$, and $a, b \in A'$ one has $h \triangleright (\text{id} \otimes \chi)(a * b) = (\text{id} \otimes \chi)(h^{(1)} \triangleright a * h^{(2)} \triangleright b) = 0$. Here we have used H -invariance of the character χ , equivariance of $*$ and the fact that \hat{H}^\perp is a bi-ideal.

Proposition 6.3. *The map $A' \otimes A' \rightarrow A' \otimes L'$ makes A' a dynamical algebra over the base (H', L') .*

Proof. By construction, A' is a module over H' and the multiplication $A' \otimes A' \rightarrow A' \otimes L'$ is H' -equivariant. Restrict the leftmost bottom vertex in (6.6) to $A' \otimes A \otimes A'$ and apply the projection ϖ to the rightmost L . This yields the commutative diagram

$$\begin{array}{ccccccc}
A \otimes L \otimes A' & \xrightarrow{\tau_A} & A \otimes A' \otimes L & \xrightarrow{\varpi} & A \otimes A' \otimes L' & \xrightarrow{*'} & A' \otimes L' \otimes L' \\
\uparrow \scriptstyle * \otimes \text{id} & & & & & & \downarrow \scriptstyle m_{L'} \\
A' \otimes A' \otimes A' & \xrightarrow{\text{id} \otimes *'} & A' \otimes A' \otimes L' & \xrightarrow{*'} & A' \otimes L' \otimes L' & \xrightarrow{m_{L'}} & A' \otimes L'
\end{array}, \quad (6.7)$$

where we suppress the obvious identity morphisms. Let τ'_A denote the canonical permutation $L' \otimes A' \rightarrow A' \otimes L'$. Obviously, $(\text{id} \otimes \varpi) \circ \tau_A = \tau'_A \circ (\varpi \otimes \text{id})$ when restricted to $L \otimes A'$. Thus we can replace all the non-dashed elements of the above diagram by their dashed counterparts. This yields the shifted associativity diagram for A' as a dynamical algebra over (H', L') . \square

Remark that the reduced dynamical algebra A' contains elements of weight α for all $\alpha \in \hat{H}$. Indeed, for every such element $a \in A$ and all $h \in \hat{H}^\perp$ we find $h \triangleright a = \alpha(h)a = 0$, so $a \in A'$.

Every invariant character defines an associative algebra I_χ on the vector space A^H of H -invariants in A . The multiplication in I_χ is set to be

$$a \overset{\chi}{*} b := (\text{id}_A \otimes \chi)(a * b).$$

In applications to deformation quantization, see [DM1], H is a quantized universal enveloping algebra $U(\mathfrak{h})$ acting on a Poisson manifold M . The dynamical algebra A is a "deformation"

of the algebra of functions on M . Then I_χ can be interpreted as a deformation of the function algebra of the corresponding quotient space. It is not realized as a subalgebra in an associative quantization of M . Rather, the function algebra on M is quantized to right and left I_χ -modules A_χ and ${}_{\tilde{\chi}}A$. Our goal is to construct a Galois map $A_\chi \otimes_{I_\chi} {}_{\tilde{\chi}}A \rightarrow A \otimes H^*$ in generalization of the standard situation when $A_\chi = {}_{\tilde{\chi}}A$ is an algebra (A is a trivial dynamical algebra). This Galois map will be also a homomorphism of bimodules over an associative algebra with the underlying vector space $A \otimes L$. This algebra is a generalization of the ordinary smash product, and it is the subject of our interest in the next section.

7 Dynamical smash product

Suppose that A is a dynamical algebra with multiplication $\ast: A \otimes A \rightarrow A \otimes L$. Define the right action of L on $A \otimes L$ by $(a \otimes \lambda)\mu = a \otimes \lambda\mu$. Introduce the associative algebra $A \rtimes L$ on the vector space $A \otimes L$ with multiplication

$$(a \otimes \lambda)(b \otimes \mu) = \left(a \ast (\lambda^{(1)} \triangleright b) \right) (\lambda^{[\infty]} \mu). \quad (7.8)$$

The last factor is an element of L acting on the term within the big parentheses, which is an element of $A \otimes L$. The algebra $A \rtimes L$ is an H -module, and the multiplication is H -equivariant.

Definition 7.1. The H -algebra $A \rtimes L$ is called dynamical smash product of A and L .

The dynamical smash product is a generalization of the usual smash product of an H -module algebra A and the base algebra L , where the dynamical multiplication $A \otimes A \rightarrow A \otimes L$ factors to the usual multiplication $A \otimes A \rightarrow A$ and the embedding of A in $A \otimes L$.

The dynamical smash product $B = A \rtimes L$ is also an L -bimodule, with the left L -action being defined through the right action by the formula $\lambda(b\mu) := (\lambda^{(1)} \triangleright b)(\lambda^{[\infty]}\mu)$, for all $\lambda, \mu \in L$ and $b \in B$. This can be also considered as a compatibility with the action of H . We call such algebras (H, L) -module algebras. In fact, they are algebras in the monoidal category of modules over the bialgebroid $L \# H$, see [Lu, DM2]. Note that B is freely generated by A over L (with respect to either actions). These conditions characterize dynamical algebras.

Proposition 7.2 ([DM1]). *Suppose A is a dynamical algebra over the base (H, L) . Then the dynamical smash product is an (H, L) -module algebra. Conversely, suppose B an (H, L) -module algebra. Suppose that B is freely generated over L by an H -submodule $A \subset B$. Then A is a dynamical algebra.*

Given an (H, L) -module algebra B we can define a new (H, L) -module algebra on the direct sum $B \oplus L$ of H -modules. The multiplication is given by the rule

$$(a \oplus \lambda)(b \oplus \mu) := (ab + \lambda b + a\mu) \oplus \lambda\mu.$$

The vector $(0, 1_L)$ is the identity in this new algebra. If B is a dynamical smash product $A \rtimes L$, so is the new algebra, and the corresponding dynamical algebra $A \oplus \mathbb{k}$ is unital. From now on we assume that all the dynamical algebras under consideration are unital. It is then convenient to identify A with $A \rtimes 1_L \subset B$ and L with $1_A \rtimes L \subset B$. The algebra $B = A \rtimes L$ is generated by A and L , with the commutation relations $ha = (h^{(1)} \triangleright a)h^{(2)}$.

For instance, put $L = H$ and suppose that H is a subalgebra in B . Then B is an H -module algebra with respect to the adjoint action. It is also an (H, H) -module algebra. If B is freely generated over H by an H -submodule A , then A is a dynamical algebra.

Definition 7.3. Suppose $(H, L) \xrightarrow{\varphi, \varpi} (H', L')$ is a homomorphism of dynamical bases and let A and A' be dynamical algebras over (H, L) and (H', L') , respectively. An equivariant mapping $\theta: A \rightarrow A' \otimes L'$ is called morphism of dynamical algebras if the diagram

$$\begin{array}{ccccccc} A \otimes A & \xrightarrow{\theta \otimes \theta} & A' \otimes L' \otimes A' \otimes L' & \xrightarrow{\tau'_{A'}} & A' \otimes A' \otimes L' \otimes L' & \xrightarrow{m_{L'}} & A' \otimes A' \otimes L' \\ \downarrow * & & & & & & \downarrow * \\ A \otimes L & \xrightarrow{\theta \otimes \varpi} & A' \otimes L' \otimes L' & \xrightarrow{m_{L'}} & A' \otimes L' & \xleftarrow{m_{L'}} & A' \otimes L' \otimes L' \end{array} \quad (7.9)$$

is commutative (here we have dropped the obvious identity maps).

Given a homomorphism $(H, L) \rightarrow (H', L')$ of dynamical bases and their module algebras B and B' we define a homomorphism $B \rightarrow B'$ as that of associative algebras which is equivariant under the left Hopf algebra actions and the two-sided base algebra actions. The following proposition is straightforward.

Proposition 7.4. *Morphisms $(H, L, A) \xrightarrow{\varphi, \varpi, \theta} (H', L', A')$ are in natural 1-1 correspondence with equivariant homomorphisms $(\text{id} \otimes m_{L'}) \circ (\theta \otimes \varpi): A \rtimes L \rightarrow A' \rtimes L'$ of associative algebras that preserve the base algebras.*

8 Linear induction from the base algebra

From now on we use the notation B for the dynamical smash product $A \rtimes L$ assuming it to be fixed once and for all. In the present section we study the modules over B that are induced from H -invariant characters of L .

Denote by \triangleleft the right H -action on A defined by $a \triangleleft h := \bar{\gamma}(h) \triangleright a$. We shall use the following symbolic presentation of the dynamical multiplication:

$$a * b = (a \overset{i}{*} b) \rho_i = \ell_i(a \overset{i}{\bullet} b) = \rho_i^{[\infty]}((a \overset{i}{*} b) \triangleleft \rho_i^{(1)}),$$

where $a \overset{i}{*} b$ and $a \overset{i}{\bullet} b$ stand for the A -component while ρ_i and ℓ_i for the L -component. The label $i = 1, 2, \dots$ is used to distinguish between different copies of these operations rather than for summation, which is implicitly understood as in the Sweedler notation.

For a fixed invariant character χ of the algebra L we regard A as the left induced B -module $B \otimes_L \mathbb{k}$ and denote it by A_χ . Similarly, the right B -module $\mathbb{k} \otimes_L B = A$ will be denoted by ${}_\chi A$. The left and right actions of H on A induce the left and right actions of L through the homomorphism $\iota_\chi: L \rightarrow H$. In particular, the right L -action on ${}_\chi A$ is given by $a \otimes \lambda = (\bar{\gamma} \circ \iota_\chi)(\lambda) \triangleright a$. Denote by $A_\chi^L = A_\chi[\chi]$ the space of right χ -generators in A under the left L -action induced by the homomorphism $\iota_\chi: L \rightarrow H$. Due to Lemma 5.1, A_χ^L coincides with the space of right $\tilde{\chi}$ -generators in the right module ${}_{\tilde{\chi}} A$.

We introduce the following H -equivariant bilinear operations on A :

$$a \overset{\times}{*} b := a \overset{1}{*} b \chi(\rho_1), \quad a \overset{\times}{\bullet} b := \chi(\ell_1)(a \overset{1}{\bullet} b).$$

The unit of the dynamical algebra A becomes the neutral element for the operations $\overset{\times}{*}$ and $\overset{\times}{\bullet}$. Note that $\overset{\times}{*}$ is the restriction to $A \subset B$ of the left action on A_χ ; similarly, $\overset{\times}{\bullet}$ is the restriction to $A \subset B$ of the right action on ${}_\chi A$.

We can characterize A_χ^L as the algebra of endomorphisms of the B -modules A_χ and ${}_{\tilde{\chi}} A$.

Theorem 8.1. *The operations $\overset{\times}{*}$ and $\overset{\times}{\bullet}$ coincide on A_χ^L . They make A_χ^L into an associative H -algebra and A into right and left A_χ^L -modules respectively. Moreover,*

$$\text{End}_B(A_\chi) \simeq (A_\chi^L)_{op} \simeq \text{End}_B({}_{\tilde{\chi}} A).$$

Proof. First we demonstrate that the operations in question coincide when restricted to A_χ^L . From the presentation $\ell_1(a \overset{1}{\bullet} b) = \rho_1^{[\infty]}(\bar{\gamma}(\rho_1^{(1)}) \triangleright (a \overset{1}{*} b)) = (\Theta_1 \triangleright \rho_1)(\bar{\gamma}(\Theta_2) \triangleright (a \overset{1}{*} b))$ we find for all $a, b \in A_\chi^L$:

$$a \overset{\times}{\bullet} b = (a \overset{1}{*} b) \triangleleft \iota_{\tilde{\chi}}(\rho_1) = \tilde{\chi}(\Theta_1 \Theta_{1'} \vartheta \triangleright \rho_1) (\bar{\gamma}(\Theta_2) \triangleright a) \overset{1}{*} (\bar{\gamma}(\Theta_{2'}) \triangleright b) = (a \overset{1}{*} b) \tilde{\chi}(\vartheta \triangleright \rho_1) = a \overset{\times}{*} b.$$

The second equality follows from the H -equivariance of the multiplication in B . In the next equality to the right we have used Lemma 5.1 allowing to drop the antipodes.

The second part of the theorem will be proved only for $\overset{\times}{*}$, as the case of $\overset{\times}{\bullet}$ is analogous. Apply the map $(\text{id}_A \otimes \chi)$ to the equality $a(bc) = (ab)c$ in the algebra $B = A \rtimes L$. The

left-hand side immediately gives $a \overset{\times}{*} (b \overset{\times}{*} c)$ for all $a, b, c \in A$. The right-hand side turns into $(a \overset{\times}{*} b) \overset{\times}{*} c$ once $c \in A_\chi^L$, because $a \overset{\times}{*} b \otimes \iota_\chi(\rho_1) \triangleright c = a \overset{\times}{*} b \otimes \chi(\rho_1)c = (a \overset{\times}{*} b) \otimes c$ in this case.

Thus we have shown that $(a \overset{\times}{*} b) \overset{\times}{*} c = a \overset{\times}{*} (b \overset{\times}{*} c)$ whenever $c \in A_\chi^L$. This proves that A_χ^L is an associative algebra and A is a right A_χ^L -module under $\overset{\times}{*}$. Let us show that $A_\chi^L \subset \text{End}_B(A_\chi)$. The algebra L acts on A_χ by multipliers $\chi(\lambda)$, $\lambda \in L$. Hence the right action of A_χ^L commutes with the left action of $L \subset B$. The action of $A \subset B$ on A_χ coincides with the operation $\overset{\times}{*}$. As the operation $\overset{\times}{*}$ is associative once the third argument is from A_χ^L , the right action of A_χ^L commutes with the left action of A . This gives an anti-algebra homomorphism $A_\chi^L \rightarrow \text{End}_B(A_\chi)$. Note that this homomorphism is an embedding, because A is assumed unital as a dynamical algebra and the operation $\overset{\times}{*}$ is also unital.

The Frobenius reciprocity gives $\text{End}_B(A_\chi) \simeq \text{Hom}_L(\mathbb{k}, A_\chi)$. The action of L on A_χ is given by $\iota_\chi(\lambda) \triangleright a$, so $\text{Hom}_L(\mathbb{k}, A_\chi) = A_\chi^L$. We have shown that $A_\chi^L \simeq \text{End}_B(A_\chi)$ as vector spaces. It is easy to see that this is also an anti-isomorphism of algebras. \square

Regarding B as an L -module through the homomorphism $\iota_\chi: L \rightarrow H$, denote by $B_\chi^L := B[\chi]$ the subspace of left χ -generators in B . By Lemma 5.1, it is at the same time the space of right $\tilde{\chi}$ -generators under the right representation $\tilde{\gamma} \circ \iota_{\tilde{\chi}}$. The vector space B_χ^L forms an H -module, and equivariance of the multiplication in B implies that B_χ^L is a subalgebra in B .

Introduce the left and, respectively, right B -module homomorphisms

$$\wp_\chi: B \rightarrow A_\chi, \quad \chi\wp: B \rightarrow \chi A, \quad (8.10)$$

obtained by evaluating χ on the L -factor in the factorizations $AL = B = LA$. It is also a homomorphism of H -modules. When restricted to B_χ^L , the map \wp_χ takes values in A_χ^L .

Proposition 8.2. *Restriction of \wp_χ to B_χ^L defines a surjective homomorphism $B_\chi^L \rightarrow A_\chi^L$ of H -algebras. Through this homomorphism, $\wp_\chi: B \rightarrow A_\chi$ becomes a right B_χ^L -bimodule map.*

Proof. As the base algebra L is unital, $A_\chi^L \subset B_\chi^L$ and the \wp_χ is identical on A_χ^L . Hence it is surjective. For all $a = \sum_i a_i \lambda_i \in B$ and $b = \sum_m b_m \mu_m \in B_\chi^L$ we find

$$\wp_\chi(ab) = \wp_\chi\left(\sum_{i,m} a_i (\lambda_i^{(1)} \triangleright (b_m \mu_m)) \lambda_i^{[\infty]}\right) = \sum_{i,m} (a_i \overset{\times}{*} b_m) \chi(\rho_1 \mu_m \lambda_i) = \wp_\chi(a) \overset{\times}{*} \wp_\chi(b).$$

In the middle equality we used the assumption $b \in B_\chi^L$ as the latter is supposed to be invariant. This proves two things: firstly, \wp_χ is an algebra homomorphism when restricted to B_χ^L and, secondly, it is a homomorphism of right B_χ^L -modules $B \rightarrow A_\chi$. \square

An analogous statement holds true for $\tilde{\chi}\wp$ with replacement of A_χ by $\tilde{\chi}A$, which becomes a left B_χ^L -module.

Proposition 8.3. *The projections \wp_χ and $\tilde{\chi}\wp$ coincide on B^H .*

Proof. An invariant element $a = \sum_i a_i \lambda_i \in B^H$ can be presented in the form

$$\sum_k (\Theta_1 \triangleright \lambda_k) \bar{\gamma}(\Theta_2) \triangleright a_k = \sum_k (\Theta_1 \bar{\gamma}^2(\Theta_2) \triangleright \lambda_k) a_k = \sum_k (\vartheta \triangleright \lambda_k) a_k.$$

This immediately implies $\wp_\chi(a) = \sum_i a_i \chi(\lambda_i) = \tilde{\chi}\wp(a)$. \square

The mapping \wp_χ induces a surjective algebra homomorphism $B^H \rightarrow I_\chi$ of H -invariants.

9 Dynamical Galois extension

In the present section we generalize the notion of Hopf-Galois extension to the case of dynamical algebras. First we recall the standard definition of the Galois map for H^* -comodule algebras.

Let B be a right H^* -comodule algebra. Recall that we use the symbolic notation $b^{[0]} \otimes b^{(1)} = \delta(b)$ for right H^* -coactions. We regard B as a left H -module algebra under the action $h \otimes b \mapsto b^{[0]} \langle b^{(1)}, h \rangle$, $h \in H$ and $b \in B$. Equip the tensor product $B \otimes H^*$ with the left and right B -actions

$$a.(b \otimes \eta).c = abc^{[0]} \otimes \eta c^{(1)},$$

for $a, b, c \in B$, $\eta \in H^*$. These actions commute with each other and make $B \otimes H^*$ a B -bimodule.

Let $B^H \subset B$ denote the subalgebra of H -invariants. Consider the map $\tilde{\Gamma}: B \otimes B \rightarrow B \otimes H^*$ defined by $a \otimes b \mapsto ab^{[0]} \otimes b^{(1)}$. This map is a homomorphism of B -bimodules, and it is factored through a homomorphism $\Gamma: B \otimes_{B^H} B \rightarrow B \otimes H^*$ called the Galois map. The algebra B is called Galois extension of B^H if Γ is a bijection and weak Galois extension if Γ is a surjection.

Lemma 9.1. *The vector subspace $J_\chi = \text{Span}(b\lambda \otimes \eta - b \otimes (\lambda \xrightarrow{\chi} \eta))$, $b \in B$, $\eta \in H^*$, $\lambda \in L$, is a B -sub-bimodule in $B \otimes H^*$.*

Proof. The subspace J_χ is obviously preserved by the left B -action:

$$a.(b\lambda \otimes \eta - b \otimes \lambda \xrightarrow{\chi} \eta) = (ab)\lambda \otimes \eta - (ab) \otimes \lambda \xrightarrow{\chi} \eta,$$

for $a, b \in B$, $\eta \in H^*$, and $\lambda \in L$. The following calculation demonstrates the right B -invariance of J_χ :

$$(b\lambda \otimes \eta - b \otimes \lambda \xrightarrow{\chi} \eta).a = b\lambda a^{[0]} \otimes \eta a^{(1)} - ba^{[0]} \otimes (\lambda \xrightarrow{\chi} \eta)a^{(1)}.$$

The first summand in the right-hand side can be rewritten as $b(\lambda^{(1)} \triangleright a^{[0]})\lambda^{[\infty]} \otimes \eta a^{(1)} = ba^{[0]}\lambda^{[\infty]} \otimes \eta(a^{(1)} \leftarrow \lambda^{(1)})$, while the second as $-ba^{[0]} \otimes \lambda^{[\infty]} \xrightarrow{\chi} (\eta(a^{(1)} \leftarrow \lambda^{(1)}))$. This proves the statement. \square

Remark that the quotient $B \otimes H^*/J_\chi$ is isomorphic to the tensor product $B \otimes_L H^*$ as a left B -module. As a vector space, it coincides with $A \otimes H^*$, and we denote it by $A \otimes_\chi H^*$. Thus the latter becomes a B -bimodule. We denote by P_χ the projection $B \otimes H^* \rightarrow A \otimes_\chi H^*$ along J_χ .

Theorem 9.2. *The map $P_\chi \circ \check{\Gamma}: B \otimes B \rightarrow A \otimes_\chi H^*$ factors through a B -bimodule homomorphism $\Gamma_\chi: A_\chi \otimes_{I_\chi} \check{\chi}A \rightarrow A \otimes_\chi H^*$.*

Proof. First of all, the map $P_\chi \circ \check{\Gamma}$ induces a B -bimodule map $\check{\Gamma}_\chi: A_\chi \otimes_{\check{\chi}} A \rightarrow A \otimes_\chi H^*$. This follows from the equality $(P_\chi \circ \check{\Gamma})(\lambda \otimes \mu) = \chi(\lambda)\check{\chi}(\mu)(1 \otimes 1)$ which holds true for all $\lambda \in L$ by the very construction and for all $\mu \in L$ by Lemma 3.2. Further, for all $a, b \in A$ and $c \in I_\chi$ one has

$$\check{\Gamma}_\chi(a \otimes c \check{\bullet} b) = \check{\Gamma}_\chi(a \otimes c).b = (a \check{*} c^{[0]} \otimes \rho_1 \xrightarrow{\chi} c^{(1)}).b = \check{\Gamma}_\chi(a \check{*} c \otimes 1).b = \check{\Gamma}_\chi(a \check{*} c \otimes b).$$

This implies that the map $\check{\Gamma}_\chi$ factors through a map $\Gamma_\chi: A_\chi \otimes_{I_\chi} \check{\chi}A \rightarrow A \otimes_\chi H^*$, which is a homomorphism of B -bimodules. \square

Definition 9.3. We call Γ_χ the χ -Galois map. The dynamical algebra A is called a (weak) χ -Galois extension of I_χ if Γ_χ is an (epimorphism) isomorphism.

Definition 9.3 reduces to the standard Galois map for an ordinary H -module algebra A upon setting $L = H$ and $\chi = \varepsilon$. Applying our construction to this situation we conclude that the standard Galois map (which is independent on χ) is a morphism of not only A -bimodules, but of $A \rtimes H$ -bimodules too. The two-sided action of $A \rtimes H$ on $A \otimes_{A \rtimes H} A$ is straightforward while it is not obvious on $A \otimes H^*$. For instance, the left H -module $A \otimes H^*$ is a tensor product of the default module A and the module H^* under the action $h.\alpha = \alpha \leftarrow \gamma(h)$. The right H -module $A \otimes H^*$ is a tensor product of the trivial module A and the module H^* with the right action $\alpha.h = \sum_{i,j} (\alpha \eta^i \gamma(\eta^j)) \leftarrow \gamma(h_i h h_j)$. Here $\{h_i\} \subset H$ is a basis and $\{\eta^i\} \subset H^*$

is its dual. Note that for either commutative or co-commutative H this action is expressed through the left co-regular action and antipode by $\alpha.h = \gamma(h)\dashv\alpha$.

Below we display the construction of Galois maps on a commutative diagram of B -bimodules.

$$\begin{array}{ccccc}
B \otimes B & \xrightarrow{\check{\Gamma}} & B \otimes H^* & & \\
\downarrow \varrho_\chi \otimes \check{\chi} \varrho & \searrow & \downarrow P_\chi & & \\
& & B \otimes_{B^H} B & \xrightarrow{\Gamma} & \\
& & \downarrow \check{\Gamma}_\chi & & \\
A_\chi \otimes \check{\chi} A & \xrightarrow{\check{\Gamma}_\chi} & A \otimes_\chi H^* & & \\
& \searrow & \downarrow \Gamma_\chi & & \\
& & A_\chi \otimes_{I_\chi} \check{\chi} A & \xrightarrow{\Gamma_\chi} &
\end{array} \tag{9.11}$$

The front downward arrow is due to Proposition 8.3. Note that A/I_χ is a weak χ -Galois extension if so is B/B^H , as follows from the diagram.

Recall from Lemma 5.1 that the subspace of χ -generators in A_χ is equal to the subspace of $\check{\chi}$ -generators in $\check{\chi}A$. By Theorem 8.1, the operations $\check{\ast}$ and $\check{\bullet}$ coincide when restricted to that subspace and define the algebra A_χ^L .

Proposition 9.4. *Restriction of the Γ_χ to $A_\chi^L \otimes_{I_\chi} \otimes A_\chi^L$ is the ordinary Galois map and takes values in $A_\chi^L \otimes (H^\chi)^* \subset A_\chi^L \otimes H^*$.*

Proof. The first statement follows from the equalities

$$a \check{\ast} b^{[0]} \otimes \rho_1 \blacktriangleright b^{(1)} = a \check{\ast} b^{[0]} \otimes \chi(\rho_1) b^{(1)} = a \check{\ast} b^{[0]} \otimes b^{(1)}$$

valid for all $a, b \in A_\chi^L$. The coaction $A_\chi^L \rightarrow A_\chi^L \otimes H^*$ takes values in $A_\chi^L \otimes (H^\chi)^*$, as the H -action on A_χ^L factors through an action of H^χ . \square

The Galois map is independent on the choice of base algebra in the following sense. Suppose we are given a homomorphism of bases $(H, L) \xrightarrow{\text{id}, \varpi} (H, L')$ identical on H . Given a dynamical algebra A over the base (H, L) we make it a dynamical algebra over (H, L') , with the multiplication

$$A \otimes A \rightarrow A \otimes L \xrightarrow{\text{id} \otimes \varpi} A \otimes L'.$$

This amounts to an H -algebra map $A \rtimes L \rightarrow A \rtimes L'$ identical on A and given by ϖ on L . A character $\chi' \in \hat{L}'$ gives rise to the character $\chi := \chi' \circ \varpi \in \hat{L}$. We have also $\check{\chi} := \check{\chi}' \circ \varpi$, as ϖ commutes with the action of $\vartheta \in \mathfrak{D}(H)$ on L and L' (because the base homomorphism is

identical on H). It is obvious that Γ_χ constructed via $A \rtimes L$ is equal to $\Gamma_{\chi'}$ constructed via $A \rtimes L'$. Below we illustrate this on the example $L' = H$.

We fix an invariant character $\chi \in \hat{L}$ and consider the homomorphism $\iota_\chi: L \rightarrow H$. As was mentioned earlier, this amounts to a homomorphism of dynamical bases $(H, L) \rightarrow (H, H)$ identical on H . The group $\hat{H} \subset H^*$ acts on the algebra H by base algebra automorphisms through the left regular action $\varpi_\alpha(h) := \alpha \cdot h = h^{(1)} \langle \alpha, h^{(2)} \rangle$, see Example 2.5. Under this action, $\alpha \circ \iota_\chi = \iota_{\alpha \cdot \chi}$. Denote by $A \rtimes_\chi H$ the dynamical smash product of A and H . For $\alpha \in \hat{H}$ the algebras $A \rtimes_\chi H$ and $A \rtimes_{\alpha \cdot \chi} H$ are isomorphic, with the isomorphism identical on A and given by ϖ_α on H .

All the characters of the base algebra H are invariant and constitute the group \hat{H} . For all $\alpha \in \hat{H}$ we construct the left and right $A \rtimes_\chi H$ -induced modules A_α and ${}_\alpha A$ as in Section 8. The homomorphism $B \rightarrow A \rtimes_\chi H$ makes A_α and ${}_\alpha A$ left and right B -modules, which are isomorphic to $A_{\alpha \cdot \chi}$ and ${}_{\alpha \cdot \chi} A$, respectively. The subspace $I_\alpha \in A$ is formed by H -invariants. We have the following isomorphisms $(I_{\alpha \cdot \chi})_{op} \simeq (I_\alpha)_{op} \simeq \text{End}_{A \rtimes_\chi H}(A_\alpha)$, where the right isomorphism is due to Theorem 8.1 applied to base algebra $L = H$.

The homomorphism $B \rightarrow B' := A \rtimes_\chi H$ gives rise to the commutative diagrams

$$\begin{array}{ccc} B \otimes B & \xrightarrow{\check{\Gamma}} & B \otimes H^* & A_{\alpha \cdot \chi} \otimes_{I_\chi} {}_{\alpha \cdot \chi} A & \xrightarrow{\Gamma_{\alpha \cdot \chi}} & A \otimes H^* \\ \downarrow & & \downarrow & \downarrow \wr & & \downarrow \wr \\ B' \otimes B' & \xrightarrow{\check{\Gamma}'} & B' \otimes H^* & A_\alpha \otimes_{I_\alpha} {}_\alpha A & \xrightarrow{\Gamma'_\alpha} & A \otimes H^* \end{array}, \quad \forall \alpha \in \hat{H}.$$

In the right square we have used the equality $\alpha \cdot \tilde{\chi} = \widetilde{\alpha \cdot \chi}$, cf. Section 3. These diagrams illustrate independence of the Galois map on the choice of base algebra.

10 Morita context for dynamical smash product

In the present section we construct a Morita context for the dynamical smash product with any invariant character $\chi \in \hat{L}$ and a left generator $e \in L$. Recall that two algebras are called Morita equivalent if their module categories are equivalent. Morita context is a weaker relation than equivalence. Given two algebras C and B , a Morita context is a quadruple $(P, Q, [\cdot, \cdot], (\cdot, \cdot))$ consisting of an $C - B$ bimodule P , an $B - C$ bimodule Q and two maps

$$[\cdot, \cdot]: P \otimes_B Q \rightarrow C, \quad (\cdot, \cdot): Q \otimes_C P \rightarrow B$$

obeying the following conditions. Firstly, they are, respectively, C - and B -bimodule maps. Secondly, for all $p, p' \in P$ and $q, q' \in Q$ they satisfy the "associativity" condition

$$p' \dashv (q, p) = [p', q] \vdash p, \quad q' \dashv [p, q] = (q', p) \vdash q.$$

Here \vdash and \dashv denote the left and right actions of C and B . Morita context defines two functors between categories of C -and B -modules and becomes equivalence if $[\cdot, \cdot]$ and (\cdot, \cdot) are surjective.

We construct a Morita context for the algebras B and A_χ^L under the assumption that the subspace L^χ of χ -generators is not zero. The construction depends on a generator $e \in L^\chi$.

Lemma 10.1. *For every $e \in L$ the map $\check{\Pi}_e: B \otimes H^* \rightarrow B$, $a \otimes \eta \mapsto a\langle \eta, e^{(1)} \rangle e^{[\infty]}$ is a homomorphism of B -bimodules. If e is a χ -generator, then this map factors through a map $\Pi_e: A \otimes_\chi H^* \rightarrow B$.*

Proof. The statement is obvious as to the left B -action. Let us demonstrate that $\check{\Pi}_e$ is a right B -homomorphism. For all $a, b \in B$ and $\eta \in H^*$ the element $ab^{[0]} \otimes \eta b^{(1)}$ goes to

$$ab^{[0]} \langle \eta b^{(1)}, e^{(1)} \rangle e^{[\infty]} = ab^{[0]} \langle \eta, e^{(1)} \rangle \langle b^{(1)}, e^{(2)} \rangle e^{[\infty]} = a(e^{(2)} \triangleright b) \langle \eta, e^{(1)} \rangle e^{[\infty]} = a \langle \eta, e^{(1)} \rangle e^{[\infty]} b,$$

as required.

Further, suppose that e is a left χ -generator. Then $\langle \eta, e^{(1)} \rangle \lambda e^{[\infty]} = \langle \lambda \xrightarrow{\chi} \eta, e^{(1)} \rangle e^{[\infty]}$ for all $\lambda \in L$ and $\eta \in H^*$. Hence

$$\check{\Pi}_e: a\lambda \otimes \eta - a \otimes (\lambda \xrightarrow{\chi} \eta) \mapsto a\lambda \langle \eta, e^{(1)} \rangle e^{[\infty]} - a \langle \lambda \xrightarrow{\chi} \eta, e^{(1)} \rangle e^{[\infty]} = 0.$$

Thus $\check{\Pi}_e(J_\chi) = \{0\}$, and $\check{\Pi}_e$ factors through a B -bimodule map $\Pi_e: A \otimes_\chi H^* \rightarrow B$. This completes the proof. \square

We call a left χ -generator e cyclic if L is a cyclic right H^* -module generated by e . In particular, in the special case $L = H$ every character of H is cyclic. This follows from the fact that the map $\eta \mapsto t\eta$, where $t \neq 0$ is a left integral of H , is a linear isomorphism.

Lemma 10.2. *Suppose that A is a weak χ -Galois extension. If the left χ -generator e is cyclic, then the map $A_\chi \otimes_{I_\chi} \bar{\chi}A \rightarrow A \otimes_\chi H^* \xrightarrow{\Pi_e} B$ induces an epimorphism $A_\chi \otimes_{A_\chi^L} \bar{\chi}A \rightarrow B$ given by the assignment $a \otimes b \mapsto aeb$.*

Proof. The map in question is given by the assignment $a \otimes b \mapsto a(e^{(1)} \triangleright b)e^{[\infty]} = aeb$. This implies the statement. \square

Put $P := A_\chi$, and $Q := \bar{\chi}A$. By construction P and Q are, respectively, $B - A_\chi^L$, and $A_\chi^L - B$ bimodules. The B -module structures are obtained by induction from the H -module \mathbb{k}_χ . Define the maps

$$[\cdot, \cdot]: P \otimes_{A_\chi^L} Q \rightarrow B, \quad (\cdot, \cdot): Q \otimes_B P \rightarrow A_\chi^L$$

by setting

$$[a, b] = aeb, \quad (a, b) = \iota_\chi(e) \triangleright (a \dot{*} b).$$

Proposition 10.3. *The quadruple $(P, Q, [., .], (., .))$ is a Morita context for B and A_χ^L . It becomes equivalence if the extension A/I_χ is weak Galois, the generator e is cyclic, and $A_\chi^L = \iota_\chi(e)(A)$. The similar statement is true if B is replaced by $A \rtimes_\chi H$ and A_χ^L by I_χ .*

Proof. It is straightforward to check that $(P, Q, [., .], (., .))$ defined as above is a Morita context indeed. The map $(., .)$ is an epimorphism if $A^L = \iota_\chi(e)(A)$. Therefore the Morita context becomes equivalence if and only if the map $[., .]$ is surjective. That is the case if A is a weak χ -Galois extension and e is cyclic. Setting $L = H$ we get $A_\chi^L = I_\chi$, so the last statement follows from the previous two. \square

The constructions of this section take an especially simple form if the generator $e = e_\chi$ is an idempotent; recall that $\tilde{\chi} = \chi$ in this case. For the algebra $B = A \rtimes L$, its left induced module A_χ is realized as the left ideal Be_χ , while the right induced module ${}_\chi A$ as the right ideal $e_\chi B$. The algebra A_χ^L is isomorphic to the subring $e_\chi B e_\chi \subset B$. The Galois map can be presented as $a \otimes b \mapsto ae_\chi b^{[0]} \otimes b^{(1)}$. The Morita context is defined through the maps

$$[ae_\chi, e_\chi b] = ae_\chi b, \quad (e_\chi a, be_\chi) = e_\chi(ab)e_\chi.$$

where $P = Be_\chi = Ae_\chi$ and $Q = e_\chi B = e_\chi A$.

11 Ring theoretical characterization of χ -Galois extension

In this last section we generalize Theorem 1.2 of [CFM] characterizing the standard Hopf-Galois extension, for the case of dynamical algebras.

Lemma 11.1 (Frobenius reciprocity). *Suppose B is a dynamical smash product $A \rtimes L$. Let M be a left B -module and fix a character $\chi \in \hat{L}$. Then $\text{Hom}_L(\mathbb{k}_\chi, M) \simeq \text{Hom}_B(A_\chi, M)$ as vector spaces. The analogous statement is true for right modules.*

This lemma is analogous to Lemma 0.6 of [CFM] and, upon replacement of L by H , plays the same role in the following theorem for $B = A \rtimes_\chi H$.

Theorem 11.2. *Let L be a base algebra over a finite dimensional Hopf algebra H and let A be a dynamical (H, L) -algebra. Fix an invariant character $\chi \in \hat{L}$. Then the following are equivalent*

- (1) A_χ/I_χ is right weak H^* -Galois.
- (2) (a) The map $A \rtimes_\chi H \rightarrow \text{End}_{I_\chi}(A_\chi)$ is an algebra isomorphism,
 (b) A_χ is a finitely generated projective right I_χ -module.
- (3) A_χ is a left $A \rtimes_\chi H$ -generator.
- (4) if $0 \neq t \in \int_l \subset H$, then the map $[\cdot, \cdot]: A_\chi \otimes_{I_\chi} {}_{\tilde{\chi}}A \rightarrow A \rtimes_\chi H$ given by $[a, b] = atb$ is surjective.
- (5) For any left $A \rtimes_\chi H$ -module the map $A_\chi \otimes_{I_\chi} M^H \rightarrow M$ given by $a \otimes m \mapsto am$ is a left $A \rtimes_\chi H$ -module isomorphism.

The proof of this theorem is literally the same as the proof of Theorem 1.2 in [CFM]. The only difference is the argumentation for the equality $(A \rtimes_\chi H)^H = tA$ (here $A \rtimes_\chi H$ is understood as a left regular H -module) in the proof of the implication (5) \Rightarrow (4). The inclusion $(A \rtimes_\chi H)^H \subset tA$ follows from $\dim \int_l = 1$ and from the fact that the left H -module $A \rtimes_\chi H$ is freely generated by A ; the assumption for A being an algebra is unnecessary.

References

- [AL] A. Alekseev and A. Lachowska: *Invariant $*$ -products on coadjoint orbits and the Shapovalov pairing*, Com. Math. Helv. **80** (2005) 795–810
- [CFM] M. Cohen, D. Fischman, and S. Montgomery: *Hopf Galois extensions, smash products, and Morita equivalence*, J. Algebra **133** (1990) 351–372.
- [Dr1] V. Drinfeld: *Quantum Groups*, in Proc. Int. Congress of Mathematicians, Berkeley, 1986, ed. A. V. Gleason, AMS, Providence (1987) 798–820.
- [Dr2] V. Drinfeld: *Almost cocommutative Hopf algebras*, Leningrad Math. J. **2** (1990), # 2, 321–342.
- [DM1] J. Donin and A. Mudrov: *Dynamical Yang-Baxter equation and quantum vector bundles*, Commun. Math. Phys. **254** (2005) 719–760.
- [DM2] J. Donin and A. Mudrov: *Quantum groupoids and dynamical categories*, J. Algebra **296** (2006) 348–384.

- [EE] B. Enriquez and P. Etingof: Quantization of classical dynamical r-matrices with non-abelian base, *Commun. Math. Phys.* **254** (2005) 603–650.
- [EEM] B. Enriquez, P. Etingof, and I. Marshall: Quantization of some Poisson-Lie dynamical r-matrices and Poisson homogeneous spaces, arXiv: math.QA/0403283.
- [KMST] E. Karolinsky, K. Muzykin, A. Stolin, and V. Tarasov: *Dynamical Yang-Baxter equations, quasi-Poisson homogeneous spaces, and quantization*, *Letters in Mathematical Physics*, **71** (2005), 179–197.
- [Lu] J.-H. Lu: *Hopf algebroids and quantum groupoids*, *Int. J. Math.*, **7** (1996) 47–70.
- [Sw] M. Sweedler: *Integrals for Hopf algebras*, *Ann. Math.* **89** (1969) 323–335.
- [Y] D. Yetter: *Quantum groups and representations of monoidal categories*, *Math. Proc. Cambridge Philos. Soc.* **108** (1990) # 2, 261–290.