

# Thermodynamic restrictions on evolutionary optimization of transcription factor proteins

Longhua Hu and Alexander Y. Grosberg  
Department of Physics, University of Minnesota,  
116 Church Street SE, Minneapolis, Minnesota 55455, USA

Robijn Bruinsma  
Department of Physics and Astronomy, University of California at Los Angeles, Los Angeles, California 90035

*Conformational fluctuations* are believed to play an important role in the process by which transcription factor proteins locate and bind their target site on the genome of a bacterium. Using a simple model, we show that the binding time can be minimized, under selective pressure, by adjusting the spectrum of conformational states so that the fraction of time spent in more mobile conformations is matched with the target recognition rate. The associated optimal binding time is then within an order of magnitude of the limiting binding time imposed by thermodynamics, corresponding to an idealized protein with instant target recognition. Numerical estimates suggest that typical bacteria operate in this regime of optimized conformational fluctuations.

The ability of bacteria to respond within minutes to changes in their environment relies on genetic switches that are controlled by *transcription factors*. Transcription factors are proteins that—following activation by an environmental change—are able to locate a specific region (the “operator sequence”) along the bacterial genome and bind to it, thereby regulating the expression of a gene (or group of genes) adjacent to that region [1]. The number of copies of a transcription factor protein associated with a specific gene varies, but typically it is in the range of  $10^2$ . Because bacterial genomes have a size in the range of  $10^7$  sites, a transcription factor must be able to “scan” the DNA for the target site at a rate of  $10^5$  sites per second or faster in order for at least one of them to reach the target site within seconds. Note that following the *search* for the target site, the transcription factor still has to *bind* to the target site to regulate the expression of the gene.

A series of classical papers on the search process [2, 3, 4] culminated in the work of Berg, Winter and von Hippel (BWH) who showed [5]—for the canonical case of the *lac* repressor protein of the bacterium *E. coli*—that the search process takes place not by straightforward 3D diffusion to the target binding site but rather by a slide-jump combination of 1D diffusional sliding along the DNA chain alternating with 3D diffusional jumps between different DNA segments. By restricting part of the search to the 1D “target space”, the binding rate is effectively enhanced with respect to a pure 3D search, while the 3D jumps reduce the repetitive visits to the same sites that characterize purely 1D diffusive searches. This scenario is made possible by a modest, non-specific electrostatic affinity between the transcription factor and duplex DNA. BWH also provided evidence that, under physiological conditions, the search time has a minimum with respect to the strength of this non-specific affinity, which may be the result of evolutionary optimization un-

der selective pressure. Subsequent structural studies [6] have shown that the DNA-binding domains of the *lac* repressor are subject to strong *conformational fluctuations* when the protein is in contact with non-operator DNA. If the binding domain is in contact with operator sequence DNA then the protein can undergo a large-scale conformational change to a stable structure with direct contacts between the amino-acid side chains and the DNA bases.

It would seem obvious that the delay time between activation and binding of a transcription factor to the operator sequence (“binding time”) is minimized by maximizing the 1D diffusion constant  $D_1$ . However, simply increasing the transport rate will impair the accuracy, or *fidelity*, with which the protein can distinguish a right from a wrong site. Specifically, if the binding of a transcription factor to the target site is characterized by a certain rate  $\Omega$ , then the protein is likely to *overshoot* the target site if the jump rate  $D_1/a^2$  between sites, with  $a$  the spacing between protein binding sites, is large compared to  $\Omega$ . Similar conflicts between process speed and process fidelity are familiar from DNA duplication and transcription where increased reaction rates increase the number of duplication and transcription errors.

Slutsky and Mirny [7] proposed that conformational fluctuations could ease the conflict between speed and fidelity. If some conformations of the transcription factor are sensitive to the DNA sequence while others are characterized by rapid transport then the transcription factor might be able to scan the genome efficiently by rapidly flipping between the two types of conformations. The aim of this paper is to analyze how close this mechanism can approach limits of search efficiency imposed by fundamental principles of thermodynamics. We will address this question by examining a simple model for the conformational fluctuations, similar to that of Ref. [7], where the transcription factor is allowed to adopt only two conformations (+ and –) when in contact with non-operator




FIG. 1: (Color online) Schematic representation of the model. A protein moving diffusively through the cell volume (a) is adsorbed on genomic DNA (b) where it adopts one of two conformations: + and -. In the + conformation it is loosely associated with the DNA and can move by one-dimensional diffusion along the DNA chain (b) while in the - conformation (c) it is tightly associated with the DNA and is immobile. After returning to the + state, it restarts the sliding motion. The protein also can desorb from the chain (d) and return to three-dimensional diffusive motion. Following a number of such cycles, the protein lands in the “antenna region” within a distance  $\lambda$  of the target binding site (e). After reaching the target site by one-dimensional diffusion it can undergo a large-scale irreversible conformational transition to the final bound state if it is in the - state (f).

DNA. As illustrated in Fig. 1, in the + state, the protein is less ordered and only loosely associated with the DNA while it can slide along the DNA chain. In the - state, the protein is more ordered, closely associated with the DNA and immobile [8]. If the transcription factor is in contact with the target operator sequence then, in addition to these two states, it also can undergo an irreversible conformational transition from the - state to the fully ordered final bound state. We will show that the *shortest possible binding time* in this model is controlled by a dimensionless binding rate  $\omega \equiv 2\Omega ab/\sqrt{KD_1D_3}$ , with  $D_3$  the protein diffusion coefficient in bulk solution,  $D_1$  the diffusion coefficient for 1D transport along the DNA in the + state,  $K$  the equilibrium constant for the non-specific protein-DNA interaction, and  $b$  the DNA-protein “capture radius” [9]. If the dimensionless binding rate is comparable to one—or larger than one—, then we can show that for a particular value of the energy difference  $\Delta E_{\pm}$  between the + and - conformations, the binding time can approach an absolute lower bound that corresponds to proteins having *infinitely fast* final binding rates. In other words, if the internal degrees of freedom of the protein in the sliding state are properly matched to the final binding rate then the binding time of a transcription factor can approach the shortest possible value allowed by thermodynamics *provided* the dimensionless binding rate is sufficiently large.

To demonstrate these claims, assume a cell of volume  $V$  containing a DNA genome of length  $L$ . The cell also contains a certain (low) concentration  $c$  of transcription factor proteins that can bind reversibly and non-specifically to the DNA. A protein whose center is located inside a

cylindrical tube of radius  $b$  surrounding the duplex DNA will be assumed to be non-specifically associated with the DNA. The fraction  $\phi$  of the total cell volume occupied by the tube is of the order of  $Lb^2/V$ . There is also a single target site on the strand where the transcription factor can bind irreversibly. We start by applying a fundamental theorem [10], which—in terms of our model—states that the mean waiting time for irreversible occupation of the target site, the quantity of interest to us, is equal to the inverse of a steady-state diffusion current of a *different* problem, namely one where the target site is replaced by a protein *sink* that constantly absorbs - state transcription factors located at the target site at a rate  $\Omega$ , while the protein concentration far from the target site is maintained at a certain fixed value  $c_3(\infty)$ . The steady-state diffusion current, denoted by  $J_{3D}$ , into the target site for this second problem can be obtained from straightforward solution of the diffusion equation, which leads to the well-known Smoluchowski relation for the reaction rate of diffusion-limited chemical reactions:

$$J_{3D} \sim D_3 c_3(\infty) \xi . \quad (1)$$

Following Ref. [11], the effective “target radius” is defined as the radius of a sphere, surrounding the target site, that determines a cross-over regime such that far outside the sphere adsorption of proteins onto the DNA chain is in equilibrium with evaporation of protein from the DNA chain while deep inside the sphere the absorption rate exceeds the evaporation rate. For the case of transcription factors obeying BWH slide-skip transport, the size of this target sphere is determined by the condition that if a protein lands on a DNA segment inside the target sphere, following a 3D diffusion step, then it typically reaches the target sink by pure 1D diffusion where it gets absorbed before there is a chance for it to “evaporate” and leave the DNA. The length  $\lambda$  of DNA chain inside this target sphere—referred to as the “antenna” length—in general depends on the spatial organization of the genome. We will assume here the simple case of a straight genome, with  $\xi$  of order  $\lambda$  [12]. This antenna length has to be determined self-consistently but first we must establish a relation between  $c_3(\infty)$  and the actual protein concentration  $c$ .

Far outside the target sphere the DNA-protein system is, by assumption, nearly in local thermal equilibrium, so one can determine the concentrations of adsorbed and free proteins purely from equilibrium considerations. If one views the association of the transcription factors with DNA as a simple chemical reaction, then the concentration  $\tilde{c}(\infty)$  of proteins adsorbed non-specifically on the DNA and the concentration  $c_3(\infty)$  of free proteins must be related to the reaction volume fraction  $\phi$  by the *Law of Mass Action* for dilute chemical systems in thermodynamic equilibrium:

$$\frac{c_3(\infty)\phi}{\tilde{c}(\infty)} \simeq K \quad (2)$$

with  $\phi \ll 1$ . The non-specific protein-DNA equilibrium constant  $K$  depends strongly on the salt concentration [4], and other thermodynamic parameters, but it is independent of the protein and DNA concentrations. Since  $c = c_3(\infty) + \tilde{c}(\infty)$ , the concentrations of free and adsorbed proteins are now determined but it will be useful to replace the bulk concentration  $\tilde{c}(\infty)$  of adsorbed proteins by the 1D concentration  $c_1(\infty) \simeq b^2 \tilde{c}(\infty)/\phi$ , the number of adsorbed proteins *per unit length* of DNA far from the target site. Solving for  $c_1(\infty)$  and  $c_3(\infty)$  gives  $c_1(\infty) \simeq cb^2/K(1 + \phi/K)$  and  $c_3(\infty) \simeq c/(1 + \phi/K)$ , still for  $\phi \ll 1$ .

Deep inside the target sphere, the system is not in thermal equilibrium, with the adsorption rate of proteins from the bulk solution to the DNA exceeding the evaporation rate. The difference is matched by a 1D diffusion current  $J_{1D}$  along the DNA chain towards the target site. In order to estimate this 1D diffusional transport, note that if the interconversion rate between the + and - states is sufficiently rapid then their respective occupancies can be approximated by the equilibrium Boltzmann distribution. The *effective* 1D diffusion constant for transport along the chain—which we will denote by  $\tilde{D}_1$ —is then proportional to the Boltzmann probability  $p(+)$  to find the protein in the + state. If  $\mu \equiv \exp(-\Delta E_{\pm}/k_B T)$ , then  $p(+)=\mu/(1+\mu)$  and  $\tilde{D}_1 \simeq D_1 \mu/(1+\mu)$ . Similarly, the *effective* target site binding rate  $\tilde{\Omega}$  is, under these same conditions, proportional to the probability  $p(-)=1-p(+)$  to find the protein in the - state and  $\tilde{\Omega} \simeq \Omega/(1+\mu)$ .

Let  $c_1(0)$  be the 1D concentration at the target site. If the final binding rate were infinitely fast, then  $c_1(0)$  would be zero but, because of the overshoot effect, this is no longer the case. If we view the surface of the target sphere as a matching region between the asymptotic regions far from the sink where the 1D concentration approaches  $c_1(\infty)$  and the region deep inside the target sphere near the sink where the 1D concentration approaches  $c_1(0)$ , then we can estimate the 1D concentration gradient as  $[c_1(\infty) - c_1(0)]/\lambda$ . It follows that the 1D diffusion current towards the sink equals:

$$J_{1D} \sim \tilde{D}_1 \frac{c_1(\infty) - c_1(0)}{\lambda}. \quad (3)$$

The number of proteins absorbed per second by the sink itself,  $J_s$ , is of the order of  $a c_1(0) \tilde{\Omega}$ , with  $a$  the spacing between protein binding sites. Conservation of the number of proteins requires the three currents  $J_{3D}$ ,  $J_{1D}$  and  $J_s$  to be equal to each other [11], so

$$J_{3D} = J_{1D} = J_s. \quad (4)$$

Equating the 1D diffusion current with the sink current allows us to eliminate  $c_1(0)$  with the result:

$$J_{1D} \sim \frac{\tilde{D}_1 c_1(\infty)}{\lambda} \left( \frac{\tilde{\Omega}}{\tilde{\Omega} + \tilde{D}_1/a\lambda} \right). \quad (5)$$




FIG. 2: (Color online) Contour plots of the transport enhancement factor  $A$  as a function of the equilibrium constant  $K$  and the occupation probability ratio  $\mu$  of the + over - states for  $D_1 = 10^{-9}$  cm<sup>2</sup>/s,  $D_3 = 3 \times 10^{-7}$  cm<sup>2</sup>/s,  $a = 0.34$  nm,  $b = 5$  nm,  $\phi = 0.01$  and  $\Omega = 3 \times 10^3$  Hz. There is a shallow maximum around  $\mu = 0.1$  and  $K = 10^3$ . The ratio of the transport enhancement factor at this maximum,  $A_{\text{opt}}$ , and the thermodynamic limiting enhancement factor  $A_{\infty}$  equals 0.193. Inset: Dependence of the ratio  $A_{\text{opt}}/A_{\infty}$  on the dimensionless binding rate  $\omega$ .

The factor in front of the square brackets is the diffusion current in the absence of overshoot. The importance of overshoot is thus determined by the dimensionless number  $a\lambda\tilde{\Omega}/\tilde{D}_1$ . Since  $\lambda^2/\tilde{D}_1$  is the typical time spent by a protein diffusing along the antenna, it follows that  $a\lambda/\tilde{D}_1$  is the typical time spent near the target site so  $a\lambda\tilde{\Omega}/\tilde{D}_1$  is the product of the typical time spent near the target site with the effective absorption rate. The term inside the square brackets can then be understood as the probability for a protein in the antenna region to be trapped by the target.

Equating the 1D and 3D currents provides us with a self-consistency condition that determines both the size of the antenna length  $\lambda$  and the reaction rate. Solving for  $\lambda$  using Eqs. (1) and (4) and using  $c_1(\infty)/c_3(\infty) \sim b^2/K$  gives the antenna length:

$$\lambda = \sqrt{\frac{b^2}{K} \left( \frac{\tilde{D}_1}{D_3} \right) + \left( \frac{\tilde{D}_1}{2\tilde{\Omega}a} \right)^2} - \frac{\tilde{D}_1}{2\tilde{\Omega}a}. \quad (6)$$

The maximum value,  $\lambda_{\infty} = \sqrt{b^2 D_1/K D_3}$ , is reached for infinite  $\Omega$  and infinite  $\mu$ .

It will be helpful to express the binding rate  $J_{3D} \sim D_3 c_3(\infty) \xi$  in dimensionless units as  $A \equiv J_{3D}/(c D_3 a)$  with  $c D_3 a$  the Smoluchowski limiting rate of a conventional 3D diffusive search for an absorber target of radius  $a$  (the spacing between binding sites), so  $A$  can be viewed

as a reaction *amplification* or enhancement factor. This enhancement factor can be expressed as a simple function of the dimensionless binding rate  $\omega = 2\Omega ab/\sqrt{KD_1D_3}$  and the Boltzmann factor  $\mu = \exp(-\Delta E_{\pm}/k_B T)$ :

$$A(\omega, \mu) \sim A_{\infty} \frac{1}{\omega} \left( \sqrt{\omega^2 \frac{\mu}{1+\mu} + \mu^2 - \mu} \right). \quad (7)$$

Here  $A_{\infty} = (b/a)\sqrt{KD_1/D_3}/(K + \phi)$  is the maximum value of the enhancement factor, corresponding to  $\lambda = \lambda_{\infty}$  with both  $\mu$  and  $\Omega$  infinite. We will examine the amplification factor  $A(\omega, \mu)$  as a function of the non-specific equilibrium constant  $K$  and the occupation ratio  $\mu$  of the + state and - state, rather than  $\omega$  and  $\mu$ , because these are physical parameters characterizing the interaction between the transcription factor and the DNA that are expected to be sensitive to specific point mutations of the transcription factor amino-acid sequence through their exponential dependence on binding and activation energies. The contour lines of constant  $A$  as a function of  $K$  and  $\mu$  in Fig. 2 show that there is a single, rather shallow maximum. The physical origin behind the maximum of  $A$  with respect to  $K$  is, as discussed earlier, the fact that a combination of 1D and 3D diffusion minimizes the search time. By contrast, the maximum of  $A$  as a function of  $\mu$  at  $\mu_{\text{opt}} = (\sqrt{1+2\omega}-1)/2$  is surprising because it might have been expected that for sufficiently long DNA, location of the target site *always* should be the “rate-limiting” step, in which case the optimal choice for  $\mu$  would be infinite since that maximizes the effective 1D diffusion constant  $\tilde{D}_1 = D_1\mu/(1+\mu)$ . It can be shown that the maximum with respect to  $\mu$  actually is a form of *impedance matching* with the effective “resistance” of the 1D diffusional search matched with the effective resistance of the binding process.

If  $\mu$  adopts the optimal value  $\mu_{\text{opt}} = (\sqrt{1+2\omega}-1)/2$ , then the ratio  $A_{\text{opt}}/A_{\infty}$  of the optimal rate amplification factor and its maximum value is a function only of the dimensionless rate  $\omega$ :

$$\frac{A_{\text{opt}}(\omega)}{A_{\infty}} = 1 - \frac{1}{\omega} (\sqrt{1+2\omega} - 1). \quad (8)$$

The dependence of  $A_{\text{opt}}/A_{\infty}$  on  $\omega$  is shown in the inset of Fig. 2:  $A_{\text{opt}}$  is of the same order of magnitude as the theoretical limit  $A_{\infty}$  already for modest values of  $\omega$ . This demonstrates our central claim: it is *possible* for the overall binding rate of a transcription factor to approach the theoretical limiting value but *only* by a suitable choice of  $\mu$ , and *only* if the dimensionless binding rate  $\omega$  is of the order of one, or larger than one.

Are these two conditions realistic for typical transcription factors? Typical values for the diffusion constants of bacterial transcription factors are [13, 14]  $D_1 \sim 10^{-9}$  cm<sup>2</sup>/s and  $D_3 \sim 3 \times 10^{-7}$  cm<sup>2</sup>/s. We can estimate the protein-DNA reaction volume fraction  $\phi$  for *E. coli* by assuming it to be comparable to the DNA volume fraction

(about 1%). The equilibrium constant can then be determined from the relation  $c_3(\infty) \simeq c/(1+\phi/K)$  and the fact that it is known that about 10% of the *lac* repressor proteins of *E. coli* are in solution [15], which means that  $K$  must be of the order of  $10^{-3}$ . If we assume  $a$  to be equal to the base-pair spacing 0.34 nm, and estimate  $b$  as 5 nm, then the dimensionless binding rate  $\omega$  is of the order of  $10^{-4}\Omega$  with the binding rate  $\Omega$  expressed in Hz. A large-scale protein conformational change typically involves millisecond to microsecond time scales, from which it follows that  $\omega$  must lie in the range of 0.1 to 100. Note, from Fig. 2 that the optimal value for  $K$  is close to  $10^{-3}$  for  $\Omega$  in the kHz range. We conclude that the second condition can be satisfied under typical conditions. Next, the optimal occupation ratio  $\mu_{\text{opt}} = (\sqrt{1+2\omega}-1)/2$  is in the range of 0.1 to 10 for  $\omega$  in the range of 0.1 to 100. The corresponding optimal energy difference  $\Delta E_{\pm}$  between the + and - states is then in the range of a few  $k_B T$ , with  $\Delta E_{\pm}$  positive for  $\omega < 4$  but negative for  $\omega > 4$ . In either case, the structure of “optimized” transcription factors bound to non-operator DNA should be subject to strong thermal fluctuations. As we saw, this is indeed the case of the *lac* repressor [6], while a recent modeling study of the *Ets*-DNA system arrives at the same conclusion [8]. The first condition can thus be satisfied as well under reasonable conditions. Finally, the measured *lac* repressor binding rates [5] are comparable to the thermodynamic limiting rate. We conclude that, under reasonable conditions, the binding rate of transcription factor proteins can be of the same order of magnitude as the thermodynamic limiting rate if the energy spectrum of conformational fluctuations is determined, under selective pressure, by minimization of the overall binding time.

We would like to thank Leonid Mirny for useful discussions and the Kavli Institute of Theoretical Physics and the Aspen Center for Physics, where this study was initiated, for their hospitality. LH and AG would like to acknowledge support by the MRSEC Program of the NSF under Award Number DMR-0212302 and RB would like to acknowledge support by the NSF under DMR Grant 0404507.

- 
- [1] B. Alberts *et al.*, *Molecular Biology of the Cell*, (Garland, New York, 1994).
  - [2] G. Adam and M. Delbrück, *Structural Chemistry and Molecular Biology*, A. Rich and N. Davidson, editors, (Freeman, New York, 1968).
  - [3] A. D. Riggs, S. Bourgeois and M. Corn, *J. Mol. Biol.* **53**, 401 (1970).
  - [4] P. H. Richter and M. Eigen, *Biophys. Chem.* **2**, 255 (1974).
  - [5] O. G. Berg, R. B. Winter and P. H. von Hippel, *Biochemistry.* **20**, 6929 (1981), R. B. Winter, O. G. Berg and P. H. von Hippel, *Biochemistry.* **20**, 6961 (1981), P. H. von Hippel and O. G. Berg, *J. Biol. Chem.* **264**, 675 (1989).

- [6] C. G. Kalodimos *et al.*, *Science*. **305**, 386 (2004).
- [7] M. Slutsky, L. A. Mirny, *Biophys. J.* **87**, 4021 (2004).
- [8] A specific realization of a conformational fluctuation spectrum of this type was recently discussed for the *Ets*-DNA system by Y. Levy, J. N. Onuchic and P. G. Wolynes, *J. Am. Chem. Soc.* **129**, 738 (2007).
- [9] More precisely,  $b$  is defined as the radius of a cylinder surrounding the DNA duplex such that a protein will be captured by the DNA, for example by electrostatic attraction, if the center of the protein is located inside the cylinder.
- [10] L. Pontryagin, A. Andronov and A. Vitt, *Zh. Eksp. Teor. Fiz.* **3**, 165 (1933); translated and reprinted in *Noise in Nonlinear Dynamical Systems*. Vol. 1, edited by F. Moss and P. V. E. McClintock, p. 329, (Cambridge University Press, Cambridge, 1989).
- [11] T. Hu, B. I. Shklovskii and A. Y. Grosberg, *Biophys. J.* **90**, 2731 (2006).
- [12] The maximum size  $\lambda_\infty$  of the antenna length is, for the parameter values used in this paper, less than the DNA persistence length.
- [13] J. Elf, G. Li and X. S. Xie, *Science*. **316**, 1191 (2007).
- [14] Y. M. Wang, R. H. Austin and E. C. Cox, *Phys. Rev. Lett.* **97**, 048302 (2006).
- [15] Y. Kao-Huang *et al.*, *Proc. Natl. Acad. Sci. USA.* **74**, 4228 (1977).