Luc Binette^{1,2} and Yair Krongold²

¹ Département de Physique, de Génie Physique et d'Optique, Université Laval, Québec, QC, G1K 7P4

 2 Instituto de Astronomía, UNAM, Ap. 70-264, 04510 México, DF, México

Received: 12th July 2007/ Accepted: 20th September 2007

ABSTRACT

Context. Emission lines in quasars are believed to originate from a photoionized plasma. There are, however, some emission features which appear to be collisionally excited, such as the Fe_{II} multiplet bands. Shortward of Ly α , there also are a few permitted lines of species from low to intermediate ionization.

Aims. Ton 34 ($z_q = 1.928$) exhibits the steepest far-UV continuum decline known ($F_\nu \propto \nu^{-5.3}$) shortward of 1050 Å. This object also emits unusually strong low to intermediate excitation permitted lines shortward of the Lyman limit.

Methods. Using archive spectra of Ton 34 from HST, IUE and Palomar, we measure the fluxes of all the lines present in the spectra and compare their relative intensities with those observed in composite quasar spectra.

Results. Our analysis reveals unusual strengths with respect to Ly α of the following low to intermediate excitation permitted lines: $\rm \tilde{O}$ II+ $\rm \rm \tilde{O}$ III (835 Å), N III+ $\rm \tilde{O}$ III (686–703 Å) and N III+N IV (765 Å). We compare the observed line spectrum with both photoionization and shock models.

Conclusions. Photoionization cannot reproduce the strengths of these far-UV lines. Shocks with $V_s \simeq 100 \,\mathrm{km \, s}^{-1}$ turn out to be extremely efficient emitters of these lines and are favored as excitation mechanism.

Key words. Line: identification — Line: formation — Atomic processes— Galaxies: quasars: emission lines — quasars: individual: Ton 34

1. Introduction

In this work, we analyze the emission lines of an unusual quasar, Ton 34, which is alternatively named PG 1017+280 or J1019+2745 with redshift $z_q = 1.928$. It is severely deficient in ionizing photons, since its Spectral Energy Distribution (SED) shows a remarkable steepening of the continuum in the rest-frame far-UV, shortward of $1100 \,\text{\AA}$ (Binette & Krongold 2007, hereafter BK07; Binette et al. 2007). If the far-UV is fitted by a powerlaw $(F_{\nu} \propto \nu^{+\alpha})$, the index^1 index^1 is as steep as $\nu^{-5.3}$. BK07 suggest that the extreme-UV flux might undergo a recovery shortward of 450\AA .

While the near-UV emission-line spectrum appears to be 'normal', the far-UV spectrum shows low to intermediate ionization species with unusual strengths. Using the UV sed constructed by BK07 from archive data, we will quantify this statement and present photoionization and shock models for comparison. The aim is to understand how the extreme deficiency of ionizing photons in Ton 34 might be impacting the emission line spectrum.

The emission-line spectrum of quasar and Seyfert I galaxies is generally believed to originate from gas photoionized by a nuclear UV source. State of the art photoionization models of the Broad Emission Line Region (BELR) such as those developed by Baldwin et al. (1995) and dubbed 'locally optimally emitting clouds' (LOC) models can successfully reproduce most of the emission lines observed in quasars. A grid of such models can be found in Korista et al. (1997, hereafter KO97) and more recently in Casebeer et al. (2006 and references therein). There are, however, a few exceptions to the success of pure photoionization. In particular, photoionization models require micro-turbulences in order to reproduce the shape and intensity of the Fe ii UV-band (Baldwin et al. 2004). A possible alternative is that the region producing Fe II is collisionally ionized, as proposed by Grandi (1981, 1982), Joly (1987) , Véron-Cetty et al. $(2004, 2006)$ and Joly et al. (2007). In this work, we present evidence that photoionization might not be sustainable in the case of some of the far-UV permitted lines reported in this paper.

2. The UV emission line spectrum of Ton 34

Below we summarize the procedure used by BK07 to derive the UV sep of Ton 34.

2.1. Description of the archival data

The current work is based on four archival or bibliographical sources. The $760-1120\text{\AA}$ spectral segment is provided by the dataset Y2IE0A0AT from the HST-FOS archives (grating G270H). To cover the extreme UV region, we borrowed from the IUE archives. The long wavelength segment (LWP) is from Tripp, Bechtold & Green (1994) and corresponds to the dataset LW0P5708. Fluxes longward of 3000 Å (observer-frame) were severely affected from reflected sunlight or moonlight (Lanzetta, Turnshek &

¹ Among the 77 quasars whose far-UV indices could be measured by Telfer et al. 2002, there were only 3 objects with a continuum steeper than ν^{-3} . continuum steeper than ν

Sandoval 1993) and have been discarded. The shorter wavelength IUE segment (SWP) was extracted directly from the archives and corresponds to the dataset SWP28188. To cover the SED behavior longward of the HST segment, we adopted the published optical spectra of Sargent, Boksenberg & Steidel (1988), which were taken at the Palomar 5.08 m Hale Telescope. Both optical spectra lacked absolute flux calibration, although the authors observed standard stars, which allowed them to provide a relative calibration.

2.2. Matching the different SED segments

We statistically corrected the UV spectral segments for the cumulated absorption caused by unresolved $Ly\alpha$ forest lines, which are responsible for the so-called far-UV "Lyman valley" (Møller & Jakobsen 1990). For that purpose, we adopted the *mean*^{[2](#page-1-0)} transmission function for $z_q = 2$ published by Zheng et al. (1997). We also applied a Galactic reddening correction assuming the Cardelli, Clayton & Mathis (1989) extinction curve corresponding to $R_V = 3.1$ and $E_{B-V} = 0.13$. The latter value corresponds to the mean extinction inferred from the 100μ maps of Schlegel et al. (1998) near Ton 34. The blue and red arm segments have been scaled to overlap smoothly with the HST-FOS segment. Both the LWP and SWP segments were multiplied by a factor 0.75. This scaling was necessary so that the LWP segment superimposes the HST-FOS spectrum as closely as possible. Continuum variability is a possible explanation for this continuum difference, since the IUE and HST observations were made in different years. Finally, all the spectral segments were shifted to rest-frame wavelengths, and F_{λ} was multiplied by $1 + z_q$. The IUE spectra have been rebinned by grouping *n* pixels together [SWP with $n = 5$ and LWP with $n = 3$) in order to improve the limited S/N. The LWP and HST-FOS spectra overlap significantly in spectral coverage. Both datasets taken nine year apart confirm the unusual steepness of the UV break in Ton 34.

2.3. Model of the ionizing SED of Ton 34

Shortward of 1100\AA , the continuum of Ton 34 undergoes a sharp fall off (see Fig. 2 in BK07), which BK07 model as dust absorption by nanodiamond grains. This resulted in a deep and broad absorption trough that fits the observed continuum reasonably well. In our photoionization calculations presented below in Sect. [3.2.1,](#page-3-0) we experiment with two ionizing seds. The first is the intrinsic 'unabsorbed' sed, which is assumed to be a powerlaw of index $+0.1$ followed by a roll-over centered on 640 Å that extends up to the Xray domain. Beyond 2 keV , sed if behaves as a powerlaw of index -1.0 , yielding an α_{OX} of -1.45 . This sep is shown in Fig. [1](#page-1-1) and, as in the work of BK07, it is labeled Model ii. The second SED used in photoionization calculations is the dust absorbed version of the same sep, which fits the ob served UV continuum of Ton 34 between 400 and $1550\,\mathrm{A}$ (la-

Fig. 1. Log-log plot of the input spectral energy distributions used in our photoionization calculations discussed in Sect. $3.2.1$. These ionizing SEDs are labeled II and IV in either F_{λ} (panel a) or F_{ν} (panel b) and are given as a function of wavelength (bottom axis) or photon energy (top axis). The distribution labeled II (solid line) is the assumed intrinsic SED while that labeled IV is the transmitted flux (dash-dotted line) assuming nanodiamond dust extinction (see BK07). Model iv is a fit of the UV continuum of Ton 34 between 400 and $1550 \text{ Å}.$

beled Model IV in Fig. [1\)](#page-1-1). Shortward of 200\AA and longward of 2000\AA , the two distributions are the same. This is because nanodiamond dust absorbs radiation over a relatively narrow domain as compared to other grain compositions.

In Fig. [2,](#page-2-0) we present the continuum subtracted spectrum of Ton 34, that is, the residual between the observed Ton 34 sed and our continuum fit represented by Model IV. The different spectral segments have been color-coded as follows, SWP: red, LWP orange, HST-FOS: blue, and Palomar: dark green.

2.4. Extraction of line fluxes and upper limits

The procedure to measure the flux of the lines was the following: we first fit a Gaussian to each observed line in the spectra. For several lines, a narrow component was required, so we added a second (narrow) Gaussian. In addition, the lines by C iv λ 1549, Si iv λ 1400, and Ly α show a clear asymmetry in the line profile, with a blue shoulder (see Fig. [2\)](#page-2-0). For these lines, we further included a third, broader, Gaussian. The FWHM of the broad component spans from \sim 3600 to 5300 km s⁻¹. It is interesting to note that the O II+O III complex at around 835\AA has a significant and strong red shoulder extending up to $\sim 850 \text{\AA}$, which is observed in both the IUE-LWP and HST-FOS spectra (see Fig. [2\)](#page-2-0). We could not find any positive identification of this shoulder with any line from a different ion/transition, and

² This correction is statistical in nature, as it relies on the average behavior with redshift of the spatial density of intervening absorbers. It cannot be used to correct small portions of the continuum, which may be coincident with a "clear patch" or an over-density in the $Ly\alpha$ forest. These inhomogeneities may generate spurious narrow features that should not be attributed to genuine emission lines.

 \circ Ć 1000 1200 600 800 1400 1600 (\AA) $\lambda_{\rm rest}$ L. Fig. 2. Residuals of the spectral energy distribution of Ton 34 after subtracting our absorbed continuum Model iv from BK07. The different spectral segments have been color coded as follows, SWP: red, LWP orange, HST-FOS: blue, and Palomar: dark green. Color-coded fiducial marks indicate the position of observed or expected (labeled with

thus we considered this feature as part of the O II+O III emission.

symbol '?') emission lines. Measurements of line intensities

and upper limits are given in Table [1.](#page-4-0)

The measured line fluxes extracted from Fig. [2](#page-2-0) as well as upper limits of other permitted lines are listed with respect to $Ly\alpha = 100$ in Col. 5 of Table [1.](#page-4-0) Note that we give the total flux under the profile, that is, the integrated flux from all the Gaussian components required to fit each emission line. A consistency check was carried out, which showed that the line fluxes measured over the original spectra or the continuum subtracted spectra were indistinguishable from each other.

In Col. 5 of Table [1,](#page-4-0) we show our error estimates, which we evaluated at a 1σ significance level. We assumed a S/N of 25 for most lines, except for N III+N IV and N III+O III where we assume a S/N of \simeq 10. The line upper limits in Table [1](#page-4-0) correspond to a significance of 2σ . As for the continuum, we estimate the errors to be $\simeq 10\%$.

Of all the emission features that we measure in the far-UV, three line systems stand out by their strengths with respect to the composite spectra: these are the O II+O III lines at 835\AA , the N III+O III lines at $686-703 \text{\AA}$ and the N III+N IV lines at 765\AA .

Many weaker features in the IUE spectrum appear to lie where other permitted lines of comparable excitation might be expected, such as O III λ 508 Å, O iv λ 554 Å, O v λ 630 Å and O IV λ 609 Å. A few of these have been reported before in other quasars (Reimers et al. 1998; Laor et al. 1995) or in composite AGN spectra (Zheng et al. 1997; Telfer et al. 2002; Scott et al. 2004). However, these line systems appear as too narrow in the IUE spectra with respect to typical

Fig. 3. Emission lines extracted from the Ton 34 spectrum plotted in velocity space. The flux scale is arbitrary for each inset. Left panels: near-UV permitted lines, right panel: far-UV permitted lines. Overall, all the lines are consistent with the rest frame system of Ton 34. Differences in the position of the lines on the right panel may be due to absorption by intergalactic gas. The narrow line of O III at 508\AA (bottom right panel) is severely affected by intergalactic absorption, and better data would be required to confirm its presence. The same applies to the other lines shown as upper limits in Table [1.](#page-4-0)

BELR line profiles (see profile comparison of Fig. [3\)](#page-2-1). They lack a broad component at their base. Given the limited S/N of the IUE spectrum at the far-UV end, we consider probable that these lines are spurious features instead. For this reason, we will consider these emission-like features as upper limits rather than real detections. The symbol '?' denotes these unconfirmed lines in our various figures.

We find little evidence of the high excitation Ne VIII line at 775 Å reported by Telfer et al. (2002) and Scott et al. (2004) in their respective composite spectrum, and we favor the identification of O iv λ 789 Å instead. Because the line spectrum of Ton 34 is of unusually low excitation as shown below in Sect. [3.1,](#page-3-1) we do not believe that the high excitation lines of Mgx and Ne viii (listed in Table [1\)](#page-4-0) are present at a detectable level.

2.5. Originality and limitations of the data

As can be gathered from Fig. [2,](#page-2-0) the strongest emission features in the far-UV coincide with the position of lines observed or expected in quasar spectra (Sect. [3.1\)](#page-3-1). However, the limited quality of the data and the possible coincidence of absorbers at inconvenient spectral positions prevent us from deriving incontrovertible conclusions. In the case of the narrower features (O III λ 508 Å, O iv λ 554 Å, O v λ 630 Å and O iv λ 609 Å), better quality data is required to confirm or discard their presence, as discussed in

Fig. 4. Line flux ratios renormalized to $Ly\alpha = 100$ of different species as a function of line wavelength (\hat{A}) . These were extracted from the the spectrum of Ton 34 (filled squares) and from the radio-loud (crosses) and radio-quiet (open lozenge) composite spectra of Telfer et al. (2002). Only ratios that have a counterpart in Ton 34 are shown. Larger squares correspond to ratios for which the difference between Ton 34 and the composites exceeds a factor 10 (also shown in bold face in Col. 5 of Table [1\)](#page-4-0). The symbol '?' denotes upper limits of unconfirmed lines in Ton 34. The gray filled triangle indicates the position of $Ly\alpha$.

Sect. [2.4.](#page-1-2) Clearly, new observations are needed in all wave bands down to the X-rays. In what follows, we will take the data at face value and present photoionization and shock models that attempt to reproduce the far-UV lines.

3. Modelling of the line spectrum

3.1. Line ratio comparison with composite quasar spectra

We now quantify to which degree the emission lines differ in Ton 34 from the 'average' quasar. To achieve this, we list the line ratios characterizing the radio-loud (Col. 3) and radio-quiet (Col. 4) composite spectra of Telfer et al. (2002) in Table [1.](#page-4-0) Comparison between Ton 34 and these two sets of ratios require some caution, since significant line ratio variations exist among quasars. For instance, Telfer et al. (2002) reported that the RMS deviation of line fluxes between the different quasars amounted to as much as 50– 70% for the strong lines of C iv λ 1549, O vi λ 1035 and Ly α . Hence, intrinsic differences of less than a factor two between the composites and Ton 34 should not be considered significant.

To facilitate the comparison of Ton 34 with the two composites, we plot their line ratios in Fig. [4.](#page-3-2) Inspection of the Table [1](#page-4-0) or Fig. [4](#page-3-2) reveals that the commonly strong BELR lines of C_{IV}, N_V and O_{VI} are all present in Ton 34. Hence the apparent sharp turndown of the ionizing UV in the range $650-912\text{\AA}$ is not affecting radically the high excitation emission lines. In particular, the $OVI \lambda 1035$ line is quite strong, although not as much as in the two composites. The C iv is substantially weaker, by more than a factor of six in Ton 34 with respect to the radio-quiet composite. Also, the line system $CIII + NIII$ near 980 Å is noticeably weaker, although the flux in this line is difficult to measure accurately due to the uncertainties introduced by the sharp continuum bent and the many $L\nu\alpha$ forest lines.

In the far-UV, we note that the intensity of the $O II + O III$ and $N III + O III$ systems in Ton 34 are a factor of \sim 14 and 18 brighter, respectively, than in the RLQ composite. There is also evidence of significant emission of N III and/or N IV at 764 and 765 Å, which are *not* detected in the composite spectra either.

3.2. Photoionization vs. shock excitation

The line spectrum of Ton 34 show peculiarities that deserve further analysis. In particular, $O II + O III$ (835 Å), N III+O III lines (686–703Å) and N III+N IV (765Å), which are measured with unusual strengths with respect to Lyα. Are these emission features necessarily genuine lines? One possibility is that extinction resonances, unaccounted for in the extinction curve used to model the deep continuum trough (BK07), may induce features that looked like broad emission lines. Another possibility is that $Ly\alpha$ absorbers at intervening redshifts might generate spurious emission features by bracketing narrow continuum regions. Although we cannot rule out either possibility with the current data, both appear unlikely to us, on the ground that the strongest emission features coincide quite well with the position of plausible atomic transitions (see Fig. [2\)](#page-2-0). The two strongest line systems of O II+O III (835 Å) and N III+O III (686– 703\AA) have previously been reported in the RLQ composite, although at a much reduced flux level. We will thus pursue our analysis under the assumption that the observed features are real and consist of low to intermediate excitation permitted lines.

3.2.1. Photoionization calculations

Can photoionization account for the strength of the far-UV permitted lines? We first establish a comparison with published BELR models, and then evaluate the impact of a strongly absorbed ionizing continuum.

Baldwin et al. (1995) showed that by integrating line fluxes over a wide range in gas density n_H and impinging ionizing flux φ_H , one obtains a much improved fit to quasar line spectra. Such models were dubbed "locally optimally emitting clouds" (LOC). Baldwin et al. (1995) also showed that by preferentially selecting the optimal slab density and impinging flux for each individual line, one can derive a line spectrum comparable (within a factor two) to that of a true LOC model. To derive an approximate LOC model, we proceeded as follows. From the grid of photoionization models published by Korista et al. (1997; hereafter KO97), we extracted the highest equivalent width found within the plane φ_H vs n_H , for each line of interest. The particular grid that we selected was labeled AGN4^{[3](#page-3-3)}. It assumes solar abundances and a SED that was defined by KO97, which

The selected grid AGN4 comprises over a 100 lines and is available at<http://www.pa.uky.edu> /∼korista/grids/grids.html

Table 1. Comparison of Ton 34 with composite SEDs and with models

Lines		Observations			Photoionization ^a			$\text{Shocks}^{a,b}$
Species	λ (Å)	RLQ	RQQ	Ton 34	$KO97^c$	$\operatorname{SED} \operatorname{II}^d$	SEDIV^d	$100^d{\rm km\,s^{-1}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
O _{III}	508	< 1	$<1\,$	≤ 5.1	$\overline{\mathcal{L}}$	$10^{-2.4}$	$0.02\,$	8.7
O _{IV}	554	$<1\,$	< 1	≤ 4.1	$\overline{\mathcal{L}}$	0.04	0.13	$5.7\,$
Ne viii	575	< 1	2.1 ^e	\equiv	$\overline{\mathcal{L}}$	$\overline{?}$	$\overline{\mathcal{L}}$	$\overline{?}$
Hel	601	$\overline{}$	$\frac{1}{2}$	≤ 0.5	γ	2.3	0.9	2.9
$O IV + Mg X$	609, 617	<1	< 1	< 4.7	$0.04 + 1.07$	$0.03 + ?$	$0.1 + ?$	$2.0 + ?$
O _V	630	$<1\,$	<1	≤ 2.8	0.6	0.13	$0.25\,$	$0.06\,$
N $III + O$ III	686, 703	1.5^e	<1	27 ± 4.9	$\overline{?}$	$0.05 + 0.35$	$0.05 + 0.78$	$6.8 + 22$
N $II1 + N$ IV	764, 765	<1	<1	11 ± 1.4	$0.02 + 0.08$	$0.04 + 0.14$	$0.04 + 0.04$	$2.2 + 3.2$
Ne VIII $+O$ IV	775, 789	4.7	$5.7\,$	≤ 2.2	$2.4 + 1.1$	$? +0.67$	$? +1.6$	$? + 5.4$
$OII+OIII$	834, 835	$1.2\,$	$1.5\,$	21 ± 0.32	$? +0.5$	$0.04 + 1.4$	$0.04 + 2.4$	$48 + 23$
$CII+NII$	906, 912	\equiv .	$\overline{}$	≤ 2.7	γ	$< 10^{-2}$	$< 10^{-2}\,$	$22 + 4.1$
$CIII+NIII$	977, 991	$8.1\,$	11.7	5.4 ± 0.64	$2.9 + 0.7$	$4.9 + 0.8$	$3.9 + 0.5$	$13 + 7.0$
$Ly\beta + OVI$	1025, 1035	19.1	18.1	14 ± 1.75	$1.1 + 20$	$0.37 + 1.5$	$0.36 + 21.5$	$2.5 + 10^{-5}$
C _{II}	1037	$\overline{}$	$-$	blended f	$\overline{?}$	0.02	0.02	4.1
N $II + He$ II	1084	$5.6\,$	$5.5\,$		$0.07 + 0.6$	$0.01 + ?$	$0.01 + ?$	$4.3 + ?$
S IV	1067	< 1	<1	5.7 ± 0.23	$\overline{?}$	0.97	0.95	$1.6\,$
Fe III	1123	0.28	2.2		0.01	0.01	$< 10^{-2}\,$	0.07
C _{III}	1176	0.44	0.43		$0.4\,$	4.8	4.9	1.7
S $III + Si$ III	1194, 1207	1.5	0.47	blended f	$0.04 + 1.0$	$0.02 + 0.11$	$0.02 + 0.06$	$0.5 + 5.0$
$_{\rm Ly\alpha}$	1216	100	100	$100^g \pm 10.2$	100	100	100	100
O _V	1218		$\frac{1}{2}$	blended f	$5.3\,$	$2.7\,$	12.8	10^{-4}
N v	1240	$22.0\,$	17.5	4.4 ± 0.2	$3.0\,$	1.5	6.7	0.03
Si II	1262	0.27	0.41	$=$	0.08	$0.03\,$	0.05	1.3
$O I + Si II$	1302, 1305	1.9	2.5	2.1 ± 0.24	$0.07 + 0.03$	$10^{-5.6} + 0.01$	$10^{-5.0} + 0.02$	$10^{-6.9} + 0.15$
C _{II}	1335	$\rm 0.35$	1.0	1.4 ± 0.13	$0.7\,$	0.63	0.7	44
$Si IV + O IV$	1397, 1402	8.6	11.9	9.2 ± 0.8	$3.5 + 2.4$	$2.9 + 1.8$	$1.1 + 0.8$	$3.9 + 0.9$
N _{IV}	1486	$2.8\,$	0.6	1.3 ± 0.2	2.6	$3.5\,$	4.5	0.3
C_{IV}	1549	52	38	12 ± 1.2	59	44	61.4	15
He II	1640	1.2	$\mathbf{1}$	$0.7 \pm ?$	$3.0\,$	4.0	9.5	2.1
O _{III}	1664	2.3	0.7	\equiv	7.8	8.1	6.3	0.7

 a Some observational entries in Cols. 3–5 corresponds to the sum of two different lines. For the corresponding models in Cols. 6–9, we list separately each line intensity using $a +$ symbol as separator.

^b Redward of 1700 Å (down to the infrared), the 100 km s^{-1} shock does not generate any strong lines. For completeness, the only other lines of significant brightness are Si III 1896 Å, Ca II 3969 Å and Ca II 3934 Å, whose intensities reach 3.6%, 1.5% and 3% of the intensity of Ly α , respectively. As for the (optical and UV) Fe II multiplet line systems, we can't say since they are not considered by MAPPINGS Ic. Shortward of 400 Å, we expect the HeII lines to be strong, with HeII 304 Å reaching 80% of $Ly\alpha$.

 \degree A crude model that approximates the optimally locally emitting BELR model described by Baldwin et al. (1995). Each line's peak emissivity was extracted from the grid AGN4 of photoionization calculations published by Korista et al. (1997)

^d These three models were computed with MAPPINGS Ic assuming an initial density $n_{\rm H}^0$ of 4×10^9 cm⁻³ and solar metallicities. The ionization parameter is 0.04 for the two photoionized models and zero for the shock model. At these densities, the $Ly\alpha$ luminosities per unit area of photoionized or shocked gas are 3.2×10^7 , 4.4×10^7 and 5.5×10^5 erg cm⁻² s⁻¹ for models shown in Cols. 7, 8 and 9, respectively. These would scale approximately in proportion to $n_{\rm H}^0$.

 e^e Measurement by one of us (YK) using the composite spectra lent by R. Telfer.

f The strong neighboring lines of Ly α or O VI makes the determination of a meaningful upper limit impossible.

⁹ The Ly α flux in Ton 34 is measured to be 6.9×10^{-13} erg cm⁻² s⁻¹ corresponding to an equivalent width of 57 Å.

peaks at 22 eV. It is the closest to our SED II with a 18.5 eV turnover (Fig. 1; see also Haro-Corzo et al. 2007).

The line ratios from this approximated LOC model are shown in Col. 6 of Table 1. Unfortunately, the $NIII+OIII$ line system $(\lambda \lambda 686 - 703 \text{Å})$ was not part of the AGN4 grid nor the O_{II} λ 834 Å line. On the other hand, the N_{III}+N_{IV} system at 765\AA and the OIII line at 835\AA were. The NIII+NIV system is a factor of a few weaker than observed while the λ 835 Å O_{III} line is predicted an order of magnitude weaker than the observed O $II + O$ III system. As we consider unlikely that the λ 834 Å O II line (absent from the AGN4 grid) is stronger than OIII, we conclude that photoionization would have great difficulties in fitting this system. Hence, even locally optimally emitting clouds would not be able to account for the intensities of at least some of the far-UV lines observed in Ton 34.

Could the peculiar shape of the Ton 34 SED be responsible for the unusual strengths of some far-UV lines? Out of curiosity, we calculated with the multipurpose code MAPPINGS Ic (Ferruit et al. 1997; Binette et al. 1989) pho-

toionization models using SED_{II} to compare with the absorbed SED IV, characterized by the deep trough. We assumed solar metallicities (Anders & Grevesse 1989) and a gas density of $4 \times 10^{9} \text{ cm}^{-3}$ $4 \times 10^{9} \text{ cm}^{-3}$. The ionization parameter⁴ was varied until a maximum in the $O III/H\beta$ (λ 1663/ λ 4861) ratio was found, which occurred at $U = 0.04$. The models were truncated at a depth where H is 10% ionized. These calculations with $U = 0.04$ using either SED II or IV (both plotted in Fig. [1\)](#page-1-1) are reported in Cols. 7 and 8 of Table [1,](#page-4-0) respectively. Because there are fewer soft ionizing photons in sed iv, we find that the mean energy of the photoelectrons is twice as high as the one given by SED II. Hence, this must result in a hotter plasma and therefore in stronger collisionally excited lines. A comparison of the calculated ratios between the two models and with Ton 34 (Col. 5) reveals that, although many metal lines in Col. 8 (SED IV) are often stronger by a factor of a few with respect to Col. 7 (SED II), the deep UV trough does not result in a sufficient increase in the strengths of the O III+N III lines at 683, 703 Å nor of the O II+O III lines at 835 Å. In conclusion, photoionization predicts far-UV line intensities much too weak in comparison with our measurements. Furthermore, making drastic changes in the shape of the ionizing continuum does not alter this conclusion.

3.2.2. Cooling shock calculations

In view of the difficulties of producing strong permitted lines of O II, O III and N III in the case of pure photoionization, we are lead to consider whether collisional ionization might not be more appropriate.

To investigate this possibility, we calculated with mappings ic a sequence of steady-state plane-parallel shock models with a preshock density of 4×10^9 cm⁻³, assuming again solar metallicities. The postshock temperatures of the different models covered the range 1.0×10^5 -8×10^5 K, corresponding to shock velocities of 75 to 235 km s⁻¹. The pre-ionization state of the shocked gas was determined self-consistently by an iterative scheme, using the ionizing radiation produced within the cooling shock that propagates upstream (Dopita, Binette & Tuohy 1984). The time evolution of the electron and ion temperatures were followed separately until they equalized, making use of the equilibration timescale as defined by Spitzer (1962). Most of the far-UV resonance lines are emitted downstream in layers of densities in the range $10^{10.6} - 10^{11.3}$ cm⁻³, well below the densities of 10^{16} where collisional de-excitation would become a concern for many resonance lines. The elapsed time for the shocked gas to cool to temperatures of 8500 K is about 10 seconds. The adiabatic cooling and recombination of the plasma was followed in time until the ionized fraction reached $\leq 2\%$. Because the integrated columns of the different ions are modest in shocks, line opacities turn out negligible with respect to those of photoionized slabs. For instance, the line center opacity of C III λ 977 and C IV λ 1549 are 20 and 1, respectively, for a 100 km s^{-1} shock, compared to $10^{5.3}$ and $10^{4.9}$ for the photoionization model of Col. 8.

Fig. 5. Line intensities from high density cooling shocks renormalized to $Ly\alpha = 100$ as a function of shock velocity. Solar metallicities have been assumed. A vertical dashed line denotes the velocity of the shock model reproduced in Table [1.](#page-4-0)

The intensities of representative far-UV lines are shown in Fig. [5](#page-5-1) as a function of shock velocity. The calculations show that shocks with gas densities appropriate to the BELR are very efficient in producing strong lines of O II+O III (λ 835 Å) and of N III+O III ($\lambda\lambda$ 686–703 Å) that reach 71% and 29% of Ly α , respectively. We also computed the intensities of many other far-UV lines that might be observable in future observations. Some high excitation lines such as O IV λ 554 Å, O IV λ 789 Å and O v λ 630 Å become intense for shock velocities exceeding 120 km s[−]¹ . By comparing in Table [1](#page-4-0) the observed upper limits for these lines with the computed intensities of O III λ 835 Å or O III λ 703 Å, we find that velocities of order 90–130 km s⁻¹ produce line intensities compatible with the estimated line ra-tios^{[5](#page-5-2)}. To be definite, we adopt the velocity of 100 km s^{-1} for the case model^{[6](#page-5-3)} presented in Col. 9 of Table [1.](#page-4-0)

Shock models by themselves predict far-UV line intensities that are too strong with respect to $Ly\alpha$ (compare Cols. 9 and 5), creating a reverse situation to that of photoionization (Sect. [3.2.1\)](#page-3-0). We are therefore lead to propose a mixed model, in which we ascribe only a fraction of the luminosity of $Ly\alpha$ to be due to shock excitation and the complementary fraction to photoionization. In this mixed model, photoionization would be responsible for the emis-

⁴ We use the customary definition of the ionization parameter $U = \varphi_H/cn_H$, which is the ratio of the density of ionizing photons impinging on the slab φ_H/c to the H density at the face of the slab $n_{\rm H}^{\rm O}$.

⁵ While the measurements for O_{IV} λ 554 Å and O v λ 630 Å formally represent only upper limits, it remains possible that the intensities of these lines are somewhat larger than evaluated given the limited S/N of the IUE-SWP spectrum and the possible presence of many intergalactic absorption lines (this would imply higher shock velocities).

 6 For completeness, Table [1](#page-4-0) includes *all* the lines that the shock model predicts to be stronger than 2% of Ly α within the reported domain of $400-1700$ Å.

sion of the strong near-UV (i.e. classical) lines, while shocks would be contributing of order a third of $L_{V\alpha}$ and (proportionally) all of the far-UV resonance lines shortward of the Lyman limit.

The preshock density $n_{\rm H}^0$ may be significantly higher than assumed above. We find similar line ratios for preshock densities up to 100 times higher. The luminosity per unit area of the shock model in this case exceeds that of the photoionization models presented in Col. 7 and 8 (see footnote a in Table [1\)](#page-4-0). Our code includes three body recombination of H, but not the process of stimulated emission, which prevents us from going beyond a preshock density of $10^{11.6}$ cm⁻³. Beyond this limit, we expect Ly α to be the first line to thermalize, which would further enhance the strengths of the metal lines with respect to $Ly\alpha$.

In summary, the far-UV lines observed in Ton 34 shortward of the Lyman limit are characterized by a much higher excitation energy than the near-UV lines. For this reason, collisional excitation (through shocks) at temperatures significantly higher than that typically provided by photoionization is strongly favored. Calculations with mappings ic show that such temperature regime is ensured when shock excitation of moderate V_s takes place. These shocks would account not only for the far-UV lines, but may contribute significantly to the Fe II multiplet lines that have been proposed to result from mechanical heating by Joly et al. (2007 and references therein).

Acknowledgements. This work was supported by the CONACyT grants J-50296 and J-49594, and the UNAM PAPIIT grant IN118905. Diethild Starkmeth helped us with proofreading.

References

- Anders, E., & Grevesse, N. 1989, Geochim. Cosmochim. Acta 53, 197 Baldwin, J., Ferland, G., Korista, K., & Verner, D. 1995, ApJ, 455, L119
- Baldwin, J. A., Ferland, G. J., Korista, K. T., Hamann, F., & LaCluyz´e, A. 2004, ApJ, 615, 610
- Binette, L., Prieto, A., Szusziewicz, E., & Zheng, W. 1989, ApJ, 343, 135
- Binette, Haro-Corzo, S., L., Krongold, Y., & Andersen, A. C. 2007, in Proc. of The nuclear region, host galaxy and environment of active galaxies, Huatulco, México, 18-20 April 2007, ed. E. Benítez, I. Crúz-Gonzalez & Y. Krongold, Rev. Mexicana Astron. Astrofis. (Conf. Ser.), in press, astroph/0706.2006
- Binette, L. & Krongold, Y. 2007, A&A, submitted (BK07)
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245
- Casebeer, D. A., Leighly, K. M., & Baron, E. 2006, ApJ, 637, 157
- Dopita, M. A., Binette, L., & Tuohy, I. R. 1984, ApJ, 282, 142 M
- Ferruit P., Binette, L., Sutherland, R. S., Pécontal, E. 1997, A&A, 322, 73
- Grandi, S. A. 1981, ApJ, 251, 451
- Grandi, S. A. 1982, ApJ, 255, 25
- Haro-Corzo, S. A. R., Binette, L., Krongold, Y., Benitez, E., Humphrey, A., Nicastro, F., & Rodriguez-Martinez, M. 2007, ApJ, 662, 145
- Joly, M. 1987, A&A, 184, 33
- Joly, M., Véron-Cetty, M. & Véron, P. 2007, in Proc. of The nuclear region, host galaxy and environment of active galaxies, Huatulco, México, 18-20 April 2007, ed. E. Benítez, I. Crúz-Gonzalez & Y. Krongold, Rev. Mexicana Astron. Astrofis. (Conf. Ser.), in press
- Korista, K., Baldwin, J., Ferland, G., & Verner, D. 1997, ApJS, 108, 401 (KO97)
- Lanzetta, K. M., Turnshek, D. A., & Sandoval, J. 1993, ApJS, 84, 109
- Laor, A., Bahcall, J. N., Jannuzi, B. T., Schneider, D. P., & Green, R. F. 1995, ApJS, 99, 1
- Møller, P., & Jakobsen, P. 1990, A&A, 228, 299
- Reimers, D., Köhler, S., Hagen, H.-J., & Wisotzki, L. 1998, ESA SP-413: Ultraviolet Astrophysics Beyond the IUE Final Archive, 579
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525 Scott, J., Kriss, G. A., Brotherton, M. S., Green, R. F., Hutchings,
- J., Shull, J. M., & Zheng, W. 2004, ApJ, 615, 135
	- Sargent, W. L. W., Boksenberg, A., & Steidel, C. C. 1988, ApJS, 68, 539
	- Spitzer, L. 1962, Physics of fully ionized gases (New York: Interscience)
	- Telfer, R. C., Zheng, W., Kriss, G. A., & Davidsen, A. F. 2002, ApJ, 565, 773
	- Tripp, T. M., Bechtold, J., & Green, R. F. 1994, ApJ, 433, 533
	- Véron-Cetty, M.-P., Joly, M., & Véron, P. 2004, A&A, 417, 515
	- Véron-Cetty, M.-P., Joly, M., Véron, P., Boroson, T., Lipari, S., & Ogle, P. 2006, A&A, 451, 851
	- Zheng, W., Kriss, G. A., Telfer, R. C., Grimes, J. P., & Davidsen, A. F. 1997, ApJ, 475, 469

