

Axion-dilaton Gauged S -duality and its Symmetry Breaking

Eiji Konishi*

Department of Physics, School of Science, Kyoto University, Sakyo-ku, Kyoto, 606-8502, Japan

(Dated: February 14, 2019)

The equation of the axion-dilaton gauge potential is derived for gauged S -duality on general type IIB string vacua. Based on this formulation of S -duality, we show how the vacua spontaneously break gauged S -duality, leading to a vanishingly small cosmological constant.

0. In this article we derive the equation for the gauge potential of gauged S -duality[1] on moduli curves in the scalar part of type IIB string universal hypermultiplet. When we consider the space of coupling constant as the Poincaré upper half plane \mathfrak{h} , the moduli curve for gauged S -duality after gauge fixing, is a torus $\Sigma = \Gamma \backslash \mathfrak{h}$ where $\Gamma = SL(2, Z)$.

The two dimensional gauge potential \mathcal{A}_i , $i = 1, 2$ on moduli curve Σ with Ramond-Ramond (R-R) fluxes has the following characteristics.

1. It has three variables: the coupling constant g and vector R-R and Neveu-Schwarz–Neveu-Schwarz (NS-NS) fluxes on Σ . The fluxes f_1 and f_2 are the integrations of the Poincaré length 1-form f along two independent geodesics C_1 and C_2 on \mathfrak{h} . These geodesics are the fundamental classes on Σ and are two circles connecting fixed points of two elements, the S -duality element γ and the identity 1, of Γ .
2. The energy conservation law for gauge modes \mathcal{A}_i is written in terms of the total static mass of an $SL(2, Z)$ multi-plet for gauge potential \mathcal{A}_i . The square of the mass is an eigenvalue of the Laplacian Δ on Σ and the gauge potentials \mathcal{A}_i for $i = 1, 2$ are eigenstates.

The Γ -gauge symmetry on the moduli curve Σ is closely related to the geometry of Σ . Indeed the fundamental group of the moduli curve Σ is determined by the gauge group Γ and we can induce a metric on it.

1. In the following, we derive the equation of the dilaton part of the Γ -gauge potential $\mathcal{A}_\varphi = \mathcal{A}_1$ on the moduli curve Σ . For the operator acting on this, we need to sum the mass operators for the dilaton φ , the axion χ , the R-R flux f_1 and the NS-NS flux f_2 because they constitute a single S -duality multi-plet.

First, we start from the Laplace equation for a massless dilaton φ induced by the gauge potential \mathcal{A}_φ on the moduli curve Σ . The dilaton vacuum expectation value $\langle \varphi \rangle$ is associated with coupling constant g by $\sqrt{-1}e^{-\langle \varphi \rangle} = g$ when we ignore the R-R field of the D-instanton, i.e., axion. The Laplace equation on Σ for the variable g

$$\Delta \mathcal{A}_\varphi(g) = 0 \quad (1)$$

is the equation for \mathcal{A}_φ when the flux variables f_1 and f_2 are fixed.

The square root of the Laplacian Δ for the dilaton part of the potential is $\partial/\partial g$ and is the total mass operator for the gauge mode \mathcal{A}_φ .

For general type IIB string vacua, we will also need to consider the mass contributions from the other components of the S -duality multiplet. So we need to introduce the R-R (or the S -dual NS-NS) flux variables f_1 and f_2 in addition to the coupling constant g in the equation for \mathcal{A}_φ . We assume that the variables f_1 and f_2 appear in the Fourier modes of gauge potential as

$$\mathcal{A}_\varphi \sim \mathcal{A}_\varphi^0 \int \frac{dQ_1 dQ_2}{2\pi} A(Q_1, Q_2) \times \exp \sqrt{-1} \left(\sum_{i=1}^2 f_i (\text{flux}_i \text{ on } \Sigma) \times Q_i (\text{mass}_i) \right) \quad (2)$$

where \mathcal{A}_φ^0 does not depend on f_1 or f_2 .

We interpret the masses Q_i as R-R and NS-NS charges of axion and dilaton in an $SL(2, Z)$ multi-plet. By assuming a gauge potential of this form, we induced the S -duality multiplet in the R-R flux part by the action of $SL(2, Z)$ on the inner product $\vec{f} \cdot \vec{Q}$.

For the operator that acts on \mathcal{A}_φ , we consider

$$\sum_{\gamma_1, \gamma_2} \gamma^{-1} \frac{\partial}{\partial f_\gamma} \gamma \quad (3)$$

where γ_1 and γ_2 are the S -duality and identity elements of the discrete S -duality gauge group Γ . The action of γ_1 is defined as follows. In the $SL(2, Z)$ multiplet we need simultaneously to transform the integral domain geodesics $C_i \in \pi_1(\Sigma, p)$ with $p \in \Sigma$ by $\gamma_1 : {}^t(C_1, C_2) \rightarrow {}^t(C_2, -C_1)$ when we change the coupling constant g to $-1/g$. As a result of this definition, γ_1 and γ_2 select R-R and NS-NS fluxes for axion and dilaton in the gauge mode \mathcal{A}_φ .

Finally, on the moduli curve Σ , we consider the mass of an axion as the soliton mass of the non-perturbative vacuum. Its mass depends on the coupling constant such that, for an axion, strong coupling gives a light mass and weak coupling gives a heavy mass. We need to add to the mass term of the $SL(2, Z)$ multi-plet the mass of an axion given by

$$-\frac{\sqrt{-1}}{g}. \quad (4)$$

*Konishi.Eiji@s04.mbox.media.kyoto-u.ac.jp

As the result, the equation (the definition) of the gauge potential $\mathcal{A}_\varphi(g, f_1, f_2)$ is extended from Eq. 1 to the following energy conservation law for the gauge mode

$$\left(\frac{\partial}{\partial g} - \sum_{i=1}^2 \sigma_{\gamma_i} \left(\frac{\partial}{\partial f_i} \right) + \frac{\sqrt{-1}}{g} \right) \mathcal{A}_\varphi(g, f_1, f_2) = 0. \quad (5)$$

Where we put $\sigma_\gamma(x) = \gamma^{-1}x\gamma$ and the actions γ_1 and γ_2 to gauge potential are defined by

$$\begin{aligned} \gamma_1 \mathcal{A}_\varphi(g, f_1, f_2) &= \mathcal{A}_\varphi(-g^{-1}, f_2, -f_1), \\ \gamma_2 \mathcal{A}_\varphi(g, f_1, f_2) &= \mathcal{A}_\varphi(g, f_1, f_2). \end{aligned} \quad (6)$$

We set the initial condition of Eq. 5 to ensure the consistency with Eq. 1

$$\mathcal{A}_\varphi(g, 0, 0) = \mathcal{A}_\varphi^{\text{Eq. 1}}(g) \quad (7)$$

where $\mathcal{A}_\varphi^{\text{Eq. 1}}(g)$ is a solution of the Laplace equation (Eq. 1). Due to the existence of γ_1 , Eq. 5 is a non-linear equation for \mathcal{A}_φ .

2. Now, based on Refs. 2 and 3, we discuss the relation between the problem of the vanishingly small cosmological constant and the above formulation of gauged S -duality. In particular we show how the type IIB string vacua break Γ -gauge symmetry spontaneously.

It is clear what kind of object we should introduce in order to spontaneously break the Γ -symmetry to the symmetry of a Fuchsian subgroup of degree one Γ_h . The subscript h labels the genus of the Riemann surface $\Sigma_h = \Gamma_h \backslash \mathfrak{h}$. The Γ -symmetry is gravitational on the Poincaré metric of \mathfrak{h} . So this object is the so-called gravitational Higgs variable on \mathfrak{h} . In our case, the Higgs variables as R-R and NS-NS fluxes f_1 and f_2 are introduced by including a cosmological constant Λ_{ptl} that is a function of Higgs variables f_1 and f_2 by analogy with Higgs potential in the action.

When the vacuum has a source term for the Lorentz metric, $\Lambda\eta_{\mu\nu}$, in the equation of motion for type IIB supergravity, the on-shell gauge group Γ is broken.[2] The Goldstone mode $\mathcal{A}_\varphi^{G(\gamma)}$ of this broken symmetry, corresponding to an element γ of the coset Γ/Γ_h , parametrizes

the space of broken symmetries. The gauge mode $\mathcal{A}_\varphi(f_1, f_2)$ that moves by the action of Γ_h around the classical field $\mathcal{A}_{\varphi,0}$ generates a non-zero, positive contribution to the cosmological constant Λ_{ptl} , which is the value of Higgs potential at the dominant variables. And two flux variables f_1 and f_2 are the lengths of two geodesics on \mathfrak{h} as fundamental classes on Σ_h . Λ_{ptl} can be considered to be the almost vanishing vacuum energy as we can see by the following argument.

The massive mode, arising from the spontaneous breakdown of the gauge group Γ to Γ_h , gives a positive contribution to the cosmological constant Λ^+

$$\Lambda^+ = \Lambda_{ptl}(f_1, f_2) + \Lambda_{mass}, \quad (8)$$

where the contribution from the gauge mode is an eigenvalue of the Laplacian Δ_{Γ_h} on Σ_h , namely the square of the mass of the gauge mode. That is

$$\Lambda_{mass} = \text{Spec} \Delta_{\Gamma_h}. \quad (9)$$

The first term in Eq. 8 (Λ_{ptl}) survives under the covariance of the remaining Γ_h -gauge symmetry.

The second term in Eq. 8 (Λ_{mass}) should cancel with the negative vacuum energy of the type IIB supergravity vacuum (for example, a de Sitter vacuum or even a broken supersymmetry vacuum) since the Goldstone mode \mathcal{A}_φ^G recovers the broken S -duality.

The scheme presented here clarifies the idea that S -duality as exact Γ (-gauge) symmetry in type IIB string vacua can be used to resolve the problem of the vanishingly small cosmological constant, as first proposed by Kar, Maharana and Singh.[3] In this interpretation, we can also see that vacua with negative vacuum energies in type IIB supergravity are in a Higgs phase of gauged S -duality.

Acknowledgments.

I wish to dedicate this article to my mentor Yoshitaka Yamamoto. I also wish to thank Professor Toshiya Kawai for valuable discussion, the anonymous referee of PRL for his constructive comments, and Professor Tohru Eguchi for encouragement in the preliminary stage of this work.

-
- [1] J. H. Schwarz, *An $SL(2, Z)$ multiplet of type IIB superstrings*, Phys. Lett. **B360**, 13 (1995); Erratum-ibid. **B364**, 252 (1995) [hep-th/9508143].
 [2] J. Maharana and H. Singh, *Symmetries of four-dimensional string effective action*, Phys. Lett. **B368**, 64

- (1996) [hep-th/9506213].
 [3] S. Kar, J. Maharana and H. Singh, *S-duality and cosmological constant in string theory*, Phys. Lett. **B374**, 43 (1996) [hep-th/9507063].