

Euler Incognito

Thomas L Curtright^{§,#} and David B Fairlie^{†,#}

[§]Department of Physics, University of Miami, Coral Gables, Florida 33124, USA

[†]Department of Mathematical Sciences, Durham University, Durham, DH1 3LE, UK

Abstract

The nonlinear flow equations discussed recently by Bender and Feinberg are all reduced to the well-known Euler equation after change of variables.

Consider for $u(x, t)$ the class of nonlinear PDEs introduced and discussed by Bender and Feinberg [1].

$$u_t = (u_x)^k u \quad (1)$$

where k is a parameter. (Including phases for the variables requires only minor changes in the discussion to follow.) We are actually interested only in $0 \neq k \neq 1$, since $k = 0$ and $k = 1$ are easily understood and well-known. So, upon changing dependent variable to

$$v = \frac{k-1}{k} u^{k/(k-1)} \quad (2)$$

equation (1) becomes

$$v_t = (v_x)^k . \quad (3)$$

Differentiating this with respect to x and making a further change of variable to

$$w = k (v_x)^{k-1} = k (u_x)^{k-1} u \quad (4)$$

we find the familiar Euler-Monge equation in canonical form

$$w_t = w w_x . \quad (5)$$

Thus for any $k \neq 0$ and $k \neq 1$ the original equation (1) is reduced to Euler's through a change of dependent variable, although for technical reasons that are more or less obvious from the explicit constructions, it is often useful to assume $k > 1$. (For $k = 1$ of course, (1) is already the Euler-Monge equation *without* any change of variable.)

As is well-known (cf. [1] or [2] for references) the general solution for w is given implicitly by

$$w = F(x + wt) \quad (6)$$

where F is an arbitrary differentiable function. By using the previous changes of variables and integrating once with respect to x , solutions for v , and hence u , follow from those for w . For example, if F is linear, $w = \frac{x-x_0}{t_0-t}$, $v = \frac{k-1}{k} \frac{(x-x_0)^{k/(k-1)}}{[k(t_0-t)]^{1/(k-1)}}$, and $u = \frac{x-x_0}{[k(t_0-t)]^{1/k}}$.

curtright@physics.miami.edu & david.fairlie@durham.ac.uk

Moreover, (1) leads to two infinite families of local conserved currents whose time and space components are powers of u and u_x , but not higher derivatives. The first family is quickly seen to be

$$\left(J_0^{(n)}, J_1^{(n)} \right) = \left(u^n u_x, u^{n+1} (u_x)^k \right) \quad (7)$$

for any n , not necessarily integer. Obviously, all these currents have simple topological charges. On the solution set of (1), $(n+1) J_\mu^{(n)} = \varepsilon_{\mu\nu} \partial^\nu u^{n+1}$, and $\partial^\mu J_\mu^{(n)} = 0$ immediately follows. The second family of currents may be obtained from the known (non-topological) conserved currents for w , namely $((n+1) w^n, n w^{n+1})$, just by changing variables. Thus¹

$$\begin{aligned} \left(K_0^{(n)}, K_1^{(n)} \right) &= \left((n+1) (v_x)^{n(k-1)}, nk (v_x)^{(n+1)(k-1)} \right) \\ &= \left((n+1) (u_x)^{n(k-1)} u^n, nk (u_x)^{(n+1)(k-1)} u^{n+1} \right). \end{aligned} \quad (8)$$

On the solution set of (1), or equivalently (3), it is straightforward to show $\partial^\mu K_\mu^{(n)} = 0$.

Finally, the linearization of (5) as given in [2] can be used to relate the spatial derivative of (3), or equivalently of (1), to a linear equation. Define

$$\psi(a, x, t) \equiv \frac{1}{a} \left(\exp \left(ak (v_x)^{k-1} \right) - 1 \right). \quad (9)$$

Technically, it is useful to assume $k > 1$ here, especially for slowly varying v . It follows that

$$\left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial a \partial x} \right) \psi = \left(v_t - (v_x)^k \right)_x \times k(k-1) (v_x)^{k-2} \exp \left(ak (v_x)^{k-1} \right). \quad (10)$$

Thus $\left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial a \partial x} \right) \psi = 0$ iff $\left(v_t - (v_x)^k \right)_x = 0$. (If the second factor on the RHS of (10) were to vanish, for both positive and negative a , this would require $k > 2$, $v_x = 0$, and hence also $\left(v_t - (v_x)^k \right)_x = 0$.) Encoding initial data for the nonlinear system in the form (9) therefore allows the data to be evolved linearly. Given a well-behaved solution to the linear equation for ψ , we may then extract the nonlinear data at other times just by taking the limit $v_x(x, t) = \left(\frac{1}{k} \lim_{a \rightarrow 0} \psi(a, x, t) \right)^{1/(k-1)}$. Integrating with respect to x modulo a function of only the time variable yields v , hence u .

Acknowledgment This material is based upon work supported by the National Science Foundation under Grant No. 0555603.

References

- [1] C Bender and J Feinberg, "Does the complex deformation of the Riemann equation exhibit shocks?" arXiv:0709.2727v1 [hep-th].
- [2] T Curtright and D Fairlie, "Extra Dimensions and Nonlinear Equations" J.Math.Phys. 44 (2003) 2692-2703, arXiv:math-ph/0207008v1.

¹In terms of u , as $k \rightarrow 1$ we note that $(n+1)(n+2) J_\mu^{(n)} \rightarrow \partial_x K_\mu^{(n+1)} \rightarrow ((n+2)(u^{n+1})_x, (n+1)(u^{n+2})_x)$.