Testing CPT Symmetry with CMB Measurements

Jun-Qing Xia¹, Hong Li², Xiulian Wang³, and Xinmin Zhang¹

1 Institute of High Energy Physics, Chinese Academy of Science, P. O. Box 918-4, Beijing 100049, P. R. China.

² Department of Astronomy, School of Physics, Peking University, Beijing, 100871, P. R. China.

Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany.

Preprint online version: November 6, 2018

ABSTRACT

Aims. In this paper we study the possibility of testing *CPT* symmetry with Cosmic Microwave Background (CMB) measurements. Methods. Working with an effective lagrangian of the photon with CPT violation $\mathcal{L} \sim p_\mu A_\nu \tilde{F}^{\mu\nu}$ which causes the polarization vectors of the propagating CMB photons rotated, we determine the rotation angle $\Delta\alpha$ using the BOOMERanG 2003 and the WMAP3 angular power spectra.

Results. In this analysis we have included the newly released TC and GC ($l < 450$) information of WMAP3 and found $\Delta \alpha = -6.2 \pm 3.8$ deg at 68% confidence level.

Conclusions. This result increases slightly the significance for the *CPT* violation obtained in our previous paper (Feng et al. 2006) $\Delta \alpha = -6.0 \pm 4.0$ deg (1 σ). Furthermore we examine the constraint on the rotation angle from the simulated polarization data with Planck precision. Our results show that the future Planck measurement will be sensitive to $\Delta \alpha$ at the level of 0.057 deg and able to test the *CPT* symmetry with a higher precision.

Key words. cosmology: theory – (cosmology:) cosmic microwave background – (cosmology:) cosmological parameters

1. Introduction

In the standard model of particle physics *CPT* is a fundamental symmetry. Probing its violation is an important way to searc h for the new physics beyond the standard model. Up to now, *CPT* symmetry has passed a number of high precision experimental tests and no definite signal of its violation has been observe d in the laboratory. So, the present *CPT* violating e ffects, if exist, should be very small to be amenable to the experimental limits.

The *CPT* symmetry could be dynamically violated the expanding universe $(Li et al. 2007)$. The cosmological *CPT* violation mechanism investigated in the literature [\(Li et al. 2002;](#page-3-1) [Li & Zhang 2003;](#page-3-2) [Li et al. 2004](#page-3-3); [Feng et al. 2005;](#page-3-4) [Li et al. 2007\)](#page-3-0) has an interesting feature that the *CPT* violating effects at present time are too small to be detected by the laboratory experiments but large enough in the early universe to account for the generation of matter-antimatter asymmetry [\(Li et al. 2002;](#page-3-1) [Li & Zhang 2003;](#page-3-2) [Li et al. 2004;](#page-3-3) [Li et al. 2007](#page-3-0)). And more importantly, this type of *CPT* violating e ffects could be accumulated to be observable in the cosmological experiments [\(Feng et al. 2005;](#page-3-4) [Li et al. 2007;](#page-3-0) [Feng et al. 2006\)](#page-3-5). With the accumulation of high quality observational data, especially those from the CMB experiments, cosmological observation becomes a powerful way to test the *CPT* symmetry.

Here we study the CMB polarizations and *CPT* violation in the photon sector with an effective lagrangian [\(Carroll et al. 1990;](#page-3-6) [Carroll & Field 1991\)](#page-3-7):

$$
\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \mathcal{L}_{cs} \,, \tag{1}
$$

where $\mathcal{L}_{cs} \sim p_{\mu} A_{\nu} \tilde{F}^{\mu\nu}$ is a Chern-Simons term, p_{μ} is an external vector and $\tilde{F}^{\mu\nu} = (1/2)\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$ is the dual of the

electromagnetic tensor. This Lagrangian is not gauge invariant, but the action is gauge independent if $\partial_{\nu}p_{\mu} = \partial_{\mu}p_{\nu}$. This may be possible if p_{μ} is constant in spacetime or the gradient of a scalar field in the quintessential baryo-/leptogenesis [\(Li & Zhang 2003;](#page-3-2) [Li et al. 2002;](#page-3-1) [De Felice et al. 2003\)](#page-3-8) or the gradient of a function of the Ricci scalar in gravitational baryo- /leptogenesis [\(Li et al. 2004;](#page-3-3) [Davoudiasl et al. 2004](#page-3-9)). The Chern-Simons term violates Lorentz and *CPT* symmetries when the background value of p_μ is nonzero.

One of the physical consequences of the Chern-Simons term is the rotation of the polarization direction of electromagnetic waves propagating over large distances [\(Carroll et al. 1990\)](#page-3-6). From the Lagrangian [\(1\)](#page-0-0), we can directly obtain the equation of motion for the electromagnetic field:

$$
\nabla_{\mu}(\nabla^{\mu}A^{\nu} - \nabla^{\nu}A^{\mu}) = -p_{\mu}\epsilon^{\mu\nu\rho\sigma}(\nabla_{\rho}A_{\sigma} - \nabla_{\sigma}A_{\rho}).
$$
\n(2)

After imposing Lorentz gauge condition $\nabla_{\mu}A^{\mu} = 0$, it becomes:

$$
\nabla_{\mu}\nabla^{\mu}A^{\nu} + R^{\nu}_{\mu}A^{\mu} = -p_{\mu}\epsilon^{\mu\nu\rho\sigma}(\nabla_{\rho}A_{\sigma} - \nabla_{\sigma}A_{\rho})\,,\tag{3}
$$

where R^{ν}_{μ} is the Ricci tensor. With the geometric optics approximation, the solution to the equation of motion is expected to be: $A^{\mu} = \text{Re}[(a^{\mu} + \epsilon b^{\mu} + \epsilon^2 c^{\mu} + ...)e^{iS/\epsilon}],$ where ϵ is a small number. With this ansatz, one can easily see that the Lorentz gauge condition implies $k_{\mu}a^{\mu} = 0$, where the wave vector $k_{\mu} \equiv \nabla_{\mu}S$ is orthogonal to the surfaces of constant phase and represents the direction which photons travel along with. The vector a^{μ} is the product of a scalar amplitude *A* and a normalized polarization vector ε^{μ} , $a^{\mu} = Ae^{\mu}$, with $\varepsilon_{\mu} \varepsilon^{\mu} = 1$. Hence in the Lorentz gauge, the wave vector k_{μ} is orthogonal to the polarization vector ε^{μ} . Substituting this solution into the modified Maxwell equation [\(3\)](#page-0-1) and neglecting the Ricci tensor we have at the leading order of ϵ the equation is $k_{\mu}k^{\mu} = 0$. It indicates that photons still propagate along the null geodesics. The e ffect of Chern-Simons term appears at the next order, $k^{\mu} \nabla_{\mu} \varepsilon^{\nu} = -p_{\mu} \epsilon^{\mu \nu \rho \sigma} k_{\rho} \varepsilon_{\sigma}$. We can see that

*Send o*ff*print requests to*: J.-Q. Xia, e-mail: xiajq@mail.ihep.ac.cn

the Chern-Simons term makes $k^{\mu} \nabla_{\mu} \varepsilon^{\nu}$ not vanished. This means that the polarization vector ε^{ν} is not parallel transported along the light-ray. It rotates as the photon propagates in spacetime.

We consider here the spacetime described by spatially flat Friedmann-Robertson-Walker (FRW) metric. The null geodesics equation is $(k^0)^2 - k^i k^i = 0$. We assume that photons propagate along the positive direction of *x* axis, *i.e.* $k^{\mu} = (k^0, k^1, 0, 0)$ and $k^1 = k^0$. Gauge invariance guarantees that the polarization vector of the photon has only two independent components which are orthogonal to the propagating direction. So, we are only interested in the changes of the components of the polarization vector, ε^2 and ε^3 . Assuming $p_\mu = p_0$ to be a non-vanishing constant, we obtain the following equations: $d\varepsilon^2/d\lambda + \mathcal{H}k^0\varepsilon^2 = p_0k^0\varepsilon^3$, $d\varepsilon^3/d\lambda + \mathcal{H}k^0\varepsilon^3 = -p_0k^0\varepsilon^2$, where we have defined the affine parameter λ which measures the distance along the light-ray, $\bar{k}^{\mu} \equiv dx^{\mu}/d\lambda$, and the reduced expansion rate $\mathcal{H} \equiv \dot{a}/a$. The polarization angle is defined as $\chi \equiv \arctan{(\varepsilon^3/\varepsilon^2)}$. It is easy to find that the rotation angle is

$$
\Delta \chi \equiv \chi_0 - \chi_z = -\int_{\eta_z}^{\eta_0} p_0 \, d\eta = \int_{t_0}^{t_z} p_0 \, \frac{dt}{a} \,, \tag{4}
$$

where the subscript ζ is the redshift of the source when the light was emitted. For CMB photons, the source is the last scattering surface with $z \approx 1100$ $z \approx 1100$ $z \approx 1100$ ¹. The subscript 0 indicates the present time. As we know, a vector rotated by an angle $\Delta \chi$ in a fixed coordinates frame is equivalent to a fixed vector observed in a coordinates frame which is rotated by $-\Delta \chi$. So, with the notion of coordinates frame rotation, the rotation angle is

$$
\Delta \alpha = -\Delta \chi = \int_{t_z}^{t_0} p_0 \frac{dt}{a} = p_0 r_z \,. \tag{5}
$$

with r_z being the comoving distance of the light source away from us. This phenomena is known as "cosmological birefringence". This rotation angle $\Delta \alpha$ can be obtained by observing polarized radiation from distant sources such as radio galaxies, quasars and CMB.

The Stokes parameters *Q* and *U* of the CMB polarization can be decomposed into a gradient-like (*G*) and a curl-like (*C*) component [\(Kamionkowski et al. 1997\)](#page-3-10). For the standard theory of CMB, the *TC* and *GC* cross-correlation power spectra vanish. With the existence of cosmological birefringence, the polarization vector of each photon is rotated by an angle $\Delta \alpha$, and one would observe nonzero *TC* and *GC* correlations, even if they are zero at the last scattering surface. Denoting the rotated quantities with a prime, one gets [\(Feng et al. 2005;](#page-3-4) [Lue et al. 1999\)](#page-3-11):

$$
C_l^{TC} = C_l^{TG} \sin(2\Delta\alpha),
$$

\n
$$
C_l^{GC} = \frac{1}{2} (C_l^{GG} - C_l^{CC}) \sin(4\Delta\alpha),
$$

\n
$$
C_l^{TG} = C_l^{TG} \cos(2\Delta\alpha),
$$

Fig. 1. One dimensional distribution on the rotation angle $\Delta \alpha$ from WMAP3 and B03 data.

$$
C_l^{GG} = C_l^{GG} \cos^2(2\Delta\alpha) + C_l^{CC} \sin^2(2\Delta\alpha),
$$

\n
$$
C_l^{CC} = C_l^{CC} \cos^2(2\Delta\alpha) + C_l^{GG} \sin^2(2\Delta\alpha),
$$
\n(6)

while the temperature power spectrum *TT* remains unchanged.

2. Method and Results

In our study we make a global analysis to the CMB data with the public available Markov Chain Monte Carlo package Cosmo MC^2 MC^2 [\(Lewid & Bridle 2002](#page-3-12)), which has been modified to allow the rotation of the power spectra discussed above, with a new free parameter $\Delta \alpha$. We assume the purely adiabatic initial conditions and impose the flatness condition motivated by inflation. Our most general parameter space is: $P \equiv$ $(\omega_b, \omega_c, \Theta_s, \tau, n_s, \log[10^{10}A_s], r, \Delta \alpha)$, where $\omega_b \equiv \Omega_b h^2$ and $\omega_c \equiv$ $\Omega_c h^2$ are the physical baryon and cold dark matter densities relative to the critical density, Θ_s is the ratio of the sound horizon to the angular diameter distance at decoupling, τ is the optical depth to re-ionization, A_s and n_s characterize the primordial scalar power spectrum, *r* is the tensor to scalar ratio of the primordial spectrum. For the pivot of the primordial spectrum we set $k_{s0} = 0.05 \text{Mpc}^{-1}$. In our calculation we have assumed that the cosmic rotation angle is not too large and imposed a conservative flat prior $-\pi/2 \leq \Delta \alpha \leq \pi/2$.

In our calculation we combine the full data of WMAP3 [\(Spergel et al. 2007;](#page-3-13) [Page et al. 2007;](#page-3-14) [Hinshaw et al. 2007;](#page-3-15) [Jarosik et al. 2007\)](#page-3-16) (including the information of *TC* and *GC* power spectra (*l* < 450)) and BOOMERanG 2003 (B03) [\(Jones et al. 2006;](#page-3-17) [Piacentini et al. 2006;](#page-3-18) [Montroy et al. 2006\)](#page-3-19). We calculate the likelihood of *TT*, *TG*, *GG* and *CC* power spectra using the routine for computing the likelihood supplied by the WMAP team. As for the *TC* and *GC* power spectra data $(l < 450)$, which are just the preliminary suboptimal results currently, we simply assume the Gaussian likelihood function^{[3](#page-1-2)}:

$$
\mathcal{L}_{TC/GC} = \exp\left(-\frac{\chi_{TC/GC}^2}{2}\right),
$$

¹ Besides the CMB photons which come from the last scattering surface, we might observed the different CMB photons which travelled different distances as well. However, we are only interested in linear perturbations of CMB photons in this paper. CMB polarizations are already linear phenomena. They are not existent at the zeroth order. When calculating the variations of polarizations due to *CPT* violation in the perturbation theory up to linear order, we may ignore the fluctuations of the travelling distances of CMB photons. Otherwise, these fluctuations combined with polarizations would give higher order result which are beyond the scope of this paper. So, in this paper, we only consider linear perturbations and we can assume that each CMB photon detected by us travelled the same distance.

² http://cosmologist.info/.

³ We are very grateful to Professor Eiichiro Komatsu (the member of WMAP group) for the email communications on the use of the *TC* and *GC* data of WMAP3. Because the current released *TC* and *GC* information of WMAP3 are not very accurate and do not include the correlations between different *l* values, our simple Gaussian likelihood function Eq.[\(7\)](#page-1-3) is accurate enough for the illustrative purpose.

$$
\chi_{\text{TC/GC}}^2 = \sum_{1} \left(\frac{C_{1,\text{th}}^{\text{TC/GC}} - C_{1,\text{obs}}^{\text{TC/GC}}}{\sigma_1} \right)^2, \tag{7}
$$

where $C_{1,\text{th}}^{\text{TC/GC}}$ and $C_{1,\text{obs}}^{\text{TC/GC}}$ denote the theoretical value and the experimental data of the *TC* or *GC* power spectra of WMAP3, σ_1 denotes the variance of estimated power spectra at each multipole.

In the computation of the CMB spectra, we have considered lensing contributions. Furthermore, we make use of the Hubble Space Telescope (HST) measurement of the Hubble parameter H_0 = 100h km s⁻¹ Mpc⁻¹ by multiplying a Gaussian likelihood function $h = 0.72 \pm 0.08$ [\(Freedman et al. 2001](#page-3-20)). We also impose a weak Gaussian prior on the baryon density $\Omega_b h^2 = 0.022 \pm 0.002$ (1 σ) from the Big Bang Nucleosynthesis [\(Burles et al. 2001\)](#page-3-21). Simultaneously we will also use a cosmic age tophat prior as $10 \text{ Gyr} < t_0 < 20 \text{ Gyr}$.

In Figure [1](#page-1-4) we plot our one-dimensional constraint on the rotation angle $\Delta \alpha$ from the combined WMAP3 and B03 data and find that the current CMB polarization data favor a nonzero rotation angle of the photons. The best fit value of the rotation angle is $\Delta \alpha = -0.122 = -7.0$ deg. Marginalizing the posterior distributions, we find that the mean value of the rotation angle is:

$$
\Delta \alpha = -0.108 \pm 0.067 = -6.2 \pm 3.8 \text{ deg } (1 \sigma) , \qquad (8)
$$

which deviates the unrotated case $\Delta \alpha = 0$ more than 1σ and gives a weak evidence for *CPT* violation. This result is frequency independent, while the Faraday Rotation [\(Scannapieco & Ferreira 1997\)](#page-3-22), which also can give nonzero *TC* and *GC* power spectra, is frequency dependent and of the much smaller order than ours. On the other hand, currently several high precision experimental tests have confirmed the *CPT* symmetry and do not detect apparent *CPT* violation in the laboratory. However, these two results are consistent. From equations [\(4,](#page-1-5)[5\)](#page-1-6) we can find that the size of the effects of *CPT* violation on CMB polarization power spectra is an integrated effect and enhanced during the propagation time, while the laboratory experiments just test the *CPT* symmetry at present. It means that the *CPT* violating effect should be too small to be detected at any specific time and enhanced during the propagation time to be detectable by the CMB measurements.

In Figure [2](#page-3-23) we illustrate the effects of the rotation angle $\Delta \alpha$ on the power spectra for two cases $\Delta \alpha = 0$ and $\Delta \alpha = -0.108$. The basic cosmological parameters we choose for these plots are the best fit values of global fitting: $\Omega_b h^2 = 0.0228$, $\Omega_m h^2 =$ 0.1320, $\tau = 0.099$, $H_0 = 73.1$, $n_s = 0.965$, $A_s = 2.2 \times 10^{-9}$ and *r* = 0.00866. One can see from Figure 2 that the *C*-mode is very sensitive to the rotation angle $\Delta \alpha$. Direct measurements of the *TC* and *GC* power spectra are crucial and would give more stringent constraints than ones of other power spectra.

From the analysis of data points of *TC* and *GC* power spectra of WMAP3 and B03, we can find that this negative rotation angle is slightly preferred. Firstly, in the B03 data, the *TC* power at *l* ∼ 250 and *l* ∼ 350 are both negative, while it is positive at *l* ∼ 450. The *GC* power at *l* ∼ 250, *l* ∼ 350 and *l* ∼ 450 are all negative. Based on the equations [\(6\)](#page-1-7) and Figure [2,](#page-3-23) we can see that the *TC* and *GC* power spectra of B03 really help to obtain this negative rotation angle. On the other hand, the quality of *TC* and *GC* power spectra of WMAP3 seems not very good. The data points are distributed around the line $\Delta \alpha = 0$ proportional and the error bars are very large. Therefore, the WMAP3 polarization data can not give much more contribution for test-

TABLE I. Assumed experimental specifications. The noise parameters Δ_T and Δ_P are given in units of μ K-arcmin.

J skv	ι _{max}	(GHz)	$\theta_{\rm fwhm}$	Δ_T	Δp
0.65	2500	100	9.5°	6.8	10.9
		143	7.1'	6.0	11.4
		217	5.0°	13.1	26.7

ing the *CPT* symmetry and not improve the previous result of rotation angle significantly.

Recently Feng et al. (2006) used the WMAP3 and B03 polarization data to constrain the rotation angle and found the similar result: $\Delta \alpha = -6.0 \pm 4.0$ deg at 1σ confidence level. As we show above, the *TC* and *GC* information are very important for this type of analysis. In their analysis they did not include the *TC* and *GC* information of WMAP3 and set the *TC* and *GC* of power spectra of WMAP3 to be zero, $C_1^{\text{TC}} = 0$ and $C_1^{\text{GC}} = 0$. This factor should affect the final result. Comparing these two results, we can find that the mean value of our result moves toward direction away from the zero and the error bar shrinks a little, which strengthens the conclusion on the *CPT* violation.

Moreover, Cabella et al. (2007) perform a wavelet analysis of the temperature and polarization maps of the CMB delivered by the WMAP3 experiment which include the information of *TC* and *GC* power spectra. They set a limit on the CMB photon rotation angle $\Delta \alpha = -2.5 \pm 3.0$ deg (1 σ). They found no evidence for the *CPT* violation from WMAP3 polarization data. However, in their analysis, they have not included the information of B03 polarization data. We also made an analysis without the B03 data and found that $\Delta \alpha = -1.6 \pm 10.3$ deg (1 σ) which means that the WMAP3 data only did not give any significant evidence for *CPT* violation. Therefore, we believe that they should obtain the similar result with ours if they also include the B03 polarization data in their analysis.

Obviously the current CMB polarization data are not good enough to verify this possible *CPT* violation. We need more ac-curate CMB data such as the Planck measurement^{[4](#page-2-0)} in the near future. In Table I we list the assumed experimental specifications of the future Planck measurement. The likelihood function is $\mathcal{L} \propto \exp(-\chi_{\text{eff}}^2/2)$ and

$$
\chi_{\text{eff}}^2 = \sum_{1} (2l+1) f_{\text{sky}} \left(\frac{A}{|\bar{C}|} + \ln \frac{|\bar{C}|}{|\hat{C}|} + 3 \right), \tag{9}
$$

where f_{sky} denotes the observed fraction of the sky in the real experiments, *A* is defined as:

$$
A = \hat{C}_{l}^{TT}(\bar{C}_{l}^{GG}\bar{C}_{l}^{CC} - (\bar{C}_{l}^{GC})^{2}) + \hat{C}_{l}^{TG}(\bar{C}_{l}^{TC}\bar{C}_{l}^{GC} - \bar{C}_{l}^{TG}\bar{C}_{l}^{CC}) + \hat{C}_{l}^{TC}(\bar{C}_{l}^{TG}\bar{C}_{l}^{GC} - \bar{C}_{l}^{TC}\bar{C}_{l}^{GG}) + \hat{C}_{l}^{TG}(\bar{C}_{l}^{TC}\bar{C}_{l}^{GC} - \bar{C}_{l}^{TG}\bar{C}_{l}^{CC}) + \hat{C}_{l}^{GG}(\bar{C}_{l}^{TT}\bar{C}_{l}^{CC} - (\bar{C}_{l}^{TC})^{2}) + \hat{C}_{l}^{GC}(\bar{C}_{l}^{TG}\bar{C}_{l}^{TC} - \bar{C}_{l}^{TT}\bar{C}_{l}^{GC}) + \hat{C}_{l}^{TC}(\bar{C}_{l}^{TG}\bar{C}_{l}^{GC} - \bar{C}_{l}^{GG}\bar{C}_{l}^{TC}) + \hat{C}_{l}^{GC}(\bar{C}_{l}^{TG}\bar{C}_{l}^{TC} - \bar{C}_{l}^{TT}\bar{C}_{l}^{GC}) + \hat{C}_{l}^{CC}(\bar{C}_{l}^{TT}\bar{C}_{l}^{GG} - (\bar{C}_{l}^{TG})^{2}), \qquad (10)
$$

and $|\bar{C}|$ and $|\hat{C}|$ denote the determinants of the theoretical and observed data covariance matrices respectively,

$$
|\bar{C}| = \bar{C}_{l}^{TT} \bar{C}_{l}^{GG} \bar{C}_{l}^{CC} + 2 \bar{C}_{l}^{TG} \bar{C}_{l}^{TC} \bar{C}_{l}^{GC} - \bar{C}_{l}^{TT} (\bar{C}_{l}^{GC})^{2} - \bar{C}_{l}^{GG} (\bar{C}_{l}^{TC})^{2} - \bar{C}_{l}^{CC} (\bar{C}_{l}^{TG})^{2} ,|\hat{C}| = \bar{C}_{l}^{TT} \bar{C}_{l}^{GG} \bar{C}_{l}^{CC} + 2 \bar{C}_{l}^{TG} \bar{C}_{l}^{TC} \bar{C}_{l}^{GC} - \bar{C}_{l}^{TT} (\bar{C}_{l}^{GC})^{2} - \bar{C}_{l}^{GG} (\bar{C}_{l}^{TC})^{2} - \bar{C}_{l}^{CC} (\bar{C}_{l}^{TG})^{2} . \tag{11}
$$

⁴ http://sci.esa.int/science-e/www/area/index.cfm?fareaid=17/.

Fig. 2. The effects of the rotation angle ∆α on the power spectra of *TT*, *TG*, *GG*, *CC*, *TG* and *GC*. The black solid line is for the case of $\Delta \alpha = 0$ and the red dashed line is for $\Delta \alpha = -0.108$.

The likelihood has been normalized with respect to the maximum likelihood $\chi_{\text{eff}}^2 = 0$, where $\bar{C}_1^{\text{XY}} = \hat{C}_1^{\text{XY}}$. For more details of this calculation we refer the readers to our previous companion paper [\(Xia et al. 2007](#page-3-24)). By the simulated data with Planck accuracy, we find that the standard deviation of the rotation angle will be significantly reduced to $\sigma = 0.057$ deg. This result is much more stringent than the current constraint. Assuming the mean value unchanged, the cosmological *CPT* violation will be confirmed around 100σ confidence level with the future Planck measurement.

In summary, probing the violation of fundamental symmetries is an important way to search for the new physics beyond the standard model. In this paper we have determined the rotation polarization angle $\Delta \alpha$ with the CMB data from the WMAP3 and BOOMERanG 2003. We find that a nonzero rotation angle of the photons is mildly favored: $\Delta \alpha = -6.2 \pm 3.8$ deg (1 σ). Our result shows a small violation of the *CPT* symmetry, albeit not conclusive with the present CMB data. With Planck data our simulations indicate that this type of *CPT* violation could be confirmed significantly, or *CPT* symmetry will be verified with a high precision.

Acknowledgements. We acknowledge the use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA). Support for LAMBDA is provided by the NASA Office of Space Science. We have performed our numerical analysis on the Shanghai Supercomputer Center (SSC). We are grateful to Eiichiro Komatsu for using of the *TC* and *GC* data of WMAP3. We thank Carlo Contaldi, Mingzhe Li and Hiranya Peiris for helpful discussions. This work is supported in part by National Natural Science Foundation of China under Grant Nos. 90303004, 10533010 and 10675136 and by the Chinese Academy of Science under Grant No. KJCX3-SYW-N2.

References

- Burles, S., et al. 2001, ApJ, 552, L1
- Cabella, P., Natoli, P. & Silk, J. 2007, Phys. Rev. D, 76, 123014
- Carroll, S. M., Field, G. B. & Jackiw, R. 1990, Phys. Rev. D, 41, 1231
- Carroll, S. M. & Field, G. B. 1991, Phys. Rev. D, 43, 3789
- Davoudiasl, H., Kitano, R., Kribs, C., Murayama, H. & Steinhardt, P. J. 2004, Phys. Rev. Lett., 93, 201301
- De Felice, A., Nasri, S. & Trodden, M. 2003, Phys. Rev. D, 67, 043509
- Feng, B., Li, H., Li, M. & Zhang, X. M. 2005, Phys. Lett. B, 620, 27
- Feng, B., Li, M., Xia, J. Q., Chen, X. & Zhang, X. M. 2006, Phys. Rev. Lett., 96, 221302
- Freedman, W., et al. 2001, ApJ, 553, 47
- Hinshaw, G., et al. 2007, ApJ, 170, 288
- Jarosik, N., et al. 2007, ApJ, 170, 263
- Jones, W. C., et al. 2006, ApJ, 647, 823
- Kamionkowski, M., Kosowsky, A. & Stebbins, A. 1997, Phys. Rev. D, 55, 7368
- Lewis, A. & Bridle, S. 2002, Phys. Rev. D, 66, 103511.
- Li, H., Li, M. & Zhang, X. M. 2004, Phys. Rev. D, 70, 047302
- Li, M., Wang, X. L., Feng, B. & Zhang, X. M. 2002, Phys. Rev. D, 65, 103511
- Li, M., Xia, J. Q., Li, H. & Zhang, X. M. 2007, Phys. Lett. B, 651, 357
- Li, M. & Zhang, X. M. 2003, Phys. Lett. B, 573, 20
- Lue, A., Wang, L. M. & Kamionkowski, M. 1999, Phys. Rev. Lett., 83, 1506 Montroy, T. E., et al. 2006, ApJ, 647, 813
-
- Page, L., et al. 2007, ApJ, 170, 335
- Piacentini, F., et al. 2006, ApJ, 647, 833 Scannapieco, E. S. & Ferreira, P. G. 1997, Phys. Rev. D, 56, 7493
- Spergel, D. N., et al. 2007, ApJ, 170, 377
- Xia, J. Q., et al. 2007, arXiv: 0708.1111