

Induced Gravity in Deconstructed Space at Finite Temperature

Nahomi KAN^{*)}

Yamaguchi Junior College, 1346-2 Daidou, Hofu-shi, Yamaguchi-ken, Japan

We study self-consistent cosmological solutions for an Einstein universe in a graph-based induced gravity model. Especially, we demonstrate specific results for cycle graphs.

§1. Pre-history

1.1. Induced Gravity

The idea of induced Gravity is, “Gravity emerges from the quantum effect of matter fields.” The one-loop effective action can be expressed as the form:

$$-\frac{1}{2} \int \frac{dt}{t} \sum_i \text{Tr} \exp [-(-\nabla^2 + M_i^2)t]. \quad (1.1)$$

In curved D-dimensional spacetime, the trace part including the D-dimensional laplacian becomes

$$\text{Tr} \exp [-(-\nabla^2)t] = \frac{\sqrt{g}}{(4\pi)^{D/2}} t^{-D/2} (a_0 + a_1 t + \dots), \quad (1.2)$$

where coefficients depend on the background fields and a_1 leads to the Einstein-Hilbert term.

1.2. Dimensional Deconstruction, Spectral Graph Theory and Our story thus far

Dimensional Deconstruction (DD)¹⁾ is equivalent to a higher-dimensional theory with discretized extra dimensions at a low energy scale. An N -sided polygon, called a “moose” diagram, is used to describe this theory. Four-dimensional fields are assigned to vertices and edges of this diagram. In $N \rightarrow \infty$, DD leads to a five-dimensional theory, where the extra space is a circle.

In general, the moose diagram does not necessarily have a continuum limit and the diagram is susceptible of a complicated connection, which is a *graph*. Therefore, DD can be generalized to *Field Theory on a Graph*.²⁾

We have constructed models of one-loop finite induced gravity by using several graphs.³⁾ With the help of knowledge in *Spectral Graph Theory*, we can easily find that the UV divergent terms concern the graph laplacian in DD.

§2. Self-consistent Einstein Universe⁴⁾

The metric of the static Einstein Universe is given by

$$ds^2 = -dt^2 + a^2(d\chi^2 + \sin^2 \chi(d\theta^2 + \sin^2 \theta d\phi^2)), \quad (2.1)$$

^{*)} E-mail: kan@yamaguchi-jc.ac.jp

where a is the scale factor. At finite temperature T , the one-loop effective action is regarded as free energy $F(a, \beta)$ and Einstein equation becomes

$$\frac{\partial(\beta F)}{\partial \beta} = \frac{\partial(\beta F)}{\partial a} = 0, \quad (2.2)$$

where $\beta \equiv 1/T$. In the first model, scalar fields are on $8 C_{N/2}$, vector fields on $4 C_N$ and Dirac fermions on $2 C_{N/2} + 3 C_N$. In the second model, scalar fields are on $16 C_{N/4} + 2 C_{N/2}$, vector fields on $5 C_N$ and Dirac fermions on $4 C_{N/4} + 3 C_N + 2 C_{N/2}$. Here C_N denotes a cycle graph with N vertices, equivalent to an N -sided polygon. In each model, Newton's constant and cosmological constant are calculable and are not given by hand.

We exhibit βF for the first model in Fig 1 and for the second in Fig 2, for large N . The horizontal axis indicates the scale factor a while the vertical one indicates the inverse of temperature T . The scale of each axis is in the unit of N/f . In the first model, cosmological constant is zero and the solution can be found at the maximum of βF , corresponding to be in Casimir regime.⁴⁾ In the second model, the solution in Casimir regime and the solution in Planck regime⁴⁾ are found.

§3. Summary

We have studied self-consistent Einstein Universe in the 'graph theory space'. The solution can be systematically obtained with the help of the graph structure.

Acknowledgements

I would like to thank K. Shiraishi for collaboration in this work, and also the organizers of ICGA8.

References

- 1) N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Rev. Lett. **86** (2001), 4757; C. T. Hill, S. Pokorski and J. Wang, Phys. Rev. D **64** (2001), 105005.
- 2) Nahomi Kan and Kiyoshi Shiraishi, J. Math. Phys. **46** (2005), 112301.
- 3) Nahomi Kan and Kiyoshi Shiraishi, Prog. Theor. Phys. **111** (2004), 745.
- 4) M. B. Altaie and J. S. Dowker, Phys. Rev. D **18** (1978), 3557; M. B. Altaie, Phys. Rev. D **65** (2001), 044028; M. B. Altaie, Class. Quantum Grav. **20** (2003), 331; M. B. Altaie and M. R. Setare, Phys. Rev. D **67** (2003), 044018; T. Inagaki, K. Ishikawa and T. Muta, Prog. Theor. Phys. **96** (1996), 847.




Fig. 1. A contour plot of βF in the first model. A solution of Einstein equation can be found at the maximum.




Fig. 2. A contour plot of βF in the second model. Two solutions of Einstein equation can be found at the maximum and at the saddle point.