

ONE-POINT REDUCTIONS OF FINITE SPACES, H-REGULAR CW-COMPLEXES AND COLLAPSIBILITY

JONATHAN ARIEL BARMAK AND ELIAS GABRIEL MINIAN

ABSTRACT. We investigate one-point reduction methods of finite topological spaces and their relationship with collapsibility of complexes. We also introduce the notion of h-regular CW-complex, generalizing the concept of regular CW-complex and prove that the h-regular CW-complexes are modeled, up to homotopy, by their associated finite spaces (or face posets). This is accomplished by generalizing a classical result of McCord on simplicial complexes.

1. INTRODUCTION

There exists a natural correspondence between topologies and preorders defined on a finite set X , which was originally investigated by Alexandroff [1]. Under this correspondence, T_0 -topologies can be regarded as order relations. On the other hand, finite spaces are closely related to simplicial complexes: Following McCord [8] one can associate to a finite simplicial complex K , the finite T_0 -space $\mathcal{X}(K)$ which corresponds to the poset of simplices of K . Moreover, there exists a weak homotopy equivalence $K \rightarrow \mathcal{X}(K)$. Conversely, one can associate to a given finite T_0 -space X the simplicial complex $\mathcal{K}(X)$ of its non-empty chains and a weak homotopy equivalence $\mathcal{K}(X) \rightarrow X$. Finite posets have become a useful tool to investigate homotopy theory of polyhedra. The finite space point of view seems to be more appropriate since it allows the development of new techniques based on the combinatorics and the topology of these objects (see for example Peter May's notes [6, 7], Stong's article [13] and our previous papers [2, 3]).

This article deals with one-point reductions. We investigate the cases in which removing a particular point of the space does not affect its homotopy, weak homotopy or simple homotopy type. Our starting point is Theorem 3.10 of [3], which relates simplicial collapses with collapses of finite spaces. More explicitly, we have proved in [3] that a collapse $X \searrow Y$ between finite spaces induces a collapse $\mathcal{K}(X) \searrow \mathcal{K}(Y)$ between their associated simplicial complexes and a simplicial collapse $K \searrow L$ induces a collapse between the associated finite spaces. One advantage of working with finite spaces is that the elementary collapses in this context are very simple to handle and describe: they consist of removing a single point of the space, which is called a *weak point*. The beat points introduced by Stong [12] and the weak points defined in [3] constitute particular cases of one-point reductions.

In this paper we introduce the notions of γ -point and γ -collapse which provide a more general method of reduction. More precisely, we prove below the following

Proposition 3.10. *If $x \in X$ is a γ -point, the inclusion $i : X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence.*

2000 *Mathematics Subject Classification.* 06A06, 52B70, 55P15, 55U05, 57Q05, 57Q10.

Key words and phrases. Finite Topological Spaces, Posets, Simplicial Complexes, Regular CW-complexes, Collapses, Weak Homotopy Types.

This also improves an old result which appears for example in [14, Proposition 5.8]. Moreover, we prove that the converse of Proposition 3.10 also holds provided x is neither maximal nor minimal (see Theorem 3.13). Therefore, the elementary γ -collapses describe almost all possible one-point reductions.

Although a γ -expansion does not induce in general a simplicial expansion, it does induce a simple homotopy equivalence. We also investigate collapsibility and γ -collapsibility of the joins $X \oplus Y$ of finite spaces in terms of the collapsibility of X and Y . This sheds some light on the analogous results for simplicial joins.

In the last section of the paper we introduce the concept of *h-regular CW-complex*. Recall that a CW-complex is called regular if the characteristic maps of its cells are homeomorphisms. It is known that if K is a regular CW-complex, its face poset $\mathcal{X}(K)$ is a finite model of K , i.e. it is weak homotopy equivalent to K . For general CW-complexes the associated finite space $\mathcal{X}(K)$ does not give relevant information about the topology of K (cf. [4]). A CW-complex is h-regular if all its closed cells are contractible subcomplexes. In particular, regular complexes are h-regular.

We prove that if K is a finite h-regular complex, there is a weak homotopy equivalence $K \rightarrow \mathcal{X}(K)$. This theorem generalizes McCord's result for finite simplicial complexes and extends substantially the class of cell complexes which can be modeled (up to homotopy) by their face posets (compare with [4]). The paper ends with the following result which relates collapses of h-regular complexes with γ -collapses.

Theorem 4.9. *Let L be a subcomplex of an h-regular complex K . If $K \searrow L$, then $\mathcal{X}(K) \searrow \mathcal{X}(L)$.*

2. PRELIMINARIES

In this section we recall the basic notions on finite spaces and their relationship with finite posets and simplicial complexes. For more details, we refer the reader to [3, 7, 8, 12].

Given a T_0 -topology τ on a finite set X , we define for each $x \in X$ the (open) set U_x as the intersection of all open sets containing x . Recall that a topological space X is called T_0 if for every pair of points in X there exists some open set containing one and only one of them. The order associated to the T_0 -topology τ is given by $x \leq y$ if $x \in U_y$. This application establishes a one to one correspondence between T_0 -topologies and order relations on the set X . Therefore we can regard finite T_0 -spaces as finite posets and viceversa. It is not hard to see that a function is continuous if and only if it is order preserving. Moreover if $f, g : X \rightarrow Y$ are two maps such that $f(x) \leq g(x)$ for every $x \in X$, then they are homotopic.

The *order complex* $\mathcal{K}(X)$ of a finite T_0 -space X is the simplicial complex whose simplices are the non-empty chains of X . It is also denoted $\Delta(X)$ by some authors. There exists a weak homotopy equivalence from the geometric realization $|\mathcal{K}(X)|$ to X , i.e. a map which induces isomorphisms in all homotopy groups [8].

Example 2.1. Consider the finite T_0 -space X represented by the Hasse diagram shown in Fig. 1. The order complex of X is homeomorphic to the Möbius Strip.




Fig. 1: Hasse diagram of X




Fig. 2: $\mathcal{K}(X)$ is a triangulation of the Möbius Strip with twelve 2-simplices

The application \mathcal{K} is also defined on maps, moreover if $f : X \rightarrow Y$ is a map between finite T_0 -spaces, there is a commutative diagram

$$\begin{array}{ccc} |\mathcal{K}(X)| & \xrightarrow{|\mathcal{K}(f)|} & |\mathcal{K}(Y)| \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

Conversely one can associate a finite space (the *face poset*) $\mathcal{X}(K)$ to each finite simplicial complex K which is the poset of simplices of K ordered by inclusion. Since $\mathcal{K}(\mathcal{X}(K)) = K'$ is the barycentric subdivision of K , there exists a weak homotopy equivalence $|K| \rightarrow \mathcal{X}(K)$.

Let X be a finite T_0 -space. A point $x \in X$ is a *down beat point* if the set $\hat{U}_x = U_x \setminus \{x\}$ of points smaller than x has a maximum, and it is an *up beat point* if the set $\hat{F}_x = F_x \setminus \{x\}$ of points which are greater than x has a minimum. Here F_x denotes the closure of $\{x\}$ in X . If x is a beat point (down or up), $X \setminus \{x\} \hookrightarrow X$ is a strong deformation retract. Moreover, X is contractible(=dismantlable poset) if and only if one can remove beat points one at the time to obtain a space of one point [12].

Following [3], we say that a point $x \in X$ is a *weak point* if \hat{U}_x or \hat{F}_x is contractible. Note that this definition generalizes the definition of a beat point since any finite space with maximum or minimum is contractible. In this case, the inclusion $X \setminus \{x\} \hookrightarrow X$ need not be a homotopy equivalence, but it is a weak homotopy equivalence. Note that in the context of finite spaces, weak homotopy equivalences are not in general homotopy equivalences.

The notion of weak point gives rise to the following notion of collapse for finite spaces.

Definition 2.2. If $x \in X$ is a weak point, we say that X *collapses* to $X \setminus \{x\}$ by an *elementary collapse*. We denote this by $X \searrow_x X \setminus \{x\}$. We say that X *collapses* to Y (or Y expands to X), and write $X \searrow Y$, if there is a sequence of elementary collapses which starts in X and ends in Y . The space X is *collapsible* if it collapses to a point. Finally, X and Y are *simply equivalent*, denoted by $X \searrow Y$, if there exists a sequence of collapses and expansions that starts in X and ends in Y .

In contrast with the classical situation, where a simple homotopy equivalence is a special kind of homotopy equivalence, homotopy equivalent finite spaces are simply equivalent. The relationship between collapses of finite spaces and simplicial complexes is given by the following

Theorem 2.3.

- (a) *Let X and Y be finite T_0 -spaces. Then, X and Y are simply equivalent if and only if $\mathcal{K}(X)$ and $\mathcal{K}(Y)$ have the same simple homotopy type. Moreover, if $X \searrow Y$ then $\mathcal{K}(X) \searrow \mathcal{K}(Y)$.*
- (b) *Let K and L be finite simplicial complexes. Then, K and L are simple homotopy equivalent if and only if $\mathcal{X}(K)$ and $\mathcal{X}(L)$ are simply equivalent. Moreover, if $K \searrow L$ then $\mathcal{X}(K) \searrow \mathcal{X}(L)$.*

The proof of this theorem can be found in [3]. In this paper we give an alternative proof of the fact that a collapse of finite spaces induces a collapse between the associated complexes.

3. γ -POINTS AND REDUCTION METHODS

In this section we delve deeper into the study of one-point reductions of finite spaces. As we pointed out above, we investigate the cases in which removing a particular point from a finite space does not affect its homotopy, weak homotopy or simple homotopy type.

Recall that the join of two topological spaces X, Y is defined as the quotient space $X * Y = X \times Y \times I / \sim$, where I denotes the unit interval, $(x, y, 0) \sim (x', y, 0)$ and $(x, y, 1) \sim (x, y', 1)$ for every $x, x' \in X$ and $y, y' \in Y$. The *simplicial join* $K * L$ of two simplicial complexes K and L is the complex

$$K * L = K \cup L \cup \{\sigma \cup \tau \mid \sigma \in K, \tau \in L\}.$$

The cone aK of a simplicial complex K is the join of K with a vertex $a \notin K$. It is well known that for finite simplicial complexes K and L , the geometric realization $|K * L|$ is homeomorphic to $|K| * |L|$.

There is an analogous construction for finite spaces.

Definition 3.1. The (*non-Hausdorff*) *join* $X \oplus Y$ of two finite T_0 -spaces X and Y is the disjoint union $X \sqcup Y$ keeping the given ordering within X and Y and setting $x \leq y$ for every $x \in X$ and $y \in Y$.

Special cases of joins are the non-Hausdorff cone $\mathbb{C}(X) = X \oplus D^0$ and the non-Hausdorff suspension $\mathbb{S}(X) = X \oplus S^0$ of any finite T_0 -space X . Here D^0 denotes the singleton (0-cell) and S^0 the discrete space on two points (0-sphere).

Remark 3.2. $\mathcal{K}(X \oplus Y) = \mathcal{K}(X) * \mathcal{K}(Y)$.

Given a point x in a finite T_0 -space X , the *star* C_x of x consists of the points which are comparable with x , i.e. $C_x = U_x \cup F_x$. Note that C_x is always contractible since $1_{C_x} \leq f \geq g$ where $f : C_x \rightarrow C_x$ is the map which is the identity on F_x and the constant map x on U_x , and g is the constant map x . The *link* of x is the subspace $\hat{C}_x = C_x \setminus \{x\}$. In case we need to specify the ambient space X , we will write \hat{C}_x^X . Note that $\hat{C}_x = \hat{U}_x \oplus \hat{F}_x$.

Proposition 3.3. *Let X and Y be finite T_0 -spaces. Then $X \oplus Y$ is contractible if and only if X or Y is contractible.*

Proof. Assume X is contractible. Then there exists a sequence of spaces

$$X = X_n \supsetneq X_{n-1} \supsetneq \dots \supsetneq X_1 = \{x_1\}$$

with $X_i = \{x_1, x_2, \dots, x_i\}$ and such that x_i is a beat point of X_i for every $2 \leq i \leq n$. Then x_i is a beat point of $X_i \oplus Y$ for each $2 \leq i \leq n$ and therefore, $X \oplus Y$ deformation retracts to $\{x_1\} \oplus Y$ which is contractible. Analogously, if Y is contractible, so is $X \oplus Y$.

Now suppose $X \oplus Y$ is contractible. Then there exists a sequence

$$X \oplus Y = X_n \oplus Y_n \supsetneq X_{n-1} \oplus Y_{n-1} \supsetneq \dots \supsetneq X_1 \oplus Y_1 = \{z_1\}$$

with $X_i \subseteq X$, $Y_i \subseteq Y$, $X_i \oplus Y_i = \{z_1, z_2, \dots, z_i\}$ such that z_i is a beat point of $X_i \oplus Y_i$ for $i \geq 2$.

Let $i \geq 2$. If $z_i \in X_i$, z_i is a beat point of X_i unless it is a maximal point of X_i and Y_i has a minimum. In the same way, if $z_i \in Y_i$, z_i is a beat point of Y_i or X_i has a maximum. Therefore, for each $2 \leq i \leq n$, either $X_{i-1} \subseteq X_i$ and $Y_{i-1} \subseteq Y_i$ are deformation retracts (in fact, one inclusion is an identity and the other inclusion is strict), or one of them, X_i or Y_i , is contractible. This proves that X or Y is contractible. \square

Corollary 3.4. *Let X be a finite T_0 -space. Then $x \in X$ is a weak point if and only if its link \hat{C}_x is contractible.*

In [3] we proved that a collapse $X \searrow Y$ of finite spaces induces a simplicial collapse $\mathcal{K}(X) \searrow \mathcal{K}(Y)$. We exhibit here an alternative proof of this result, using Corollary 3.4 and the following easy lemma whose proof we omit.

Lemma 3.5. *Let aK be a simplicial cone of a finite complex K . Then K is collapsible if and only if $aK \searrow K$.*

We study first a particular case (cf. [10, Theorem 3.3]).

Theorem 3.6. *If x is a beat point of a finite T_0 -space X , then $\mathcal{K}(X) \searrow \mathcal{K}(X \setminus \{x\})$. In particular, if X is contractible, $\mathcal{K}(X)$ is collapsible.*

Proof. Since x is a beat point, there exists $x' \in X$ and subspaces $Y, Z \subseteq X$ such that $\hat{C}_x = Y \oplus \{x'\} \oplus Z$. Then the link $lk(x)$ of the vertex x in $\mathcal{K}(X)$ is collapsible, since $lk(x) = \mathcal{K}(\hat{C}_x) = x' \mathcal{K}(Y \oplus Z)$. By the previous lemma, the star $st(x) = xlk(x)$ collapses to $lk(x) = \mathcal{K}(X \setminus \{x\}) \cap st(x)$. Thus, $\mathcal{K}(X) = \mathcal{K}(X \setminus \{x\}) \cup st(x) \searrow \mathcal{K}(X \setminus \{x\})$. \square

Theorem 3.7. *If $X \searrow Y$, then $\mathcal{K}(X) \searrow \mathcal{K}(Y)$.*

Proof. We may assume that $Y = X \setminus \{x\}$, where $x \in X$ is a weak point. By Corollary 3.4, \hat{C}_x is contractible and then $\mathcal{K}(\hat{C}_x)$ is collapsible. Now the result follows as in the proof of Theorem 3.6. \square

Note that if \hat{C}_x is collapsible (but not necessarily contractible), we also have that $\mathcal{K}(X) \searrow \mathcal{K}(X \setminus \{x\})$.

It is known that if K and L are finite simplicial complexes and one of them is collapsible, then $K * L$ is also collapsible. As far as we know the converse of this result is an open problem (see [15, (4.1)]). In the setting of finite spaces, the analogous result and its converse hold.

Proposition 3.8. *Let X and Y be finite T_0 -spaces. Then $X \oplus Y$ is collapsible if and only if X or Y is collapsible.*

Proof. We proceed as in Proposition 3.3, replacing beat points by weak points and deformation retractions by collapses. Note that if x_i is a weak point of X_i , then x_i is also a weak point of $X_i \oplus Y$, since $\hat{C}_{x_i}^{X_i \oplus Y} = \hat{C}_{x_i}^{X_i} \oplus Y$ is contractible by Proposition 3.3.

On the other hand, if z_i is a weak point of $X_i \oplus Y_i$ and $z_i \in X_i$, then by Proposition 3.3, z_i is a weak point of X_i or Y_i is contractible. \square

Corollary 3.4 motivates the following definition.

Definition 3.9. A point x of a finite T_0 -space X is a γ -point if \hat{C}_x is homotopically trivial (i.e. if all its homotopy groups are trivial).

Note that weak points are γ -points. It is not difficult to see that both notions coincide in spaces of height less than or equal to 3. This is because any space of height 2 is contractible if and only if it is homotopically trivial. However, this is false for spaces of height greater than 3.

Let x be a γ -point of a finite T_0 -space X . We will show that the inclusion $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence. Note that since \hat{U}_x and \hat{F}_x need not be homotopically trivial, we cannot proceed as we did in [3]. However, in this case, one has the following pushout

$$\begin{array}{ccc} |\mathcal{K}(\hat{C}_x)| & \longrightarrow & |\mathcal{K}(C_x)| \\ \downarrow & & \downarrow \\ |\mathcal{K}(X \setminus \{x\})| & \longrightarrow & |\mathcal{K}(X)| \end{array}$$

Where $|\mathcal{K}(\hat{C}_x)| \rightarrow |\mathcal{K}(C_x)|$ is a homotopy equivalence and $|\mathcal{K}(\hat{C}_x)| \rightarrow |\mathcal{K}(X \setminus \{x\})|$ satisfies the homotopy extension property. Therefore $|\mathcal{K}(X \setminus \{x\})| \rightarrow |\mathcal{K}(X)|$ is a homotopy equivalence. This proves the following

Proposition 3.10. *If $x \in X$ is a γ -point, the inclusion $i : X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence.*

As we mentioned in the introduction, this result improves an old result which appears for example in Walker's Thesis [14, Proposition 5.8], which asserts, in the language of finite spaces, that $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence provided \hat{U}_x or \hat{F}_x is homotopically trivial. By Proposition 3.17 below, it is clear that a point x is a γ -point if \hat{U}_x or \hat{F}_x is homotopically trivial, but the converse is false.

We will show that the converse of Proposition 3.10 is true in most cases. First, we need some results.

Proposition 3.11. *Let x be a point of a finite T_0 -space X . The inclusion $i : X \setminus \{x\} \hookrightarrow X$ induces isomorphisms in all homology groups if and only if the subspace \hat{C}_x is acyclic.*

Proof. Apply the Mayer-Vietoris sequence to the triple $(\mathcal{K}(X); \mathcal{K}(C_x), \mathcal{K}(X \setminus \{x\}))$. \square

Remark 3.12. If X and Y are non-empty finite T_0 -spaces with n and m connected components respectively, the fundamental group $\pi_1(X \oplus Y)$ is a free product of $(n-1)(m-1)$ copies of \mathbb{Z} . In particular if $x \in X$ is neither maximal nor minimal, the fundamental group of $\hat{C}_x = \hat{U}_x \oplus \hat{F}_x$ is a free group.

Theorem 3.13. *Let X be a finite T_0 -space, and $x \in X$ a point which is neither maximal nor minimal and such that $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence. Then x is a γ -point.*

Proof. If $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence, \hat{C}_x is acyclic by Proposition 3.11. Then $\pi_1(\hat{C}_x)$ is a perfect group and therefore trivial, by Remark 3.12. Now the result follows from the Hurewicz Theorem. \square

The theorem fails if x is maximal or minimal as the next example shows.

Example 3.14. Let X be an acyclic finite T_0 -space with non-trivial fundamental group. Let $\mathbb{S}(X) = X \cup \{-1, 1\}$ be its non-Hausdorff suspension. Then $\mathbb{S}(X)$ is also acyclic and $\pi_1(\mathbb{S}(X)) = 0$. Therefore it is homotopically trivial. Hence, $X \cup \{1\} \hookrightarrow \mathbb{S}(X)$ is a weak homotopy equivalence. However -1 is not a γ -point of $\mathbb{S}(X)$.

Using the relativity principle of simple homotopy theory [5, (5.3)] one can prove that if x is a γ -point, $|\mathcal{K}(X \setminus \{x\})| \rightarrow |\mathcal{K}(X)|$ is a simple homotopy equivalence. In fact this holds whenever $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence.

Theorem 3.15. *Let X be a finite T_0 -space and let $x \in X$. If the inclusion $i : X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence, it induces a simple homotopy equivalence $|\mathcal{K}(X \setminus \{x\})| \rightarrow |\mathcal{K}(X)|$. In particular $X \setminus \{x\} \frown X$.*

Proof. Since $|\mathcal{K}(X \setminus \{x\})|$ is a strong deformation retract of $|\mathcal{K}(X)|$ and the open star of x ,

$$\overset{\circ}{st}(x) = |\mathcal{K}(X)| \setminus |\mathcal{K}(X \setminus \{x\})|$$

is contractible, then by [5, (20.1)], the Whitehead Torsion $\tau(|\mathcal{K}(X)|, |\mathcal{K}(X \setminus \{x\})|) = 0$. \square

This result essentially shows that one-point reductions are not sufficient to describe all weak homotopy types of finite spaces. Of course they are sufficient to reach all finite models of spaces with trivial Whitehead group. On the other hand, note that the fact that $X \setminus \{x\}$ and X have the same weak homotopy type does not imply that the inclusion $X \setminus \{x\} \hookrightarrow X$ is a weak homotopy equivalence.

Definition 3.16. If $x \in X$ is a γ -point, we say that there is an *elementary γ -collapse* from X to $X \setminus \{x\}$. A finite T_0 -space X *γ -collapses* to Y if there is a sequence of elementary γ -collapses that starts in X and ends in Y . We denote this by $X \searrow Y$. If X γ -collapses to a point, we say that it is γ -collapsible.

In contrast to collapses, a γ -collapse does not induce in general a collapse between the associated simplicial complexes. For example, if K is any triangulation of the Dunce hat, $\mathbb{C}(\mathcal{X}(K)) \searrow \mathcal{X}(K)$, but $aK' \not\searrow K'$ since K' is not collapsible (see Lemma 3.5).

We finish this section analyzing the relationship between γ -collapsibility and joins.

Proposition 3.17. *Let X and Y be finite T_0 -spaces. Then*

- (i) $X \oplus Y$ is homotopically trivial if X or Y is homotopically trivial.
- (ii) $X \oplus Y$ is γ -collapsible if X or Y is γ -collapsible.

Proof. If X or Y is homotopically trivial, $|\mathcal{K}(X)|$ or $|\mathcal{K}(Y)|$ is contractible and then so is $|\mathcal{K}(X)| * |\mathcal{K}(Y)| = |\mathcal{K}(X \oplus Y)|$. Therefore $X \oplus Y$ is homotopically trivial.

The proof of (ii) follows as in Proposition 3.3. If $x_i \in X_i$ is a γ -point, $\hat{C}_{x_i}^{X_i \oplus Y} = \hat{C}_{x_i}^{X_i} \oplus Y$ is homotopically trivial by item (i) and then x_i is a γ -point of $X_i \oplus Y$. \square

There is an analogous result for acyclic spaces that follows from the Künneth formula for joins [9].

Note that the converse of these results are false. To see this, consider two finite simply connected simplicial complexes K, L such that $H_2(|K|) = \mathbb{Z}_2$, $H_2(|L|) = \mathbb{Z}_3$ and $H_n(|K|) = H_n(|L|) = 0$ for every $n \geq 3$. Then $\mathcal{X}(K)$ and $\mathcal{X}(L)$ are not acyclic, but $\mathcal{X}(K) \oplus \mathcal{X}(L)$, which is weak homotopy equivalent to $|K| * |L|$, is acyclic by the Künneth formula and, since it is simply connected (see [9] or Remark 3.12), it is homotopically trivial.

A counterexample for the converse of item (ii) is the following.

Example 3.18. Let K be a triangulation of the Dunce hat. Then, $\mathcal{X}(K)$ is a homotopically trivial finite space of height 3. The non-Hausdorff suspension $\mathbb{S}(\mathcal{X}(K)) = \mathcal{X}(K) \cup \{-1, 1\}$ is γ -collapsible since 1 is a γ -point and $\mathbb{S}(\mathcal{X}(K)) \setminus \{1\}$ has maximum. However $\mathcal{X}(K)$ is not collapsible and then $\mathbb{S}(\mathcal{X}(K))$ is not collapsible by Proposition 3.8. Moreover $\mathcal{X}(K)$ and S^0 are not γ -collapsible either because their heights are less than or equal to 3.

4. H-REGULAR COMPLEXES

Recall that a CW-complex K is regular if for each (open) cell e^n , the characteristic map $D^n \rightarrow \overline{e^n}$ is a homeomorphism, or equivalently, the attaching map $S^{n-1} \rightarrow K$ is a homeomorphism onto its image \dot{e}^n , the boundary of e^n . As usual, D^n denotes the closed n -disk and S^{n-1} , the $(n-1)$ -sphere. In this case, it can be proved that the closure $\overline{e^n}$ of each cell is a subcomplex, which is equivalent to say that \dot{e}^n is a subcomplex.

Let e and e' be two cells of a regular complex. We say that e is a *face* of e' if $e \subseteq \overline{e'}$. This will be denoted by $e \leq e'$. If K is a regular complex, its barycentric subdivision K' is the simplicial complex whose vertices are the cells of K and whose simplices are the sets $\{e_1, e_2, \dots, e_n\}$ such that e_i is a face of e_{i+1} .

We can define, as in the case of simplicial complexes, the face poset $\mathcal{X}(K)$ of a regular complex K , which is the set of cells ordered by \leq . Note that $\mathcal{K}(\mathcal{X}(K)) = K'$, which is homeomorphic to K and therefore $\mathcal{X}(K)$ is a finite model of K , i.e. it has the same weak homotopy type as K .

Example 4.1. The following figure (Fig. 3) shows a regular structure for the real projective plane $\mathbb{R}P^2$. The edges are identified in the way indicated by the arrows. It has three 0-cells, six 1-cells and four 3-cells. Therefore its face poset has 13 points (Fig. 4).




Fig. 3




Fig. 4

In this section we introduce the concept of *h-regular complex*, generalizing the notion of regular complex. Given an h-regular complex K , one can define $\mathcal{X}(K)$ as before. In general, K and $\mathcal{K}(\mathcal{X}(K))$ are not homeomorphic. However we prove that $\mathcal{X}(K)$ is a finite model of K . We also study the relationship between collapses of h-regular complexes and of finite spaces.

Definition 4.2. A CW-complex K is *h-regular* if the attaching map of each cell is a homotopy equivalence with its image and the closed cells \bar{e}^n are subcomplexes of K .

Thus, regular complexes are h-regular.

Proposition 4.3. Let $K = L \cup e^n$ be a CW-complex such that \dot{e}^n is a subcomplex of L . Then \bar{e}^n is contractible if and only if the attaching map $\varphi : S^{n-1} \rightarrow \dot{e}^n$ of the cell e^n is a homotopy equivalence.


Proof. Suppose $\varphi : S^{n-1} \rightarrow \dot{e}^n$ is a homotopy equivalence. Since $S^{n-1} \hookrightarrow D^n$ has the homotopy extension property, the characteristic map $\psi : D^n \rightarrow \bar{e}^n$ is also a homotopy equivalence.

Suppose now that \bar{e}^n is contractible. The map $\bar{\psi} : D^n/S^{n-1} \rightarrow \bar{e}^n/\dot{e}^n$ is a homeomorphism and therefore it induces isomorphisms in homology and, since \bar{e}^n is contractible, by the long exact sequence of homology it follows that $\varphi_* : H_k(S^{n-1}) \rightarrow H_k(\dot{e}^n)$ is an isomorphism for every k .

If $n \geq 3$, $\pi_1(\dot{e}^n) = \pi_1(\bar{e}^n) = 0$ and by a theorem of Whitehead, φ is a homotopy equivalence. If $n = 2$, \dot{e}^n is just a graph and since $\varphi_* : H_1(S^1) \rightarrow H_1(\dot{e}^n)$ is an isomorphism, the attaching map φ is a homotopy equivalence. Finally, if $n = 1$, since the cell is contractible, φ is one-to-one and therefore a homeomorphism. \square

Corollary 4.4. A CW-complex is h-regular if and only if the closed cells are contractible subcomplexes.

Example 4.5. The following are four different h-regular structures for the Dunce hat which are not regular structures. In each example the edges are identified in the way indicated by the arrows.



For h-regular complexes we define the notion of a face in the same way as we did in the regular case. We define also the *associated finite space* (or *face poset*) $\mathcal{X}(K)$ as the poset of cells of K ordered by the face relation \leq .

The proof of the following lemma is standard.

Lemma 4.6. *Let $K \cup e$ be a CW-complex, let $\psi : D^n \rightarrow \bar{e}$ be the characteristic map of the cell e and let A be a subspace of \dot{e} . We denote $C_e(A) = \{\psi(x) \mid x \in D^n \setminus \{0\}, \psi(\frac{x}{\|x\|}) \in A\} \subseteq \bar{e}$. Then*

- (1) *If $A \subseteq \dot{e}$ is open, $C_e(A) \subseteq \bar{e}$ is open.*
- (2) *$A \subseteq C_e(A)$ is a strong deformation retract.*

Theorem 4.7. *If K is a finite h-regular complex, $\mathcal{X}(K)$ is a finite model of K .*

Proof. We define recursively a weak homotopy equivalence $f_K : K \rightarrow \mathcal{X}(K)$.

Assume $f_{K^{n-1}} : K^{n-1} \rightarrow \mathcal{X}(K^{n-1}) \subseteq \mathcal{X}(K)$ is already defined and let $x = \psi(a)$ be a point in an n -cell e^n with characteristic map $\psi : D^n \rightarrow \bar{e}^n$. If $a = 0 \in D^n$, define $f_K(x) = e^n$. Otherwise, define $f_K(x) = f_{K^{n-1}}(\psi(\frac{a}{\|a\|}))$.

In particular note that if $e^0 \in K$ is a 0-cell, $f_K(e^0) = e^0 \in \mathcal{X}(K)$. Notice also that if L is a subcomplex of K , $f_L = f_K|_L$.

We will show by induction on the number of cells of K , that for every cell $e \in K$, $f_K^{-1}(U_e)$ is open and contractible. This will prove that f_K is continuous and, by McCord's Theorem [8, Theorem 6], a weak homotopy equivalence.

Let e be a cell of K . Suppose first that there exists a cell of K which is not contained in \bar{e} . Take a maximal cell e' (respect to the face relation \leq) with this property. Since K is h-regular, $L = K \setminus e'$ is a subcomplex. By induction, $f_L^{-1}(U_e)$ is open in L . Then $f_L^{-1}(U_e) \cap e' \subseteq e'$ is open and by the previous lemma, $C_{e'}(f_L^{-1}(U_e) \cap e') \subseteq \bar{e}'$ is open. Therefore,

$$f_K^{-1}(U_e) = f_L^{-1}(U_e) \cup C_{e'}(f_L^{-1}(U_e) \cap e')$$

is open in K .

Moreover, since $f_L^{-1}(U_e) \cap e' \subseteq C_{e'}(f_L^{-1}(U_e) \cap e')$ is a strong deformation retract, so is $f_L^{-1}(U_e) \subseteq f_K^{-1}(U_e)$. By induction $f_K^{-1}(U_e)$ is contractible.

In the case that every cell of K is contained in \bar{e} , $f_K^{-1}(U_e) = \bar{e} = K$, which is open and contractible. \square

As an application we deduce that the finite spaces associated to the h-regular structures of the Dunce hat considered in Example 4.5 are all homotopically trivial. The first one is a contractible space of 5 points, the second one is a collapsible and non-contractible space of 13 points and the last two are non-collapsible spaces of 15 points since they do not have weak points. Here we exhibit the Hasse diagram of the space associated to the third h-regular structure of the Dunce hat.






Fig. 5: A homotopically trivial non-collapsible space of 15 points.


Example 4.8. Let K be the space which is obtained from a square by identifying all its edges as indicated.



We verify that K is homotopy equivalent to S^2 using techniques of finite spaces. Consider the following h-regular structure of K



which consists of three 0-cells, three 1-cells and two 2-cells. The Hasse diagram of the associated finite space $\mathcal{X}(K)$ is



The 0-cell b is an up beat point of $\mathcal{X}(K)$ and the 1-cell ab is a down beat point of $\mathcal{X}(K) \setminus \{b\}$. Therefore K is weak homotopy equivalent to $\mathcal{X}(K) \setminus \{b, ab\}$ which is a (minimal) finite model of S^2 (see [2]). In fact $\mathcal{X}(K) \setminus \{b, ab\} = S^0 \oplus S^0 \oplus S^0$ is weak homotopy equivalent to $S^0 * S^0 * S^0 = S^2$.

In [3] we proved that a collapse $K \searrow L$ of finite simplicial complexes induces a collapse $\mathcal{X}(K) \searrow \mathcal{X}(L)$ between the associated finite spaces. This is not true when K and L are regular complexes. Consider $L = \mathcal{K}(W)$ the associated simplicial complex to the Wallet W (see Fig. 6 below), and K the CW-complex obtained from L by attaching a regular 2-cell e^2 with boundary $\mathcal{K}(\{a, b, c, d\})$ and a regular 3-cell e^3 with boundary $L \cup e^2$.




Fig. 6: W

Note that the complex K is regular and collapses to L , but $\mathcal{X}(K) = \mathcal{X}(L) \cup \{e^2, e^3\}$ does not collapse to $\mathcal{X}(L)$ because $\hat{U}_{e^3}^{\mathcal{X}(K) \setminus \{e^2\}} = \mathcal{X}(L) = W'$ is not contractible. However, one can prove that a collapse $K \searrow L$ between h -regular CW-complexes induces a γ -collapse $\mathcal{X}(K) \searrow \mathcal{X}(L)$.

Theorem 4.9. *Let L be a subcomplex of an h -regular complex K . If $K \searrow L$, then $\mathcal{X}(K) \searrow \mathcal{X}(L)$.*

Proof. Assume $K = L \cup e^n \cup e^{n+1}$. Then e^n is an up beat point of $\mathcal{X}(K)$. Since $K \searrow L$, $\overline{e^{n+1}} \searrow L \cap \overline{e^{n+1}} = \dot{e}^{n+1} \searrow e^n$. In particular $\dot{e}^{n+1} \searrow e^n$ is contractible and then

$$\hat{C}_{e^{n+1}}^{\mathcal{X}(K) \setminus \{e^n\}} = \mathcal{X}(\dot{e}^{n+1} \searrow e^n)$$

is homotopically trivial. Therefore

$$\mathcal{X}(K) \searrow \mathcal{X}(K) \setminus \{e^n\} \searrow \mathcal{X}(L).$$

□

REFERENCES

- [1] P.S. Alexandroff. *Diskrete Räume*. *Mathematiceski Sbornik* (N.S.) 2 (1937), 501-518.
- [2] J.A. Barmak and E.G. Minian. *Minimal finite models*. *Journal of Homotopy and Related Structures* 2 (2007), 127-140.
- [3] J.A. Barmak and E.G. Minian. *Simple homotopy types and finite spaces*. *Advances in Math.* (to appear).
- [4] A. Björner. *Posets, Regular CW Complexes and Bruhat Order*. *Europ. J. Combinatorics* 5 (1984), 7-16.
- [5] M.M. Cohen. *A Course in Simple Homotopy Theory*. Springer-Verlag New York, Heidelberg, Berlin (1970).
- [6] J.P. May. *Finite topological spaces*. Notes for REU (2003). Available at <http://www.math.uchicago.edu/~may/MISCMaster.html>
- [7] J.P. May. *Finite spaces and simplicial complexes*. Notes for REU (2003). Available at <http://www.math.uchicago.edu/~may/MISCMaster.html>
- [8] M.C. McCord. *Singular homology groups and homotopy groups of finite topological spaces*. *Duke Mathematical Journal* 33 (1966), 465-474.
- [9] J. Milnor. *Construction of universal bundles, II*. *Ann. of Math.* 63 (1956), 430-436.
- [10] T. Osaki. *Reduction of finite topological spaces*. *Interdisciplinary Information Sciences* 5 (1999), 149-155.
- [11] D. Quillen. *Homotopy properties of the poset of non-trivial p -subgroups of a group*. *Advances in Math.* 28 (1978), 101-128.

- [12] R.E. Stong. *Finite topological spaces*. Trans. Amer. Math. Soc. 123 (1966), 325-340.
- [13] R.E. Stong. *Group actions on finite spaces*. Discrete Mathematics 49 (1984), 95-100.
- [14] J.W. Walker. *Topology and combinatorics of ordered sets* Thesis, M.I.T., Cambridge, MA (1981).
- [15] V. Welker. *Constructions preserving evasiveness and collapsibility*. Discrete Math. 207 (1999), 243-255.
- [16] J.H.C Whitehead. *Simplicial spaces, nuclei and m-groups*. Proc. London Math. Soc. 45 (1939), 243-327.
- [17] J.H.C Whitehead. *Simple homotopy types*. Amer. J. Math. 72 (1950), 1-57.
- [18] E.C. Zeeman *On the dunce hat*. Topology 2 (1964), 341-358.

DEPARTAMENTO DE MATEMÁTICA, FCEyN, UNIVERSIDAD DE BUENOS AIRES, BUENOS AIRES, ARGENTINA

E-mail address: `jbarmak@dm.uba.ar`

E-mail address: `gminian@dm.uba.ar`