

EXAMPLES OF PARA-COCYCLIC OBJECTS INDUCED BY BD -LAWS

GABRIELLA BÖHM AND DRAGOȘ ȘTEFAN

Dedicated to Freddy Van Oystaeyen on the occasion of his 60th birthday.

ABSTRACT. In a recent paper [BS], we gave a general construction of a para-cocyclic structure on a cosimplex, associated to a so called admissible septuple – consisting of two categories, three functors and two natural transformations, subject to compatibility relations. The main examples of such admissible septuples were induced by algebra homomorphisms. In this note we provide more general examples coming from appropriate (‘locally braided’) morphisms of monads.

2000 *Mathematics Subject Classification*: 16E40 (primary), 16W30 (secondary)

INTRODUCTION

History of cyclic homology started in the early eighties of the last century. The seminal works of the pioneers A Connes, B Tsygan, D Quillen and J-L Loday were motivated by looking for non-commutative generalizations of de Rham cohomology on one hand, and Lie algebra homology of matrices on the other hand.

In the subsequent decades cyclic homology has been extensively studied and became an important tool in diverse areas of mathematics, such as homological algebra, algebraic topology, Lie algebras, algebraic K-theory and so non-commutative differential geometry. Thus by most various motivations, lots of examples have been constructed. In order to study general features of the examples, and also to be able to construct new ones, it was desirable to find a unifying general description. A fundamental first step in this direction was made by A Kaygun in [Kay], who gave a construction of para-(co)cyclic objects in symmetric monoidal categories in terms of (co)monoids. In particular, in this way he managed to describe in a universal form all examples arising from Hopf cyclic theory (upto cyclic duality, cf. [KR]). Motivated by a generalization to bialgebroids (over non-commutative rings, in which case the underlying bimodule categories are not symmetric), in [BS] we made a further step of generalization and constructed para-(co)cyclic objects in arbitrary categories, in terms of (co)monads. Kaygun’s construction can be recovered as a particular case when the (co)monads in question are induced by (co)monoids.

Mean examples of Kaygun’s construction are induced by algebras over a commutative ring. By analogy, in this paper we show that appropriate monad morphisms (which are ‘locally braided’ in a sense to be described) induce examples of para-cocyclic objects in [BS].

The paper is organized as follows. In Section 1 we recall some facts about monads and BD -laws that are used in the paper. In Section 2 we introduce the notion of a *locally braid preserving morphism* of monads, generalizing a homomorphism of algebras, and we investigate their basic properties that are needed to state and prove our main result. In Section 3 we show that any such morphism determines an ‘admissible septuple’ in the sense of [BS], hence can be used to construct para-cocyclic objects. Here we also illustrate how this construction works in the example of a morphism of algebras in a braided monoidal category (i.e. when there is a *global* braiding). Particular examples will be provided by appropriate homomorphisms of (co)module algebras of a (co)quasitriangular Hopf algebra.

Acknowledgment. It is our pleasure to thank the organizers of the conference ‘*Non-commutative Rings and Geometry*’, Almería, Spain, 18-22 September 2007, held in the honour of the 60th birthday of Freddy Van Oystaeyen. The first author was partially supported by the Hungarian Scientific Research Fund OTKA K 68195 and the Bolyai János Research Scholarship. The second author was supported by Contract 2-CEX06-11-20 of the Romanian Ministry of Education and Research.

1. MONADS AND THE CATEGORY OF THEIR (BI)MODULES

Throughout the paper, we use the notations introduced in [BS]. That is, in the 2-category CAT horizontal composition (of functors) is denoted by juxtaposition, while \circ is used for vertical composition (of natural transformations). For example, for two functors $F : \mathcal{C} \rightarrow \mathcal{C}'$, $G : \mathcal{C}' \rightarrow \mathcal{C}''$ and an object X in \mathcal{C} , instead of $G(F(X))$ we write $GF X$. For two natural transformations $\mu : F \rightarrow F'$ and $\nu : G \rightarrow G'$ we write $G'\mu X \circ \nu F X : GF X \rightarrow G'F' X$ instead of $G'(\mu_X) \circ \nu_{F(X)}$. In equalities of natural transformations we shall omit the object X in our formulae.

We shall also use a graphical representation of morphisms in a category. For functors F_1, \dots, F_n , G_1, \dots, G_m , which can be composed to $F_1 F_2 \dots F_n : \mathcal{D}_1 \rightarrow \mathcal{C}$ and $G_1 G_2 \dots G_m : \mathcal{D}_2 \rightarrow \mathcal{C}$, and objects X in \mathcal{D}_1 and Y in \mathcal{D}_2 , a morphism $f : F_1 F_2 \dots F_n X \rightarrow G_1 G_2 \dots G_m Y$ will be represented vertically, with the domain up, as in Figure 1(a). Furthermore, for a functor $T : \mathcal{C} \rightarrow \mathcal{C}'$, the morphism Tf will be drawn as in (b). Keeping the notation from the first paragraph of this section, the picture representing $\mu G X$ is shown in diagram (c). The composition $g \circ f$ of the morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ will be represented as in diagram (d). For the multiplication t and the unit τ of a monad T on \mathcal{C} (see Definition 1.1), and an object X in \mathcal{C} , to draw tX and τX we shall use the diagrams (e) and (f), while for a distributive law $\natural : RT \rightarrow TR$ (see Definition 1.9) $\natural X$ will be drawn as in the picture (g). If \natural is invertible, the representation of $\natural^{-1} X$ is shown in diagram (h).




FIGURE 1. Diagrammatic representation of morphisms in a category

For simplifying the diagrams containing only natural transformations, we shall always omit the last string, that corresponds to an object in the category.

DEFINITION 1.1. A *monad* on a category \mathcal{C} is a triple (R, r, ρ) , where $R : \mathcal{C} \rightarrow \mathcal{C}$ is a functor, $r : R^2 \rightarrow R$ and $\rho : \text{Id}_{\mathcal{C}} \rightarrow R$ are natural transformations such that the first two diagrams in Figure 2, expressing associativity and unitality, are commutative. We call r and ρ the *multiplication*




FIGURE 2. Monads and morphisms of monads.

and the *unit* of the monad R , respectively.

For two monads (R, r, ρ) and (T, t, τ) on \mathcal{C} , we say that a natural transformation $\varphi : R \rightarrow T$ is a *morphism of monads* if the last two diagrams in Figure 2 are commutative.

EXAMPLE 1.2. Let $(\mathcal{C}, \otimes, \alpha, \iota_l, \iota_r, \mathbf{1})$ be a monoidal category with *tensor product* $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and *unit object* $\mathbf{1} \in \text{OB}(\mathcal{C})$. Recall that the associativity constraint $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$

and the unit constraints $\iota_r X : X \otimes \mathbf{1} \rightarrow X$ and $\iota_l X : \mathbf{1} \otimes X \rightarrow X$ are natural isomorphisms that obey *Pentagon Axiom* and *Triangle Axioms*, cf. [Kas, Chapter XI].

Algebras in monoidal categories can be defined as in the classical case, of algebras over a commutative ring, see [AMS, §1]. To such an algebra \mathbf{R} in \mathcal{C} , with multiplication $\mathbf{r} : \mathbf{R} \otimes \mathbf{R} \rightarrow \mathbf{R}$ and unit $\boldsymbol{\rho} : \mathbf{1} \rightarrow \mathbf{R}$, one associates a monad $R := \mathbf{R} \otimes (-)$ on \mathcal{C} . Its multiplication and unit are respectively defined by the morphisms $rX := (\mathbf{r} \otimes X) \circ \alpha_{\mathbf{R}, \mathbf{R}, X}^{-1}$ and $\rho X := \boldsymbol{\rho} \otimes X$, where X is an arbitrary object in \mathcal{C} . An algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ in \mathcal{C} induces a monad morphism $\varphi \otimes (-) : \mathbf{R} \otimes (-) \rightarrow \mathbf{T} \otimes (-)$ that will be denoted by φ .

DEFINITION 1.3. For a monad (R, r, ρ) on \mathcal{C} , a pair (X, x) is said to be an R -module if X is an object in \mathcal{C} and $x : RX \rightarrow X$ is a morphism such that the first two diagrams in Figure 3, expressing associativity and unitality, are commutative. A *morphism of R -modules* from (X, x) to (Y, y) is a morphism $f : X \rightarrow Y$ in \mathcal{C} such that the third diagram in Figure 3 is commutative. The category of R -modules is denoted by ${}_R\mathcal{C}$.




FIGURE 3. Modules over a monad and morphisms of modules.

EXAMPLE 1.4. For a monad $R = \mathbf{R} \otimes (-)$, induced by an algebra \mathbf{R} in a monoidal category \mathcal{C} as in Example 1.2, ${}_R\mathcal{C}$ is isomorphic to the category $\mathbf{R}\text{-Mod}$ of left modules for the algebra \mathbf{R} . For the definition of modules over an algebra in a monoidal category, see [AMS, §1].

For a monad (R, r, ρ) on a category \mathcal{C} , there is a faithful *forgetful functor* $\rho^* : {}_R\mathcal{C} \rightarrow \mathcal{C}$. It is given by the object map $(X, x) \mapsto X$ and it acts on the morphisms as the identity map. The forgetful functor possesses a left adjoint, the *free functor* $\rho_* : \mathcal{C} \rightarrow {}_R\mathcal{C}$. It has the object map $X \mapsto (RX, rX)$ and it acts on the morphisms by $f \mapsto Rf$.

Assuming that coequalizers exist in \mathcal{C} , it is natural to ask if the category of modules over a monad T on \mathcal{C} has the same property. In what follows we will frequently use the standard result that this question has a positive answer if T preserves coequalizers in \mathcal{C} . Recall that a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ preserves coequalizers if, for any coequalizer (Z, π) in \mathcal{C}

$$X \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} Y \xrightarrow{\pi} Z, \quad (1.1)$$

$(FZ, F\pi)$ is the coequalizer of (Ff, Fg) . Note that in this case the canonical morphisms π and $F\pi$ are epimorphisms (while for an arbitrary epimorphism p in \mathcal{C} , the morphism Fp is not necessarily an epimorphism in \mathcal{D}).

PROPOSITION 1.5. Let (T, t, τ) be a monad on a category \mathcal{C} that preserves coequalizers. Assume that (Z, π) is the coequalizer of a parallel pair of morphisms (f, g) in \mathcal{C} , as in (1.1). If, in addition, X and Y are T -modules such that f and g are morphisms of T -modules, then there is a unique T -module structure on Z such that (Z, π) is the coequalizer of (f, g) in ${}_T\mathcal{C}$. In particular, if any pair of parallel morphisms in \mathcal{C} has a coequalizer then any pair of parallel morphisms in ${}_T\mathcal{C}$ has a coequalizer, too.

Proof. Let x and y denote the actions of T on X and Y , respectively. Since $(TZ, T\pi)$ is the coequalizer of (Tf, Tg) in \mathcal{C} , and $\pi \circ y$ coequalizes (Tf, Tg) , it follows that there is a unique morphism $z : TZ \rightarrow Z$ such that

$$z \circ T\pi = \pi \circ y. \quad (1.2)$$

Our aim is to prove that (Z, z) is a T -module. Since T is a functor, in view of (1.2), associativity of y and of the fact that t is natural, we get

$$z \circ Tz \circ TT\pi = z \circ T\pi \circ Ty = \pi \circ y \circ Ty = \pi \circ y \circ tY = z \circ T\pi \circ tY = z \circ tZ \circ TT\pi.$$

Since T preserves coequalizers, $TT\pi$ is an epimorphism, so the associativity condition in the definition of T -modules is verified. In a similar way one can show that τZ is a right inverse of z . Thus (Z, z) is a T -module and, in view of (1.2), π is a morphism of T -modules.

Let h be a morphism in ${}_T\mathcal{C}$ that coequalizes (f, g) . As (Z, π) is the coequalizer of (f, g) in \mathcal{C} , there is a unique morphism \bar{h} in \mathcal{C} such that $\bar{h} \circ \pi = h$. Obviously, \bar{h} is a morphism of T -modules, so the proposition is proved. \square

Let $\varphi : R \rightarrow T$ be a morphism of monads on \mathcal{C} . We show that one can associate to φ “forgetful” and “free” functors connecting the categories of modules over R and T . The construction of $\varphi^* : {}_T\mathcal{C} \rightarrow {}_R\mathcal{C}$ is quite obvious, hence the proof is left to the reader.

PROPOSITION 1.6. *There exists a functor $\varphi^* : {}_T\mathcal{C} \rightarrow {}_R\mathcal{C}$ such that*

$$\varphi^*(X, x) := (X, x \circ \varphi X) \quad (1.3)$$

and $\varphi^*f = f$, for every T -module (X, x) and every T -module morphism f .

PROPOSITION 1.7. *Let \mathcal{C} be a category with coequalizers. If (R, τ, ρ) and (T, t, τ) are monads on \mathcal{C} , such that T that preserves coequalizers, and $\varphi : R \rightarrow T$ is a morphism of monads then the functor φ^* in Proposition 1.6 has a left adjoint.*

Proof. Let (X, x) be an R -module. It is not difficult to see that (TRX, tRX) and (TX, tX) are T -modules, and that $tX \circ T\varphi X$ and Tx are morphism of T -modules. Let

$$(\varphi_*(X, x), \pi X) := \text{Coeq}(tX \circ T\varphi X, Tx). \quad (1.4)$$

By Proposition 1.5, $\varphi_*(X, x)$ is a T -module such that $\pi X : (TX, tX) \rightarrow \varphi_*(X, x)$ is a morphism of T -modules. By the universal property of coequalizers, for any R -module morphism $f : X \rightarrow Y$, there is a T -module morphism φ_*f such that

$$\varphi_*f \circ \pi X = \pi Y \circ Tf. \quad (1.5)$$

In conclusion, we have constructed a functor φ_* from ${}_R\mathcal{C}$ to ${}_T\mathcal{C}$. Moreover, one can compose the forgetful functor $\rho^* : {}_R\mathcal{C} \rightarrow \mathcal{C}$ with the free functor $\tau_* : \mathcal{C} \rightarrow {}_T\mathcal{C}$ to obtain a functor from ${}_R\mathcal{C}$ to ${}_T\mathcal{C}$, given by $(X, x) \mapsto (TX, tX)$ and $f \mapsto Tf$. Therefore, (1.5) means that $(\pi X)_{X \in \text{Ob } {}_R\mathcal{C}}$ defines a natural transformation $\pi : \tau_*\rho^* \rightarrow \varphi_*$.

It remains to show that φ_* is a left adjoint of φ^* . For objects (X, x) in ${}_R\mathcal{C}$ and (Y, y) in ${}_T\mathcal{C}$ we define

$$\Phi_{X,Y} : \text{Hom}_{{}_T\mathcal{C}}(\varphi_*(X, x), (Y, y)) \rightarrow \text{Hom}_{{}_R\mathcal{C}}((X, x), \varphi^*(Y, y)), \quad \Phi_{X,Y}(f) = f \circ \pi X \circ \tau X.$$

In order to prove that $\Phi_{X,Y}(f)$ is a morphism of R -modules, recall that, by the proof of Proposition 1.5, the module structure on $\varphi_*(X, x)$ is given by the unique morphism \bar{x} satisfying

$$\bar{x} \circ T\pi X = \pi X \circ tX. \quad (1.6)$$

As $\pi X \circ tX \circ T\varphi X = \pi X \circ Tx$, by a simple but tedious computation, one can show that

$$y \circ \varphi Y \circ R\Phi_{X,Y}(f) = \Phi_{X,Y}(f) \circ x,$$

i.e. $\Phi_{X,Y}(f)$ is a morphism of R -modules from (X, x) to $\varphi^*(Y, y)$.

We are going to construct an inverse of $\Phi_{X,Y}$. Let g be a morphism of R -modules from (X, x) to $\varphi^*(Y, y)$. Then $y \circ Tg$ is a morphism of T -modules and coequalizes $(Tx, tX \circ T\varphi X)$. Since $\varphi_*(X, x)$ is the coequalizer in ${}_T\mathcal{C}$ of this pair, there exists a unique T -module morphism $\Theta_{X,Y}(g)$ such that

$$\Theta_{X,Y}(g) \circ \pi X = y \circ Tg.$$

Furthermore, by construction of $\Phi_{X,Y}$ and $\Theta_{X,Y}$, and the facts that τ is natural and y is unital, we get

$$\Phi_{X,Y}(\Theta_{X,Y}(g)) = \Theta_{X,Y}(g) \circ \pi X \circ \tau X = y \circ Tg \circ \tau X = y \circ \tau Y \circ g = g.$$

Thus we deduce that $\Theta_{X,Y}$ is a right inverse of $\Phi_{X,Y}$. Let $f : \varphi_*(X, x) \rightarrow (Y, y)$ be a morphism of T -modules. To conclude, we must prove

$$\Theta_{X,Y}(\Phi_{X,Y}(f)) \circ \pi X = f \circ \pi X.$$

The left hand side of this relation can be rewritten as

$$y \circ T\Phi_{X,Y}(f) = y \circ Tf \circ T\pi X \circ T\tau X = f \circ \bar{x} \circ T\pi X \circ T\tau X = f \circ \pi X \circ tX \circ T\tau X = f \circ \pi X,$$

where for the first two equalities we used the definitions of $\Phi_{X,Y}$ and $\Theta_{X,Y}$. Since f and πX are morphisms of T -modules we obtained the third and the fourth relations, while for the last one we used the definition of monads. \square

EXAMPLE 1.8. Let $(\mathcal{C}, \otimes, \alpha, \iota_l, \iota_r, \mathbf{1})$ be an abelian monoidal category, that is \mathcal{C} be abelian, and $X \otimes (-) : \mathcal{C} \rightarrow \mathcal{C}$ and $(-) \otimes X : \mathcal{C} \rightarrow \mathcal{C}$ be additive and right exact functors, for any $X \in \mathcal{C}$. For details on abelian monoidal categories, the reader is referred to [AMS, §1.]. Following [AMS, (1.11)], for a right \mathbf{R} -module $\mu_X : X \otimes \mathbf{R} \rightarrow X$ and a left \mathbf{R} -module $\mu_Y : \mathbf{R} \otimes Y \rightarrow Y$ one defines $X \otimes_{\mathbf{R}} Y$ by

$$X \otimes_{\mathbf{R}} Y := \text{Coker}(\mu_X \otimes Y - X \otimes \mu_Y).$$

Take another algebra \mathbf{T} in \mathcal{C} . Proceeding as in [AMS, (1.4)] one defines the category $\mathbf{T}\text{-Mod-}\mathbf{R}$ of \mathbf{T} - \mathbf{R} bimodules in \mathcal{C} . By definition, (X, μ_X^l, μ_X^r) is an \mathbf{T} - \mathbf{R} bimodule if $\mu_X^l : \mathbf{T} \otimes X \rightarrow X$ and $\mu_X^r : X \otimes \mathbf{R} \rightarrow X$ define compatible left and right module structures such that

$$\mu_X^l \circ (T \otimes \mu_X^r) = \mu_X^r \circ (\mu_X^l \otimes R) \circ \alpha_{\mathbf{T}, X, \mathbf{R}}^{-1}. \quad (1.7)$$

If (X, μ_X^l, μ_X^r) is a \mathbf{T} - \mathbf{R} bimodule and (Y, μ_Y^l, μ_Y^r) is an \mathbf{R} - \mathbf{S} bimodule, then $X \otimes_{\mathbf{R}} Y$ is a \mathbf{T} - \mathbf{S} bimodule, for any algebra \mathbf{S} in \mathcal{C} . Its left \mathbf{T} -action and right \mathbf{S} -action are respectively induced by $\mu_X^l \otimes Y$ and $X \otimes \mu_Y^r$. As in [AMS, Theorem 1.12], one can prove that $\mathbf{T}\text{-Mod-}\mathbf{R}$ is abelian. Moreover, $\mathbf{R}\text{-Mod-}\mathbf{R}$ is monoidal with respect to $(-) \otimes_{\mathbf{R}} (-)$, see [AMS, Theorem 1.12].

For a monad morphism $\varphi \otimes (-) : \mathbf{R} \otimes (-) \rightarrow \mathbf{T} \otimes (-)$, induced by an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$, the functors in Propositions 1.6 and 1.7 are $\varphi^* : \mathbf{T}\text{-Mod} \rightarrow \mathbf{R}\text{-Mod}$, the ‘restriction of scalars’, and $\varphi^* = \mathbf{T} \otimes_{\mathbf{R}} (-) : \mathbf{R}\text{-Mod} \rightarrow \mathbf{T}\text{-Mod}$, the ‘extension of scalars’.

Distributive laws were introduced by J. Beck [Be]. As we shall see later, they give a way to compose two monads in order to obtain a monad.

DEFINITION 1.9. A *distributive law* between two monads (R, r, ρ) and (T, t, τ) is a natural transformation $\iota : RT \rightarrow TR$ satisfying the four conditions in Figure 4.




FIGURE 4. The definition of distributive laws.

REMARK 1.10. Since we are using for the first time the diagrammatic representation of morphisms, we note that the first and the third relations in Figure 4 can be explicitly written as follows:

$$\iota \circ rT = Tr \circ \iota R \circ R\iota, \quad \iota \circ \rho T = T\rho.$$

EXAMPLE 1.11. Let \mathbf{R} and \mathbf{T} be two algebras in a braided monoidal category $(\mathcal{C}, \otimes, \alpha, \iota_l, \iota_r, \mathbf{1}, \chi)$, with braiding $\chi : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$. Let $R = \mathbf{R} \otimes (-)$ and $T = \mathbf{T} \otimes (-)$ be the monads induced by \mathbf{R} and \mathbf{T} , as in Example 1.2. For an object X in \mathcal{C} , we put

$$\begin{aligned} \mathfrak{l}X &:= \alpha_{\mathbf{T}, \mathbf{R}, X} \circ (\chi_{\mathbf{R}, \mathbf{T}} \otimes X) \circ \alpha_{\mathbf{R}, \mathbf{T}, X}^{-1}, \\ \mathfrak{r}X &:= \alpha_{\mathbf{R}, \mathbf{R}, X} \circ (\chi_{\mathbf{R}, \mathbf{R}} \otimes X) \circ \alpha_{\mathbf{R}, \mathbf{R}, X}^{-1}, \\ \mathfrak{t}X &:= \alpha_{\mathbf{T}, \mathbf{T}, X} \circ (\chi_{\mathbf{T}, \mathbf{T}} \otimes X) \circ \alpha_{\mathbf{T}, \mathbf{T}, X}^{-1}. \end{aligned}$$

One can see easily that $\mathfrak{l} : RT \rightarrow TR$, $\mathfrak{r} : RR \rightarrow RR$ and $\mathfrak{t} : TT \rightarrow TT$ are distributive laws.

DEFINITION 1.12. Consider a monad (T, t, τ) on a category \mathcal{C} . A distributive law $\mathfrak{t} : TT \rightarrow TT$ is said to be a *BD-law* if it satisfies the *YB-equation*:

$$\mathfrak{t}T \circ T\mathfrak{t} \circ \mathfrak{t}T = T\mathfrak{t} \circ \mathfrak{t}T \circ T\mathfrak{t}.$$

For a diagrammatic representation of the *YB-equation* see the first picture in Figure 5.

The distributive laws \mathfrak{t} and \mathfrak{r} in Example 1.11 are, in fact, *BD-laws*.

2. BRAIDED PAIRS OF MONADS AND THEIR MORPHISMS

Our starting point of a construction of para-cocyclic objects will be the following notion.

DEFINITION 2.1. The sextuple $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ is said to be a *braided pair of monads* if and only if the following conditions are satisfied.

- \mathcal{C} is a category, in which any pair of parallel morphisms has a coequalizer.
- (R, r, ρ) and (T, t, τ) are monads on \mathcal{C} , both of which preserve coequalizers.
- $\mathfrak{l} : RT \rightarrow TR$ is an *invertible* distributive law.
- $\mathfrak{t} : TT \rightarrow TT$ and $\mathfrak{r} : RR \rightarrow RR$ are *invertible* *BD-laws* such that conditions (A 1) and (A 2) in Figure 5 hold.




FIGURE 5. *YB-equation* and conditions (A 1)–(A 4).

A morphism of monads $\varphi : R \rightarrow T$ is said to be *braid preserving* if it satisfies conditions (A 3) and (A 4) in Figure 5.

REMARK 2.2. In the proof of several results we do not need all assumptions from the definition of a braid preserving monad morphism in Definition 2.1. Still, for reader's convenience, we prefer to state all of them at the same time.

EXAMPLE 2.3. Let $\varphi : (\mathbf{R}, \mathbf{r}, \rho) \rightarrow (\mathbf{T}, \mathbf{t}, \tau)$ be a morphism of algebras in a braided monoidal category $(\mathcal{C}, \otimes, \alpha, \iota_l, \iota_r, \mathbf{1}, \chi)$. We take the monads R and T as in Example 1.2, and the distributivity laws \mathfrak{l} , \mathfrak{r} and \mathfrak{t} as in Example 1.11. Clearly, $(\mathcal{C}, R, T, \mathfrak{l}, \mathfrak{r}, \mathfrak{t})$ is a braided pair of monads. Moreover, by pushing φ over \mathfrak{r} in the second diagram in (A 3) and under \mathfrak{t} in the second picture in (A 4), it follows that $\varphi := \varphi \otimes (-)$ is a braid preserving monad morphism if, and only if,

$$(\varphi \otimes \mathbf{T}) \circ \chi_{\mathbf{T}, \mathbf{R}} = (\varphi \otimes \mathbf{T}) \circ \chi_{\mathbf{R}, \mathbf{T}}^{-1}, \quad \text{and} \quad \chi_{\mathbf{R}, \mathbf{T}} \circ (\mathbf{R} \otimes \varphi) = \chi_{\mathbf{T}, \mathbf{R}}^{-1} \circ (\mathbf{R} \otimes \varphi). \quad (2.1)$$

For example, if $\chi_{\mathbf{T},\mathbf{R}} = \chi_{\mathbf{R},\mathbf{T}}^{-1}$, then φ is such a morphism. In a similar manner one can show that φ is a braid preserving monad morphism, provided that $\chi_{\mathbf{R},\mathbf{R}} = \chi_{\mathbf{R},\mathbf{R}}^{-1}$ and $\chi_{\mathbf{T},\mathbf{T}} = \chi_{\mathbf{T},\mathbf{T}}^{-1}$. For, it is enough to rewrite the second equalities in (A 3) and (A 4) by pushing φ under \lrcorner^{-1} .

Distributive laws, as we have already remarked, were introduced by Beck in order to construct a monad on the composite of two monads. For two monads (R, r, ρ) and (T, t, τ) , a distributive law $\lrcorner : RT \rightarrow TR$ induces a monad TR on \mathcal{C} , with multiplication and unit

$$Tr \circ tR^2 \circ T\lrcorner R = tR \circ T^2r \circ T\lrcorner R : TRTR \rightarrow TR \quad \text{and} \quad T\rho \circ \tau = \tau R \circ \rho : \text{Id}_{\mathcal{C}} \rightarrow TR.$$

Therefore, it makes sense to speak about ${}_{TR}\mathcal{C}$, the category TR -modules. Its objects can be described equivalently as follows. Let (X, x) be an object in ${}_{TR}\mathcal{C}$. We set $x_T := x \circ T\rho X$ and $x_R := x \circ \tau RX$. One can prove that (X, x_R) is an R -module, (X, x_T) is a T -module and these structures commute in the sense that

$$x_T \circ Tx_R \circ \lrcorner X = x_R \circ Rx_T. \quad (2.2)$$

Conversely, if (X, x_R) is an R -module and (X, x_T) is a T -module that obey relation (2.2) then X is an TR -module with respect to $x := x_T \circ Tx_R$. Moreover, a morphism in \mathcal{C} is a morphism of TR -modules if, and only if, it is a morphism of R -modules and T -modules with respect to the above defined structures. Thus, ${}_{TR}\mathcal{C}$ is isomorphic to the category of triples (X, x_R, x_T) such that x_R and x_T define commuting module structures over R and T , respectively. For details see [KLV].

For any BD -law $\mathfrak{r} : RR \rightarrow RR$, one can deform the monad (R, r, ρ) to obtain a new monad

$$R^{\mathfrak{r}} := R, \quad r^{\mathfrak{r}} := r \circ \mathfrak{r}, \quad \rho^{\mathfrak{r}} := \rho.$$

For a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \lrcorner, t, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$, it follows by the definition of distributive laws and (A 1) that $\lrcorner : R^{\mathfrak{r}}T \rightarrow TR^{\mathfrak{r}}$ is a distributive law too. Hence $TR^{\mathfrak{r}}$ admits a monad structure. By the above considerations, the category of $TR^{\mathfrak{r}}$ -modules is isomorphic to the category of triples (X, x_R^0, x_T) , where (X, x_R^0) is an $R^{\mathfrak{r}}$ -module and (X, x_T) is a T -module such that the compatibility condition

$$x_T \circ Tx_R^0 \circ \lrcorner X = x_R^0 \circ R^{\mathfrak{r}}x_T \quad (2.3)$$

holds. Their morphisms are both T -module, and $R^{\mathfrak{r}}$ -module morphisms. We call the triple (X, x_R^0, x_T) a (T, R) -bimodule and denote their category by ${}_{T}\mathcal{C}_R$.

REMARK 2.4. By the above considerations, in particular, for any monad R and BD -law $\mathfrak{r} : RR \rightarrow RR$, also $\mathfrak{r} : R^{\mathfrak{r}}R \rightarrow RR^{\mathfrak{r}}$ is a distributive law. Hence, as above, $R^e := RR^{\mathfrak{r}}$ is a monad, that is called the *enveloping monad of R* . The category ${}_{R^e}\mathcal{C}$ is isomorphic to the category of (R, R) -bimodules. Recall that an (R, R) -bimodule is an object X in \mathcal{C} , together with two morphisms $x : RX \rightarrow X$ and $x^0 : R^{\mathfrak{r}}X \rightarrow X$ such that (X, x) is an R -module, (X, x^0) is an $R^{\mathfrak{r}}$ -module and these structures commute in the sense that

$$x \circ Rx^0 \circ \mathfrak{r}X = x^0 \circ R^{\mathfrak{r}}x. \quad (2.4)$$


The category of (R, R) -bimodules will be denoted by ${}_{R}\mathcal{C}_R$. In order to simplify notations, the forgetful functor $U : {}_{R}\mathcal{C}_R \rightarrow \mathcal{C}$ (with object map $(X, x, x^0) \mapsto X$) will be omitted in our formulae whenever it causes no danger.

For an *invertible* BD -law $\mathfrak{r} : RR \rightarrow RR$, also $\mathfrak{r}^{-1} : R^{\mathfrak{r}}R^{\mathfrak{r}} \rightarrow R^{\mathfrak{r}}R^{\mathfrak{r}}$ is an (invertible) BD -law. Hence the previous construction can be repeated with $\mathfrak{r} : RR \rightarrow RR$ replaced by $\mathfrak{r}^{-1} : R^{\mathfrak{r}}R^{\mathfrak{r}} \rightarrow R^{\mathfrak{r}}R^{\mathfrak{r}}$. The objects of the resulting category ${}_{R^{\mathfrak{r}}}\mathcal{C}_{R^{\mathfrak{r}}}$ are triples (X, x^0, x) , where (X, x^0) and (X, x) are $R^{\mathfrak{r}}$ and $R = (R^{\mathfrak{r}})^{\mathfrak{r}^{-1}}$ -modules, respectively, such that

$$x^0 \circ R^{\mathfrak{r}}x \circ \mathfrak{r}^{-1}X = x \circ Rx^0.$$

Hence the object map $(X, x, x^0) \mapsto (X, x^0, x)$ induces an isomorphism ${}_{R}\mathcal{C}_R \rightarrow {}_{R^{\mathfrak{r}}}\mathcal{C}_{R^{\mathfrak{r}}}$.

EXAMPLE 2.5. Let R and T be the monads coming from algebras \mathbf{R} and \mathbf{T} in a braided monoidal \mathcal{C} , as in Example 1.2. We denote by \lrcorner, \mathfrak{r} and \mathfrak{t} the distributive laws defined in Example 1.11. In this case, the category ${}_{T}\mathcal{C}_R$ is isomorphic to the category $\mathbf{T}\text{-Mod-}\mathbf{R}$, defined in Example 1.8. Indeed,

FIGURE 6. The categories $T\mathcal{C}_R$ and $\mathbf{T}\text{-Mod-}\mathbf{R}$ are equivalent.

an object in $T\mathcal{C}_R$ is a triple (X, x, x^0) , such that (X, x) is a T -module, (X, x^0) is an R^r -module and these structures satisfy relation (2.3). We define a functor


$$F : T\mathcal{C}_R \rightarrow \mathbf{T}\text{-Mod-}\mathbf{R}, \quad F(X, x, x^0) := (X, x, x^0 \circ \chi_{\mathbf{R}, X}^{-1}).$$

Let us first prove that F is well defined, that is $(X, x, x^0 \circ \chi_{\mathbf{R}, X}^{-1})$ is an object in $\mathbf{T}\text{-Mod-}\mathbf{R}$. We have to check that $(X, x^0 \circ \chi_{\mathbf{R}, X}^{-1})$ is a right \mathbf{R} -module and that the structure maps obey (1.7). The graphical proofs of these properties are given in Figure 6, where for simplicity we use the notation $\chi_1 := \chi_{\mathbf{R}, X}$, $\chi_2 := \chi_{\mathbf{R}, \mathbf{R}}$ and $\chi_3 := \chi_{\mathbf{R}, \mathbf{T}}$. In the first frame we prove that X is a right \mathbf{R} -module. The first and the fourth equalities follow by the fact that the braiding χ is a natural transformation. The second identity is trivial, while for the third one we used that (X, x^0) is an \mathbf{R}^r -module. In the second frame we show that the module structures of X satisfy condition (1.7). The first relation in that frame is obvious. The second one follows by (2.3) and the last one is a consequence of naturality of the braiding. Obviously, $F^{-1} : \mathbf{T}\text{-Mod-}\mathbf{R} \rightarrow T\mathcal{C}_R$ is given by $F^{-1}(X, \mu_X^l, \mu_X^r) := (X, \mu_X^l, \mu_X^r \circ \chi_{\mathbf{R}, X})$.

In view of this example, from now on, we shall always regard an object in $T\mathcal{C}_R$ as a $\mathbf{T}\text{-R}$ bimodule.

PROPOSITION 2.6. *Let (R, r, ρ) be a monad and $\tau : RR \rightarrow RR$ be a BD-law. For an (R, R) -bimodule (X, x, x^0) , also the triple $(RX, Rx \circ \tau X, Rx^0 \circ \tau X)$ is an R -bimodule. This construction defines a lifting $\tilde{R} : R\mathcal{C}_R \rightarrow R\mathcal{C}_R$ of R .*

Proof. Let us denote the actions of R and R^r on RX by y and y^0 , respectively. In the first frame of Figure 7 we prove that RX is an R -module. The first equality follows by the definition of y . The second equality is a consequence of the definition of distributive laws, cf. Figure 4, while the third one results by the definition of R -modules. For the last relation we use the definition of y . In

FIGURE 7. RX is an (R, R) -bimodule.

the second frame we prove that RX is an R^r -module too. We proceed as above using, in addition, YB -equation. The fact that y and y^0 commute, i.e. they satisfy relation (2.4), is proved in the third frame, where for the first relation we use the definition of the actions and YB -equation. Since x and x^0 commute, we obtain the second relation. We conclude the proof of the proposition by applying the definition of y and y^0 . \square

REMARK 2.7. If $\mathfrak{r} : RR \rightarrow RR$ is an invertible BD -law, Proposition 2.6 can be applied to the BD -law $\mathfrak{r}^{-1} : R^{\mathfrak{r}}R^{\mathfrak{r}} \rightarrow R^{\mathfrak{r}}R^{\mathfrak{r}}$ and the $R^{\mathfrak{r}}$ -bimodule (X, x^0, x) . Using the isomorphism ${}_R\mathcal{C}_R \cong {}_{R^{\mathfrak{r}}}\mathcal{C}_{R^{\mathfrak{r}}}$ in Remark 2.4, we obtain a functor $\tilde{R}^0 : {}_R\mathcal{C}_R \rightarrow {}_R\mathcal{C}_R$.

EXAMPLE 2.8. For a monad $R = \mathbf{R} \otimes (-)$, induced by an algebra \mathbf{R} as in Example 1.2, the functor $\tilde{R} : \mathbf{R}\text{-Mod-}\mathbf{R} \rightarrow \mathbf{R}\text{-Mod-}\mathbf{R}$, maps an \mathbf{R} -bimodule (X, μ_X^l, μ_X^r) to $\mathbf{R} \otimes X$, with \mathbf{R} -actions

$$\mu_{\mathbf{R} \otimes X}^l := (\mathbf{R} \otimes \mu_X^l) \circ \alpha_{\mathbf{R}, \mathbf{R}, X} \circ (\chi_{\mathbf{R}, \mathbf{R}} \otimes X) \circ \alpha_{\mathbf{R}, \mathbf{R}, X}^{-1}, \quad \mu_{\mathbf{R} \otimes X}^r := (\mathbf{R} \otimes \mu_X^r) \circ \alpha_{\mathbf{R}, X, \mathbf{R}}. \quad (2.5)$$

REMARK 2.9. Note that, for a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and any braid preserving monad morphism $\varphi : R \rightarrow T$ is braid preserving also for the braided pair of monads $\mathcal{S}^0 := (\mathcal{C}, T^{\mathfrak{t}^{-1}}, R^{\mathfrak{r}}, \mathfrak{l}, \mathfrak{t}^{-1}, \mathfrak{r}^{-1})$.

PROPOSITION 2.10. Consider a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$. For an R -module (X, x) and an $R^{\mathfrak{r}}$ -module (Y, y^0) , the following assertions hold.

- (1) The triple (TX, x_i, x_i^0) defines an (R, R) -bimodule, where

$$x_i := Tx \circ \mathfrak{l}X \quad \text{and} \quad x_i^0 := tX \circ T\varphi X \circ \mathfrak{l}X. \quad (2.6)$$

It is called the inside structure on TX . This construction defines a functor ${}_R\mathcal{C} \rightarrow {}_R\mathcal{C}_R$.

- (2) The triple (TY, y_o, y_o^0) defines an (R, R) -bimodule, where

$$y_o := tY \circ \varphi TY \quad \text{and} \quad y_o^0 := Ty^0 \circ \mathfrak{l}Y. \quad (2.7)$$

It is called the outside structure on TY . This construction defines a functor ${}_{R^{\mathfrak{r}}}\mathcal{C} \rightarrow {}_{R^{\mathfrak{r}}}\mathcal{C}_R$.

Proof. Claim (1) is verified by straightforward computation, which is left to the reader. Part (2) is obtained from (1) by replacing the braided pair of monads \mathcal{S} with \mathcal{S}^0 in Remark 2.9, and replacing the R -module (X, x) with the $R^{\mathfrak{r}}$ -module (Y, y^0) . \square

EXAMPLE 2.11. Consider the braid preserving monad morphism coming from an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ in a braided monoidal category, as in Example 2.3. For a left \mathbf{R} -module (X, μ) , the inner actions on $\mathbf{T} \otimes X$ in Proposition 2.10 (1) come out as

$$\begin{aligned} \mu_i^l &:= (\mathbf{T} \otimes \mu) \circ \alpha_{\mathbf{T}, \mathbf{R}, X} \circ (\chi_{\mathbf{R}, \mathbf{T}} \otimes X) \circ \alpha_{\mathbf{R}, \mathbf{T}, X}^{-1} \\ \mu_i^r &:= [\mathfrak{t} \circ (\mathbf{T} \otimes \varphi) \otimes X] \circ \alpha_{\mathbf{T}, \mathbf{R}, X}^{-1} \circ (\mathbf{T} \otimes \chi_{\mathbf{R}, X}^{-1}) \circ \alpha_{\mathbf{T}, X, \mathbf{R}}. \end{aligned}$$

For a right \mathbf{R} -module (Y, ν) , the outer actions on $\mathbf{T} \otimes Y$ in Proposition 2.10 (2) come out as

$$\nu_o^l := [\mathfrak{t} \circ (\varphi \otimes \mathbf{T}) \otimes Y] \circ \alpha_{\mathbf{R}, \mathbf{T}, Y}^{-1}, \quad \nu_o^r := (\mathbf{T} \otimes \nu) \circ \alpha_{\mathbf{T}, Y, \mathbf{R}}.$$

Composition of the forgetful functor ${}_R\mathcal{C}_R \rightarrow {}_R\mathcal{C}$ with the functor in Proposition 2.10 (1) yields a functor $T_i : {}_R\mathcal{C}_R \rightarrow {}_R\mathcal{C}_R$. Symmetrically, composition of the forgetful functor ${}_{R^{\mathfrak{r}}}\mathcal{C}_R \rightarrow {}_{R^{\mathfrak{r}}}\mathcal{C}$ with the functor in Proposition 2.10 (2) yields a functor $T_o : {}_{R^{\mathfrak{r}}}\mathcal{C}_R \rightarrow {}_{R^{\mathfrak{r}}}\mathcal{C}_R$. In the following proposition some natural transformations between various composites of these endofunctors on ${}_R\mathcal{C}_R$, and the functors \tilde{R} and \tilde{R}^0 in Proposition 2.6, are studied.

PROPOSITION 2.12. Let $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ be a braided pair of monads and $\varphi : R \rightarrow T$ be a braid preserving monad morphism. Consider the above endofunctors \tilde{R} , \tilde{R}^0 , T_i and T_o on ${}_R\mathcal{C}_R$. Then the following hold.

- (1) The mappings $\text{Ob}({}_R\mathcal{C}_R) \rightarrow \text{Mor}({}_R\mathcal{C}_R)$, $(X, x, x^0) \mapsto x_i$ and $(X, x, x^0) \mapsto x_i^0$, defined in terms of the inner actions in Proposition 2.10 (1), determine natural transformations $\tilde{R}T_o \rightarrow T_o$.
- (2) The mappings $\text{Ob}({}_R\mathcal{C}_R) \rightarrow \text{Mor}({}_R\mathcal{C}_R)$, $(X, x, x^0) \mapsto x_o$ and $(X, x, x^0) \mapsto x_o^0$, defined in terms of the outer actions in Proposition 2.10 (2), determine natural transformations $\tilde{R}^0T_i \rightarrow T_i$.
- (3) The BD -law \mathfrak{t} defines a natural transformation $T_iT_o \rightarrow T_oT_i$.
- (4) The BD -law \mathfrak{l} defines a natural transformations $\tilde{R}T_o \rightarrow T_o\tilde{R}$ and $\tilde{R}^0T_o \rightarrow T_o\tilde{R}^0$.

Proof. Note that part (2) is obtained from part (1) by replacing the braided pair of monads \mathcal{S} with \mathcal{S}^0 in Remark 2.9, and using the isomorphism ${}_R\mathcal{C}_R \cong {}_{R^r}\mathcal{C}_{R^r}$.

Claims (1), (3) and (4) are proven by straightforward but somewhat lengthy computations. We illustrate the main steps on the example of part (3). We use the graphical representation of morphisms in a category again. We denote the morphisms defining the (R, R) -bimodule structure of $T_i T_o X$ by x_{io} and x_{io}^0 , respectively. Similarly, for the actions of R and R^r on $T_o T_i X$ we use the notation x_{oi} and x_{oi}^0 . The diagrammatic representation of these morphisms is given in the first frame of Figure 8. In the second frame we prove that $\mathfrak{t}X$ is a morphism of R -modules. Note that for the first equality we used that \mathfrak{l} is a distributive law and φ satisfies (A 4). For the graphical proof of the fact $\mathfrak{t}X$ is a morphism of R^r -modules see the third frame of Figure 8. Note that, besides the properties of \mathfrak{l} and φ that we already used, we also need condition (A 2) in Definition 2.1. \square




FIGURE 8. $\mathfrak{t}X$ is a morphism of (R, R) -bimodules.

PROPOSITION 2.13. *For a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$, consider an (R, R) -bimodule (X, x, x^0) and a (T, R) -bimodule (Y, y, y^0) . Then the following assertions hold.*

- (1) *The actions x^0 and y^0 induce R^r -module structures respectively on $\varphi_*(X, x)$ and $\varphi^*(Y, y)$.*
- (2) *With respect to the above actions, φ_* in Proposition 1.7 and φ^* in Proposition 1.6 can be regarded as functors $\varphi_* : {}_R\mathcal{C}_R \rightarrow {}_T\mathcal{C}_R$ and $\varphi^* : {}_T\mathcal{C}_R \rightarrow {}_R\mathcal{C}_R$.*
- (3) *The functors constructed in part (2) define a pair of adjoint functors (φ_*, φ^*) .*

Proof. (1) and (2). Obviously, y^0 induces an R^r -action on $\varphi^*(Y, y) = Y$. By construction, $\varphi^*(Y, y) = (Y, y \circ \varphi Y)$. We claim that $(Y, y \circ \varphi Y, y^0)$ is an (R, R) -bimodule. We have to check that the module structures commute. Since φ is a natural morphism, condition (A 3) holds true and y and y^0 commute, we get

$$y \circ \varphi Y \circ R y^0 \circ \mathfrak{r} Y = y \circ T y^0 \circ \varphi R^0 Y \circ \mathfrak{r} Y = y \circ T y^0 \circ \mathfrak{l} Y \circ R^0 \varphi Y = y^0 \circ R^0 y \circ R^0 \varphi Y.$$

Hence $y \circ \varphi Y$ and y^0 commute too. Consequently, φ^* can be seen as a functor from ${}_T\mathcal{C}_R$ to ${}_R\mathcal{C}_R$.

Recall from the proof of Proposition 1.7 that the T -module $(\varphi_* X, \bar{x})$ is the coequalizer of $(T x, t X \circ T \varphi X)$. In terms of the inner actions in (2.6), the T -module morphisms $T x$ and $t X \circ T \varphi X$ are equal to $x_i \circ \mathfrak{l}^{-1} X$ and $x_i^0 \circ \mathfrak{l}^{-1} X$, respectively. Hence they are R^r -module morphisms $T_o \tilde{R} X \rightarrow T_o X$ by Proposition 2.12 (1) and (4). Thus $T x$ and $t X \circ T \varphi X$ are morphisms of (T, R) -bimodules from $(T R X, t R X, T R x^0 \circ T \mathfrak{r} X \circ \mathfrak{l} R X)$ to $(T X, t X, x_o^0)$. Hence, by Proposition 1.5, there is an action $\bar{x}^0 : R^r \varphi_* X \rightarrow \varphi_* X$ such that $(\varphi_* X, \bar{x}, \bar{x}^0)$ is a (T, R) -bimodule and πX is a morphism of (T, R) -bimodules from $(T X, t X, x_o^0)$ to $(\varphi_* X, \bar{x}, \bar{x}^0)$.

Let $f : (X, x, x^0) \rightarrow (Z, z, z^0)$ be a morphism of (R, R) -bimodules. We already know that $\varphi_* f$ is a morphism of T -modules. By Proposition 2.10 (2), $T_o f$ is an R^r -module morphism. Hence so is $\varphi_* f$ by Proposition 1.5. Thus φ_* can be regarded as a functor from ${}_R\mathcal{C}_R$ to ${}_T\mathcal{C}_R$.

(3). Let (X, x, x^0) be an (R, R) -bimodule and let (Y, y, y^0) be a (T, R) -bimodule. It is sufficient to show that $\Phi_{X, Y}(f)$ and $\Theta_{X, Y}(g)$ are morphisms of R^r -modules, for any $f : \varphi_* X \rightarrow Y$ in ${}_T\mathcal{C}_R$

and any $g : X \rightarrow \varphi^*Y$ in ${}_R\mathcal{C}_R$ (for the definition of $\Phi_{X,Y}$ and $\Theta_{X,Y}$ see the proof of Proposition 1.7). The fact that $\Phi_{X,Y}(f)$ is a morphism of R^τ -modules is proved in the following computation:

$$y^0 \circ Rf \circ R\pi X \circ R^0\tau X = f \circ \bar{x}^0 \circ R\pi X \circ R\tau X = f \circ \pi X \circ Tx^0 \circ \mathfrak{L}X \circ R\tau X = f \circ \pi X \circ \tau X \circ x^0.$$

Since $R\pi X$ is an epimorphism, the relation meaning that $\Theta_{X,Y}(g)$ is a morphism of R^τ -modules follows from the computation below:

$$\begin{aligned} y^0 \circ R^\tau\Theta_{X,Y}(g) \circ R^\tau\pi X &= y^0 \circ R^\tau y \circ R^\tau Tg \stackrel{(A)}{=} y \circ Ty^0 \circ \mathfrak{L}Y \circ R^\tau Tg \stackrel{(B)}{=} y \circ Tg \circ Tx^0 \circ \mathfrak{L}X \\ &= \Theta_{X,Y}(g) \circ \pi X \circ Tx^0 \circ \mathfrak{L}X \stackrel{(C)}{=} \Theta_{X,Y}(g) \circ \bar{x}^0 \circ R^\tau\pi X. \end{aligned}$$

To get the equalities (A), (B) and (C) above we used the commutation relation between the module structures of Y and, respectively, that $T_o g$ and π are morphisms of R^τ -modules. \square

COROLLARY 2.14. *For a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \tau)$ and a braid preserving monad morphism $\varphi : R \rightarrow T$, consider the adjoint functors (φ_*, φ^*) in Proposition 2.13.*

- (1) *The unit of the adjunction $\sigma : \text{Id}_{{}_R\mathcal{C}_R} \rightarrow \varphi^*\varphi_*$ is given, for any (R, R) -bimodule (X, x, x^0) , by*

$$\sigma X := \pi X \circ \tau X. \quad (2.8)$$

- (2) *The counit of the adjunction $\xi : \varphi_*\varphi^* \rightarrow \text{Id}_{T\mathcal{C}_R}$ satisfies the following relation for any (T, R) -bimodule (Y, y, y^0) .*

$$\xi Y \circ \pi\varphi^*(Y, y, y^0) = y. \quad (2.9)$$

Proof. By definition, $\sigma X := \Phi_{X, \varphi_*(X, x, x^0)}(\text{Id}_{\varphi_*(X, x, x^0)})$ and $\xi Y = \Theta_{\varphi^*(Y, y, y^0), Y}(\text{Id}_{\varphi^*(Y, y, y^0)})$. \square

EXAMPLE 2.15. For the braid preserving monad morphism, coming from an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ in a braided monoidal category as in Example 2.3, \mathbf{T} is a \mathbf{T} - \mathbf{R} bimodule via the left regular \mathbf{T} -action and the right \mathbf{R} -action induced by φ . If (as in Example 1.8) the functors $X \otimes (-)$ and $(-) \otimes X$ are right exact, for any object X in \mathcal{C} , then the pair of adjoint functors in Proposition 2.13 (3) consists of the ‘restriction of scalars’ functor $\varphi^* : \mathbf{T}\text{-Mod-}\mathbf{R} \rightarrow \mathbf{R}\text{-Mod-}\mathbf{R}$ and the induction functor $\varphi_* = \mathbf{T} \otimes_{\mathbf{R}} (-) : \mathbf{R}\text{-Mod-}\mathbf{R} \rightarrow \mathbf{T}\text{-Mod-}\mathbf{R}$.

3. THE PARA-COCYCLIC OBJECT ASSOCIATED TO A BRAID PRESERVING HOMOMORPHISM

In this section we show, using the main result in [BS], that to every braid preserving monad morphism there corresponds an admissible septuple, and hence a certain para-cocyclic object.

Recall that, for every pair (F, G) of adjoint functors, with $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$, the triple $(GF, G\xi F, \sigma)$ is a monad on \mathcal{C} , where ξ and σ are respectively the counit and the unit of the adjunction. For details the reader is referred to [We, Main Application 8.6.2, p. 280]. In particular, the pair of adjoint functors (φ_*, φ^*) , constructed in Proposition 2.13 (3), determines a monad structure on $\varphi^*\varphi_* : {}_R\mathcal{C}_R \rightarrow {}_R\mathcal{C}_R$ that will be denoted by $(\bar{T}_o, \bar{t}_o, \bar{\tau}_o)$.

Since \mathfrak{l} is invertible, the object in \mathcal{C} , underlying the (R, R) -bimodule $\bar{T}_o(X, x, x^0)$, is the coequalizer of the morphisms x_i and x_i^0 in (2.6). Since x_i and x_i^0 are (R, R) -bimodule morphisms by Proposition 2.12 (1), it follows by Proposition 1.5 that

$$\bar{T}_o(X, x, x^0) = \text{Coeq}(x_i, x_i^0)$$

in ${}_R\mathcal{C}_R$. That is, φ^* takes the canonical epimorphism $\pi X : (TX, tX, x_o^0) \rightarrow \varphi_*(X, x, x^0)$ to a canonical (R, R) -bimodule epimorphism $\pi_o X : T_o(X, x, x^0) \rightarrow \bar{T}_o(X, x, x^0)$. In other words, the respective actions \bar{x}_o and \bar{x}_o^0 of R and R^τ on $\bar{T}_o X$ are uniquely determined such that $\pi_o X : T_o X \rightarrow \bar{T}_o X$ is a morphism of (R, R) -bimodules, in the sense that

$$\bar{x}_o \circ R\pi_o X = \pi_o X \circ x_o \quad \text{and} \quad \bar{x}_o^0 \circ R^\tau\pi_o X = \pi_o X \circ x_o^0. \quad (3.1)$$

The unit of \bar{T}_o is

$$\bar{\tau}_o = \pi_o \circ \tau. \quad (3.2)$$

In order to compute the multiplication \bar{t}_o of \bar{T}_o , substitute $Y = \varphi_*(X, x, x^0)$ in (2.9), apply φ^* on it, and compose the resulting relation on the right by $T\pi_o X$. Since $T\pi_o X$ is a coequalizer and the

module structure \bar{x} satisfies (1.6), we deduce that $\bar{t}_o X$ is the unique morphism in ${}_R \mathcal{C}_R$ satisfying relation

$$\bar{t}_o X \circ \pi_o \bar{T}_o X \circ T \pi_o X = \pi_o X \circ t X. \quad (3.3)$$

Summarizing, we prove the following proposition.

PROPOSITION 3.1. *Associated to a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$, there is a monad $(\bar{T}_o, \bar{t}_o, \bar{\tau}_o)$ on ${}_R \mathcal{C}_R$ such that, for every (R, R) -bimodule (X, x, x^0) ,*

$$(\bar{T}_o X, \pi_o X) = \text{Coeq}(x_i, x_i^0),$$

(cf. Proposition 2.10 (1)). *That is, the R , and R^r -actions \bar{x}_o and \bar{x}_o^0 on $\bar{T}_o X$ are uniquely defined such that $\pi_o X : T_o X \rightarrow \bar{T}_o X$ is a morphism of (R, R) -bimodules. The multiplication $\bar{t}_o : \bar{T}_o \bar{T}_o \rightarrow \bar{T}_o$ satisfies (3.3) and the unit $\bar{\tau}_o : {}_R \mathcal{C}_R \rightarrow \bar{T}_o$ satisfies (3.2).*

A symmetrical version of Proposition 3.1 is obtained by replacing the braided pair of monads \mathcal{S} with \mathcal{S}^0 , introduced in Remark 2.9.

PROPOSITION 3.2. *Associated to a braided pair of monads $\mathcal{S} := (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$, there is a monad $(\bar{T}_i, \bar{t}_i, \bar{\tau}_i)$ on ${}_R \mathcal{C}_R$ such that, for every (R, R) -bimodule (X, x, x^0) ,*

$$(\bar{T}_i X, \pi_i X) = \text{Coeq}(x_o, x_o^0), \quad (3.4)$$

(cf. Proposition 2.10 (2)). *That is, the actions \bar{x}_i and \bar{x}_i^0 on $\bar{T}_i X$ are uniquely defined such that $\pi_i X : T_i X \rightarrow \bar{T}_i X$ is a morphism of (R, R) -bimodules, i.e.*

$$\bar{x}_i \circ R \pi_i X = \pi_i X \circ x_i \quad \text{and} \quad \bar{x}_i^0 \circ R^r \pi_i X = \pi_i X \circ x_i^0. \quad (3.5)$$

The unit $\bar{\tau}_i$ and the multiplication \bar{t}_i satisfy

$$\bar{\tau}_i = \pi_i \circ \tau \quad \text{and} \quad \bar{t}_i \circ \pi_i \bar{T}_i \circ T \pi_i = \pi_i \circ t \circ \mathfrak{t}^{-1}. \quad (3.6)$$

EXAMPLE 3.3. Let \mathcal{C} be a braided monoidal category. For the braid preserving monad morphism coming from an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ as in Example 2.3, \mathbf{T} has a natural \mathbf{R} -bimodule structure via φ . If (as in Example 1.8) the functors $X \otimes (-)$ and $(-) \otimes X$ are right exact, for any object X in \mathcal{C} , then the functor underlying the monad in Proposition 3.1 is induced by the \mathbf{R} -bimodule \mathbf{T} , i.e. $\bar{T}_o = \mathbf{T} \otimes_{\mathbf{R}} (-) : \mathbf{R}\text{-Mod-}\mathbf{R} \rightarrow \mathbf{R}\text{-Mod-}\mathbf{R}$. Multiplication is given by $(\bar{t}_o \otimes_{\mathbf{R}} (-)) \circ \bar{\alpha}_{\mathbf{T}, \mathbf{T}, (-)}^{-1}$, where $\bar{\alpha}$ is the associator isomorphism in $\mathbf{R}\text{-Mod-}\mathbf{R}$ and the \mathbf{R} -bimodule morphism $\bar{t}_o : \mathbf{T} \otimes_{\mathbf{R}} \mathbf{T} \rightarrow \mathbf{T}$ is defined as the projection of $\mathfrak{t} : \mathbf{T} \otimes \mathbf{T} \rightarrow \mathbf{T}$. The unit is $\varphi \otimes_{\mathbf{R}} (-)$ (where we used that $\mathbf{R} \otimes_{\mathbf{R}} (-)$ is naturally isomorphic to the identity functor on $\mathbf{R}\text{-Mod-}\mathbf{R}$).

Symmetrically, on an \mathbf{R} -bimodule (X, μ_X^l, μ_X^r) , the functor \bar{T}_i is defined as the coequalizer of $x_o = (\mathfrak{t} \otimes X) \circ [(\varphi \otimes \mathbf{T}) \otimes X] \circ \alpha_{\mathbf{R}, \mathbf{T}, X}^{-1}$ and $x_o^0 = (T \otimes \mu_X^r) \circ \alpha_{\mathbf{T}, X, \mathbf{R}} \circ \chi_{\mathbf{R}, \mathbf{T} \otimes X}$. Equivalently, composing both x_o and x_o^0 by the isomorphism $\alpha_{\mathbf{R}, \mathbf{T}, X} \circ (\chi_{\mathbf{R}, \mathbf{T}}^{-1} \otimes X)$, as a coequalizer

$$(\mathbf{T} \otimes \mathbf{R}) \otimes X \begin{array}{c} \xrightarrow{[\mathbf{T} \otimes (\mu_X^r \circ \chi_{\mathbf{R}, X})] \circ \alpha_{\mathbf{T}, \mathbf{R}, X}} \\ \xrightarrow{[\mathfrak{t} \circ \chi_{\mathbf{T}, \mathbf{T}} \circ (\mathbf{T} \otimes \varphi)] \otimes X} \end{array} \mathbf{T} \otimes X \xrightarrow{\pi_i} \bar{T}_i X.$$

In order to obtain the form of the morphism corresponding to the lower one of the parallel arrows, we used the first identity in (2.1). Note that $\mu_X^r \circ \chi_{\mathbf{R}, X}$ is a left action, and $\mathfrak{t} \circ \chi_{\mathbf{T}, \mathbf{T}} \circ (\mathbf{T} \otimes \varphi)$ is a right action for the algebra $\mathbf{R}^X := (\mathbf{R}, \mathfrak{r} \circ \chi_{\mathbf{R}, \mathbf{R}}, \rho)$. Therefore, $\bar{T}_i X = \mathbf{T} \otimes_{\mathbf{R}^X} X$. Since φ can be regarded as an algebra homomorphism $\mathbf{R}^X \rightarrow \mathbf{T}^{X^{-1}}$, the unit of the monad $\bar{T}_i(-) = \mathbf{T} \otimes_{\mathbf{R}^X} (-)$ is given by $\varphi \otimes_{\mathbf{R}^X} (-)$. Multiplication is induced by the projection $\bar{t}_i : \mathbf{T} \otimes_{\mathbf{R}^X} \mathbf{T} \rightarrow \mathbf{T}$ of the \mathbf{R}^X -bimodule morphism $\mathfrak{t} \circ \chi_{\mathbf{T}, \mathbf{T}}^{-1} : \mathbf{T} \otimes \mathbf{T} \rightarrow \mathbf{T}$. Note that, if χ is a symmetry, then $\mathbf{R}^X = \mathbf{R}^{op}$, hence $\bar{T}_i = \mathbf{T}^{op} \otimes_{\mathbf{R}^{op}} (-) \cong (-) \otimes_{\mathbf{R}} \mathbf{T}$.

Next we introduce a functor $\Pi : {}_R \mathcal{C}_R \rightarrow \mathcal{C}$, that will be used to construct a para-cocyclic module associated to a braid preserving monad morphism.

DEFINITION 3.4. For a BD -law $\tau : RR \rightarrow RR$, the functor $\Pi : {}_R\mathcal{C}_R \rightarrow \mathcal{C}$ is defined, for an (R, R) -bimodule (X, x, x^0) , by

$$(\Pi X, pX) := \text{Coeq}(x, x^0).$$

For every morphism $f : X \rightarrow Y$ of (R, R) -bimodules, Πf is the unique morphism in \mathcal{C} such that $pY \circ f = \Pi f \circ pX$. Hence p can be interpreted as a natural epimorphism from the forgetful functor $U : {}_R\mathcal{C}_R \rightarrow \mathcal{C}$ to Π .

In the graphical notation we do not denote the forgetful functor $U : {}_R\mathcal{C}_R \rightarrow \mathcal{C}$. That is, a box representing the natural transformation p in Definition 3.4 has only a lower leg, corresponding to the functor Π .

REMARK 3.5. In the context of Definition 3.4, the functor $\Pi T_i : {}_R\mathcal{C}_R \rightarrow \mathcal{C}$ is equal to the composite of $\overline{T}_o : {}_R\mathcal{C}_R \rightarrow {}_R\mathcal{C}_R$ and the forgetful functor $U : {}_R\mathcal{C}_R \rightarrow \mathcal{C}$. Symmetrically, $\Pi T_o = U\overline{T}_i$.

EXAMPLE 3.6. For the braid preserving monad morphism coming from an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ as in Example 2.3, the functor Π maps an \mathbf{R} -bimodule (X, μ_X^l, μ_X^r) to

$$\Pi X = \text{Coker}(\mu_X^l - \mu_X^r \circ \chi_{\mathbf{R}, X}).$$

For two \mathbf{R} -bimodules X and Y we shall use the notation $X \widehat{\otimes}_{\mathbf{R}} Y := \Pi(X \otimes_{\mathbf{R}} Y)$ and we shall say that this object in \mathcal{C} is the *braided cyclic tensor product* of X and Y .

In the next theorem we prove that the BD -law \mathfrak{t} , in a braided pair of monads $(\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$, lifts to a distributive law of the monads \overline{T}_o and \overline{T}_i in Propositions 3.1 and 3.2.

THEOREM 3.7. *Let $\mathcal{S} = (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ be a braided pair of monads and $\varphi : R \rightarrow T$ be a braid preserving monad morphism. Consider the associated functors T_i, T_o and $\overline{T}_i, \overline{T}_o$, and the natural epimorphisms $\pi_i : T_i \rightarrow \overline{T}_i$ and $\pi_o : T_o \rightarrow \overline{T}_o$ in Propositions 3.1 and 3.2.*

- (1) *There is a natural transformation $\mathfrak{t}' : T_i \overline{T}_o \rightarrow \overline{T}_o T_i$, between endofunctors on the category of (R, R) -bimodules, such that*

$$\mathfrak{t}' \circ T_i \pi_o = \pi_o T_i \circ \mathfrak{t}. \quad (3.7)$$

- (2) *There is a natural transformation $\bar{\mathfrak{t}} : \overline{T}_i \overline{T}_o \rightarrow \overline{T}_o \overline{T}_i$, between endofunctors on the category of (R, R) -bimodules, such that*

$$\pi_o \overline{T}_i \circ T_o \pi_i \circ \mathfrak{t} = \bar{\mathfrak{t}} \circ \pi_i \overline{T}_o \circ T_i \pi_o. \quad (3.8)$$

- (3) $\bar{\mathfrak{t}} : \overline{T}_i \overline{T}_o \rightarrow \overline{T}_o \overline{T}_i$ *is a distributive law.*

Proof. (1). Let (X, x, x^0) be a given (R, R) -bimodule. Since T preserves coequalizers, the sequence in the top row of the diagram in Figure 9 is a coequalizer. Since \mathfrak{l} is invertible, the sequence in

$$\begin{array}{ccccc} T_i T_o \widetilde{R}X & \xrightarrow[\quad g_1 := T_i T_o x \quad]{f_1 := T_i \mathfrak{t} X \circ T_i T_o \varphi X} & T_i T_o X & \xrightarrow{T_i \pi_o X} & T_i \overline{T}_o X \\ \mathfrak{t} \widetilde{R}X \downarrow & & \mathfrak{t} X \downarrow & & \downarrow \mathfrak{t}' X \\ T_o T_i \widetilde{R}X & \xrightarrow[\quad g_2 := T_o T_i x \quad]{f_2 := \mathfrak{t} T_i X \circ T_o \varphi T_i X \circ T_o \Gamma^{-1} X} & T_o T_i X & \xrightarrow{\pi_o T_i X} & \overline{T}_o T_i X \end{array}$$

FIGURE 9. Construction of \mathfrak{t}' .

the bottom row is a coequalizer by the definition of π_o . By Proposition 2.12 (3) the square whose horizontal edges are g_1 and g_2 is commutative. Using that \mathfrak{t} is a BD -law and taking into account condition (A 4), it results that the square with horizontal arrows f_1 and f_2 is also commutative. Hence $\pi_o T_i X \circ \mathfrak{t} X$ coequalizes (f_1, g_1) . Furthermore, by Proposition 2.12 (3), $\mathfrak{t} X : T_i T_o X \rightarrow T_o T_i X$ is a morphism of (R, R) -bimodules. By construction of \overline{T}_o , the canonical epimorphism $\pi_o X : T_o X \rightarrow \overline{T}_o X$ is a morphism of (R, R) -bimodules. Hence $\pi_o T_i X : T_o T_i X \rightarrow \overline{T}_o T_i X$ and

$T_i\pi_oX : T_iT_oX \rightarrow T_i\bar{T}_oX$ are also morphisms in ${}_R\mathcal{C}_R$. Moreover, $tX \circ T\varphi X$ and Tx are (R, R) -bimodule morphisms by Proposition 2.12 (1) and (4). Hence so are f_1 and g_1 by Proposition 2.10 (1). In conclusion, we can apply Proposition 1.5 in the category ${}_R\mathcal{C}_R$ to show that there exists an (R, R) -bimodule morphism $\mathfrak{t}'X$, rendering commutative the right hand side square on Figure 9. By construction, \mathfrak{t}' is natural.

(2). It follows by condition (A 4), naturality of φ and the fact that \mathfrak{t} is a distributive law that the morphisms x_o and x_o^0 in (2.7) satisfy

$$T_o x_o \circ \mathfrak{t}_i X \circ RtX = \mathfrak{t}X \circ tT_oX \circ \varphi T_i T_oX \quad \text{and} \quad T_i x_o^0 \circ \mathfrak{t}_i X \circ RtX = \mathfrak{t}X \circ T_i x_o^0 \circ \mathfrak{t}_o X. \quad (3.9)$$

Taking into account (3.7), the identities in (3.9) imply

$$\begin{aligned} \mathfrak{t}'X \circ T_i\pi_oX \circ tT_oX \circ \varphi T_i T_oX &= \pi_o T_i X \circ T_o x_o \circ \mathfrak{t}_i X \circ RtX \quad \text{and} \\ \mathfrak{t}'X \circ T_i\pi_oX \circ T_i x_o^0 \circ \mathfrak{t}_o X &= \pi_o T_i X \circ T_o x_o^0 \circ \mathfrak{t}_i X \circ RtX. \end{aligned} \quad (3.10)$$

Using the naturality of t , φ and π_o together with the (R, R) -bimodule morphism property (3.1) of π_o , the equations in (3.10) can be seen to be equivalent to the commutativity of the left hand side squares on Figure 10. Since $\pi_i X$ coequalizes (x_o, x_o^0) , the morphism $\bar{T}_o\pi_i X$ coequalizes $\bar{T}_o x_o \circ$


FIGURE 10. Construction of $\bar{\mathfrak{t}}$.

$\pi_o \tilde{R}^0 T_i X \circ \mathfrak{t}_i X \circ \tilde{R}^0 \mathfrak{t}X$ and $\bar{T}_o x_o^0 \circ \pi_o \tilde{R}^0 T_i X \circ \mathfrak{t}_i X \circ \tilde{R}^0 \mathfrak{t}X$. Hence $\bar{T}_o \pi_i X \circ \mathfrak{t}'X$ coequalizes $t\bar{T}_o X \circ \varphi T_i \bar{T}_o X \circ \tilde{R}^0 T_i \pi_o X$ and $T_i x_o^0 \circ \bar{\mathfrak{t}}_o X \circ \tilde{R}^0 T_i \pi_o X$. Since $\tilde{R}^0 T_i \pi_o X$ is an epimorphism, this implies the existence of a morphism $\bar{\mathfrak{t}}X$ in \mathcal{C} , rendering commutative the right hand side square on Figure 10. Furthermore, the parallel arrows $t\bar{T}_o X \circ \varphi T_i \bar{T}_o X$ and $T_i x_o^0 \circ \bar{\mathfrak{t}}_o X$ on Figure 10 are morphisms of (R, R) -bimodules by Proposition 2.12 (2) and $\mathfrak{t}'X$ is a morphism of (R, R) -bimodules by part (1). By applying Proposition 1.5 in ${}_R\mathcal{C}_R$, we conclude that $\bar{\mathfrak{t}}X$ is an (R, R) -bimodule morphism. Naturality of $\bar{\mathfrak{t}}$ follows obviously from the naturality of \mathfrak{t}' .

(3). The first axiom in Definition 1.9 is proven in the left frame in Figure 11. The first and fourth equalities follow by (3.8) and the fact that π_i and π_o are natural transformations. The second and fifth equalities are consequences of the second identity in (3.6). For the third relation we used YB -equation on \mathfrak{t} and that \mathfrak{t} is a distributive law. Since $\pi_i \bar{T}_i \bar{T}_o \circ T\pi_i \bar{T}_o \circ TT\pi_o$ is epi, this proves the first axiom in the definition of distributivity laws.

Similarly, taking into account the first identity in (3.6) we get the first and last equalities in the right frame of Figure 11. By (3.8) and the fact that \mathfrak{t} is a distributive law we deduce the second and the third identities. Note that in the first, second and last equalities we also used naturality which, in this case, means that e.g. π_o can be pushed up and down along a string. Since π_o is an epimorphism, this proves that $\bar{\mathfrak{t}}$ satisfies the third axiom of a distributive law in Definition 1.9. The remaining two axioms are verified similarly. \square


THEOREM 3.8. *Take a braided pair of monads $\mathcal{S} = (\mathcal{C}, T, R, \mathfrak{l}, \mathfrak{t}, \mathfrak{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$. Consider the functors \bar{T}_o and \bar{T}_i , constructed in Propositions 3.1 and 3.2, respectively, and the functor Π introduced in Definition 3.4. For these data there are mutually*


 FIGURE 11. $\bar{\tau}$ is a distributivity law.

inverse natural isomorphisms $i : \Pi \bar{T}_o \rightarrow \Pi \bar{T}_i$ and $j : \Pi \bar{T}_i \rightarrow \Pi \bar{T}_o$, between functors ${}_R \mathcal{C}_R \rightarrow \mathcal{C}$, such that

$$i \circ p\bar{T}_o \circ \pi_o = p\bar{T}_i \circ \pi_i, \quad j \circ p\bar{T}_i \circ \pi_i = p\bar{T}_o \circ \pi_o. \quad (3.11)$$

Proof. In the first diagram of Figure 12 the squares are commutative, for any (R, R) -bimodule (X, x, x^0) , as $\pi_i X : T_i X \rightarrow \bar{T}_i X$ is a morphism of (R, R) -bimodules. Hence $p\bar{T}_i X \circ \pi_i X$ coequalizes (x_i, x_i^0) . Then there is a morphism $i' X : \bar{T}_o X \rightarrow \Pi \bar{T}_i X$ such that $i' X \circ \pi_o X = p\bar{T}_i X \circ \pi_i X$. In


 FIGURE 12. The construction of i .

the second diagram of Figure 12, the squares are also commutative, as $\pi_o X$ is a morphism of (R, R) -bimodules. Therefore $i' X$ coequalizes (\bar{x}_o, \bar{x}_o^0) , as $\pi_i X$ coequalizes (x_o, x_o^0) and $R\pi_o X$ is an epimorphism. Thus there exists a morphism $i X : \Pi \bar{T}_o X \rightarrow \Pi \bar{T}_i X$ satisfying the required relation. To construct j one proceeds analogously. The natural morphisms i and j are mutual inverses, as $p\bar{T}_i X \circ \pi_i X$ and $p\bar{T}_o X \circ \pi_o X$ are epimorphisms. Naturality of i follows by (3.11) and naturality of p , π_o and π_i . \square

EXAMPLE 3.9. Let \mathcal{C} be an abelian braided monoidal category such that the functors $X \otimes (-)$ and $(-) \otimes X$ are right exact, for any object X in \mathcal{C} (cf. Example 1.8). For the monad morphism coming from an algebra homomorphism $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ as in Example 2.3, the natural transformation $\bar{\tau}$ in Theorem 3.7 (2) is the projection of $\alpha_{\mathbf{T}, \mathbf{T}, X} \circ (\chi_{\mathbf{T}, \mathbf{T}} \otimes X) \circ \alpha_{\mathbf{T}, \mathbf{T}, X}^{-1}$. That is, it is the unique morphism satisfying

$$\pi_o(\mathbf{T} \otimes_{\mathbf{R}X} X) \circ (\mathbf{T} \otimes \pi_i X) \circ \alpha_{\mathbf{T}, \mathbf{T}, X} \circ (\chi_{\mathbf{T}, \mathbf{T}} \otimes X) \circ \alpha_{\mathbf{T}, \mathbf{T}, X}^{-1} = \bar{\tau} X \circ \pi_i(\mathbf{T} \otimes_{\mathbf{R}X} X) \circ (T \otimes \pi_o X),$$

where notations in Example 3.3 are used.

For an \mathbf{R} -bimodule X , the natural transformation $i X : \mathbf{T} \widehat{\otimes}_{\mathbf{R}X} X \rightarrow \Pi(\mathbf{T} \otimes_{\mathbf{R}X} X)$, constructed in the previous theorem, is the projection of the identity map $\mathbf{T} \otimes X \rightarrow \mathbf{T} \otimes X$.

As it is explained in [We, page 281], for any monad (\mathbf{T}, μ, η) on a category \mathcal{M} and an object X in \mathcal{M} , there is an associated cosimplex in \mathcal{M} , given at degree n by $\mathbf{T}^{n+1} X$. Coface and codegeneracy maps are given, for $k = 0, \dots, n$, by

$$\mathbf{T}^k \eta \mathbf{T}^{n-k} X : \mathbf{T}^n X \rightarrow \mathbf{T}^{n+1} X \quad \text{and} \quad \mathbf{T}^k \mu \mathbf{T}^{n-k} X : \mathbf{T}^{n+2} X \rightarrow \mathbf{T}^{n+1} X,$$

respectively. Clearly, application of any functor $\mathbf{\Pi} : \mathcal{M} \rightarrow \mathcal{C}$ yields a cosimplex in \mathcal{C} . In [BS], paracocyclic structures on the resulting cosimplex in \mathcal{C} were studied. Recall from [BS] the following construction.

DEFINITION 3.10. [BS, Definitions 1.7 and 1.8] An *admissible septuple* \mathcal{A} consists of the data $(\mathcal{M}, \mathcal{C}, \mathbf{T}_o, \mathbf{T}_i, \mathbf{\Pi}, \mathbf{t}, \mathbf{i})$, where

- \mathcal{C} and \mathcal{M} are categories,
- $(\mathbf{T}_o, \boldsymbol{\mu}_o, \boldsymbol{\eta}_o)$ and $(\mathbf{T}_i, \boldsymbol{\mu}_i, \boldsymbol{\eta}_i)$ are monads on \mathcal{M} ,
- $\mathbf{\Pi}$ is a functor $\mathcal{M} \rightarrow \mathcal{C}$,
- $\mathbf{t} : \mathbf{T}_i \mathbf{T}_o \rightarrow \mathbf{T}_o \mathbf{T}_i$ is a distributive law,
- $\mathbf{i} : \mathbf{\Pi} \mathbf{T}_o \rightarrow \mathbf{\Pi} \mathbf{T}_i$ is a natural transformation,

subject to the conditions

$$\mathbf{i} \circ \mathbf{\Pi} \boldsymbol{\eta}_o = \mathbf{\Pi} \boldsymbol{\eta}_i \quad \text{and} \quad \mathbf{i} \circ \mathbf{\Pi} \boldsymbol{\mu}_o = \mathbf{\Pi} \boldsymbol{\mu}_i \circ \mathbf{i} \mathbf{T}_i \circ \mathbf{\Pi} \mathbf{t} \circ \mathbf{i} \mathbf{T}_o. \quad (3.12)$$

A *transposition morphism* for the admissible septuple \mathcal{A} is a pair (X, w) , consisting of an object X and a morphism $w : \mathbf{T}_i X \rightarrow \mathbf{T}_o X$ in \mathcal{M} , satisfying

$$w \circ \boldsymbol{\eta}_i X = \boldsymbol{\eta}_o X \quad \text{and} \quad w \circ \boldsymbol{\mu}_i X = \boldsymbol{\mu}_o X \circ \mathbf{T}_o w \circ \mathbf{t} X \circ \mathbf{T}_i w. \quad (3.13)$$

THEOREM 3.11. [BS, Theorem 1.10] Consider an admissible septuple $(\mathcal{M}, \mathcal{C}, \mathbf{T}_o, \mathbf{T}_i, \mathbf{\Pi}, \mathbf{t}, \mathbf{i})$ and a transposition morphism (X, w) for it. The associated cosimplex $Z^* := \mathbf{\Pi} \mathbf{T}_o^{*+1} X$ is paracocyclic with paracocyclic morphism

$$w_n := \mathbf{\Pi} \mathbf{T}_o^n w \circ \mathbf{\Pi} \mathbf{T}_o^{n-1} \mathbf{t} X \circ \mathbf{\Pi} \mathbf{T}_o^{n-2} \mathbf{t} \mathbf{T}_o X \circ \dots \circ \mathbf{\Pi} \mathbf{T}_o \mathbf{t} \mathbf{T}_o^{n-2} X \circ \mathbf{\Pi} \mathbf{t} \mathbf{T}_o^{n-1} X \circ \mathbf{i} \mathbf{T}_o^n X. \quad (3.14)$$

The next theorem is our main result.

THEOREM 3.12. Consider a braided pair of monads $\mathcal{S} = (\mathcal{C}, T, R, \mathbf{l}, \mathbf{t}, \mathbf{r})$ and a braid preserving monad morphism $\varphi : R \rightarrow T$. Then the data, consisting of

- the categories \mathcal{C} and $\mathcal{M} := {}_R \mathcal{C}_R$,
- the monads \overline{T}_o in Proposition 3.1 and \overline{T}_i in Proposition 3.2,
- the functor $\mathbf{\Pi} : {}_R \mathcal{C}_R \rightarrow \mathcal{C}$ in Definition 3.4,
- the natural transformations $\overline{\mathbf{t}} : \overline{T}_i \overline{T}_o \rightarrow \overline{T}_o \overline{T}_i$, constructed in Theorem 3.7 (2), and $\mathbf{i} : \mathbf{\Pi} \overline{T}_o \rightarrow \mathbf{\Pi} \overline{T}_i$, constructed in Theorem 3.8,

constitute an admissible septuple in the sense of Definition 3.10. Moreover, a transposition morphism for it consists of an (R, R) -bimodule X and an (R, R) -bimodule map $w : \overline{T}_i X \rightarrow \overline{T}_o X$, satisfying

$$\tilde{w} \circ \tau X = \overline{\tau}_o X, \quad \text{and} \quad \tilde{w} \circ \mathbf{t} X = \overline{\mathbf{t}}_o X \circ \overline{T}_o \tilde{w} \circ \mathbf{t}' X \circ T \tilde{w} \circ \mathbf{t} X, \quad (3.15)$$

where \tilde{w} is defined in terms of the natural morphism π_i in Proposition 3.2 as $\tilde{w} := w \circ \pi_i X$ and the natural transformation $\mathbf{t}' : T_i \overline{T}_o \rightarrow \overline{T}_o T_i$ is defined in Theorem 3.7 (1).

Proof. \overline{T}_o and \overline{T}_i are monads on ${}_R \mathcal{C}_R$ by Proposition 3.1 and Proposition 3.2, respectively. $\mathbf{\Pi}$ is a functor ${}_R \mathcal{C}_R \rightarrow \mathcal{C}$ by Definition 3.4. $\overline{\mathbf{t}}$ is a distributive law $\overline{T}_i \overline{T}_o \rightarrow \overline{T}_o \overline{T}_i$ by Theorem 3.7 (3). Thus in order to verify that they constitute an admissible septuple, we have to prove that the natural transformation $\mathbf{i} : \mathbf{\Pi} \overline{T}_o \rightarrow \mathbf{\Pi} \overline{T}_i$ in Theorem 3.8 satisfies conditions (3.12). In terms of the natural epimorphism p from the forgetful functor ${}_R \mathcal{C}_R \rightarrow \mathcal{C}$ to $\mathbf{\Pi}$,

$$\mathbf{i} \circ \mathbf{\Pi} \overline{\tau}_o \circ p = \mathbf{i} \circ \mathbf{\Pi} \pi_o \circ \mathbf{\Pi} \tau \circ p = \mathbf{i} \circ \mathbf{\Pi} \pi_o \circ p T \circ \tau = p \overline{T}_i \circ \pi_i \circ \tau = p \overline{T}_i \circ \overline{\tau}_i = \mathbf{\Pi} \overline{\tau}_i \circ p. \quad (3.16)$$

In the first equality we used (3.2) and in the penultimate equality we used the first identity in (3.6). The third equality is a consequence of the first identity in (3.11). The other equalities follow by naturality. Since p is epi, (3.16) proves the first condition in (3.12).

The second condition in (3.12) is proven in Figure 13. The first, fourth and seventh equalities follow by the first identity in (3.11). The third equality is a consequence of (3.8). In the fifth equality we used the second condition in (3.6) and in the seventh one we used (3.3). The other equalities follow by naturality. Since $p \overline{T}_o \overline{T}_o \circ \pi_o \overline{T}_o \circ T \pi_o$ is epi, we have the second condition in (3.12) proven.




FIGURE 13. The proof of Theorem 3.12 (3).

The last claim about transposition morphisms is proven by showing the equivalence of conditions (3.15) and (3.13). In view of the first condition in (3.6), the first conditions in (3.15) and (3.13) are obviously equivalent. In the second frame of Figure 13 it is shown that the other conditions in (3.15) and (3.13) are equivalent. Composition on the right of both sides of the second condition in (3.13) with the epimorphism $\pi_i \bar{T}_i X \circ T\pi_i X$ yields an equivalent condition, namely the first equality in the above mentioned picture. This is equivalent to the second equality, in view of (3.6) together with the fact that, by (3.7) and (3.8), $\bar{T}_o \pi_i \circ \iota' = \bar{t} \circ \pi_i \bar{T}_o$. To conclude the proof we use $\tilde{w} = w \circ \pi_i X$. \square

Combining Theorem 3.12 with Theorem 3.11, we obtain the following

COROLLARY 3.13. *Let $\mathcal{S} = (\mathcal{C}, T, R, \iota, \iota', \tau)$ be a braided pair of monads and $\varphi : R \rightarrow T$ be a braid preserving monad morphism. Consider the monads \bar{T}_o on ${}_R \mathcal{C}_R$ in Proposition 3.1 and \bar{T}_i in Proposition 3.2 and the functor $\Pi : {}_R \mathcal{C}_R \rightarrow \mathcal{C}$ in Definition 3.4. Then any (R, R) -bimodule morphism $w : \bar{T}_i X \rightarrow \bar{T}_o X$, satisfying (3.15), determines a para-cocyclic structure on the cosimplex $\Pi \bar{T}_o^{\rightarrow s+1} X$. The para-cocyclic morphism is given in terms of the natural transformation \mathfrak{i} , constructed in Theorem 3.8, as*

$$\Pi \bar{T}_o^n w \circ \Pi \bar{T}_o^{n-1} \bar{t} \circ \Pi \bar{T}_o^{n-2} \bar{t} \bar{T}_o \circ \dots \circ \Pi \bar{T}_o \bar{t} \bar{T}_o^{n-2} \circ \Pi \bar{t} \bar{T}_o^{n-1} \circ \mathfrak{i} \bar{T}_o^n : \Pi \bar{T}_o^{n+1} X \rightarrow \Pi \bar{T}_o^{n+1} X.$$

An immediate example of the situation in Corollary 3.13 is induced by an algebra homomorphism in a braided monoidal category.

EXAMPLE 3.14. In this example, in order to simplify formulae, we do not write the associativity constraints. Let $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ be a morphism of algebras in a braided monoidal category, in which the functors $X \otimes (-)$ and $(-) \otimes X$ are right exact, for any object X in \mathcal{C} . Assume that φ satisfies (2.1) and consider the corresponding braid preserving monad morphism in Example 2.3. It determines an admissible septuple $(\mathcal{C}, \mathbf{R}\text{-Mod-}\mathbf{R}, \mathbf{T} \otimes_{\mathbf{R}} (-), \mathbf{T} \otimes_{\mathbf{R}^\times} (-), \Pi, \bar{t}, \mathfrak{i})$, where the functors $\mathbf{T} \otimes_{\mathbf{R}} (-)$, $\mathbf{T} \otimes_{\mathbf{R}^\times} (-)$ and Π are described in Example 3.3 and Example 3.6, respectively, and the natural transformations \bar{t} and \mathfrak{i} can be found in Example 3.9. For any \mathbf{R} -bimodule X there is a corresponding cosimplex, given at degree n by

$$Z^n(X, \varphi) := \mathbf{T}^{\widehat{\otimes}_{\mathbf{R}} n+1} \widehat{\otimes}_{\mathbf{R}} X,$$

known as a (braided) cyclic \mathbf{R} -module tensor product. A transposition morphism is an \mathbf{R} -bimodule map $w : \mathbf{T} \otimes_{\mathbf{R}^\times} X \rightarrow \mathbf{T} \otimes_{\mathbf{R}} X$, satisfying the conditions

$$\begin{aligned} w \circ (\varphi \otimes_{\mathbf{R}^\times} X) &= \varphi \otimes_{\mathbf{R}} X \\ w \circ (\bar{t}_i \otimes_{\mathbf{R}^\times} X) &= (\bar{t}_o \otimes_{\mathbf{R}} X) \circ (\mathbf{T} \otimes_{\mathbf{R}} w) \circ \bar{t} X \circ (\mathbf{T} \otimes_{\mathbf{R}^\times} w), \end{aligned}$$

where the multiplication maps $\bar{t}_o : \mathbf{T} \otimes_{\mathbf{R}} \mathbf{T} \rightarrow \mathbf{T}$ and $\bar{t}_i : \mathbf{T} \otimes_{\mathbf{R}^\times} \mathbf{T} \rightarrow \mathbf{T}$ are induced by $\mathfrak{t} : \mathbf{T} \otimes \mathbf{T} \rightarrow \mathbf{T}$ and $\mathfrak{t} \circ \chi_{\mathbf{T}, \mathbf{T}}^{-1} : \mathbf{T} \otimes \mathbf{T} \rightarrow \mathbf{T}$, respectively, (cf. Example 3.3). In degree n , the para-cocyclic operator is given by

$$w_n = (\mathbf{T}^{\widehat{\otimes}_{\mathbf{R}} n} \widehat{\otimes}_{\mathbf{R}} w) \circ \Pi \bar{t}^n X \circ \mathfrak{i}(\mathbf{T}^{\otimes_{\mathbf{R}^\times} n} \otimes_{\mathbf{R}} X),$$

where $\bar{\tau}^n X$ is the projection of $\chi_{\mathbf{T}, \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes X}$, i.e. the unique map $\mathbf{T} \otimes_{\mathbf{R}^\times} (\mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes_{\mathbf{R}} X) \rightarrow \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes_{\mathbf{R}} (\mathbf{T} \otimes_{\mathbf{R}^\times} X)$, for which

$$P_{oi} \circ (\chi_{\mathbf{T}, \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes X) = \bar{\tau}^n X \circ P_{io},$$

where $P_{oi} : \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes \mathbf{T} \otimes X \rightarrow \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes_{\mathbf{R}} (\mathbf{T} \otimes_{\mathbf{R}^\times} X)$ and $P_{io} : \mathbf{T} \otimes \mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes X \rightarrow \mathbf{T} \otimes_{\mathbf{R}^\times} (\mathbf{T}^{\otimes_{\mathbf{R}} n} \otimes_{\mathbf{R}} X)$ denote the canonical epimorphisms.

Examples of cosimplices in Hopf cyclic theory, corresponding to (co)module algebras of a bialgebroid (or in particular a bialgebra) over \mathbf{R} , are of this kind [BS]. Further examples are presented in Example 3.15 and Example 3.16 below.

EXAMPLE 3.15. Let H be a Hopf algebra with comultiplication $\Delta : H \rightarrow H \otimes H$ and counit $\varepsilon : H \rightarrow \mathbb{K}$. For simplifying computations, we shall use the Sweedler-Heynemann Σ -notation $\Delta(h) = \sum h_{(1)} \otimes h_{(2)}$.

An important class of braided monoidal categories is provided by the categories \mathcal{M}^H of right comodules over a coquasitriangular Hopf algebra H . Cf. [Mo, Chapter 10], a Hopf algebra H is *coquasitriangular* if there is an invertible (with respect to the convolution product) \mathbb{K} -linear map $\langle -, - \rangle : H \otimes H \rightarrow \mathbb{K}$ such that, for $h, k, l \in H$,

$$\sum \langle h_{(1)}, k_{(1)} \rangle k_{(2)} h_{(2)} = \sum h_{(1)} k_{(1)} \langle h_{(2)}, k_{(2)} \rangle, \quad (3.17)$$

$$\langle h, kl \rangle = \sum \langle h_{(1)}, k \rangle \langle h_{(2)}, l \rangle, \quad (3.18)$$

$$\langle hk, l \rangle = \sum \langle h, l_{(2)} \rangle \langle k, l_{(1)} \rangle. \quad (3.19)$$

If, in addition, the inverse in convolution of $\langle -, - \rangle$ is equal to $\langle -, - \rangle \circ \tau$, we shall say that H is *cotriangular*. Here $\tau : H \otimes H \rightarrow H \otimes H$ denotes the usual flip map. By definition, $\langle -, - \rangle$ is called the coquasitriangular map of H .

We fix a coquasitriangular Hopf algebra $(H, \langle -, - \rangle)$ over a field \mathbb{K} . Let us briefly recall the braided monoidal structure of \mathcal{M}^H , that corresponds to $\langle -, - \rangle$. First, for (M, ρ_M) and (N, ρ_N) in \mathcal{M}^H , the tensor product of M and N in \mathcal{M}^H is $M \otimes_{\mathbb{K}} N$, regarded as a comodule with respect to the right diagonal coaction

$$\rho(m \otimes n) = \sum m_{(0)} \otimes n_{(0)} \otimes m_{(1)} n_{(1)}.$$

In this formula, for a right comodule (M, ρ_M) , we used the Σ -notation $\rho_M(m) = \sum m_{(0)} \otimes m_{(1)}$. The associativity and unity constraints in \mathcal{M}^H are induced by the corresponding structures in the monoidal category of \mathbb{K} -linear spaces. The unit object is \mathbb{K} , regarded as a trivial H -comodule. To define the braiding in \mathcal{M}^H we use the coquasitriangular map as follows. For two right H -comodules M and N we define the natural map

$$\chi_{M,N} : M \otimes_{\mathbb{K}} N \rightarrow N \otimes_{\mathbb{K}} M, \quad \chi_{M,N}(m \otimes n) = \sum \langle n_{(1)}, m_{(1)} \rangle n_{(0)} \otimes m_{(0)}.$$

It is well-known that \mathcal{M}^H is a braided monoidal category with braiding χ . Furthermore, \mathcal{M}^H is a symmetric monoidal category with respect to χ (that is $\chi_{N,M} \circ \chi_{M,N} = \text{Id}_{M \otimes N}$, for $N, M \in \mathcal{M}^H$), if and only if H is *cotriangular*.

Commutative Hopf algebras are the simplest examples of cotriangular Hopf algebras. In this case the cotriangular structure $\langle -, - \rangle$ may be taken the trivial \mathbb{K} -linear map $\langle k, k \rangle := \varepsilon(h)\varepsilon(k)$, for any $h, k \in H$. Thus, the braiding is induced by τ , the canonical flip map.

Group Hopf algebras are not necessarily coquasitriangular. Indeed, let G denote a group and assume that $\langle -, - \rangle$ is a coquasitriangular map on $\mathbb{K}G$. Let $\gamma : G \times G \rightarrow \mathbb{K}$ be the map

$$\gamma(h, k) := \langle h, k \rangle, \quad \text{for } h, k \in G.$$

Since the coquasitriangular map is invertible in convolution, it follows easily that $\langle h, k \rangle \in \mathbb{K}^*$, for any $h, k \in G$. Hence γ can be regarded as a map to \mathbb{K}^* . Clearly, then the condition (3.17) is equivalent to the fact that G is abelian. On the other hand, relations (3.18) and (3.19) are equivalent to

$$\gamma(hk, g) = \gamma(h, g)\gamma(k, g) \quad \text{and} \quad \gamma(g, hk) = \gamma(g, h)\gamma(g, k), \quad \forall h, g, k \in G.$$

A map $\gamma : G \times G \rightarrow K^*$, $\gamma(h, k) = \langle h, k \rangle$ satisfying the above identities is called *bi-character* of G . In conclusion, $\mathbb{K}G$ is coquasitriangular if, and only if, G is abelian and there is a bi-character $\gamma : G \times G \rightarrow \mathbb{K}$ on G . As a matter of fact, we have also proved that, for an abelian group G , there is an one-to-one correspondence between the set of coquasitriangular structures on $\mathbb{K}G$ and the set of bi-characters on G .

The category $\mathcal{M}^{\mathbb{K}G}$ is equivalent to the category of G -graded vector spaces. Therefore, an object in $\mathcal{M}^{\mathbb{K}G}$ is a vector space V together with a decomposition $V := \bigoplus_{g \in G} V_g$. The tensor product of V and W in $\mathcal{M}^{\mathbb{K}G}$ is $V \otimes_{\mathbb{K}} W$ on which we take the decomposition

$$(V \otimes_{\mathbb{K}} W)_g = \bigoplus_{hk=g} V_h \otimes_{\mathbb{K}} W_k.$$

The braiding $\chi_{V,W} : V \otimes_{\mathbb{K}} W \rightarrow W \otimes_{\mathbb{K}} V$, for $v \in V_h$ and $w \in W_k$, is given by

$$\chi_{V,W}(v \otimes w) = \gamma(k, h)w \otimes v.$$

Note that the bi-character $\gamma : G \times G \rightarrow \mathbb{K}^*$ defines a cotriangular structure on $\mathbb{K}G$ if, and only if, γ is *symmetric*, that is, for $h \in G$ and $k \in G$,

$$\gamma(h, k)^{-1} = \gamma(k, h).$$

Recall that the category of super vector spaces can be seen as the symmetric monoidal category of \mathbb{Z}_2 -graded vector spaces, whose braiding is induced by the bi-character $\gamma : \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{K}^*$, $\gamma(g, h) = (-1)^{gh}$.

Our aim now is to specialize the construction in Example 3.14 to the case when $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ is a braid preserving homomorphism of algebras in \mathcal{M}^H , for a coquasitriangular Hopf algebra H . In this case, the conditions in (2.1) take the form

$$\begin{aligned} \varphi(r) \otimes_{\mathbb{K}} t &= \sum \langle r_{(1)(1)}, t_{(1)(1)} \rangle \langle t_{(1)(2)}, r_{(1)(2)} \rangle \varphi(r_{(0)}) \otimes_{\mathbb{K}} t_{(0)} && \text{and} && (3.20) \\ \varphi(r) \otimes_{\mathbb{K}} r' &= \sum \langle r_{(1)(1)}, r'_{(1)(1)} \rangle \langle r'_{(1)(2)}, r_{(1)(2)} \rangle \varphi(r_{(0)}) \otimes_{\mathbb{K}} r'_{(0)}, && \text{for } r, r' \in \mathbf{R}, t \in \mathbf{T}. \end{aligned}$$

Clearly, \overline{T}_o is the functor $\mathbf{T} \otimes_{\mathbf{R}} (-)$. The multiplication in \mathbf{R}^X is defined, for r' and r'' in \mathbf{R} , by

$$r' \cdot r'' := \sum \langle r''_{(1)}, r'_{(1)} \rangle r''_{(0)} r'_{(0)}.$$

We have already noticed that \mathbf{R}^X is an algebra in \mathcal{M}^H and that \mathbf{T} is a right \mathbf{R}^X -module with respect to the action

$$t \cdot r := \sum \langle r_{(1)}, t_{(1)} \rangle \varphi(r_{(0)}) t_{(0)}.$$

Furthermore, every \mathbf{R} -bimodule X can be seen as a left \mathbf{R}^X -module, with respect to the action

$$r \cdot x := \sum \langle x_{(1)}, r_{(1)} \rangle x_{(0)} r_{(0)},$$

where $x \in X$ and $r \in \mathbf{R}$. Hence $\overline{T}_i := \mathbf{T} \otimes_{\mathbf{R}^X} (-)$. For X as above, ΠX is the quotient of X with respect to the vector space generated by the commutators

$$[x, r] := rx - \sum \langle x_{(1)}, r_{(1)} \rangle x_{(0)} r_{(0)},$$

where r and x run arbitrarily in \mathbf{R} and X , respectively. For $t', t'' \in \mathbf{T}$ and $x \in X$, the morphisms $\overline{i}X : \mathbf{T} \otimes_{\mathbf{R}^X} (\mathbf{T} \otimes_{\mathbf{R}} X) \rightarrow \mathbf{T} \otimes_{\mathbf{R}} (\mathbf{T} \otimes_{\mathbf{R}^X} X)$ and $iX : \mathbf{T} \widehat{\otimes}_{\mathbf{R}^X} X \rightarrow \Pi(\mathbf{T} \otimes_{\mathbf{R}^X} X)$ are given by the following formulae

$$\begin{aligned} \overline{i}X(t' \otimes_{\mathbf{R}^X} t'' \otimes_{\mathbf{R}} x) &= \sum \langle t''_{(1)}, t'_{(1)} \rangle t''_{(0)} \otimes_{\mathbf{R}} t'_{(0)} \otimes_{\mathbf{R}^X} x, \\ iX(t \widehat{\otimes}_{\mathbf{R}^X} x) &= p(t \otimes_{\mathbf{R}^X} x), \end{aligned}$$

where $p : \text{Id}_{\mathbf{R}\text{-Mod-}\mathbf{R}} \rightarrow \Pi$ is the canonical projection and $\widehat{\otimes}_{\mathbf{R}}$ denotes the braided cyclic tensor product introduced in Example 3.6.

Let X be an \mathbf{R} -bimodule in \mathcal{M}^H . For a transposition morphism $w : \mathbf{T} \otimes_{\mathbf{R}^X} X \rightarrow \mathbf{T} \otimes_{\mathbf{R}} X$, we use the notation

$$w(t \otimes_{\mathbf{R}^X} x) = \sum t_w \otimes_{\mathbf{R}} x_w.$$

Using this notation, the defining conditions of a transposition morphism are equivalent to

$$w(1 \otimes_{\mathbf{R}^x} x) = 1 \otimes_{\mathbf{R}} x, \quad \text{and}$$

$$\sum \left\langle t''_{\langle 1 \rangle}, t'_{\langle 1 \rangle} \right\rangle \left(t''_{\langle 0 \rangle} t'_{\langle 0 \rangle} \right)_w \otimes_{\mathbf{R}} x_w = \sum \left\langle (t''_w)_{\langle 1 \rangle}, t'_{\langle 1 \rangle} \right\rangle (t''_{\langle 0 \rangle})_{w'} \left(t'_{\langle 0 \rangle} \right)_{w'} \otimes_{\mathbf{R}} (x_w)_{w'}.$$

In degree n , the corresponding para-cocyclic object is given by $Z^n(X, \varphi, w) := \mathbf{T}^{\widehat{\otimes}_{\mathbf{R}}} n+1 \widehat{\otimes}_{\mathbf{R}} X$. Its para-cocyclic operator is

$$\begin{aligned} w_n(t^0 \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t^n \widehat{\otimes}_{\mathbf{R}} x) &= \sum \left\langle t^1_{\langle 1 \rangle}, t^0_{\langle n \rangle} \right\rangle \cdots \left\langle t^n_{\langle 1 \rangle}, t^0_{\langle 1 \rangle} \right\rangle t^1_{\langle 0 \rangle} \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t^n_{\langle 0 \rangle} \widehat{\otimes}_{\mathbf{R}} \left(t^0_{\langle 0 \rangle} \right)_w \widehat{\otimes}_{\mathbf{R}} x_w \\ &= \sum \left\langle t^1_{\langle 1 \rangle} \cdots t^n_{\langle 1 \rangle}, t^0_{\langle 1 \rangle} \right\rangle t^1_{\langle 0 \rangle} \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t^n_{\langle 0 \rangle} \widehat{\otimes}_{\mathbf{R}} \left(t^0_{\langle 0 \rangle} \right)_w \widehat{\otimes}_{\mathbf{R}} x_w. \end{aligned}$$

In the particular case when $H := \mathbb{K}G$ and the braiding is defined by a bi-character γ , conditions (3.20) reduce to

$$\varphi(r) \otimes_{\mathbb{K}} t = \gamma(k, h) \gamma(h, k) \varphi(r) \otimes_{\mathbb{K}} t \quad \text{and} \quad \varphi(r) \otimes_{\mathbb{K}} r' = \gamma(k, h) \gamma(h, k) \varphi(r) \otimes_{\mathbb{K}} r',$$

for $h, k \in G$, $r \in \mathbf{R}_h$, $r' \in \mathbf{R}_k$ and $t \in \mathbf{T}_k$. Note that, if some component \mathbf{R}_h lies within the kernel of φ , then these conditions may hold also for a non-trivial braiding between copies of \mathbf{R} and \mathbf{T} . The second condition in the definition of a transposition map becomes

$$\sum \gamma(k, h) (t'' t')_w \otimes_{\mathbf{R}} x_w = \gamma(g, h) (t''_u t'_{w'}) \otimes_{\mathbf{R}} (x_w)_{w'},$$

where $h, k, g \in G$ and $t' \in \mathbf{T}_h$, $t'' \in \mathbf{T}_k$ and x belongs to the component X_g of the G -graded vector space X . In this particular case, for $t^i \in \mathbf{T}_{g_i}$, the para-cocyclic operator satisfies

$$w_n(t^0 \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t^n \widehat{\otimes}_{\mathbf{R}} x) = \sum \gamma(g_1 \cdots g_n, g_0) t^1 \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t^n \widehat{\otimes}_{\mathbf{R}} t^0 \widehat{\otimes}_{\mathbf{R}} x_w.$$

EXAMPLE 3.16. Dually, the category ${}_H\mathcal{M}$ of left modules over a \mathbb{K} -Hopf algebra H is braided monoidal if, and only if, H is quasitriangular. Cf. [Mo, Chapter 10], a Hopf algebra H is *quasitriangular* if there is an invertible element $R := \sum_i a_i \otimes b_i$ in $H \otimes H$ such that

$$\begin{aligned} \sum h_{(2)} \otimes h_{(1)} &= R \left(\sum h_{(1)} \otimes h_{(2)} \right) R^{-1} \\ (\Delta \otimes \text{Id}_H)(R) &= R^{13} R^{23}, \\ (\text{Id}_H \otimes \Delta)(R) &= R^{13} R^{12}, \end{aligned}$$

where $R^{12} = \sum_i a_i \otimes b_i \otimes 1$ and R^{13} and R^{23} are defined analogously. If $R^{-1} := \sum_i b_i \otimes a_i$, then we say that H is *triangular*.

The braided monoidal structure that corresponds to a quasitriangular element R is defined as follows. For M and N in ${}_H\mathcal{M}$, the tensor product of M and N in ${}_H\mathcal{M}$ is $M \otimes_{\mathbb{K}} N$, regarded as a module with respect to the left diagonal action

$$h \triangleright (m \otimes n) = \sum h_{(1)} \triangleright m \otimes h_{(2)} \triangleright n.$$

The associativity and unity constraints in ${}_H\mathcal{M}$ are induced by the corresponding structures in the monoidal category of \mathbb{K} -linear spaces. The unit object is \mathbb{K} , regarded as a trivial H -module. The functorial morphism of left H -modules

$$\chi_{M,N} : M \otimes_{\mathbb{K}} N \rightarrow N \otimes_{\mathbb{K}} M, \quad \chi_{M,N}(m \otimes n) = \sum_j (c_j \triangleright n) \otimes (d_j \triangleright m)$$

is a braiding on ${}_H\mathcal{M}$, where $R^{-1} = \sum_j c_j \otimes d_j$.

Let us specialize the para-cocyclic object constructed in Example 3.14 to a homomorphism of algebras $\varphi : \mathbf{R} \rightarrow \mathbf{T}$ in the braided monoidal category ${}_H\mathcal{M}$. In this case, conditions (2.1) read as

$$\varphi(r) \otimes_{\mathbb{K}} t = \sum_{j,k} (c_k d_j \triangleright \varphi(r)) \otimes_{\mathbb{K}} (d_k c_j \triangleright t) \quad \text{and} \quad \varphi(r) \otimes_{\mathbb{K}} r' = \sum_{j,k} (c_k d_j \triangleright \varphi(r)) \otimes_{\mathbb{K}} (d_k c_j \triangleright r'),$$

for $r, r' \in \mathbf{R}$ and $t \in \mathbf{T}$, where $\sum_j c_j \otimes d_j = R^{-1} = \sum_k c_k \otimes d_k$. Clearly, \overline{T}_o is the functor $\mathbf{T} \otimes_{\mathbf{R}} (-)$. Moreover, the multiplication in \mathbf{R}^x is defined, for r' and r'' in \mathbf{R} , by

$$r' \cdot r'' := \sum_j (c_j \triangleright r'') (d_j \triangleright r').$$

We have already noticed that \mathbf{R}^X is an algebra in ${}_H\mathcal{M}$ and \mathbf{T} is a right \mathbf{R}^X -module with respect to the action

$$t \cdot r := \sum (c_j \triangleright \varphi(r)) (d_j \triangleright t).$$

Furthermore, if X is an \mathbf{R} -bimodule then, for $x \in X$ and $r \in \mathbf{R}$, the following formula

$$r \cdot x := \sum_j (c_j \triangleright x) (d_j \triangleright r)$$

defines a left \mathbf{R}^X -module structure. Hence $\bar{T}_i := \mathbf{T} \otimes_{\mathbf{R}^X} (-)$. For X as above, ΠX is the quotient of X with respect to the vector space generated by the commutators

$$[x, r] := rx - \sum_j (c_j \triangleright x) (d_j \triangleright r),$$

where r and x run arbitrarily in \mathbf{R} and X , respectively. For $t', t'' \in \mathbf{T}$ and $x \in X$, the morphisms $\bar{i}X : \mathbf{T} \otimes_{\mathbf{R}^X} (\mathbf{T} \otimes_{\mathbf{R}} X) \rightarrow \mathbf{T} \otimes_{\mathbf{R}} (\mathbf{T} \otimes_{\mathbf{R}^X} X)$ and $iX : \mathbf{T} \widehat{\otimes}_{\mathbf{R}} X \rightarrow \Pi(\mathbf{T} \otimes_{\mathbf{R}^X} X)$ are given by the following formulae

$$\begin{aligned} \bar{i}X(t' \otimes_{\mathbf{R}^X} t'' \otimes_{\mathbf{R}} x) &= \sum_j c_j \triangleright t'' \otimes_{\mathbf{R}} d_j \triangleright t' \otimes_{\mathbf{R}^X} x, \\ iX(t \widehat{\otimes}_{\mathbf{R}} x) &= p(t \otimes_{\mathbf{R}^X} x), \end{aligned}$$

where $p : \text{Id}_{\mathbf{R}\text{-Mod-}\mathbf{R}} \rightarrow \Pi$ is the canonical projection and $\widehat{\otimes}_{\mathbf{R}}$ denotes the braided cyclic tensor product introduced in Example 3.6.

Let X be an \mathbf{R} -bimodule in ${}_H\mathcal{M}$. For a transposition morphism $w : \mathbf{T} \otimes_{\mathbf{R}^X} X \rightarrow \mathbf{T} \otimes_{\mathbf{R}} X$, we use the notation

$$w(t \otimes_{\mathbf{R}^X} x) = \sum t_w \otimes_{\mathbf{R}} x_w.$$

Using this notation, the defining conditions of a transposition map are equivalent to

$$\begin{aligned} w(1 \otimes_{\mathbf{R}^X} x) &= 1 \otimes_{\mathbf{R}} x, \quad \text{and} \\ \sum_j [(c_j \triangleright t'') (d_j \triangleright t)]_w \otimes_{\mathbf{R}} x_w &= \sum_j (c_j \triangleright t''_w) (d_j \triangleright t'_w) \otimes_{\mathbf{R}} (x_w)_{w'}. \end{aligned}$$

In degree n , the corresponding para-cocyclic object is given by $Z^n(X, \varphi, w) := \mathbf{T} \widehat{\otimes}_{\mathbf{R}}^{n+1} \widehat{\otimes}_{\mathbf{R}} X$ and its para-cocyclic operator is

$$\begin{aligned} w_n(t_0 \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} t_n \widehat{\otimes}_{\mathbf{R}} x) &= \sum_{j_1, \dots, j_n} \left(c_{j_1}^{(1)} \triangleright t_1 \right) \widehat{\otimes}_{\mathbf{R}} \cdots \widehat{\otimes}_{\mathbf{R}} \left(c_{j_n}^{(n)} \triangleright t_n \right) \widehat{\otimes}_{\mathbf{R}} \\ &\quad \widehat{\otimes}_{\mathbf{R}} \left[d_{j_n}^{(n)} \triangleright \left(\cdots \triangleright \left(d_{j_1}^{(1)} \triangleright t_0 \right) \cdots \right) \right]_w \widehat{\otimes}_{\mathbf{R}} x_w, \end{aligned}$$

where $\sum_{j_k} c_{j_k}^{(k)} \otimes d_{j_k}^{(k)} = R^{-1}$, for every $k = 1, \dots, n$.

REFERENCES

- [AMS] A. Ardizzoni, C. Menini and D. Ştefan, *Hochschild cohomology and “smoothness” in monoidal categories*, J. Pure Appl. Algebra **208** (2007), 297-330.
- [Be] J. Beck, *Distributive laws*, Lecture Notes in Mathematics **80**, 119–140, Springer Verlag, 1969.
- [BŞ] G. Böhm and D Ştefan, (Co)cyclic (co)homology of bialgebroids: An approach via (co)monads, preprint arXiv:arXiv:0705.3190.
- [KLV] S. Kasangian, S. Lack and E. Vitale, *Coalgebras, braidings, and distributive laws*, Theory and Applications of Categories **13** (2004), 129-146.
- [Kas] C. Kassel, *Quantum groups*, Graduate Text in Mathematics, **155**, Springer-Verlag, New York, 1995.
- [Kay] A. Kaygun, *The universal Hopf cyclic theory*, preprint arXiv:math.KT/0609311.
- [KR] M. Khalkhali and B. Rangipour, *A note on cyclic duality and Hopf algebras*, Comm. Algebra **33** (2005), 763–773.
- [LPvO] J. Lopez, F. Panaite and F. van Oystaeyen, *General twisting of algebras*, to appear in Adv. Mathematics.
- [Mo] S. Montgomery, “Hopf Algebras and their actions on rings”, CMBS Regional Conference Series in Mathematics **82**, 1993.
- [We] C. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics **38**, Cambridge, University Press, 1994.

RESEARCH INSTITUTE FOR PARTICLE AND NUCLEAR PHYSICS, BUDAPEST, H-1525 BUDAPEST 114, P.O.B.49,
HUNGARY.

UNIVERSITY OF BUCHAREST, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, 14 ACADEMIEI STREET,
BUCHAREST, RO-010014, ROMANIA.