General Single Field Inflation with Large Positive Non-Gaussianity

Miao Li^{1,2}, Tower Wang^{2,1}, Yi Wang^{2,1}

 ¹ Interdisciplinary Center of Theoretical Studies, USTC, Hefei, Anhui 230026, P.R.China
 ² Institute of Theoretical Physics, CAS, Beijing 100080, P.R.China

Recent analysis of the WMAP three year data suggests $f_{NL}^{local} \simeq 86.8$ in the WMAP convention. It is necessary to make sure whether general single field inflation can produce a large positive f_{NL} before turning to other scenarios. We give some examples to generate a large positive f_{NL}^{equil} in general single field inflation. Our models are different from ghost inflation. Due to the appearance of non-conventional kinetic terms, $f_{NL}^{equil} \gg 1$ can be realized in single field inflation.

1 Introduction

As modern cosmological observations become more and more precise, study of the non-Gaussianity of CMB temperature fluctuations has become a more and more pressing issue in recent years. In the WMAP convention [1, 2], the primordial non-Gaussianity is parameterized by f_{NL} assuming the ansatz

$$\zeta = \zeta_L + \frac{3}{5} f_{NL} \zeta_L^2,\tag{1}$$

where ζ is the scalar perturbation and ζ_L is its linear Gaussian part¹. However, this ansatz only corresponds to a restricted shape of non-Gaussianity. Theoretically, during inflation, the non-Gaussianity is usually produced in different shapes, and the estimator f_{NL} can be defined in terms of the 3-point function $\langle \zeta_{k_1} \zeta_{k_2} \zeta_{k_3} \rangle$. Two limits of f_{NL} are of most interest. One is the local, squeezed limit $k_1 \ll k_2 \simeq k_3$, for which we will use the notation f_{NL}^{local} . The other is the non-local, equilateral limit $k_1 \simeq k_2 \simeq k_3$, for which the notation will be f_{NL}^{equil} .

In a recent work [7], utilizing the fast estimator of primordial non-Gaussianity [8], Yadav and Wandelt claimed that data from two channels of WMAP3 reject $f_{NL}^{local} = 0$ at the 2.89 σ level, or 99.6% significance. They also showed that 26.91 $< f_{NL}^{local} <$ 146.71 at 95% C.L., with a central value of $f_{NL}^{local} =$ 86.8. If this result is confirmed by future observations, it will have a great impact on our study of the early universe, because a large class of inflation models will be ruled out. For example, the simplest model of inflation is slow rolling, driven by a single scalar field. Ignoring the non-Gaussianity, previous observational data fit well with single-field slow-roll inflation [2, 9]. On the other hand, in the conventional single-field slow-roll inflation, it has been found for both local and equilateral forms that $|f_{NL}| < 1$, which is too small to detect in the near future [3, 10]. Therefore, the confirmation of a large f_{NL}^{local} observationally will exclude almost all models of single-field slow-roll inflation.

Confronted with the evidence for $f_{NL}^{local} \gg 1$, is single-field slow-roll inflation dying? Not necessarily. There are at least two ways to save it. First, a large non-Gaussianity may arise in stochastic inflation due to non-linear effects between inflaton and metric

¹Note that Maldacena's convention is also popularly used in the literature [3, 4], which assumes $\zeta = \zeta_L - \frac{3}{5} f_{NL} \zeta_L^2$ instead of (1). That convention is different in sign of f_{NL} from the WMAP convention. Please refer to [5, 6] for clarification. In this paper, we will use the WMAP convention.

perturbations [11]. Second, the curvaton mechanism provides an elegant way to produce a large positive f_{NL}^{local} [12].

Although there is no evidence for $|f_{NL}^{equil}|$ to be large at present, several inflation models with $|f_{NL}^{equil}| \gg 1$ have already appeared in the past few years. In single-field slow-roll inflation, this is realized by the introduction of non-canonical kinetic terms, such as k-inflation [13, 14], ghost inflation [15] DBI inflation [16, 17, 18, 19], and some other mechanisms [20]. Translated into the WMAP convention, most of the models predict $f_{NL}^{equil} \ll -1$ and a small f_{NL}^{local} . It is remarkable that the value of f_{NL}^{local} favored by [7] is of positive sign in the WMAP convention. For f_{NL}^{equil} , the present constraint is not stringent enough to make a conclusion. So one expects that there are three possibilities in the future:

- 1. $f_{NL}^{local} \gg 1$, $-1 < f_{NL}^{equil} < 1$. This can be explained by the conventional single-field slow-roll inflation + curvaton mechanism.
- 2. $f_{NL}^{local} \gg 1$, $f_{NL}^{equil} < -1$. This corresponds to DBI inflation (or most known k-inflation/ghost inflation) + curvaton mechanism.
- 3. $f_{NL}^{local} \gg 1$, $f_{NL}^{equil} > 1$. This is more challenging to explain.

For both the second and the third possibility, one should keep in mind that additional fine-tuning is needed in order that both the curvaton and the inflaton produce perturbations of comparable magnitude.

The purpose of this paper is to search for k-inflation models with $f_{NL}^{equil} \gg 1$. If this type of models are constructed, one can combine them with the curvaton mechanism to meet the challenge posed by the third possibility.

Indeed, starting from the action (2), we have found several examples of general single field inflation in which $f_{NL}^{equil} \gg 1$. As will be shown in section 4, even if the inflation is driven by the potential of inflaton, a large positive f_{NL}^{equil} can be generated by non-conventional kinetic terms. In some other models, the non-conventional kinetic terms not only give rise to the desired f_{NL}^{equil} , but also drive the inflation. Examples for this type of models are constructed in power-law k-inflation. In all our models, typically the desired non-Gaussianity stems from high order terms in X, with X defined in (3).

The paper is organized as follows. In section 2, we review briefly the general single field inflation. In section 3, we prove a no-go theorem for the p(X) models. We show that if the matter Lagrangian depends only on X, but not on ϕ directly, then one can not obtain a large and positive f_{NL}^{equil} . In section 4, we construct generalized slow roll inflation models with $f_{NL}^{equil} \gg 1$. In section 5, we construct power-law inflation models with $f_{NL}^{equil} \gg 1$. We conclude in section 6.

2 A Brief Review of General Single Field Inflation

In this section, we give a brief review on the general single field inflation. For further details, please refer to [4, 13, 14, 21]. We consider the action [13]

$$S = \int d^4x \sqrt{-g} [\frac{1}{16\pi G} R + p(\phi, X)],$$
 (2)

where

$$X = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\phi\partial_{\nu}\phi, \qquad (3)$$

and the signature of metric is (-1, 1, 1, 1). We henceforth set the reduced Planck mass $M_{pl} = (8\pi G)^{-\frac{1}{2}}$ to unity.

In terms of the pressure $p(\phi, X)$ and its derivatives with respect to X (denoted by p_{X} etc.), we can write down the energy density

$$\rho(\phi, X) = 2Xp_{,X} - p \tag{4}$$

of the inflaton as well as the speed of sound

$$c_s^2 = \frac{p_{,X}}{p_{,X} + 2Xp_{,XX}}.$$
 (5)

The Friedmann equation and the continuity equation are given by

$$3H^2 = \rho,$$

$$\dot{\rho} = -3H(\rho + p). \tag{6}$$

It proves useful to define two quantities

$$\Sigma = X p_{,X} + 2X^2 p_{,XX}, \lambda = X^2 p_{,XX} + \frac{2}{3} X^3 p_{,XXX},$$
(7)

and some slow-variation parameters

$$\epsilon = -\frac{H}{H^2} = \frac{3Xp_{,X}}{2Xp_{,X} - p},$$

$$\eta = \frac{\dot{\epsilon}}{H\epsilon}, \quad s = \frac{\dot{c}_s}{Hc_s}, \quad l = \frac{\dot{\lambda}}{H\lambda}$$
(8)

following [4, 21]. We make note that one of the "slow-variation" parameters η here is different from one of the "slow-roll" parameters frequently used in ordinary slow-roll inflation. On this point please see [21] for clarification. Throughout our discussion, we will be interested only in the the slow-variation case $\epsilon, \eta, s, l \ll 1$ with $\dot{H} \leq 0$, $\epsilon \geq 0$.

According to [14], to the leading order, the power spectra for scalar and tensor perturbations are

$$P_k^{\zeta} = \frac{H^2}{8\pi^2 c_s \epsilon},$$

$$P_k^h = \frac{2H^2}{\pi^2},$$
(9)

which lead to the tensor-to-scalar ratio

$$r = \frac{P_k^h}{P_k^{\zeta}} = 16c_s \epsilon. \tag{10}$$

While their spectral indices are

$$n_s - 1 = -2\epsilon - \eta - s,$$

$$n_T = -2\epsilon.$$
(11)

In accordance with the WMAP convention (1), the non-Gaussianity parameter f_{NL} in the equilateral triangle limit is [4]

$$f_{NL}^{equil} = -\frac{10}{81}\frac{\lambda}{\Sigma} + \left(\frac{5}{81} - \frac{35}{108}\right)\left(\frac{1}{c_s^2} - 1\right) + \frac{5(3-2\gamma)}{81}\frac{l\lambda}{\Sigma} + \mathcal{O}\left(\frac{\epsilon}{c_s^2}, \frac{\epsilon\lambda}{\Sigma}\right) + \mathcal{O}(\epsilon).$$
(12)

Here γ is the Euler-Mascheroni constant, which is denoted by \mathbf{c}_1 in [4]. Numerically $\gamma \simeq 0.577$. Please always keep in mind that we follow the WMAP convention hence the f_{NL}^{equil} here is opposite in sign with respect to that in [3, 4]. From (12) it is clear that in order to get a large positive f_{NL}^{equil} , we should have $-\frac{\lambda}{\Sigma} \gg 1$. Neglecting the sub-leading terms, one can naively estimate $-\frac{10}{81\Sigma} \frac{\lambda}{\Sigma}$ with f_{NL}^{equil} , thus we have $-\frac{\lambda}{\Sigma} \sim$

 $\mathcal{O}(10^3)$ in order to get $f_{NL}^{equil} \sim 100$. This observation will be useful in our model reconstruction below.

Note that in the case $-\frac{\lambda}{\Sigma} \gg 1$, the $\mathcal{O}(\epsilon \lambda / \Sigma)$ correction also gives a $\mathcal{O}(1)$ contribution. So to be more precise, we have

$$f_{NL}^{equil} = -\frac{10}{81}\frac{\lambda}{\Sigma} + \left(\frac{5}{81} - \frac{35}{108}\right)\left(\frac{1}{c_s^2} - 1\right) + \frac{5(3 - 2\gamma)}{81}\frac{l\lambda}{\Sigma} - \frac{5}{243}\frac{\lambda}{\Sigma}\left\{\left(2\epsilon + \eta + s\right)\left(-3\gamma - 48 + \frac{252}{2}\ln\frac{3}{2}\right) - \eta\left(6\gamma - \frac{33}{2}\right) + s(3\gamma - 12)\right\} + \mathcal{O}\left(\epsilon, \frac{\epsilon}{c_s^2}, \frac{\epsilon^2\lambda}{\Sigma}\right), \qquad (13)$$

where we have performed the integral in $R(k_1, k_2, k_3)$ [4] in the equilateral limit $k_1 = k_2 = k_3$. The details of the integration is given in Appendix A. In (13), we neglect $\mathcal{O}(\frac{\epsilon}{c_s^2})$ terms by concentrating on the parameter space $c_s^2 \gg \epsilon$. If one would like to consider models with a smaller c_s^2 , those terms should be taken into account.

3 A No-Go Result for p(X) Model

In the simplest case, Lagrangian p takes the form p = p(X) independent of ϕ . This model mimics a de Sitter space, in which the cosmological perturbations are "ill-defined" [13]. This model cannot give large positive f_{NL}^{equil} , as we now show by applying the general results in section 2 to it.

Making use of (4-8), it is not hard to check for p = p(X) that

$$s = \frac{\dot{X}}{2HX} \left(\frac{X^2 p_{,XX}}{X p_{,X}} - \frac{3X^2 p_{,XX} + 2X^3 p_{,XXX}}{X p_{,X} + 2X^2 p_{,XX}} \right)$$

$$= -\frac{3X p_{,X}}{X p_{,X} + 2X^2 p_{,XX}} \left(\frac{X^2 p_{,XX}}{X p_{,X}} - \frac{3X^2 p_{,XX} + 2X^3 p_{,XXX}}{X p_{,X} + 2X^2 p_{,XX}} \right)$$

$$= 9c_s^2 \left(\frac{\lambda}{\Sigma} + \frac{1}{6} \right) - \frac{3}{2}.$$
 (14)

It is obvious from (12) that a large positive f_{NL}^{equil} implies $\frac{\lambda}{\Sigma} \ll -1$. This requirement leads to s < -1 in (14), violating the slow-variation condition.

So a large positive f_{NL}^{equil} cannot appear in this simple model.

For a more general Lagrangian $p = p(\phi, X)$, terms like $(2Xp_{X\phi} - p_{\phi})\dot{\phi}$ will show up in the continuity equation (6), hence the above relation does not hold anymore, and one should study case by case. For some special forms of $p(\phi, X)$, one may get a relation similar to (14) and a no-go theorem likewise. For other cases, as we will investigate in the following sections, such a no-go theorem does not exist and we can construct slow-variation inflation models with $\frac{\lambda}{\Sigma} \ll -1$.

4 Reconstruction of the Generalized Slow Roll Inflation

In this section, we investigate the non-Gaussianity estimator f_{NL}^{equil} of the generalized slow roll inflation. We will show that a large and positive f_{NL}^{equil} can be obtained in relatively simple models of this class. By generalized slow roll inflation models, we mean that inflation is still driven by the potential energy of inflaton, while the inflaton has generalized kinetic terms, which can generate large non-Gaussianities. For this purpose, we study the Lagrangian

$$p(\phi, X) = g(\phi)f(X) - V(\phi) .$$
(15)

Using (4), the energy density can be written as

$$\rho = 2gXf_{,X} - gf + V . \tag{16}$$

For our purpose, we look for solutions with $|2gXf_{X}| \ll V$ and $|gf| \ll V$. The validity of this ansatz will be checked later in this section. Then to the leading order approximation, we have

$$\rho \simeq V \ . \tag{17}$$

The equation of motion of ϕ can be written as

$$\partial_t (gf_{,X}\dot{\phi}) + 3Hgf_{,X}\dot{\phi} - g_{,\phi}f + V_{,\phi} = 0$$
 (18)

For simplicity, we study solutions with $\ddot{\phi} \simeq 0$. This is the direct generalization of the $\frac{1}{2}m^2\phi^2$ model with the standard kinetic term. We expect this simplification does not lose much generality for two reasons. Firstly, for slow roll inflation, $\ddot{\phi}/(H\dot{\phi}) \ll 1$, so they commonly behave the way we assume. Secondly, we are mainly interested in the non-Gaussianity, which is generated mainly by f(X). As we will show, we can choose g and V so that the assumption $\ddot{\phi} \simeq 0$ does not lose generality for f(X).

After this approximation, and using the Friedmann equation, (18) takes the form

$$g_{,\phi}\left(f - 2Xf_{,X} - (\operatorname{sgn}\dot{\phi})\sqrt{6VX}f_{,X}\frac{g}{g_{,\phi}}\right) = V_{,\phi} , \qquad (19)$$

where $(\operatorname{sgn}\dot{\phi})$ denotes the sign of $\dot{\phi}$, which comes from the square root $\dot{\phi} = (\operatorname{sgn}\dot{\phi})\sqrt{2X}$. We demand the equation (19) to boil down to an equation of only X, so

$$\frac{g_{,\phi}}{g} \propto \sqrt{V} , \quad g_{,\phi} \propto V_{,\phi} .$$
 (20)

The solution of (20) takes the form

$$g = -\frac{1}{\cosh^2(\alpha\phi)}$$
, $V = \frac{\alpha^2\beta^2}{3} \tanh^2(\alpha\phi)$, (21)

where α and β are constants. Without losing generality, we set $\alpha > 0$, $\beta > 0$, and $\phi > 0$. In this case, ϕ rolls backwards, so $\dot{\phi} < 0$.

Inserting the solution (21) into (19), we have the equation for X

$$f - 2Xf_{,X} - \beta \sqrt{\frac{X}{2}}f_{,X} = \frac{\alpha^2 \beta^2}{3}$$
 (22)

We can take this equation either as a differential equation, which is valid for all X, or as an algebraic equation, which is valid for some certain X. It can be shown that the former possibility leads to

$$f = \frac{\alpha^2 \beta^2}{3} + C\left(\frac{\beta}{\sqrt{2}} + 2\sqrt{X}\right) , \qquad (23)$$

where C is a constant. In this case, $c_s \to \infty$ and $\lambda/\Sigma \to 0/0$, so the next to leading order contribution in the slow roll approximation must be taken into consideration. In the remainder of this section, we consider the latter possibility, and treat (22) as an algebraic equation.

Now let us verify the slow roll conditions. Compare (22) with the energy density, one can find that if $f - 2Xf_{,X}$ and $\beta\sqrt{\frac{X}{2}}f_{,X}$ do not cancel at leading order of slow roll approximation, then the condition $\alpha\phi \gg 1$ leads to the slow roll condition $\epsilon \ll 1$. This condition can be satisfied by imposing proper initial conditions.

It can be shown that

$$\eta \simeq l \simeq \frac{6\sqrt{2X}}{\beta}.$$
(24)

So the condition $\eta, l \ll 1$ can be satisfied by requiring that β is large enough.

The slow roll condition for s is automatically satisfied, because we have assumed $\dot{X} \propto \ddot{\phi} \simeq 0$, which is verified in (22) where X is a constant.

To solve (22), we need to give an explicit expression for f(X). As an illustration, we consider the simplest polynomial case

$$f = c_1 X + c_2 X^2 , (25)$$

the calculation can be generalized to other models of f(X) straightforwardly.

Note that our model has a rescaling invariance. Suppose the solution of (22) is $X = X_0$, then we can always redefine $\alpha \to \sqrt{X_0}\alpha$, $\beta \to \beta/\sqrt{X_0}$, $c_1 \to X_0c_1$ and $c_2 \to X_0^2c_2$, so that we get the solution X = 1 after performing the rescaling. So we set X = 1 in the following calculation.

One can show that λ / Σ and c_s^2 can be expressed as

$$\frac{\lambda}{\Sigma} = \frac{2c_2}{c_1 + 6c_2} , \quad c_s^2 = 1 - 2\frac{\lambda}{\Sigma} .$$
 (26)

Combining with (22), we can express the coefficients c_1 and c_2 as functions of λ/Σ as

$$c_1 = -\frac{\frac{\alpha^2 \beta^2}{3} \left(\frac{2}{-\frac{\lambda}{\Sigma}} + 6\right)}{\frac{24}{\eta} + 3 + \left(\frac{6}{\eta} + 1\right) \left(\frac{2}{-\frac{\lambda}{\Sigma}}\right)} , \quad c_2 = \frac{\frac{\alpha^2 \beta^2}{3}}{\frac{24}{\eta} + 3 + \left(\frac{6}{\eta} + 1\right) \left(\frac{2}{-\frac{\lambda}{\Sigma}}\right)} . \tag{27}$$

From the equation (12), we see when $\lambda/\Sigma \ll -1$, we can get a large and positive f_{NL}^{equil} . Note that this case corresponds to $c_s \gg 1$. In order not to generate too large tensor mode perturbation, we need $\epsilon \ll 1$, and mainly use η to generate a red spectrum for the scalar perturbations.

The parameter region $c_s \gg 1$ seems exotic, because this leads to a superluminal propagation of the inflaton perturbations. However, as discussed in [22], during inflation, the inflaton field provides a time dependent background, which determines a preferable coordinate frame. The superluminal propagation occurs only in this special frame. So causality is not violated. This causality issue is discussed in more detail in [23].

Comparing with data, from $n_s \simeq 0.96$, and the observation that ϵ is very small, we get $\eta \simeq 0.04$, so $\beta \simeq 150\sqrt{2}$. If we assume $f_{NL}^{equil} \simeq 100$, then we get $\lambda/\Sigma \simeq -810$, and $c_s \simeq 40$. Finally, from the COBE normalization $P^{\zeta} \simeq 2.5 \times 10^{-9}$, we get $\alpha^2/\epsilon \simeq 1.6 \times 10^{-9}$. Note that ϵ can still be chosen arbitrarily within the experimental range. A different choice of ϵ leads to a different tensor-to-scalar ratio r.

For example, when $r \simeq 0.3$, we have $\epsilon = 4.7 \times 10^{-4}$, $\alpha = 8.7 \times 10^{-7}$, $\phi = 3.0 \times 10^{6}$, $c_1 = -1.1 \times 10^{-10}$, and $c_2 = 1.9 \times 10^{-11}$.

At the first sight, these parameters seem to be rather unnatural. While note that for simplicity, we have rescaled the parameters to have X = 1. If we rescale back the parameters such that $-c_1/\cosh^2(\alpha\phi) \simeq 1$, then the parameters become $\alpha = 0.53$, $\beta = 0.00034$, $\phi = 4.8$, $X = 2.6 \times 10^{-12}$, $c_1 = -43$, and $c_2 = 2.7 \times 10^{12}$. Note that c_2 still seems to be too large in the Planck units. The largeness of c_2 implies the existence of a new scale, for example, the string scale M_s . If we recover the Planck mass, then

$$f = -43X + \left(2.7 \times 10^{12} \times \frac{M_s^4}{M_{pl}^4}\right) \frac{X^2}{M_s^4} .$$
 (28)

If the string scale is $M_s \sim 10^{-2} M_{pl}$ or $M_s \sim 10^{-3} M_{pl}$, then c_2 becomes of order 1. The above action assumes the form of an effective action, with the mass scale M_s playing the role of a physical cut-off.

As a matter of fact, the physical inflaton should be a "nearly canonical" field

$$\tilde{\phi} = \tilde{\phi}_0 + \frac{2\sqrt{-c_1}}{\alpha}\arctan(e^{\alpha\phi}), \qquad (29)$$

with a first order canonical kinetic term

$$\tilde{X} = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\tilde{\phi}\partial_{\nu}\tilde{\phi} = -\frac{c_1 X}{\cosh^2(\alpha\phi)}$$
(30)

when the Lagrangian is expanded. Here $\tilde{\phi}_0$ is a free parameter, which can be fixed by hand. The normalization $-c_1/\cosh^2(\alpha\phi) \simeq 1$ we have chosen facilitates our discussion greatly. It implies $\tilde{X} = X$, hence the result (28) still holds if we replace X with \tilde{X} . Numerically the inflaton $\tilde{\phi} = \tilde{\phi}_0 + 37$ in this case.

As another example, when $r \simeq 10^{-3}$, we have $\alpha = 5.0 \times 10^{-8}$, $\phi = 1.1 \times 10^8$, $c_1 = -3.8 \times 10^{-13}$, and $c_2 = 6.3 \times 10^{-14}$. After rescaling to $-c_1/\cosh^2(\alpha\phi) \simeq 1$, we have $\alpha = 9.2$, $\beta = 1.2 \times 10^{-6}$, $\phi = 0.59$, $X = 2.9 \times 10^{-17}$, $c_1 = -1.3 \times 10^4$, $c_2 = 7.3 \times 10^{19}$ and $\tilde{\phi} = \tilde{\phi}_0 + 39$.

Before proceeding to the next section, we discuss some physical issues in the models we studied above. First we note that the coefficient c_1 is negative, this is nice,

since the function g in equation (21) is negative, thus the leading kinetic term X in energy is always positive and the relation (29) is well-defined. However, the coefficient c_2 is positive, this leads to a negative X^2 term in energy, and if X is sufficiently large, this negative term will cause instability. This problem can be eased by introducing a positive higher order term in X.

We have introduced a mass scale M_s above to indicate that these models may be treated as an effective field theory arising in string theory. Just as in the DBI action, high order terms in X can be regarded as stringy correction at the tree level. In an effective action, operators with larger scaling dimensions are suppressed by power of $1/M_s$. Here X has dimension 4, so with each extra factor X, a factor $1/M_s^4$ is introduced. As we have seen, these high order terms are certainly important during inflation, as a merely X^2 can help to produce a large f_{NL} . Nevertheless, these terms become less important when the universe evolves to regimes of low energy.

In the first model discussed above, the "nearly canonical" scalar field assumes a value $\tilde{\phi} = \tilde{\phi}_0 + 37$ in the reduced Planck unit. Unless we fine-tune $\tilde{\phi}_0$, this value lies in the trans-Planckian regime. Since the potential $V(\phi)$ is proportional to the square of the hyperbolic tangent function, the scalar field ϕ rolls down towards smaller values (the linear term X dominates slightly over the quadratic term X^2 in the kinetic energy). The second model still gives a $\tilde{\phi}$ above the Planck scale without fine tuning, although ϕ is smaller than the reduced Planck scale. Again, ϕ rolls towards smaller values too. It is interesting to study carefully whether $\tilde{\phi} > 1$ will cause a disaster to our models, following the arguments in [24, 25].

5 Reconstruction of Power-Law k-Inflation

As has been discussed in [13], the Lagrangian of power-law k-inflation takes the form

$$p(\phi, X) = \frac{1}{\phi^2} g(X) \tag{31}$$

in general.

In power-law k-inflation models, the equations of motion (6) are solved by

$$\phi = \sqrt{2X}t,$$

$$X = \frac{1}{2}\dot{\phi}^2 = \text{constant},$$

$$a \propto t\sqrt{(2Xg_{,X}-g)/X},$$

$$g_{,X} = \sqrt{\frac{2(2Xg_{,X}-g)}{3X}}.$$
(32)

While the slow-variation parameters in (8) and the spectral indices are reduced to

$$\epsilon = \frac{3Xp_{,X}}{2Xp_{,X} - p},$$

$$\eta = s = 0, \quad l = -2\epsilon,$$

$$n_s - 1 = -2\epsilon = n_T.$$
(33)

The equality of $n_s - 1$ and n_T in (33), and the no-running condition

$$\alpha_s = \frac{dn_s}{d\ln k} = 0 \tag{34}$$

are important features of power-law k-inflation. These features can be used to test or rule out this class of models by future experiments. Another feature is that the spectral indices depend on ϵ exclusively, due to the vanishing of η and s in this class of models.

As we have argued, a large positive f_{NL}^{equil} in (12) requires that $-\frac{\lambda}{\Sigma} \sim \mathcal{O}(10^3)$. On the other hand, the experimentally favorite value $n_s \simeq 0.958 \pm 0.016$ indicates $\epsilon \sim \mathcal{O}(10^{-2})$ by (33). So we can see $-\frac{\epsilon\lambda}{\Sigma} \gtrsim 1$. The value of c_s can be experimentally determined by the aid of (10). Theoretically we cannot exclude the possibility that models might exist with both $c_s^2 \lesssim \epsilon$ and a large positive f_{NL}^{equil} . But from now on we will concentrate on models with $c_s^2 \gg \epsilon$, which is much simpler. Using (13), up to $\mathcal{O}(1)$, the equilateral non-Gaussianity estimator becomes

$$f_{NL}^{equil} = -\frac{10}{81}\frac{\lambda}{\Sigma} - \frac{85}{324}\left(\frac{1}{c_s^2} - 1\right) + \frac{10}{81}\left(29 + 4\gamma - 84\ln\frac{3}{2}\right)\frac{\epsilon\lambda}{\Sigma} + \mathcal{O}(\frac{\epsilon}{c_s^2}, \epsilon).$$
(35)

In this section we will reconstruct the power-law k-inflation with g(X) of some specific forms. The input parameters for reconstruction are ϵ , c_s and f_{NL}^{equil} . The sound speed c_s may be translated by relation (10) into tensor-to-scalar ratio r, which is constrained by experiment. The last line of (33) translates ϵ into n_s or n_T , so an experimental constraint on spectral indices is equivalent to a constraint on ϵ in power-law k-inflation models.

It is useful to note that

$$p_{,X} = \frac{1}{\phi^2} g_{,X}, \qquad \frac{Xp_{,X}}{p} = \frac{Xg_{,X}}{g},$$
 (36)

etc. in power-law k-inflation models (31). The expressions of c_s^2 and ϵ in (5) and (8) can be rewritten in the form

$$\frac{Xg_{,X}}{g} = \frac{\epsilon}{2\epsilon - 3} \equiv \xi_1,$$

$$\frac{X^2g_{,XX}}{Xg_{,X}} = \frac{1 - c_s^2}{2c_s^2} \equiv \xi_2.$$
 (37)

On the other hand, combining (5), (7) and (35), we have

$$\frac{X^{3}g_{,XXX}}{X^{2}g_{,XX}} = -\frac{3}{2} + \frac{3}{1-c_{s}^{2}}\frac{\lambda}{\Sigma} \\
= -\frac{3}{2} - 3\left[1 - \left(29 + 4\gamma - 84\ln\frac{3}{2}\right)\epsilon\right]^{-1}\left[\frac{81f_{NL}^{equil}}{10(1-c_{s}^{2})} + \frac{17}{8c_{s}^{2}}\right] \\
\equiv \xi_{3}.$$
(38)

Please note here

$$29 + 4\gamma - 84\ln\frac{3}{2} \simeq -2.75 \tag{39}$$

numerically. Thanks to (37), the last equation of (32) may take a simple form as follows:

$$g_{,X} = \frac{2}{\epsilon} \equiv \xi_4. \tag{40}$$

We have defined four parameters $\xi_1, \xi_2, \xi_3, \xi_4$ for later use. In addition, it is necessary to check the condition

$$2Xg_{,X} - g > 0 \tag{41}$$

dictated by $\rho > 0$, otherwise the solution (32) would break down. After a few calculations, one can quickly confirm that

$$\rho = \frac{1}{\phi^2} (2Xg_{,X} - g) = \frac{6X}{\phi^2 \epsilon^2} = \frac{3}{\epsilon^2 t^2} > 0$$
(42)

So the condition $\rho > 0$ is always satisfied in power-law k-inflation.

Equations (37-40) will be our main starting point. We will reconstruct g(X) of polynomial form

$$g(X) = c_1 X + c_2 X^2 + c_3 X^3 + c_4 X^4$$
(43)

in subsection 5.1, and that of DBI-like plus constant form

$$g(X) = -(c_0 + c_1 X + c_2 X^2)^{\frac{1}{2}} + c_3$$
(44)

in subsection 5.2. Subsection 5.3 will concern the DBI-like form

$$g(X) = -(c_0 + c_1 X + c_2 X^2 + c_3 X^3)^{\frac{1}{2}}.$$
(45)

5.1 Power-Law Model *I*: $p = \frac{1}{\phi^2}(c_1X + c_2X^2 + c_3X^3 + c_4X^4)$

For polynomial form (43), the Lagrangian

$$p = \frac{1}{\phi^2} (c_1 X + c_2 X^2 + c_3 X^3 + c_4 X^4).$$
(46)

By substituting (43) into (37-40), one immediately gets

$$c_{1} = -\frac{\xi_{4}}{6\xi_{1}} (\xi_{1}\xi_{2}\xi_{3} - 6\xi_{1}\xi_{2} + 18\xi_{1} - 24),$$

$$c_{2} = \frac{\xi_{4}}{2X\xi_{1}} (\xi_{1}\xi_{2}\xi_{3} - 5\xi_{1}\xi_{2} + 12\xi_{1} - 12),$$

$$c_{3} = -\frac{\xi_{4}}{2X^{2}\xi_{1}} (\xi_{1}\xi_{2}\xi_{3} - 4\xi_{1}\xi_{2} + 8\xi_{1} - 8),$$

$$c_{4} = \frac{\xi_{4}}{6X^{3}\xi_{1}} (\xi_{1}\xi_{2}\xi_{3} - 3\xi_{1}\xi_{2} + 6\xi_{1} - 6).$$
(47)

During the reconstruction, $\xi_1, \xi_2, \xi_3, \xi_4$ may be traded for ϵ , c_s and f_{NL}^{equil} by (37-40), while the value of X is put by hand, contingent on the scale of ϕ as will be shown in (53). Please note here c_1 is independent of X.

In principle, the Lagrangian (46) can be reckoned as the Lagrangian of a massless scalar with higher order corrections. To see this, we have to redefine the scalar field as

$$\tilde{\phi} = \sqrt{c_1} \ln \frac{\phi}{\phi_0},\tag{48}$$

whose first order kinetic term

$$\tilde{X} = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\tilde{\phi}\partial_{\nu}\tilde{\phi} = \frac{c_1X}{\phi^2}$$
(49)

is canonical as we intend to show now. In terms of $\tilde{\phi}$ and \tilde{X} , the Lagrangian (46) takes a "nearly canonical" form

$$p = \frac{1}{\phi^2} (c_1 X + c_2 X^2 + c_3 X^3 + c_4 X^4)$$

$$= \tilde{X} + \frac{c_2}{c_1^2} \phi^2 \tilde{X}^2 + \frac{c_3}{c_1^3} \phi^4 \tilde{X}^3 + \frac{c_4}{c_1^4} \phi^6 \tilde{X}^4$$

$$= \tilde{X} + \tilde{c}_2 \exp\left(\frac{2\tilde{\phi}}{\sqrt{c_1}}\right) \tilde{X}^2 + \tilde{c}_3 \exp\left(\frac{4\tilde{\phi}}{\sqrt{c_1}}\right) \tilde{X}^3 + \tilde{c}_4 \exp\left(\frac{6\tilde{\phi}}{\sqrt{c_1}}\right) \tilde{X}^4$$

$$= M_s^4 \left(\frac{\tilde{X}}{M_s^4} + c_2^* \frac{\tilde{X}^2}{M_s^8} + c_3^* \frac{\tilde{X}^3}{M_s^{12}} + c_4^* \frac{\tilde{X}^4}{M_s^{16}}\right).$$
(50)

Using (6), (9), (10) and (32), (42), one can confirm that

$$P^{\zeta} = \frac{4X}{r\pi^2 \epsilon^2 \phi^2}.$$
(51)

Through this relation, given a normalization of power spectrum, the scale of \tilde{X} is dictated by

$$\tilde{X} = \frac{c_1 r \pi^2 \epsilon^2}{4} P^{\zeta},\tag{52}$$

while the scale of X depends on ϕ_0 and $\tilde{\phi}$ as

$$X = \frac{r\pi^2 \epsilon^2}{4} P^{\zeta} \phi_0^2 \exp\left(\frac{2\tilde{\phi}}{\sqrt{c_1}}\right).$$
(53)

In the rest of this subsection, we will set the "canonical" scalar field $\tilde{\phi} \simeq 0.01$, which is of the same order as $\tilde{X}^{\frac{1}{4}}$ approximately. Since $\sqrt{c_1} \gg 1$ in (53), our result will not change significantly if the scale of $\tilde{\phi}$ is lowered down. That is to say, with $\tilde{\phi}$ at a sub-Planckian energy scale, X is determined by ϕ_0 but insensitive to $\tilde{\phi}$. We will not fix X or ϕ_0 here, since they do not appear in our final result.

We choose parameters

$$P^{\zeta} = 2.5 \times 10^{-9}, \quad n_s \simeq 0.97, \quad f_{NL}^{equil} \simeq 100, \quad r \simeq 0.1,$$
 (54)

and set

$$\tilde{\phi} \simeq 0.01, \quad \frac{M_s^4}{M_{pl}^4} \simeq 5 \times 10^{-9},$$
(55)

from (47), (48), (52) and (53), we can reconstruct the coefficients in Lagrangian (50) as

$$c_{1} \simeq 4.6 \times 10^{4}, \quad c_{2} \simeq -2.1 \times 10^{18} \phi_{0}^{-2},$$

$$c_{3} \simeq 1.8 \times 10^{31} \phi_{0}^{-4}, \quad c_{4} \simeq -4.7 \times 10^{43} \phi_{0}^{-6},$$

$$\tilde{c}_{2} \simeq -1.0 \times 10^{9}, \quad \tilde{c}_{3} \simeq 1.9 \times 10^{17}, \quad \tilde{c}_{4} \simeq -1.1 \times 10^{25},$$

$$c_{2}^{*} \simeq -5.1, \quad c_{3}^{*} \simeq 4.8, \quad c_{4}^{*} \simeq -1.3,$$
(56)

and

$$\phi \simeq \phi_0, \quad X \simeq 1.4 \times 10^{-13} \phi_0^2, \quad \tilde{X} \simeq 6.3 \times 10^{-9} M_{pl}^4 \simeq 1.3 M_s^4.$$
 (57)

Note that selecting a value of X is equivalent to choosing a normalization of ϕ . In the last line of (50), we have tuned c_2^*, c_3^*, c_4^* to be of order unity by choosing the string energy scale M_s . The first term in (50) recovers a canonical form apparently. One should also note that contributions from each term in (50) are comparable. This is reasonable for the Lagrangian of a scalar field with higher order corrections.

From (10) and (33), one can show that for this set of parameters, ϵ and c_s^2 take the values

$$\epsilon \simeq 0.015, \quad c_s^2 \simeq 0.174. \tag{58}$$

So we are sure that for the choice of parameters (54), the $\mathcal{O}(\frac{\epsilon}{c_s^2})$ contribution to f_{NL}^{equil} is indeed negligible in (35).

As we have mentioned, by relations (10) and (33), the parameters ϵ and c_s can be translated into n_s and r, which are experimentally constrained. The WMAP3 data alone [2] gives

$$n_s = 0.958 \pm 0.016 \text{ (at } 68\% \text{ C.L.)}, \quad r_{0.002} < 0.65 \text{ (at } 95\% \text{ C.L.)}.$$
 (59)

Therefore, more generally, given a reasonable value of n_s , we can plot the ratio $\frac{\epsilon}{c_s^2}$ as a function of r,

$$\frac{\epsilon}{c_s^2} = \frac{256\epsilon^3}{r^2} = \frac{32(1-n_s)^3}{r^2}.$$
(60)

This has been done using solid blue lines in figure 1. The figure tells us that to neglect the $\mathcal{O}(\frac{\epsilon}{c_s^2})$ contribution we should treat our power-law models in large r region (typically $r \gtrsim \mathcal{O}(10^{-2})$), otherwise $\mathcal{O}(\frac{\epsilon}{c_s^2})$ terms will be involved.



Figure 1: The (logarithmic) ratio $\log_{10}(\frac{\epsilon}{c_s^2})$ as a function of r. The solid blue lines are plotted according to relation (60). We have set $n_s \simeq 0.97$ in the left plot, and $n_s \simeq 0.96$ in the right one. This figure is valid for all of the power-law models considered in section 5. The dashed red lines are used to highlight the black points corresponding to $c_s = 1$, since we have the additional constraint $c_s^2 > 1$ for power-law model II in subsection 5.2.



Figure 2: The allowed f_{NL} -r region (blue shaded) for nearly canonical powerlaw model *I*. This figure is plotted to depict the condition (61).

One should be aware that we choose $n_s = 0.97$ instead of the best-fit value $n_s = 0.96$ in this section. This choice will be essential in subsection 5.2, because models with $n_s = 0.96$ and r < 0.3 can not be constructed in that subsection. In subsections 5.1 and 5.3, models with $n_s = 0.96$ can nevertheless be constructed as well.

If we assume $r \simeq 0.3$ in (54), we are not fortunate enough to get a reasonable model here, because the reconstruction gives values of ϕ with non-vanishing imaginary parts. This indicates terms like $c_5 X^5$ will be important for large r.

Let us see it in more details. In fact, the definition (48) implies $c_1 > 0$, namely

$$f_{NL}^{equil} \gtrsim -0.059 - \frac{0.015}{r^2} + 7103r^2$$
 (61)

assuming the same choice of parameters P^{ζ} and n_s as in (54). Clearly this condition is broken down when $f_{NL}^{equil} \lesssim 100$ and $r \simeq 0.3$. The allowed region for f_{NL} as a function of r is shaded in figure 2 according to this condition.

For large r, although it is difficult to reconstruct the nearly canonical Lagrangian (50), such a model may be available if we introduce a $c_5 X^5$ term. In deed, when this term is introduced, the condition (61) will be relaxed, with the form

$$f_{NL}^{equil} \gtrsim -0.059 - \frac{0.015}{r^2} + 7103r^2 - 2.5 \times 10^{-49} c_5 \phi^8 r^6, \tag{62}$$

which is not hard to meet with reasonable values of parameters even if r is large.

Note that we introduced a mass scale M_s for this class of models again. All the coefficients look reasonable in this scale, and the scalar field itself is in the sub-Planckian regime. This tells us that this class of model as a candidate for explaining a large f_{NL} is attractive. Note also the highest term X^4 has a negative coefficient, once again signaling instability. This may be a general feature of all models accommodating a large positive non-Gaussianity. Numerically, all the terms in p are comparable in magnitude.

5.2 Power-Law Model *II*: $p = \frac{1}{\phi^2} (-\sqrt{c_0 + c_1 X + c_2 X^2} + c_3)$

For the DBI-like plus constant form

$$p = \frac{1}{\phi^2} \left(-\sqrt{c_0 + c_1 X + c_2 X^2} + c_3\right),\tag{63}$$

a similar computation gives

$$c_{0} = \frac{X^{2}\xi_{4}^{2}}{\xi_{3}^{2}}(\xi_{3}^{2} - 3\xi_{2}\xi_{3} + 6\xi_{3} + 9),$$

$$c_{1} = \frac{2X\xi_{4}^{2}}{\xi_{3}}(3\xi_{2} - \xi_{3} - 3),$$

$$c_{2} = \frac{\xi_{4}^{2}}{\xi_{3}}(\xi_{3} - 3\xi_{2}),$$

$$c_{3} = \frac{X\xi_{4}}{\xi_{3}}\left(\frac{\xi_{3}}{\xi_{1}} + 3\right),$$
(64)

and

$$\sqrt{c_0 + c_1 X + c_2 X^2} = \frac{3\xi_4 X}{\xi_3} = \frac{6X}{\epsilon\xi_3}.$$
(65)

Since we concentrate on the case $\epsilon \geq 0$, the result (65) tells us that $\xi_3 > 0$. According to this requirement and $f_{NL}^{equil} \gg 1$, the definition of ξ_3 in (38) suggests $c_s^2 > 1$ in this model. According to (10) and (33), this bound implies a constraint on this model

$$c_s = \frac{r}{8(1 - n_s)} > 1 \tag{66}$$

if $n_s < 1$ and $f_{NL}^{equil} \gg 1$.

Now let us try to recast the model into a more realistic form

$$p = \frac{1}{\phi^{2}} \left(-\sqrt{c_{0} + c_{1}X + c_{2}X^{2}} + c_{3} \right)$$

$$= -\frac{\sqrt{c_{0}}}{\phi^{2}} \sqrt{1 - \frac{2}{\sqrt{c_{0}}} \phi^{2} \tilde{X} + \frac{4c_{2}}{c_{1}^{2}} \phi^{4} \tilde{X}^{2}} + \frac{c_{3}}{\phi^{2}}$$

$$= -\tilde{c}_{0} \exp(-2\alpha\tilde{\phi}) \sqrt{1 - \frac{2}{\tilde{c}_{0}}} \exp(2\alpha\tilde{\phi}) \tilde{X} + \tilde{c}_{2} \exp(4\alpha\tilde{\phi}) \tilde{X}^{2} + \tilde{c}_{3} \exp(-2\alpha\tilde{\phi})$$

$$= M_{s}^{4} \left(-c_{0}^{*} \sqrt{1 - \frac{2}{c_{0}^{*}} \frac{\tilde{X}}{M_{s}^{4}} + c_{2}^{*} \frac{\tilde{X}^{2}}{M_{s}^{8}}} + c_{3}^{*} \right), \qquad (67)$$

which to the first order recovers a DBI action. This may be accomplished by introducing

$$\tilde{\phi} = \sqrt{\frac{-c_1}{2\sqrt{c_0}}} \ln \frac{\phi}{\phi_0} = \frac{1}{\alpha} \ln \frac{\phi}{\phi_0}, \quad \alpha = \sqrt{\frac{2\sqrt{c_0}}{-c_1}},\tag{68}$$

and correspondingly

$$\tilde{X} = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\tilde{\phi}\partial_{\nu}\tilde{\phi} = \frac{-c_1X}{2\sqrt{c_0}\phi^2}.$$
(69)

Comparing (68), (69) with (51), one quickly writes down

$$\tilde{X} = -\frac{c_1 r \pi^2 \epsilon^2}{8\sqrt{c_0}} P^{\zeta},$$

$$X = \frac{r \pi^2 \epsilon^2}{4} P^{\zeta} \phi_0^2 \exp(2\alpha \tilde{\phi}).$$
(70)

Again with a sub-Planckian inflaton $\tilde{\phi}$, we have $\alpha \tilde{\phi} \ll 1$ and thus X is insensitive to the scale of $\tilde{\phi}$. In the following we will assume the "canonical" scalar field and $\tilde{X}^{\frac{1}{4}}$ are roughly of the same order, *e.g.* $\tilde{\phi} \simeq 10^{-3}$, but we will leave X and ϕ_0 undetermined.

This time the condition $c_s^2 > 1$ put a tighter constraint on r, see inequality (66) and figure 1. In each plot of the figure, this condition is satisfied only for the solid blue line below the black dot. So we choose $r \simeq 0.3$. The other input parameters of reconstruction are chosen to be the same as in (54). In addition we take

$$\tilde{\phi} \simeq 10^{-3}, \quad \frac{M_s^4}{M_{pl}^4} \simeq 5 \times 10^{-11}.$$
 (71)

With (64), (68) and (70) at hand, we can perform a computation similar to the previous subsection, resulted in coefficients

$$c_{0} \simeq 3.1 \times 10^{-21} \phi_{0}^{4}, \quad c_{1} \simeq -1.5 \times 10^{-8} \phi_{0}^{2},$$

$$c_{2} \simeq 1.8 \times 10^{4}, \quad c_{3} \simeq -1.1 \times 10^{-8} \phi_{0}^{2},$$

$$\tilde{c}_{0} \simeq 5.6 \times 10^{-11}, \quad \tilde{c}_{2} \simeq 3.2 \times 10^{20}, \quad \tilde{c}_{3} \simeq -1.1 \times 10^{-8},$$

$$c_{0}^{*} \simeq 1.1, \quad c_{2}^{*} \simeq 0.81, \quad c_{3}^{*} \simeq -220$$
(72)

in Lagrangian (67), and other quantities

$$\phi \simeq \phi_0, \quad X \simeq 4.2 \times 10^{-13} \phi_0^2, \quad \tilde{X} \simeq 5.55 \times 10^{-11} M_{pl}^4 \simeq 1.1 M_s^4,$$

$$\epsilon = 0.015, \quad c_s^2 = 1.56. \tag{73}$$

Hence this class of models can be considered as a DBI action with higher order corrections.

Note that the coefficient c_1 has the desired sign, while the sign of c_2 is positive and this leads to the kinetic energy unbounded from below when X is large. The last term c_3/ϕ^2 dominates the Lagrangian numerically, but other terms are important to drive the inflation with appropriate spectral indices, otherwise we can only have a de Sitter space.

5.3 Power-Law Model III: $p = -\frac{1}{\phi^2}\sqrt{c_0 + c_1 X + c_2 X^2 + c_3 X^3}$

The Lagrangian for the DBI-like form (45) is

$$p = -\frac{1}{\phi^2}\sqrt{c_0 + c_1 X + c_2 X^2 + c_3 X^3}.$$
(74)

We still start from (37-40). After some calculations, we obtain

$$c_{0} = X^{2}\xi_{4}^{2}(1-\xi_{2}) - \frac{X^{2}\xi_{4}^{2}}{3\xi_{1}^{2}}(\xi_{1}\xi_{2}\xi_{3} - 3\xi_{1}\xi_{2} + 6\xi_{1} - 3),$$

$$c_{1} = X\xi_{4}^{2}(3\xi_{2} - 2) + \frac{X\xi_{4}^{2}}{\xi_{1}}(\xi_{2}\xi_{3} - 2\xi_{2} + 2),$$

$$c_{2} = \xi_{4}^{2}(1-3\xi_{2}) + \frac{\xi_{2}\xi_{4}^{2}}{\xi_{1}}(1-\xi_{3}),$$

$$c_{3} = \frac{\xi_{2}\xi_{4}^{2}}{3X\xi_{1}}(3\xi_{1} + \xi_{3}),$$
(75)

and

$$\sqrt{c_0 + c_1 X + c_2 X^2 + c_3 X^3} = -\frac{3X\xi_4}{\xi_1} = \frac{2X(3 - 2\epsilon)}{\epsilon^2}.$$
(76)

This model is all right if we do not demand the kinetic term be canonical, with the value of X or ϕ to be put by hand. For instance, if we take $\phi \simeq 0.1$ and (54), by using equations (51) and (75), we can get

$$c_0 \simeq -1.4 \times 10^{-20}, \quad c_1 \simeq 3.3 \times 10^{-5}, \quad c_2 \simeq -2.4 \times 10^{10}, \quad c_3 \simeq 5.8 \times 10^{24},$$

 $X \simeq 1.4 \times 10^{-15}, \quad \epsilon = 0.015, \quad c_s^2 = 0.174.$
(77)

This naive model gives $f_{NL}^{equil} \simeq 100$ as we desired.

Now turn to reconstruction of canonical models with Lagrangian (74). Different from the previous subsection, for this class of models, without introducing more terms, we cannot recover DBI action to the first order. The key point is as follows.

Formally, employing (68-70), one may rewrite Lagrangian (74) as

$$p = -\tilde{c}_0 \exp(-2\alpha\tilde{\phi}) \sqrt{1 - \frac{2}{\tilde{c}_0} \exp(2\alpha\tilde{\phi})\tilde{X} + \tilde{c}_2 \exp(4\alpha\tilde{\phi})\tilde{X}^2 + \tilde{c}_3 \exp(6\alpha\tilde{\phi})\tilde{X}^3}.$$
 (78)

However, the validity of (68) indicates $c_1 < 0$, namely

$$f_{NL}^{equil} \lesssim 0.11 - \frac{0.015}{r^2} + 5.6r^2$$
 (79)

when we make the same choice of parameters P^{ζ} and n_s as in (54). Clearly it is impossible to arrange parameters obeying the condition (79) with $f_{NL}^{equil} \gg 1$ and r < 1. In other words, even by tuning M_s and the normalization of ϕ , it is impossible to rewrite (74) in the form of a conventional DBI action with higher order corrections.

Nevertheless, we can still construct other models of the form (74), whose leading order kinetic term is not canonical when expanded. If we consider that class of models which are not "nearly canonical", the model (77) is able to reproduce the desired f_{NL} .

If one prefers being restricted to "nearly canonical" models, another $c_4 X^4$ term will lend a hand, just as what happened in subsection 5.1. Actually, when we take roughly $c_4 \simeq 10^{36} \phi^{-4}$ and choose the assumption (54), a model

$$p = -\frac{1}{\phi^2} \sqrt{c_0 + c_1 X + c_2 X^2 + c_3 X^3 + c_4 X^4}$$

$$= -\tilde{c}_0 e^{-2\alpha\tilde{\phi}} \sqrt{1 - \frac{2}{\tilde{c}_0} e^{2\alpha\tilde{\phi}} \tilde{X} + \tilde{c}_2 e^{4\alpha\tilde{\phi}} \tilde{X}^2 + \tilde{c}_3 e^{6\alpha\tilde{\phi}} \tilde{X}^3 + \tilde{c}_4 e^{8\alpha\tilde{\phi}} \tilde{X}^4}$$

$$= -c_0^* M_s^4 \sqrt{1 - \frac{2}{c_0^*} \frac{\tilde{X}}{M_s^4} + c_2^* \frac{\tilde{X}^2}{M_s^8} + c_3^* \frac{\tilde{X}^3}{M_s^{12}} + c_4^* \frac{\tilde{X}^4}{M_s^{16}}}$$
(80)

is constructed, with the main results

$$c_{0} \simeq 2.3 \times 10^{-16} \phi_{0}^{4}, \quad c_{1} \simeq -7.4 \times 10^{-3} \phi_{0}^{2},$$

$$c_{2} \simeq 9.2 \times 10^{10}, \quad c_{3} \simeq -5.0 \times 10^{23} \phi_{0}^{-2}, \quad c_{4} \simeq 10^{36} \phi_{0}^{-4},$$

$$\tilde{c}_{0} \simeq 1.5 \times 10^{-8}, \quad \tilde{c}_{2} \simeq 6.75 \times 10^{15}, \quad \tilde{c}_{3} \simeq -1.5 \times 10^{23}, \quad \tilde{c}_{4} \simeq 1.3 \times 10^{30},$$

$$c_{0}^{*} \simeq 0.51, \quad c_{2}^{*} \simeq 6.1, \quad c_{3}^{*} \simeq -4.1, \quad c_{4}^{*} \simeq 1.0$$
(81)

and

$$\begin{split} \tilde{\phi} &\simeq 0.01, \quad \frac{M_s^4}{M_{pl}^4} \simeq 3 \times 10^{-8}, \quad \alpha \simeq 2.03 \times 10^{-3}, \\ \phi &\simeq \phi_0, \quad X \simeq 1.4 \times 10^{-13} \phi_0^2, \quad \tilde{X} \simeq 3.4 \times 10^{-8} M_{pl}^4 \simeq 1.1 M_s^4, \\ \epsilon &= 0.015, \quad c_s^2 = 0.174. \end{split}$$
(82)

This model is more complicated than (74), but it is canonical when expanded in terms of \tilde{X} to the first order. What is more, it does not suffer from trans-Planckian effects, because the inflaton $\tilde{\phi}$ is well below the Planck scale.

6 Conclusion

In this paper, we have constructed several models with a large positive f_{NL} in the WMAP convention [1, 2]. These models are general single field inflation with higher

order kinetic terms. In one class of models, the inflation is driven by the potential term of the inflaton, as shown in section 4. In section 5, we have given another class of models, in which inflation is driven by non-conventional kinetic terms [13, 14]. In both classes of models, due to the appearance of non-conventional kinetic terms, we can arrange the parameters to produce $f_{NL}^{equil} \gg 1$. These are first examples to generate $f_{NL}^{equil} \gg 1$ in general single field k-inflation. They are different from ghost inflation [15] since we restricted our discussion in k-inflation and did not introduce terms like $\nabla^2 \phi$ in our Lagrangian.

The common features of these models are as following:

- 1. Typically we need to introduce four parameters in a model, although there are only three data to fit, the COBE normalization of the two point function, the spectral index n_s and the non-Gaussianity parameter f_{NL} . Introduction of four parameters is not strictly necessary. For example, for models studied in section 4, we introduced two parameters α and β in order to have a model easy to solve. In section 5, we introduced four parameters c_i since we try to also fit a parameter r whose value is not fixed by experiments yet. However, a upper bound on r is set, so it is necessary to make sure that our model does not violate this bound.
- 2. High order terms in X are absolutely necessary, and luckily these terms can be viewed as operators of high dimensions in an effective field theory with a mass cut-off lower than the Planck scale. The bad news is that, the highest order term in general triggers an instability if we do not introduce more high order terms.

Acknowledgments

We thank Qing-Guo Huang for discussions. We are grateful to KITPC for a fruitful workshop on string cosmology, during which this work was initiated. We thank all the speakers on the topic of non-Gaussianity for the education, especially Komatsu for his informative talk. We thank Henry Tye especially, for his emphasis on the importance of the non-Gaussianity in discriminating among inflation models. This work was supported by grants from NSFC, a grant from Chinese Academy of Sciences and a grant from USTC.

A Calculation of $R^{equil}(k)$

In the equilateral triangle case, the function $R(k_1, k_2, k_3)$ defined in [4] takes the form

$$R^{equil}(k) = R(k_1, k_2, k_3)|_{k_1 = k_2 = k_3 = k}$$

= $3k^3 \operatorname{Re}\left[\int_0^\infty (1 - 2ix)e^{-2ix}h^*(x)dx\right]$ (83)

with

$$h(x) = -2ie^{ix} + ie^{-ix}(1+ix)[\operatorname{Ci}(2x) + i\operatorname{Si}(2x)] - i\pi\sin x + i\pi x\cos x.$$
(84)

With the help of the relation

$$Ci(2x) - iSi(2x) = -\int_{2x}^{\infty} \frac{\cos t dt}{t} - i \int_{0}^{2x} \frac{\sin t dt}{t}$$
$$= -\int_{2x}^{\infty} \frac{e^{-it} dt}{t} - \frac{i\pi}{2}$$
(85)

and the equality

$$\int_{-2ix}^{\infty} \frac{e^{-it}dt}{t} = \int_{2x}^{\infty} \frac{e^{-u}du}{u} \quad \text{for } x \ge 0,$$
(86)

after an analytical continuation $x \to -ix$, we can transform it into

$$R^{equil}(k) = 3k^{3} \operatorname{Re} \left\{ -i \int_{0}^{\infty} (1 - 2x) e^{-2x} \times \left[\left(2i - \frac{\pi}{2} (1 + x) \right) e^{-x} + i(1 - x) e^{x} \left(\int_{-2ix}^{\infty} \frac{e^{-it} dt}{t} \right) \right] dx \right\}$$

$$= 3k^{3} \int_{0}^{\infty} 2(1 - 2x) e^{-3x} dx$$

$$+ 3k^{3} \int_{0}^{\infty} (1 - x)(1 - 2x) e^{-x} \left(\int_{2x}^{\infty} \frac{e^{-u} du}{u} \right) dx.$$
(87)

The first term can be easily evaluated with a result $\frac{2}{3}k^3$. To calculate the second term, let us note that

$$\int_{0}^{\infty} \dots dx = \lim_{\epsilon \to 0^{+}} \int_{\epsilon}^{\infty} \dots dx$$
(88)

and consider the integral

$$\lim_{\epsilon \to 0^{+}} \left[\int_{\epsilon}^{\infty} (1-x)(1-2x)e^{-x} \left(\int_{2x}^{\infty} \frac{e^{-u}du}{u} \right) dx \right] \\
= \lim_{\epsilon \to 0^{+}} \left[(2+\epsilon+2\epsilon^{2})e^{-\epsilon} \int_{2\epsilon}^{\infty} \frac{e^{-u}du}{u} - 2 \int_{\epsilon}^{\infty} \frac{e^{-3x}dx}{x} - \left(\frac{5}{9} + \frac{2}{3}\epsilon\right) e^{-3\epsilon} \right] \\
= \lim_{\epsilon \to 0^{+}} \left(2 \int_{2\epsilon}^{\infty} \frac{e^{-u}du}{u} - 2 \int_{\epsilon}^{\infty} \frac{e^{-3x}dx}{x} - \frac{5}{9} \right) \\
= -\frac{5}{9} + \lim_{\epsilon \to 0^{+}} \left(2 \int_{2\epsilon}^{\infty} \frac{e^{-u}du}{u} - 2 \int_{3\epsilon}^{\infty} \frac{e^{-u}du}{u} \right) \\
= -\frac{5}{9} + \lim_{\epsilon \to 0^{+}} 2 \int_{2\epsilon}^{3\epsilon} \frac{(1-u)du}{u} \\
= -\frac{5}{9} + 2\ln\frac{3}{2}.$$
(89)

Here we have integrated by parts and used the equalities

$$\frac{d}{dx}\int_{2x}^{\infty}\frac{e^{-u}du}{u} = -\frac{e^{-2x}}{x}.$$

As the last result,

$$R^{equil}(k) = \frac{2}{3}k^3 + 3k^3\left(-\frac{5}{9} + 2\ln\frac{3}{2}\right) = \left(-1 + 6\ln\frac{3}{2}\right)k^3 \tag{90}$$

References

- [1] E. Komatsu and D. N. Spergel, Phys. Rev. D 63, 063002 (2001)
 [arXiv:astro-ph/0005036].
- [2] D. N. Spergel et al. [WMAP Collaboration], Astrophys. J. Suppl. 170, 377 (2007)
 [arXiv:astro-ph/0603449].
- [3] J. M. Maldacena, JHEP **0305**, 013 (2003) [arXiv:astro-ph/0210603].
- [4] X. Chen, M. X. Huang, S. Kachru and G. Shiu, JCAP 0701, 002 (2007)
 [arXiv:hep-th/0605045].
- [5] E. Komatsu, arXiv:astro-ph/0206039.
- [6] E. Komatsu, 2007, talk at String Theory and Cosmology Program in KITPC, http://www.kitpc.ac.cn/Activities/program/63574646/fnl.pdf

- [7] A. P. S. Yadav and B. D. Wandelt, arXiv:0712.1148 [astro-ph].
- [8] A. P. S. Yadav, E. Komatsu, B. D. Wandelt, M. Liguori, F. K. Hansen and S. Matarrese, arXiv:0711.4933 [astro-ph].
- [9] L. Alabidi and D. H. Lyth, JCAP **0608**, 013 (2006) [arXiv:astro-ph/0603539].
- [10] V. Acquaviva, N. Bartolo, S. Matarrese and A. Riotto, Nucl. Phys. B 667, 119 (2003) [arXiv:astro-ph/0209156].
- [11] A. Gangui, F. Lucchin, S. Matarrese and S. Mollerach, Astrophys. J. 430, 447 (1994) [arXiv:astro-ph/9312033].
- [12] D. H. Lyth, C. Ungarelli and D. Wands, Phys. Rev. D 67, 023503 (2003)
 [arXiv:astro-ph/0208055].
- [13] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, Phys. Lett. B 458, 209 (1999) [arXiv:hep-th/9904075].
- [14] J. Garriga and V. F. Mukhanov, Phys. Lett. B 458, 219 (1999)
 [arXiv:hep-th/9904176].
- [15] N. Arkani-Hamed, P. Creminelli, S. Mukohyama and M. Zaldarriaga, JCAP 0404, 001 (2004) [arXiv:hep-th/0312100].
- [16] E. Silverstein and D. Tong, Phys. Rev. D 70, 103505 (2004)
 [arXiv:hep-th/0310221].
- [17] M. Alishahiha, E. Silverstein and D. Tong, Phys. Rev. D 70, 123505 (2004)
 [arXiv:hep-th/0404084].
- [18] X. Chen, Phys. Rev. D 71, 063506 (2005) [arXiv:hep-th/0408084].
- [19] X. Chen, JHEP **0508**, 045 (2005) [arXiv:hep-th/0501184].
- [20] S. Gupta, A. Berera, A. F. Heavens and S. Matarrese, Phys. Rev. D 66, 043510 (2002) [arXiv:astro-ph/0205152]. I. G. Moss and C. Xiong, JCAP 0704, 007 (2007) [arXiv:astro-ph/0701302]. I. G. Moss and C. M. Graham, JCAP 0711, 004 (2007) [arXiv:0707.1647 [astro-ph]].

- [21] D. Seery and J. E. Lidsey, JCAP 0506, 003 (2005) [arXiv:astro-ph/0503692].
- [22] V. F. Mukhanov and A. Vikman, JCAP 0602, 004 (2006) [arXiv:astro-ph/0512066].
- [23] E. Babichev, V. Mukhanov and A. Vikman, arXiv:0708.0561 [hep-th].
- [24] Q. G. Huang, JHEP 0705, 096 (2007) [arXiv:hep-th/0703071].
- [25] Q. G. Huang, Phys. Rev. D 76, 061303 (2007) [arXiv:0706.2215 [hep-th]].