

NON-EXISTENCE OF ABSOLUTELY CONTINUOUS INVARIANT PROBABILITIES FOR EXPONENTIAL MAPS

NEIL DOBBS AND BARTŁOMIEJ SKORULSKI

ABSTRACT. We show that for entire maps of the form $z \mapsto \lambda \exp(z)$ such that the orbit of zero is bounded and such that Lebesgue almost every point is transitive, no absolutely continuous invariant probability measure can exist. This answers a long-standing open problem.

In this paper we introduce a new method to deal with the problem of existence of invariant measures for entire maps. To illustrate this method, avoiding uninteresting technical difficulties, we show the following theorem.

Theorem 1. *Let $\lambda \in \mathbb{C} \setminus \{0\}$ be such that the Julia set of $f : z \mapsto \lambda \exp(z)$ is equal to \mathbb{C} , the forward orbit of 0 is bounded and such that there is a set of positive Lebesgue measure of points $z \in \mathbb{C}$ such that $\omega(z) \not\subset \mathcal{P}(f)$. Then f has a σ -finite absolutely continuous invariant measure, but it does not have an absolutely continuous invariant probability measure.*

We denote, as usual, the ω -limit set of $z \in \mathbb{C}$ by $\omega(z) \subset \overline{\mathbb{C}}$ and the post-singular set by $\mathcal{P}(f)$, here equal to the union of $\{\infty\}$ with the closure of the orbit of 0.

Theorem 1 implies, in particular, that the map $z \mapsto 2\pi i \exp(z)$ has no absolutely continuous invariant probability measure, which was a long-standing open problem (see [9]).

For a class of unimodal maps f of the interval with non-recurrent critical point, Benedicks and Misiurewicz in [1] showed that there exists an absolutely continuous invariant probability measure if and only if $\int \log |f'(x)| dx > -\infty$. The necessity of the integrability condition was later extended by the first author (in Theorem 1 of [3]) to all $C^{1+\epsilon}$ interval maps without any hypothesis on the critical orbits, but under the assumption that the measure has positive Lyapunov exponent. As an example, no unimodal map of the form $x \mapsto C_1 - C_2 \exp(-|x|^{-\alpha})$ with $\alpha \geq 1$ has an absolutely continuous invariant probability measure with positive Lyapunov exponent.

This paper extends the main result of [1] to the holomorphic setting. The strategy of the proof has two elements: construct a *nice set* on which the density of a hypothetical measure must be bounded away from zero; show that the return time to the nice set is not integrable. This is similar in philosophy to the proof of Benedicks and Misiurewicz, although the settings differ.

Date: January 30, 2008.

The authors were partially supported by Research Network on Low Dimensional Dynamics, PBCT ACT 17, CONICYT, Chile and by the EU Research Training Network “Conformal Structures and Dynamics”. The second author was also supported by Chilean FONDECYT Grant No. 11060538.

Existence of a σ -finite measure is not new; it was shown in [7] under weaker hypotheses but with a considerably more difficult proof. On the other hand, existence of absolutely continuous invariant probability measures for transcendental entire maps has been an interesting and open question for some time with a response in only one situation: J-M Hemke in [5] proved that, for a class of entire maps for which the orbits of all asymptotic values converge to infinity sufficiently fast, $\omega(z) \subset \mathcal{P}(f)$. For these maps, $\mathcal{P}(f)$ has zero measure and it follows from the Poincaré Recurrence Theorem that no absolutely continuous invariant probability measure can exist. Hemke's work generalised a result proved independently by M. Rees and M. Lyubich for $z \mapsto \exp(z)$ ([11], [8]).

For non-entire maps the second author, in [13], has another negative result for some postcritically finite tangent maps. For a large class of transcendental non-entire maps which satisfy a Misiurewicz-type condition J. Kotus and G. Świątek in [6] showed that absolutely continuous invariant probability measures can exist.

The mathematics involved in the proof have the merit of being surprisingly elementary. An important and somewhat magical technique is Juan Rivera-Letelier's construction of nice sets for rational dynamics (see [12]) which we adapt to the entire setting.

An open set U is called *nice* if $f^n(\partial U) \cap U = \emptyset$ for all $n > 0$. This implies that every pair of pullbacks (connected components of $f^{-n}(U), f^{-n'}(U)$ for some $n, n' \geq 0$) is either nested or disjoint. Let us fix some $D > 0$ such that $\mathcal{P}(f) \subset B(0, D)$.

Lemma 2. *For each sufficiently large $x \in \mathbb{R}$, there exists a connected nice set $U \subset \mathbb{C}$ satisfying $B(x, 4\pi) \subset U \subset B(x, 8\pi)$.*

Proof. There exists a $K > 1$ such that for any r and any holomorphic function g , univalent on $B(x, Kr)$, one has

$$\left| \frac{g'(z)}{g'(z')} \right| \leq 2$$

for all $z, z' \in B(x, r)$, by the Koebe distortion theorem.

Let x satisfy $x > 8K\pi + D$. Let W be a (connected) pullback of $B(x, 8\pi)$ and let $n > 0$ be such that f^n maps W univalently onto $B(x, 8\pi)$. Since $f^n|_W$ extends to map univalently onto $B(x, 8K\pi)$, it follows that the distortion of f^n restricted to W is bounded by 2. Thus there is $r > 0$ such that $B((f^n|_W)^{-1}(x), r) \subset W \subset B((f^n|_W)^{-1}(x), 2r)$. But $B((f^n|_W)^{-1}(x), r)$ must lie in a horizontal strip of height 2π , so we have that $|W| < 4\pi$.

We shall use this to construct nice sets exactly as per [12]. We include the proof for the reader's convenience: Let $U_0 := B(x, 4\pi)$ and define U_n as the connected component of $\bigcup_{i=0}^n f^{-i}(U_0)$ containing U_0 and $U = \bigcup_{n \geq 0} U_n$. We prove by induction that $U_n \subset B(x, 8\pi)$ for all $n \geq 0$. This is clearly true for $n = 0$. So suppose it is true for all $n \leq k$. We must show it holds for $n = k + 1$.

Let X be a connected component of $U_{k+1} \setminus U_0$. Then there is a minimal $m \geq 0$ such that $f^m(z) \in U_0$ for some $z \in X$, and necessarily $m \geq 1$. Consider $f^m(X)$. This set is contained in U_{k+1-m} , and so by hypothesis is contained in $B(x, 8\pi)$. But then X , being connected, is contained in some pullback W with $|W| < 4\pi$. The result follows. \square

Lemma 3. *There exists a $c > 0$ such that if $f^n(z) \notin B(0, 2D)$ then $n > -c \log |z|$.*

Proof. Let $M > 1$ be such that $|f'(z)| < M$ for all $z \in B(0, 2D)$. Suppose $f^n(z) \notin B(0, 2D)$. Then $|f^n(z) - f^n(0)| > D > 1$. This implies that $|z - 0| = |z| > M^{-n}$. Thus $\log |z| > -n \log M$ and $n > (-1/\log M) \log |z|$. \square

In what follows, let U be a nice set given by Lemma 2 for some $x > 8\pi + 2D$, we fix x too. In particular, $U \cap B(0, 2D) = \emptyset$. We denote by $r_U(z)$ the first return time of z to U . Also let $r, \phi \in \mathbb{R}$ such that $\lambda = re^{i\phi}$. \square

Lemma 4. *There exists $C \in \mathbb{R}$ and $c > 0$ with the following property. Suppose $z \in U$ and $\operatorname{Re}(f^k(z)) \leq -K$ for some $0 < k < r_U(z)$ and $K > 0$. Then $r_U(z) > C + cK$.*

Proof. Let c be given by Lemma 3. We have $|f^{k+1}(z)| \leq re^{-K}$. Then the time it takes for $f^{k+1}(z)$ to leave $B(0, 2D)$ is greater than $-c \log(re^{-K}) = -c(-K + \log r)$ by Lemma 3. Take $C := -c \log r$. \square

Lemma 5. *Denote by m Lebesgue measure. Then $\int_U r_U(z) dm = \infty$.*

Proof. Define $h : \mathbb{R} \rightarrow \mathbb{R}$ by $h(y) = (r/2) \exp(y)$ and let

$$S_R := \{z : \operatorname{Re}(z) > x \text{ and } \arg f(z) \in [-\pi/4, \pi/4]\}$$

and

$$S_L := \{z : \operatorname{Re}(z) > x \text{ and } \arg f(z) \in [3\pi/4, 5\pi/4]\}.$$

Note that each connected component of $\{z : \arg f(z) \in [-\pi/4, \pi/4]\}$ is a horizontal strip of height $\pi/2$ and the components are periodic of period $2i\pi$. A similar statement holds for $\{z : \arg f(z) \in [3\pi/4, 5\pi/4]\}$. Let

$$P_n := \{z \in B(x, 4\pi) : f^k(z) \in S_R \text{ for all } 0 \leq k \leq n\},$$

and let $Q_n := P_{n-1} \cap f^{-n}(S_L)$. For $z \in S_R$, $\operatorname{Re}(f(z)) \geq h(\operatorname{Re}(z))$, so by induction, for all $z \in P_n$, $\operatorname{Re}(f^n(z)) \geq h^n(x)$. Then distortion arguments like in [10] give that $m(Q_n)/m(P_n)$ tends to one and that

$$\lim_{n \rightarrow \infty} m(P_n)/m(P_{n+1}) = 1/4.$$

Thus there exists a $\gamma \in (0, 1/4)$ such that for all $n \geq 1$,

$$m(Q_n) \geq \gamma^n.$$

Now for $z \in Q_n$, $\operatorname{Re}(f^{n+1}(z)) < -h^{n+1}(x)$, so we have $r_U(z) > C + ch^{n+1}(x)$ where the constants c, C are given by Lemma 4. But $h^n(x)$ grows with n faster than any exponential so

$$\lim_{n \rightarrow \infty} m(Q_n) \inf\{r_U(z) : z \in Q_n\} = \infty.$$

\square

Proof of Theorem 1. Let ψ denote the first return map to U . Since U is nice and disjoint from $\mathcal{P}(f)$, every connected component of the domain of ψ is mapped univalently onto U by ψ . Moreover the branches of ψ are uniformly extendible, so the Koebe distortion theorem gives a uniform distortion bound for all branches of all iterates of ψ . Note that by Corollary 2 of [2], Lebesgue almost every point has a transitive orbit.

Denote by V_k the set of points from U whose first return time to U is k . By the Folklore Theorem (see for example [4]), there exists a unique absolutely continuous invariant probability ν for ψ and its density is bounded below by some $\varepsilon > 0$. Then

the spread measure $\mu := \sum_{k=1}^{\infty} \sum_{n=0}^{k-1} f_*^n \nu|_{V_k}$ is a σ -finite absolutely continuous invariant measure for f . This gives the easy proof of its existence.

Suppose now μ is an absolutely continuous f -invariant probability measure. By transitivity of Lebesgue almost every point, $\mu(U) > 0$. Then μ is also a finite invariant measure for ψ , since ψ is a first return map. By uniqueness, the density of μ is then bounded from below on U by $\mu(U)\varepsilon > 0$. Thus

$$1 = \int_U r_U(z) d\mu \geq \mu(U)\varepsilon \int_U r_U(z) dm,$$

the first equality being Kac' Lemma. This contradicts Lemma 5, so no absolutely continuous invariant probability measure can exist. \square

Acknowledgements. We would like to thank J. Rivera-Letelier and the referee for helpful suggestions that improved the final version of the paper.

REFERENCES

- [1] Michael Benedicks and Michał Misiurewicz. Absolutely continuous invariant measures for maps with flat tops. *Inst. Hautes Études Sci. Publ. Math.*, (69):203–213, 1989.
- [2] Heinrich Bock. On the dynamics of entire functions on the Julia set. *Results Math.*, 30(1-2):16–20, 1996.
- [3] Neil Dobbs. On cusps and flat tops. 2007, <http://arxiv.org/abs/0801.3815>.
- [4] Sébastien Gouëzel. *Vitesse de décorrélation et théorèmes limites pour les applications non uniformément dilatantes*. PhD thesis, Université de Paris-Sud, 2004.
- [5] Jan-Martin Hemke. Recurrence of entire transcendental functions with simple post-singular sets. *Fund. Math.*, 187(3):255–289, 2005.
- [6] Janina Kotus and Grzegorz Świątek. Invariant measures for meromorphic Misiurewicz maps. *to appear in Math. Proc. Cambridge Philos. Soc.*, 2007.
- [7] Janina Kotus and Mariusz Urbański. Existence of invariant measures for transcendental subexpanding functions. *Math. Z.*, 243(1):25–36, 2003.
- [8] M. Yu. Lyubich. The measurable dynamics of the exponential. *Sibirsk. Mat. Zh.*, 28(5):111–127, 1987.
- [9] V. Mayer and M Urbański. Ergodic properties of sub-hyperbolic functions with polynomial Schwarzian derivative. Preprint 2007, <http://www.math.unt.edu/~urbanski/papers.html>.
- [10] Curt McMullen. Area and Hausdorff dimension of Julia sets of entire functions. *Trans. Amer. Math. Soc.*, 300(1):329–342, 1987.
- [11] Mary Rees. The exponential map is not recurrent. *Math. Z.*, 191(4):593–598, 1986.
- [12] Juan Rivera-Letelier. A connecting lemma for rational maps satisfying a no-growth condition. *Ergodic Theory Dynam. Systems*, 27(2):595–636, 2007.
- [13] B. Skorulski. Non-ergodic maps in the tangent family. *Indag. Math. (N.S.)*, 14(1):103–118, 2003.

INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8 P.O. BOX 21, 00-956 WARSZAWA, POLAND

E-mail address: neil.dobbs@gmail.com

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD CATÓLICA DEL NORTE, AVENIDA ANGAMOS 0610, CASILLA 1280, ANTOFAGASTA, CHILE

E-mail address: bskorulski@ucn.cl