

# Quantum gravitational corrections to the stress-energy tensor around the BTZ black hole

Dawood A. Kothawala\*

*IUCAA, Post Bag 4, Ganeshkhind, Pune 411 007, India.*

S. Shankaranarayanan†

*Institute of Cosmology and Gravitation, University of Portsmouth, Mercantile House, Portsmouth P01 2EG, U.K..*

L. Sriramkumar‡

*Harish-Chandra Research Institute, Chhatnag Road, Jhansi, Allahabad 211 019, India.*

(Dated: May 28, 2019)

Modes emerging out of a collapsing black hole are red-shifted to such an extent that Hawking radiation at future null infinity consists of modes that have energies beyond the Planck scale at past null infinity. This indicates that physics at the Planck scale may modify the spectrum of Hawking radiation and the associated stress-energy tensor of the quantum field. Recently, it has been shown that, the T-duality symmetry of string fluctuations along compact extra dimensions leads to a modification of the standard propagator of point particles in quantum field theory. At low energies, the modified propagator is found to behave as though the spacetime possesses a minimal length, say,  $L_P$ , which we shall assume to be of the order of the Planck length. We utilize the duality approach to evaluate the modified propagator around the BTZ black hole and show that the propagator is finite in the coincident limit. We compute the stress-energy tensor associated with the modified Green's function and briefly comment on the results we have obtained.

PACS numbers: 04.70.Dy, 04.60.Kz, 04.62.+v, 04.60.-m

## I. WHY DO WE NEED TO CONSIDER PLANCK SCALE PHYSICS?

Hawking radiation primarily arises due to the asymmetry in the extent of the redshift and the blueshift of the modes of a quantum field as they propagate through matter that is collapsing gravitationally and, eventually, goes on to become a black hole [1]. Consider a typical mode that constitutes Hawking radiation at the future null infinity (i.e. at  $\mathcal{I}^+$ ), say, the mode where the intensity of the radiation is the maximum and whose wavelength can be identified, for instance, using Wein's law. When one traces such a mode back to the past null infinity (i.e. to  $\mathcal{I}^-$ ) where the initial conditions are imposed on the quantum field, one finds that the energy of the mode turns out to be way beyond the Planck scale. (This feature seems to have been originally noticed in Ref. [2]; in this context, also see Ref. [3].) As Hawking radiation mostly consists of modes that leave the future event horizon just before its formation, such a phenomenon essentially occurs due to the enormous red-shifting of the modes near the horizon. This behavior then raises the question as to whether Planck scale effects will modify the spectrum of Hawking radiation and the associated stress-energy tensor of the quantum field.

There has been a sufficient amount of effort in the literature towards understanding the effects of Planck scale physics on Hawking radiation (for the earliest discussions, see Refs. [4–7], and, for relatively recent efforts, see Refs. [8, 9]). In the absence of a workable quantum theory of gravity, to study the Planck scale effects, most of these efforts (apart from one notable exception, see Ref. [6]) consider phenomenological models constructed by hand—models which purportedly contain one or more features of the actual effective theory obtained by integrating out the gravitational degrees of freedom. These models either introduce new features in the standard dispersion relation [4, 5], or work with a classical fluctuating geometry [7], or assume that the spacetime coordinates are non-commutative [8]. Often—though, we should hasten to add—but, not always, these high energy models do not preserve local Lorentz invariance. Moreover, some of them either consider a simpler model of Hawking radiation (say, the popular model of a moving mirror in flat space-time [8]) rather than the actual situation, or just consider the spherically symmetric (i.e. the  $\ell = 0$ ) mode in higher dimensional cases, which, essentially, reduces to studying the effects in  $(1 + 1)$ -dimensions.

---

\*E-mail: dawood@iucaa.ernet.in

†E-mail: Shanki.Subramaniam@port.ac.uk

‡E-mail: sriram@mri.ernet.in

Our aim in this work, is to use a locally Lorentz invariant approach to evaluate the Planck scale modifications to stress-energy tensor around a black hole without resorting such approximations.

An approach that introduces a fundamental length scale (say, the Planck scale  $L_p$ ) into standard quantum field theory while preserving local Lorentz invariance is the so-called principle of path integral duality (for the original discussion, see Refs. [10]; for various applications of the approach, see Refs. [11–13]). Interestingly, it has been shown that, at low energies, the modified propagator of matter fields obtained through such an approach is equivalent to taking into account the string fluctuations propagating along compact extra dimensions [14–16]. Effectively, the path-integral duality approach can be said to provide a prescription to evaluate the modified Green's function of a free quantum field that is propagating in a given classical background (for further details, see the next section.) In this work, we shall use this prescription to evaluate the modified Green's function and the Planck scale corrections to the stress-energy tensor around a specific black hole.

In  $(3 + 1)$ -dimensions, it proves to be difficult to evaluate the two-point function exactly around even the simplest of black holes. As a result, the stress-energy tensor of a quantum field around, say, the Schwarzschild black hole has been evaluated only under an approximation (see, for example, Refs. [17–19]). In contrast, the spacetime around the  $(2 + 1)$ -dimensional Banados-Teitelboim-Zanelli (BTZ, hereafter) black hole [20–22] provides a situation wherein it is possible to calculate the two-point function in a closed form [23–29]. We shall utilize this feature to compute the modified propagator and the Planck scale corrections to the standard stress-energy tensor around the BTZ black hole.

The remainder of this paper is organized as follows. In the following section, we shall briefly outline as to how the principle of path integral duality modifies the two-point function of a free quantum field evolving in a given spacetime. In Section III, we evaluate the modified propagator around the BTZ black hole, and show that the path integral duality approach regulates the ultra-violet behavior of the two-point function. In Section IV, we evaluate the stress-energy tensor associated with the modified two-point function and illustrate the form of the Planck scale corrections to the stress-energy tensor. Finally, in Section V, we close with a brief discussion on the results we have obtained.

Before we proceed, a few words on our notations and conventions are in order. We shall work in  $(2 + 1)$ -dimensions and adopt the metric signature of  $(-, +, +)$ . Also, for convenience, we shall denote the set of three coordinates  $x^\mu$  as  $\tilde{x}$ , and use natural units such that  $\hbar = c = 1$ .

## II. STRING FLUCTUATIONS, DUALITY AND THE MODIFIED GREEN'S FUNCTION

Recently, it was shown that, when the fluctuations of closed strings along compact extra dimensions are taken into account, one arrives at an effective propagator for point particles that is regular at high energies [16]. It was found that, the  $T$ -duality symmetry between the topologically non-trivial excitations and the winding modes of the strings around the compact dimensions leads to a modified propagator in the Minkowski vacuum wherein the original spacetime interval  $(\tilde{x} - \tilde{x}')^2$  between the two spacetime events  $\tilde{x}$  and  $\tilde{x}'$  is replaced by  $[(\tilde{x} - \tilde{x}')^2 + L_p^2]$ , where, as we mentioned above,  $L_p$  denotes the Planck length. Thus, effectively, the string fluctuations introduce a zero point length  $L_p$  into standard field theory.

A similar modification of the Minkowski propagator has been obtained earlier by invoking the principle of path integral duality [10]. Recall that, in standard quantum field theory, the path integral amplitude for a path connecting events  $\tilde{x}$  and  $\tilde{x}'$  in a given spacetime is proportional to the proper length, say,  $\mathcal{R}(\tilde{x}, \tilde{x}')$  between the two events. The duality principle proposes that the path integral amplitude should be invariant under the transformation  $\mathcal{R} \rightarrow (L_p^2/\mathcal{R})$ . Operationally, it turns out to be convenient to express the effect of the duality approach and the string fluctuations on the propagator in terms of the Schwinger's proper time formulation [10]. One finds that, in such a formulation, the string fluctuations and the duality approach effectively modify the weightage given to a point particle of mass  $m$ , from  $\exp(i m^2 s)$  to  $\exp[i(m^2 s + (L_p^2/s))]$ .

Consider a free scalar field of mass  $m$  that is propagating in a given classical gravitational background described by the metric tensor  $g_{\mu\nu}$ . Let us further assume that the field is non-minimally coupled to gravity. In Schwinger's proper time formalism, the two-point function corresponding to such a scalar field can be expressed as [30, 31]

$$G(\tilde{x}, \tilde{x}') = \int_0^\infty ds K(\tilde{x}, \tilde{x}'; s), \quad (1)$$

where  $K(\tilde{x}, \tilde{x}'; s)$  is defined as

$$K(\tilde{x}, \tilde{x}'; s) \equiv \langle \tilde{x} | e^{-i(\square - m^2 - \xi R)s} | \tilde{x}' \rangle, \quad (2)$$

and  $R$  is the curvature of the background spacetime with  $\xi$  denoting the coefficient of the non-minimal coupling. In other words, the quantity  $K(\tilde{x}, \tilde{x}'; s)$  is the path integral amplitude for a quantum mechanical system described by the following Hamiltonian:

$$\hat{H} = (\square - m^2 - \xi R) \equiv \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu) - m^2 - \xi R. \quad (3)$$

As we mentioned, the effects due to path integral duality or, equivalently, the string fluctuations, correspond to modifying the expression (1) above for the two point function to [10, 16]

$$G^M(\tilde{x}, \tilde{x}') = \int_0^\infty ds e^{iL_P^2/4s} K(\tilde{x}, \tilde{x}'; s). \quad (4)$$

In the following section, we shall make use of this prescription to evaluate the modified two-point function around the BTZ black hole.

### III. THE MODIFIED GREEN'S FUNCTION AROUND THE BTZ BLACK HOLE

The (non-rotating) BTZ black hole is a spherically symmetric vacuum solution of the Einstein's equations in three-dimensional anti-de Sitter spacetime (AdS<sub>3</sub>, hereafter). It can be represented as AdS<sub>3</sub> identified under a discrete subgroup of its isometry group [20–22].

Recall that AdS<sub>3</sub> is a maximally symmetric space sourced by a *negative* cosmological constant, say,  $-\ell^{-2}$ . AdS<sub>3</sub> can be described by the line-element

$$ds^2 = - \left( \frac{\hat{r}^2}{\ell^2} - 1 \right) d\hat{t}^2 + \left( \frac{\hat{r}^2}{\ell^2} - 1 \right)^{-1} d\hat{r}^2 + \hat{r}^2 d\hat{\phi}^2. \quad (5)$$

where  $-\infty < (\hat{t}, \hat{\phi}) < \infty$  and  $0 < \hat{r} < \infty$ . It is important to note that the coordinate  $\hat{\phi}$  in the above line-element has an infinite range and, hence, is *not* periodic. The non-rotating BTZ black hole of mass  $M$  can be obtained from the above AdS<sub>3</sub> line-element by the following periodic identification of the coordinate  $\hat{\phi}$  [20–22]

$$\hat{\phi} \equiv \hat{\phi} + (2\pi n) \sqrt{M}, \quad (6)$$

where  $n$  is an integer. On redefining the coordinates  $(\hat{t}, \hat{r}, \hat{\phi})$  as follows (see, for instance, Ref. [23]):

$$\hat{t} = \sqrt{M} t, \quad \hat{r} = (r/\sqrt{M}) \quad \text{and} \quad \hat{\phi} = \sqrt{M} \phi, \quad (7)$$

we obtain the metric around the BTZ black hole to be

$$ds^2 = - \left( \frac{r^2}{\ell^2} - M \right) dt^2 + \left( \frac{r^2}{\ell^2} - M \right)^{-1} dr^2 + r^2 d\phi^2 \quad (8)$$

with  $-\infty < t < \infty$ ,  $0 < r < \infty$  and  $0 < \phi \leq (2\pi)$ . Evidently, the event horizon of the black hole is located at  $r_H = (\sqrt{M} \ell)$ .

It is clear from the above discussion that, if we can evaluate the Green's function, say,  $G_{\text{AdS}_3}(\tilde{x}, \tilde{x}')$ , in AdS<sub>3</sub>, then the Green's function, say,  $G_{\text{BTZ}}(\tilde{x}, \tilde{x}')$  around the BTZ black hole can be obtained on imposing by hand the periodicity condition (6) of the angular coordinate, and transforming to the black holes coordinates using the relations (7) [23–29]. Therefore, we need to first evaluate the Green's function in AdS<sub>3</sub>. We shall do so by evaluating the quantum mechanical kernel  $K(\tilde{x}, \tilde{x}'; s)$  and using the Schwinger's proper time expression (1) to arrive at the Green's function.

In order to calculate the kernel  $K(\tilde{x}, \tilde{x}'; s)$  in AdS<sub>3</sub>, it turns out to be more convenient to work with the following set of coordinates [21]:

$$x_2 = \hat{r}^{-1} \sqrt{(\hat{r}^2 - \ell^2)} e^\phi \sinh(\hat{t}/\ell), \quad y = (\ell/\hat{r}) e^\phi, \quad x_1 = \hat{r}^{-1} \sqrt{(\hat{r}^2 - \ell^2)} e^\phi \cosh(\hat{t}/\ell). \quad (9)$$

In terms of these new coordinates, the AdS<sub>3</sub> line-element (5) reduces to

$$ds^2 = (\ell^2/y^2) (-dx_2^2 + dy^2 + dx_1^2), \quad (10)$$

which, evidently, is conformal to flat spacetime. The kernel of a massive and non-minimally coupled scalar field propagating in the above conformally flat line-element can be easily evaluated using the method of spectral decomposition (see, for instance, the Appendix in Ref. [29]). We find that the kernel can be expressed as [27–29]

$$K(\tilde{x}, \tilde{x}'; s) = \left( \frac{1}{(4\pi i s)^{3/2}} \right) \left( \frac{\sigma(\tilde{x}, \tilde{x}')/\ell}{\sinh(\sigma(\tilde{x}, \tilde{x}')/\ell)} \right) \exp [i\sigma^2(\tilde{x}, \tilde{x}')/4s - i b s/\ell^2], \quad (11)$$

where  $b = \ell^2 (1/\ell^2 + m^2 + \xi R)$ ,  $R = -(6/\ell^2)$  being the Ricci scalar (a constant in AdS<sub>3</sub>), and the quantity  $\sigma(\tilde{x}, \tilde{x}')$  is given by

$$\cosh(\sigma(\tilde{x}, \tilde{x}')/\ell) = \left[ \left( (y^2 + y'^2) + (x_1 - x'_1)^2 - (x_2 - x'_2)^2 \right) / 2 y y' \right]. \quad (12)$$

In terms of the original coordinates  $(\hat{t}, \hat{r}, \hat{\phi})$ , the quantity  $\sigma(\tilde{x}, \tilde{x}')$  is given by

$$\sinh[\sigma(\tilde{x}, \tilde{x}')/2\ell] = \left( \sqrt{2} \ell \right)^{-1} \left[ \hat{r} \hat{r}' \cosh(\hat{\phi} - \hat{\phi}') - \ell^2 - \sqrt{(\hat{r}^2 - \ell^2)(\hat{r}'^2 - \ell^2)} \cosh[(\hat{t} - \hat{t}')/\ell] \right]^{1/2}. \quad (13)$$

On using the expression (1), the Green's function corresponding to the kernel (11) can then immediately be evaluated to be [27]

$$G_{\text{AdS}}(\tilde{x}, \tilde{x}') = \left( \frac{1}{4\pi\ell} \right) \left( \frac{1}{\sinh(\sigma(\tilde{x}, \tilde{x}')/\ell)} \right) \exp - \left[ \sqrt{b} \sigma(\tilde{x}, \tilde{x}')/\ell \right]. \quad (14)$$

We should mention here that this Green's function corresponds to a particular choice of boundary condition (actually, the Dirichlet condition [27–29]) that are required to be imposed at spatial infinity in AdS space-times (see, for example Refs. [32]).

The Green's function around the BTZ black hole can now be obtained by imposing the periodicity condition (6) in the AdS<sub>3</sub> Green's function (14), and transforming into the black holes coordinates using the relations (7). The BTZ Green's function is given by [23–29]

$$G_{\text{BTZ}}(\tilde{x}, \tilde{x}') = \left( \frac{1}{4\pi\ell} \right) \sum_{n=-\infty}^{\infty} \left( \frac{1}{\sinh(\sigma_n(\tilde{x}, \tilde{x}')/\ell)} \right) \exp - \left[ \sqrt{b} \sigma_n(\tilde{x}, \tilde{x}')/\ell \right], \quad (15)$$

where  $\sigma_n(\tilde{x}, \tilde{x}')$  is defined as

$$\sinh[\sigma_n(\tilde{x}, \tilde{x}')/2\ell] = \left( \sqrt{2M}\ell \right)^{-1} \left( r r' \cosh \left[ \sqrt{M} (\phi - \phi' + 2\pi n) \right] - M \ell^2 - \sqrt{(\hat{r}^2 - \ell^2)(\hat{r}'^2 - \ell^2)} \cosh \left[ \sqrt{M} (t - t')/\ell \right] \right)^{1/2}. \quad (16)$$

The time translational invariance clearly indicates that the above Green's function corresponds to the Hawking-Hartle state of the black hole (see, for instance, Ref. [33]).

The duality modified Green's function in AdS<sub>3</sub> can now be obtained by substituting the kernel (11) in the expression (4) and carrying out the integral. We obtain the duality modified Green's function in AdS<sub>3</sub> to be

$$G_{\text{AdS}}^{\text{M}}(\tilde{x}, \tilde{x}') = \left( \frac{1}{4\pi\ell} \right) \left( \frac{\sigma(\tilde{x}, \tilde{x}')}{\sinh(\sigma(\tilde{x}, \tilde{x}')/\ell)} \right) \left( \frac{1}{\sqrt{\sigma^2(\tilde{x}, \tilde{x}') + L_{\text{P}}^2}} \right) \exp - \left( \sqrt{\frac{b}{\ell^2} [\sigma^2(\tilde{x}, \tilde{x}') + L_{\text{P}}^2]} \right). \quad (17)$$

with  $\sigma(\tilde{x}, \tilde{x}')$  given by Eq. (13). In the coincidence limit (i.e. when  $\tilde{x} \rightarrow \tilde{x}'$ ),  $\sigma(\tilde{x}, \tilde{x}') \rightarrow 0$ , and the above modified Green's function reduces to

$$G_{\text{AdS}}^{\text{M}}(\tilde{x}, \tilde{x}') = \left( \frac{1}{4\pi L_{\text{P}}} \right) \exp - \left( \sqrt{b} L_{\text{P}}/\ell \right). \quad (18)$$

Note that the Green's function is finite in the coincident limit independent of the nature of the coupling and the mass of the scalar field. This behavior clearly illustrates that the string theory inspired modification regulates the theory at the Planck scale.

The modified Green's function around the BTZ black can be obtained from the above modified Green's function in AdS<sub>3</sub> as in the standard case. It can be expressed as

$$G_{\text{BTZ}}^{\text{M}}(\tilde{x}, \tilde{x}') = \left(\frac{1}{4\pi\ell}\right) \sum_{n=-\infty}^{\infty} \left(\frac{\sigma_n(\tilde{x}, \tilde{x}')}{\sinh(\sigma_n(\tilde{x}, \tilde{x}')/\ell)}\right) \left(\frac{1}{\sqrt{\sigma_n^2(\tilde{x}, \tilde{x}') + L_{\text{P}}^2}}\right) \exp - \left(\sqrt{\frac{b}{\ell^2} [\sigma_n^2(\tilde{x}, \tilde{x}') + L_{\text{P}}^2]}\right). \quad (19)$$

with  $\sigma_n(\tilde{x}, \tilde{x}')$  given by Eq. (16). The  $n \neq 0$  terms are finite in the coincident limit even in the standard (i.e. unmodified) case, and the  $n = 0$  term corresponds to AdS<sub>3</sub>. Obviously, the modified Green's function around the BTZ black hole is finite in the coincident limit as well.

#### IV. PLANCK SCALE MODIFICATIONS TO THE STRESS-ENERGY TENSOR

In this section, using the modified Green function (19), we shall evaluate the leading Planck scale corrections to the stress tensor around the BTZ black hole. For convenience in calculation, we shall restrict ourselves to the case of a massless and a conformally coupled scalar field [i.e. when  $\xi = (1/8)$ ,  $m = 0$  and, hence,  $b = (1/4)$ ]. In such a situation, given the Green's function, say,  $G(\tilde{x}_1, \tilde{x}_2)$ , the corresponding mean value of the stress-energy tensor can be expressed as [23–25]

$$\langle \hat{T}_{\mu\nu} \rangle = \lim_{2 \rightarrow 1} \mathcal{T}_{\mu\nu}^{(1,2)} G(\tilde{x}_1, \tilde{x}_2). \quad (20)$$

The quantity  $\mathcal{T}_{\mu\nu}^{(1,2)}$  is a differential operator and is given by

$$\mathcal{T}_{\mu\nu}^{(1,2)} = \left[ \left(\frac{3}{4}\right) (\nabla_{\mu}^1 \nabla_{\nu}^2) - \left(\frac{1}{4}\right) g_{\mu\nu} g^{\alpha\beta} (\nabla_{\alpha}^1 \nabla_{\beta}^2) - \left(\frac{1}{8}\right) (\nabla_{\mu}^1 \nabla_{\nu}^1 + \nabla_{\mu}^2 \nabla_{\nu}^2) - \left(\frac{1}{16\ell^2}\right) g_{\mu\nu} \right]. \quad (21)$$

with the covariant derivatives  $\nabla_{\mu}^1$  and  $\nabla_{\mu}^2$  acting on the points  $\tilde{x}_1$  and  $\tilde{x}_2$ , respectively.

For the case of the Dirichlet boundary condition imposed at spatial infinity, the mean value of the stress-energy tensor associated with a massless and conformally coupled scalar field can be obtained by substituting the Green's function (15) [with  $b$  set to  $(1/2)$  in the expression (20) above. We find that the stress-energy tensor is given by [24, 25]

$$\begin{aligned} \langle \hat{T}_{\nu}^{\mu} \rangle = & \left(\frac{M^{3/2}}{32\pi r^3}\right) \sum_{n=1}^{\infty} \left[ 2 + \left(1 - \left[1 + \left(\frac{M\ell^2}{r^2 \sinh^2(\sqrt{M}\pi n)}\right)\right]^{-3/2}\right) \sinh^2(\sqrt{M}\pi n) \right] \\ & \times \left(\frac{1}{\sinh^3(\sqrt{M}\pi n)}\right) \text{diag. } (1, 1, -2) \\ & + \left(\frac{3(r^2 - M\ell^2)}{r^2 \sinh^3(\sqrt{M}\pi n)}\right) \left[1 + \left(\frac{M\ell^2}{r^2 \sinh^2(\sqrt{M}\pi n)}\right)\right]^{5/2} \text{diag. } (1, 0, -1). \quad (22) \end{aligned}$$

In order to arrive at this finite result, we have regularized the stress-energy tensor by simply dropping the  $n = 0$  term which corresponds to the stress-energy tensor in AdS<sub>3</sub>. It is well-known that the stress-energy associated with a massless and conformally coupled scalar field vanishes in AdS<sub>3</sub> because of the absence of the trace anomaly in odd dimensions [33].

The Planck modifications to the stress-energy tensor can now be arrived at upon using the modified Green's function (19) instead corrections. We find that the resulting stress-energy tensor is again diagonal, but the expressions turn out to be fairly long to be displayed. As an illustration of the behavior of the stress-energy tensor, we have instead plotted the the  $n = 2$  term in energy density (viz the  $\langle \hat{T}_t^t \rangle$  component) in the accompanying figure. It is evident from the figure that the Planck scale modifications to the stress-energy tensor are rather small.

#### V. DISCUSSION

In this work, using the T-duality symmetry of the string fluctuations, we have evaluated the modified two-point function and the resulting stress-energy tensor around the BTZ black hole. We should emphasize here that we have not made any approximation whatsoever in our calculations and the results we have obtained around the BTZ black hole are exact. This is important, since, as we had mentioned in the introductory section, much of the earlier analysis




FIG. 1: The  $n = 2$  term in the standard energy density (red continuous line) and the energy density obtained using the modified propagator (blue dashed line) has been plotted on the left. The difference between these two quantities has been plotted on the right. We have assumed that  $M = 0.1$ ,  $\ell = 1$ , and to illustrate the modification (or, rather, the lack thereof) due to the Planck scale effects, we have assumed a very large value for  $L_P$ —we have set  $L_P = 10^{-3}$ ! Clearly, the Planck scale modifications are negligibly small.

in the literature, had either worked with the moving mirror model of Hawking radiation or had just considered the spherically symmetric mode in  $(3 + 1)$ -dimensions, which, effectively, simplifies to the  $(1 + 1)$ -dimensional model.

Interestingly, we find that the modified Green's function remains finite in the coincident limit. Actually, such a result could have been expected based on the Schwinger-DeWitt expansion [31]. According to the expansion, for small separations, the kernel  $K(\tilde{x}, \tilde{x}'; s)$  in an arbitrary spacetime has the same form as in the Minkowski vacuum. Therefore, if the modified Green function is ultra-violet regulated in flat spacetime, then it can be expected to remain finite in the coincident limit in any curved spacetime as well. Further, we find the Planck scale modification to the stress-energy tensor are negligibly small, which is a corroboration of the results that have been arrived at earlier in the literature.

Ideally, rather than evaluate the Planck scale modifications to the stress-energy tensor, one would have liked to evaluate the corrections to Hawking radiation itself. This in turn requires that one considers a dynamical situation, and evaluates either the effective Lagrangian or the in-out Bogoliubov coefficient, while taking into the Planck scale modifications. We are currently investigating these issues.

### Acknowledgments

The authors would like to thank T. Padmanabhan for discussions. DAK and LS wish to thank the Harish-Chandra Research Institute, Allahabad, India, and the Inter University Centre for Astronomy and Astrophysics, Pune, India, respectively, for hospitality, where part of this work was carried out. DAK is supported by the Senior Research Fellowship of the Council for Scientific and Industrial Research, India. SS is supported by the Marie Curie Incoming International Grant IIF-2006-039205.

- 
- [1] S. W. Hawking, *Nature* **248**, 30 (1974); *Commun. Math. Phys.* **43**, 199 (1975).
  - [2] R. Wald, *Phys. Rev. D* **13**, 3176 (1976).
  - [3] R. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984), Footnote on p. 406.
  - [4] T. Jacobson, *Phys. Rev. D* **48**, 728 (1993); *ibid.* **53**, 7082 (1996); W. G. Unruh, *ibid.* **51**, 2827 (1995); S. Corley and T. Jacobson, *ibid.* **54**, 1568 (1996); T. Jacobson, *Prog. Theor. Phys. Suppl.* **136**, 1 (1999).
  - [5] R. Brout, S. Massar, R. Parentani and Ph. Spindel, *Phys. Rev. D* **52** 4559 (1995); R. Brout, Cl. Gabriel, M. Lubo, Ph. Spindel, *ibid.* **59**, 044005 (1999).
  - [6] N. Hambli, C. P. Burgess *Phys. Rev. D* **53**, 5717 (1996).
  - [7] C. Barrabes, V. Frolov and R. Parentani, *Phys. Rev. D* **59**, 124010 (1999); *ibid.* **62**, 044020 (2000); R. Parentani, *ibid.* **63** 041503 (2001).
  - [8] R. Casadio, P. H. Cox, B. Harms and O. Micu, *Phys. Rev. D* **73**, 044019 (2006).

- [9] I. Agullo, J. Navarro-Salas, G. J. Olmo and L. Parker, Phys. Rev. D **76**, 044018 (2007).
- [10] T. Padmanabhan, Phys. Rev. Letts. **78**, 1854 (1997); T. Padmanabhan, Phys. Rev. D **57**, 6206 (1998).
- [11] K. Srinivasan, L. Sriramkumar and T. Padmanabhan, Phys. Rev. D **58**, 044009(1998).
- [12] S. Shankaranarayanan and T. Padmanabhan, Int. J. Mod. Phys. D **10**, 351 (2001).
- [13] L. Sriramkumar and S. Shankaranarayanan, JHEP **0612**, 050 (2006).
- [14] A. Smailagic, E. Spallucci and T. Padmanabhan, arXiv:hep-th/0308122;
- [15] E. Spallucci and M. Fontanini, arXiv:gr-qc/0508076.
- [16] M. Fontanini, E. Spallucci and T. Padmanabhan, Phys. Lett. B **633**, 627 (2006).
- [17] D. Page, Phys. Rev. D **25**, 1499 (1982).
- [18] A. Campos and B. L. Hu, Phys. Rev. D **58**, 125021 (1998).
- [19] N. Phillips and B. L. Hu, Phys. Rev. D **67**, 104002 (2003).
- [20] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69**, 1849 (1992); M. Bañados, M. Henneaux, C. Teitelboim and J. Zanelli, Phys. Rev. D **48**, 1506 (1993).
- [21] S. Carlip and C. Teitelboim, Phys. Rev. D **51**, 622 (1995).
- [22] S. Carlip, *Quantum Gravity in (2 + 1)-Dimensions* (Cambridge University Press, 1998), Secs. 3.2 and 12.2.
- [23] A. R. Steif, Phys. Rev. D **49**, R585 (1994).
- [24] G. Lifschytz and M. Ortiz, Phys. Rev. D **49**, 1929 (1994).
- [25] K. Shiraishi and T. Maki, Phys. Rev. D **49**, 5286 (1994).
- [26] I. Ichinose and Y. Satoh, Nucl. Phys. B **447**, 340 (1995).
- [27] R. B. Mann and S. N. Solodukhin, Phys. Rev. D **55**, 3622 (1997).
- [28] A. A. Bytsenko, L. Vanzo and S. Zerbini, Phys. Rev. D **57**, 4917 (1998).
- [29] D. Binosi, V. Moretti, L. Vanzo and S. Zerbini, Phys. Rev. D **59**, 104017 (1999).
- [30] J. Schwinger, Phys. Rev. **82** (1951), 664.
- [31] B. S. DeWitt, Phys. Rep. **19** (1975), 297.
- [32] S. J. Avis, C. J. Isham and D. Storey, Phys. Rev. D **18**, 3565 (1978); C. J. C. Burges, D. Z. Freedman, S. Davis and G. W. Gibbons, Ann. Phys. (N.Y.) **167**, 285 (1986).
- [33] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982).