

# Data Tables for Lorentz and CPT Violation

V. Alan Kostelecký<sup>a</sup> and Neil Russell<sup>b</sup>

<sup>a</sup>*Physics Department, Indiana University, Bloomington, IN 47405*

<sup>b</sup>*Physics Department, Northern Michigan University, Marquette, MI 49855*

IUHET 608; February 2016 update to *Reviews of Modern Physics* **83**, 11 (2011) [arXiv:0801.0287]

This work tabulates measured and derived values of coefficients for Lorentz and CPT violation in the Standard-Model Extension. Summary tables are extracted listing maximal attained sensitivities in the matter, photon, neutrino, and gravity sectors. Tables presenting definitions and properties are also compiled.

## CONTENTS

I. Introduction	1
II. Summary tables	2
III. Data tables	3
IV. Properties tables	5
A. Minimal QED extension	6
B. Minimal SME	7
C. Nonminimal sectors	8
References	10
Table 1: List of tables	14
Summary tables S2–S5	15
Data tables D6–D38	20
Properties tables P39–P52	73

## I. INTRODUCTION

Recent years have seen a renewed interest in experimental tests of Lorentz and CPT symmetry. Observable signals of Lorentz and CPT violation can be described in a model-independent way using effective field theory [1].

The general realistic effective field theory for Lorentz violation is called the Standard-Model Extension (SME) [2, 3]. It includes the Standard Model coupled to General Relativity along with all possible operators for Lorentz violation. Both global [2] and local [3] Lorentz violation are incorporated. Since CPT violation in realistic field theories is accompanied by Lorentz violation [4], the SME also describes general CPT violation. Reviews of the SME can be found in Refs. [5–7].

Each Lorentz-violating term in the Lagrange density of the SME is constructed as the coordinate-independent product of a coefficient for Lorentz violation with a Lorentz-violating operator. The Lorentz-violating physics associated with any operator is therefore controlled by the corresponding coefficient, and so any experimental signal for Lorentz violation can be expressed in terms of one or more of these coefficients.

The Lorentz-violating operators in the SME are systematically classified according to their mass dimension, and operators of arbitrarily large dimension can appear. At any fixed dimension, the operators are finite in number and can in principle be enumerated. A limiting case of particular interest is the minimal SME, which can be viewed as the restriction of the SME to include only Lorentz-violating operators of mass dimension four or less. The corresponding coefficients for Lorentz violation are dimensionless or have positive mass dimension.

The results summarized here concern primarily but not exclusively the coefficients for Lorentz violation in the minimal SME. We compile 33 *data tables* for these SME coefficients, including both existing experimental measurements and theory-derived limits. Each of these data tables provides information about the results of searches for Lorentz violation for a specific sector of the SME. For each measurement or constraint, we list the relevant coefficient or combination of coefficients, the result as presented in the literature, the context in which the search was performed, and the source citation. The tables include results available from the literature up to January 31, 2016.

The scope of the searches for Lorentz violation listed in the data tables can be characterized roughly in terms of depth, breadth, and refinement. Deep searches yield great sensitivity to a small number of SME coefficients. Broad searches cover substantial portions of the coefficient space, usually at a lesser sensitivity. Searches with high refinement disentangle combinations of coefficients. In the absence of a compelling signal for Lorentz violation, all types of searches are necessary to obtain complete coverage of the possibilities.

As a guide to the scope of the existing searches, we extract from the data tables four *summary tables* covering the sectors for matter (electrons, protons, neutrons, and their antiparticles), photons, neutrinos, and gravity. These summary tables list our best estimates for the maximal attained sensitivities to the relevant SME

coefficients in the corresponding sectors. Each entry in the summary tables is obtained under the assumption that only one coefficient is nonzero. The summary tables therefore provide information about the overall search depth and breadth, at the cost of masking the search refinement.

In addition to the data tables and the summary tables, we also provide 14 *properties tables* listing some features and definitions of the SME and the coefficients for Lorentz violation. The Lagrange densities for the minimal QED extension in Riemann spacetime, for the minimal SME in Riemann-Cartan spacetime, for a non-minimal Dirac fermion in Minkowski spacetime, and for the nonminimal photon sector in Minkowski spacetime are provided in tabulated form. The mass dimensions of the operators for Lorentz violation and their properties under the various discrete spacetime transformations are displayed. Standard combinations of SME coefficients that appear in the literature are listed. Along with the data tables and the summary tables, the properties tables can be used to identify open directions for future searches. Among these are first measurements of unconstrained coefficients, improved sensitivities to constrained coefficients, and studies disentangling combinations of coefficients.

The organization of the tables is as follows. Table 1 contains a list of all tables. The four summary tables are presented next, Tables S2–S5. These are followed by the 33 data tables, Tables D6–D38. The 14 properties tables appear last, Tables P39–P52.

A description of the summary tables is given in Sec. II. Information about the format and content of the data tables is presented in Sec. III, while Sec. IV provides an overview of the properties tables. The bibliography for the text and all the tables follows Sec. IV.

## II. SUMMARY TABLES

The four summary tables (Tables S2–S5) list maximal experimental sensitivities attained for coefficients in the matter, photon, neutrino, and gravity sectors of the minimal SME. To date, there is no confirmed experimental evidence supporting Lorentz violation. A few measurements suggest nonzero coefficients at weak confidence levels. These latter results have been excluded in constructing the summary tables but are listed in the data tables. Also excluded are results based on the reported  $6\sigma$  difference between the speeds of muon neutrinos and light in the OPERA experiment [8], which has since been identified as a systematic effect [9].

In the four summary tables, each displayed sensitivity value represents our conservative estimate of a  $2\sigma$  limit, given to the nearest order of magnitude, on the modulus of the corresponding coefficient. Our rounding convention is logarithmic: a factor greater than or equal to  $10^{0.5}$




FIG. 1: Standard Sun-centered inertial reference frame [10].

rounds to 10, while a factor less than  $10^{0.5}$  rounds to 1. In a few cases, tighter results may exist when suitable theoretical assumptions are adopted; these results can be found in the data tables that follow.

Where observations involve a linear combination of the coefficients appearing in the summary tables, the displayed sensitivity for each coefficient assumes for definiteness that no other coefficient contributes. Some caution is therefore advisable in applying the results in these summary tables to situations involving two or more nonzero coefficient values. Care in applications is also required because under some circumstances certain coefficients can be intrinsically unobservable or can be absorbed into others by field or coordinate redefinitions, as described in Sec. IV A.

In presenting the physical sensitivities, we adopt natural units with  $\hbar = c = \epsilon_0 = k_B = 1$  and express mass units in GeV. Our values are reported in the standard Sun-centered inertial reference frame [10] widely used in the literature. This frame is illustrated in Fig. 1. The origin of the time coordinate  $T$  is at the 2000 vernal equinox. The  $Z$  axis is directed north and parallel to the rotational axis of the Earth at  $T = 0$ . The  $X$  axis points from the Sun towards the vernal equinox, while the  $Y$  axis completes a right-handed system. Some further details about this frame, including transformations to other standard frames, can be found in Section III A and Appendix C of Ref. [11].

Table S2 lists the maximal attained sensitivities involving electrons, protons, neutrons, and their antiparticles. For each distinct massive spin-half Dirac fermion in the minimal SME in Minkowski spacetime, there are 44 independent observable combinations of coefficients for Lorentz violation in the nonrelativistic limit. Of these,

20 also control CPT violation. The 44 combinations are conventionally chosen as the tilde coefficients shown. The definitions of these 44 tilde coefficients in terms of coefficients in the minimal SME are listed in Table P41. All the definitions appear elsewhere in the literature [10] except the four combinations  $\tilde{b}_J^*$  and  $\tilde{c}_{TT}$ . The three tilde coefficients  $\tilde{b}_J^*$  are the antimatter equivalent of the tilde coefficients  $\tilde{b}_J$ . They appear in nonrelativistic studies of antimatter properties, such as the hyperfine transitions of antihydrogen [12]. The tilde coefficient  $\tilde{c}_{TT}$  is a simple scaling of the coefficient  $c_{TT}$  in the minimal SME, introduced here to ensure completeness of the set of tilde coefficients. All tilde coefficients have dimensions of GeV in natural units. In Table S2, a superscript indicating the particle species of relevance is understood on all coefficients. For example, the first line of the table presents limits on three different tilde coefficients,  $\tilde{b}_X^e$ ,  $\tilde{b}_X^p$ ,  $\tilde{b}_X^n$ . In the table, a dash indicates that no sensitivity to the coefficient has been identified to date. A few maximal sensitivities are obtained by applying the inverse of the definitions in Table P41 to data in Tables D6 and D9. These inverses are presented in Table P42.

Table S3 displays the maximal attained sensitivities to coefficients for Lorentz violation in the photon sector. There are 23 observable coefficient combinations for photons in the minimal SME, of which four also control CPT violation. The first four rows of the table concern combinations of these four coefficients, which have dimensions of GeV in natural units. The second part of the table displays maximal sensitivities to 19 tilde coefficients, which are conventional combinations of the 19 dimensionless coefficients in the minimal SME. The definitions of all 23 combinations of coefficients in the minimal SME are taken from the literature [11, 13] and are provided in Table P43. The final section of Table S3 concerns those few coefficients in the photon sector of the full SME that control isotropic Lorentz violation. Exactly one such coefficient exists for each operator of mass dimension  $d$  [14]. The operators also violate CPT when  $d$  is odd. The table displays maximal attained sensitivities to isotropic Lorentz violation from operators of dimension  $3 \leq d \leq 9$ .

Table S4 lists the maximal attained sensitivities to a subset of the coefficients controlling Lorentz-violating mixing and propagation of the three flavors  $\nu_e$ ,  $\nu_\mu$ ,  $\nu_\tau$  of left-handed neutrinos and antineutrinos in the SME. In this table, each numerical value gives the limit on the modulus of the corresponding coefficient. A dash indicates that no sensitivity to the coefficient has been identified as yet. The first 14 rows of the table concern cartesian coefficients governing neutrino mixing for operator dimensions  $d = 3$  and  $d = 4$ . All these flavor-mixing coefficients are complex, so bounds on both real and imaginary parts are provided. A subscript  $e\mu$ ,  $e\tau$ , or  $\mu\tau$  indicating the specific flavor mixing is understood on each coefficient and can be identified by the column head-

ing. For example, the entry in the first row and second column provides the constraint  $|\text{Re}(a_L)_{e\mu}^T| < 10^{-20}$  GeV. The last eight rows of Table S4 concern spherical coefficients in the nonminimal neutrino sector, which are defined in the literature [15] and listed in Table P52. These eight rows list maximal attained sensitivities to isotropic oscillation-free operators of dimension  $3 \leq d \leq 10$  controlling propagation, along with isotropic mixing operators of dimension  $3 \leq d \leq 10$  controlling  $\nu_e \leftrightarrow \nu_\mu$  oscillations. One isotropic coefficient of each type exists for each  $d$ , and the corresponding operators violate CPT when  $d$  is odd.

Table S5 displays the maximal attained sensitivities to certain coefficients for Lorentz violation involving the gravity sector of the minimal SME. Two classes of coefficients can be distinguished in this context: ones appearing in the matter sector, and ones appearing in the pure-gravity sector. For the first class, Table S5 contains results for the 24 coefficients  $\bar{a}_\mu^e$ ,  $\bar{a}_\mu^p$ ,  $\bar{a}_\mu^n$  and  $\bar{c}_\mu^e$ ,  $\bar{c}_\mu^p$ ,  $\bar{c}_\mu^n$  involving the electron, proton, and neutron sectors. These observables are associated with CPT-odd operators. The prefactor  $\alpha$  is a model-dependent number [16]. For the second class, the table displays nine combinations of the nine dimensionless coefficients for Lorentz violation  $\bar{\mathcal{A}}^{\mu\nu}$ . Additional coefficients exist in the minimal gravity sector, but no sensitivities to them have been identified to date (the ‘t puzzle’ [17]).

### III. DATA TABLES

We present 33 data tables compiled from the existing literature. Some of these include results restricted to the minimal SME, others list bounds on nonminimal coefficients, and a few contain data involving both minimal and nonminimal sectors. The 33 tables cover the electron sector (Tables D6–D8), the proton sector (Tables D9 and D10), the neutron sector (Table D11), the photon sector (Tables D12–D18), the charged-lepton sector (Tables D19–D22), the neutrino sector (Tables D23–D30), the quark sector (Tables D31 and D32), the electroweak sector (Table D33), the gluon sector (Table D34), and the gravity sector (Tables D35–D38).

Each of these 33 data tables contains four columns. The first column lists the coefficients for Lorentz violation or their relevant combinations. Results for coefficients of the same generic type are grouped together. Certain results involve combinations of coefficients across more than one sector; each of these has been listed only once in the table deemed most appropriate. Some minor changes in notation or format have been introduced as needed, but for the most part the results are quoted as they appear in the cited references. Definitions for standard combinations of coefficients are provided in the properties tables that follow. A few authors use unconventional notation; where immediate, the match to the

standard notation is shown. Parentheses enclosing a pair of indices on a coefficient indicate symmetrization without a factor of two.

The second column contains the measurements and bounds, presented in the same form as documented in the literature. For each generic type of coefficient, the results are listed in reverse chronological order. If no significant figures appear in the quoted limit on an absolute value, the order of magnitude of the limit is given as a power of 10. Where both statistical and systematic errors appear in a given result, they are quoted in that order.

The third column contains a succinct reminder of the physical context in which the bound is extracted, while the fourth column contains the source citations. The reader is referred to the latter for details of experimental and theoretical procedures, assumptions underlying the results, definitions of unconventional notations, and other relevant information. Results deduced on theoretical grounds are distinguished from those obtained via direct experimental measurement by an asterisk placed after the citation.

Tables D6–D11 contain data for the electron, proton, and neutron sectors. These data are organized sequentially, beginning with combinations involving the minimal coefficients  $b_\nu$ ,  $c_{\mu\nu}$ ,  $H_{\mu\nu}$ ,  $d_{\mu\nu}$ ,  $g_{\mu\nu\lambda}$  and followed by combinations involving nonminimal coefficients. For the electron and proton sectors, minimal and nonminimal coefficients are displayed in distinct tables. A superscript indicating the particle species of relevance is understood on the coefficients in all these tables. Standard definitions for the minimal coefficients and their combinations are provided in Tables P39 and P41, while definitions for the nonminimal coefficients are given in Tables P48 and P49. Many entries in Tables D6–D11 use coefficients for a single Dirac fermion written in a cartesian basis, but some adopt instead coefficients decomposed using spin-weighted spherical harmonics, which provide a convenient means of enumerating Lorentz-violating fermion operators of arbitrary mass dimension  $d$  [18]. Some entries for the nonminimal sectors appearing in these tables involve nonrelativistic spherical coefficients identified by the superscript NR, which are formed as special linear combinations of spherical coefficients for all dimensions  $d$ , while others involve isotropic ultrarelativistic coefficients denoted by a ring diacritic and the superscript UR. Some results depend on  $\eta \simeq 23.5^\circ$ , which is the angle between the equatorial and ecliptic planes in the solar system. Note that the existing bounds on observables involving  $a_\nu^e$ ,  $a_\nu^p$ ,  $a_\nu^n$  and  $e_\nu^e$ ,  $e_\nu^p$ ,  $e_\nu^n$  are obtained from gravitational experiments and are listed with the gravity-sector results in Table D35.

Tables D12 and D13 present the photon-sector data for operator dimensions  $d = 3$  and  $d = 4$ , respectively. Most of the combinations of coefficients for Lorentz violation appearing in the first columns of these tables are de-

finied in Tables P43 and P51. The combinations  $k_{(V)jm}^{(3)}$ ,  $k_{(E)jm}^{(4)}$ , and  $k_{(B)jm}^{(4)}$  arise from analyses [13, 14, 19] using spin-weighted spherical harmonics. The factor of  $\beta_\oplus$  appearing in some places is the speed of the Earth in the standard Sun-centered reference frame, which is about  $10^{-4}$  in natural units.

Tables D14–D18 contain a compilation of some measurements and bounds on coefficients for Lorentz violation in the nonminimal photon sector of the SME. Results are available for a variety of operators of dimensions greater than 4, with the coefficients for  $d = 5$ ,  $d = 6$ ,  $d = 7$ ,  $d = 8$ , and  $d \geq 9$  being placed in distinct tables. The spin-weighted spherical harmonics also provide a convenient basis for classifying photon operators of arbitrary mass dimension  $d$  [14]. The corresponding coefficients in the photon sector are listed in Table P51. Some constraints have been obtained for the vacuum coefficients for Lorentz violation, which are  $c_{(I)jm}^{(d)}$ ,  $k_{(E)jm}^{(d)}$ ,  $k_{(B)jm}^{(d)}$  for even  $d$  and  $k_{(V)jm}^{(d)}$  for odd  $d$ , where the subscripts  $jm$  label the angular-momentum quantum numbers. In the first columns of Tables D14–D18, the various spherical harmonics  ${}_s Y_{jm}(\theta, \phi)$  of spin weight  $s$  are evaluated at specified angles, which are the celestial coordinates of certain astrophysical sources. The conventional spherical harmonics of spin weight 0 are denoted as  $Y_{jm}(\theta, \phi)$ . Measurements are also listed of some vacuum-orthogonal coefficients for Lorentz violation, which are distinguished by a negation diacritic  $\bar{\cdot}$ .

Tables D19–D22 list measurements and bounds on coefficients for Lorentz violation involving the second- and third-generation charged leptons. Results for the muon and its antiparticle for operator dimensions  $d = 3$ ,  $d = 4$ , and  $d \geq 5$  are collected in Tables D19, D20, and D21, respectively. These results are expressed either using the cartesian coefficients specified in the lepton sector of Table P44 or via the spherical basis for Dirac fermions [18] using the spherical coefficients summarized in Table P49. Some nonrelativistic experiments are sensitive to special linear combinations of spherical coefficients involving all dimensions  $d$ , which are identified by the superscript NR and have properties listed in Table P49. The results for these nonrelativistic coefficients are placed in the tables according to their mass dimension and are separated from other coefficients of definite  $d$  by a horizontal line. A few rows of the muon data tables present limits on nonminimal operators governed by isotropic ultrarelativistic coefficients, denoted by a ring diacritic and the superscript UR, which are also listed in Table P49. Experiments with storage rings studying the muon anomalous magnetic moment are sensitive to combinations of spherical coefficients at each  $d$  that involve boost factors and are denoted with a háček diacritic. These háček coefficients are defined in Table XI of Ref. [20]. The few existing results for the tau and its antiparticle can be found in Table D22.

Tables D23–D30 collect results involving coefficients in the neutrino sector. Limits for neutrino operators of renormalizable dimension appear in Tables D23–D25, while Tables D26–D30 contain results for coefficients in the nonminimal neutrino sector. Most results are for operators of dimension  $2 \leq d \leq 10$ , but a few results in Table D30 concern operators of arbitrary dimension. In compiling the results, a separate table is used for each value of  $d$  in the range  $2 \leq d \leq 8$  and for  $d \geq 9$ . Many of the cartesian coefficients appearing in the first columns of these tables are provided in Table P47. For a general treatment, in parallel with the fermion and photon sectors, a basis using spin-weighted spherical harmonics is useful to classify the neutrino operators [15]. The corresponding sets of spherical coefficients are listed in Table P52. In Tables D23–D30, oscillation-free coefficients are listed before those controlling neutrino mixing, with a horizontal line clarifying the separation where useful. In Table D30, horizontal lines also separate coefficients of different  $d$ . The last ten lines of Table D30 concern more general models and present constraints involving combinations of an infinite number of coefficients. In some entries in these tables, orientation information is encoded in the factors  ${}_s\mathcal{N}_{jm}$ , which are defined in Eq. (93) of Ref. [15] and evaluated at specific angles determined by the neutrino propagation direction and the laboratory location. In all these tables, many of the results listed are obtained in the context of specific neutrino models, as described in the cited papers.

Tables D31 and D32 present experimental sensitivities to coefficients for operators in the quark sector. Some coefficients appearing in these tables are obtained from experiments with mesons and are composite quantities defined in the corresponding references. They are effective coefficients for which complete analytical expressions are as yet unknown, formed from certain quark-sector coefficients appearing in Table P44 and from other quantities arising from the quark binding in the mesons. A few rows of Table D32 contain bounds on isotropic ultra-relativistic spherical coefficients [18], denoted by a ring diacritic and the superscript UR. Properties of these coefficients are provided in Table P49.

Tables D33 and D34 concern coefficients in the gauge sectors of the minimal SME. Results for the electroweak sector are listed in Table D33, while those for the gluon sector are in Table D34. The coefficients for the electroweak sector are defined in the gauge and Higgs sections of Table P45. The gluon-sector coefficient is the analogue of the corresponding photon-sector coefficient defined in Table P43.

Tables D35–D38 present measurements and bounds concerning operators in the minimal and nonminimal gravity sector of the SME. Table D35 includes limits on the matter-sector coefficients  $a_\nu^e$ ,  $a_\nu^p$ ,  $a_\nu^n$ , and  $e_\nu^e$ ,  $e_\nu^p$ ,  $e_\nu^n$ , which arise at  $d = 3$  and  $d = 4$  and are detectable in gravitational experiments. Table D36 involves coefficients in

the pure-gravity sector of the minimal SME, all of which occur at  $d = 4$ . The specific combinations appearing in some entries in the first column are defined in the references and are expressed in terms of the coefficients for Lorentz violation listed in the gravity section of Table P45. Tables D37 and D38 list existing experimental constraints on coefficients in the nonminimal pure-gravity sector. They include measurements of combinations of coefficients with  $d = 6$  that are observable in nonrelativistic experiments. These coefficients are defined in Eq. (7) of Ref. [17]. Tables D37 and D38 also display constraints on all effective spherical coefficients  $\bar{s}_{jm}^{(d)}$  with even values of  $d \leq 8$ , which govern Lorentz violation affecting the propagation of gravitational waves. These effective spherical coefficients are defined in Eq. (45) of Ref. [207].

#### IV. PROPERTIES TABLES

The 14 properties tables list various features and definitions related to Lorentz violation. Five tables concern the terms in the restriction of the minimal SME to quantum electrodynamics (QED) in Riemann spacetime. For this theory, which is called the minimal QED extension, the tables include information about the operator structure (Table P39), the action of discrete symmetries (Table P40), and some useful coefficient combinations (Tables P41, P42, and P43). Two tables contain information about the matter sector (Table P44) and the gauge and gravity sectors (Table P45) of the minimal SME in Riemann-Cartan spacetime. Another two tables summarize some features of the coefficients for Lorentz violation concerning operators of renormalizable dimension in the neutrino sector. One (Table P46) lists the cartesian coefficients and some of their properties, while the other (Table P47) provides the connection between cartesian and spherical coefficients. The five remaining tables (Tables P48–P52) provide information about the operator structure and the spherical coefficients for Lorentz violation in the nonminimal fermion, photon, and neutrino sectors.

For these properties tables, our primary conventions are those of Ref. [3]. Greek indices  $\mu, \nu, \lambda, \dots$  refer to curved-spacetime coordinates and Latin indices  $a, b, c, \dots$  to local Lorentz coordinates. The vierbein formalism [21], which relates the two sets of coordinates, is adopted to facilitate the description of spinors on the spacetime manifold. The determinant  $e$  of the vierbein  $e_\mu^a$  is related to the determinant  $g$  of the metric  $g_{\mu\nu}$  by  $e = \sqrt{-g}$ . The conventions for the Dirac matrices  $\gamma^a$  are given in Appendix A of Ref. [3]. The Newton gravitational constant  $G_N$  enters as the combination  $\kappa \equiv 8\pi G_N$ , and it has dimensions of inverse mass squared.

In the Minkowski-spacetime limit, the metric  $g_{\mu\nu}$  is written  $\eta_{\mu\nu}$  with diagonal entries  $(-1, 1, 1, 1)$ . For

decompositions into time and space components, we adopt the Sun-centered frame of Fig. 1 and use indices  $J, K, L, \dots$  to denote the three spatial components  $X, Y, Z$ . The sign of the antisymmetric tensor  $\epsilon_{\kappa\lambda\mu\nu}$  is fixed via the component  $\epsilon_{TXYZ} = +1$ , and the antisymmetric symbol in three spatial dimensions is defined with  $\epsilon_{XYZ} = +1$ . Note that some of the literature on the SME in Minkowski spacetime adopts a metric  $\eta_{\mu\nu}$  of opposite sign, following the common present usage in quantum physics instead of the one in relativity. Under this alternative convention, terms in the Lagrange density with an odd number of index contractions have opposite signs to those appearing in this work. The numerical results for the SME coefficients in the tables are unaffected by the convention.

### A. Minimal QED extension

Table P39 concerns the minimal QED extension, for which the basic nongravitational fields are a Dirac fermion  $\psi$  and the photon  $A_\mu$ . The electromagnetic field-strength tensor is  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . The pure-gravity sector involves the Riemann tensor  $R_{\kappa\lambda\mu\nu}$ , the Ricci tensor  $R_{\mu\nu}$ , the curvature scalar  $R$ , and the cosmological constant  $\Lambda$ . The spacetime covariant derivative  $D_\mu$  corrects local Lorentz indices using the spin connection, corrects spacetime indices using the Cartan connection, and contains the usual gauge field  $A_\mu$  for the photon. The notation  $\overset{\leftrightarrow}{D}_\mu$  is an abbreviation for the difference of two terms, the first with derivative acting to the right and the second acting to the left. Note that Table P39 is restricted to the zero-torsion limit of the minimal SME. The general case [3] involves additional operators constructed with the torsion tensor  $T^\alpha{}_{\mu\nu}$ . The Minkowski-spacetime limit of QED with nonzero torsion contains terms that mimic Lorentz violation, so searches for Lorentz violation can be used to bound components of the torsion tensor [22, 23].

In Table P39, each line specifies one term in the Lagrange density for the QED extension in Riemann spacetime. Both conventional QED terms and ones with Lorentz violation are included. The first column indicates the sector to which the term belongs. The second column lists the coefficient controlling the corresponding operator. Note the standard use of an upper-case letter for the coefficient  $H_{\mu\nu}$ , which distinguishes it from the metric fluctuation  $h_{\mu\nu}$ . The third column shows the number of components for the coefficient. The next three columns list the operator, its mass dimension, and the vierbein factor contracting the coefficient and the operator. The final two columns list the properties of the term under CPT and Lorentz transformations. A CPT-even operator is indicated by a plus sign and a CPT-odd one by a minus sign, while terms violating Lorentz invariance

are identified by a check mark.

As an example, consider the fourth row of Table P39. This concerns the term in the fermion sector with coefficient  $a_\mu$  for Lorentz violation. The coefficient has four independent components, which control the four Lorentz-violating operators  $\bar{\psi}\gamma^a\psi$ . The gravitational couplings of this operator are contained in the vierbein product  $ee^\mu{}_a$ . The corresponding term in the Lagrange density for the minimal QED extension in Riemann spacetime is  $\mathcal{L}_a = -ea_\mu e^\mu{}_a \bar{\psi}\gamma^a\psi$ . It has mass dimension 3 and is CPT odd. The Minkowski-spacetime limit of this term can be obtained by the vierbein replacement  $e_\mu{}^a \rightarrow \delta_\mu{}^a$ . The number of index contractions in  $\mathcal{L}_a$  is two, one each for the  $\mu$  and  $a$  indices, so the overall sign of  $\mathcal{L}_a$  is unaffected by the choice of convention for the Minkowski metric.

The properties listed in Table P39 are those of the operators in the Lagrange density rather than those associated with observables. The issue of observability of a given coefficient can be subtle because experiments always involve comparisons of at least two quantities. The point is that in certain tests a given coefficient may produce the same effect on two or more quantities and so may be unobservable, or it may produce effects indistinguishable from those of other coefficients. This situation can often be theoretically understood via a field redefinition that eliminates the coefficient from the relevant part of the Lagrange density without affecting the dynamics of the experiment in question. For example, a constant coefficient  $a_\mu$  in the minimal QED extension in Minkowski spacetime is unobservable in any experiment involving a single fermion flavor because it can be absorbed as a phase shift in the fermion field [2]. The situation changes in Riemann spacetime, where three of the four components of  $a_\mu$  become observables affecting the gravitational properties of the fermion [3]. Another example is provided by the coefficient  $f_\mu$  in the minimal QED extension in Minkowski spacetime, which can be converted into a coefficient of the  $c_{\mu\nu}$  type via a change of spinor basis [24, 25]. Additional subtleties arise because any experiment must always choose definitions of clock ticking rates, clock synchronizations, rod lengths, and rod isotropies. This involves 10 free coordinate choices and implies the unobservability of 10 combinations of coefficients for Lorentz violation [26].

Table P40 lists the properties under discrete-symmetry transformations of the Lorentz-violating operators in the minimal QED extension [27]. The seven transformations considered are charge conjugation C, parity inversion P, time reversal T, and their combinations CP, CT, PT, and CPT. The first column specifies the operator by indicating its corresponding coefficient. Each of the other columns concerns one of the seven transformations. An even operator is indicated by a plus sign and an odd one by a minus sign. The table contains eight rows, one for each of the eight possible combinations of signs under C,

P, and T.

Tables P41 and P42 concern the 44 combinations of coefficients for Lorentz violation that frequently appear in experimental analyses involving the fermion sector of the minimal QED extension in Minkowski spacetime in the nonrelativistic limit. These combinations are conventionally denoted by tilde coefficients, and they are defined by the first two columns of Table P41. Note that six of these combinations,  $\tilde{c}_X$ ,  $\tilde{c}_Y$ ,  $\tilde{c}_Z$ ,  $\tilde{g}_{TX}$ ,  $\tilde{g}_{TY}$ , and  $\tilde{g}_{TZ}$ , are denoted as  $\tilde{c}_{Q,Y}$ ,  $\tilde{c}_{Q,X}$ ,  $\tilde{c}_{XY}$ ,  $\tilde{g}_{Q,Y}$ ,  $\tilde{g}_{Q,X}$ , and  $\tilde{g}_{XY}$ , respectively, in some early publications. The definitions in the table are given for a generic fermion of mass  $m$ . Most applications in the literature involve electrons, protons, neutrons, and their antiparticles, for which the corresponding mass is understood. The final column lists the number of independent components of each coefficient. For matter involving electrons, protons, neutrons, and their antiparticles, there are therefore 132 independent observable coefficients for Lorentz violation in the minimal QED sector of the SME in Minkowski spacetime.

The 44 tilde coefficients in Table P41 are formed from 44 linear combinations of the 76 available coefficients in the minimal SME restricted to a single free fermion in flat spacetime. In this limit, the other 32 degrees of freedom can be removed by field redefinitions. Table P42 lists each of these 44 linear combinations and gives its equivalent in terms of tilde coefficients. For brevity, some of the 44 linear combinations are expressed using the definitions

$$\begin{aligned} d_{\mu\nu}^{\pm} &= \frac{1}{2}(d_{\mu\nu} \pm d_{\nu\mu}), & g^{(A)\mu} &= \frac{1}{6}\epsilon^{\alpha\beta\gamma\mu}g_{\alpha\beta\gamma}, \\ g_{\mu\nu\lambda}^{(M)} &= \frac{1}{3}(g_{\mu\nu\lambda} + g_{\mu\lambda\nu} + \eta_{\mu\lambda}g_{\nu\alpha\beta}\eta^{\alpha\beta}) - (\mu \leftrightarrow \nu). \end{aligned}$$

The map between the two coefficient sets can be block-diagonalized [28], and this is reflected in Table P42 by horizontal lines separating each block. The number of independent degrees of freedom in each block is provided in the third column of the table.

Table P43 presents definitions for certain combinations of the 23 coefficients for Lorentz violation in the photon sector of the minimal QED extension in Minkowski spacetime. This table has three sections. The first section consists of five rows listing 19 widely used combinations of the 19 coefficients for CPT-even Lorentz violation. The second section provides 10 alternative combinations involving the 10 CPT-even Lorentz-violating operators relevant to leading-order birefringence [11]. The third section lists four combinations of the four coefficients for CPT-odd Lorentz violation. These combinations appear when a basis of spin-weighted spherical harmonics is adopted.

## B. Minimal SME

Table P44 concerns the fermion-sector terms in the Lagrange density of the minimal SME in Riemann-Cartan spacetime. The column headings are similar to those in Table P39. In the lepton sector, the left- and right-handed leptons are denoted by  $L_A$  and  $R_A$ , where  $A$  is the generation index. The  $SU(2)$  doublet  $L_A$  includes the three neutrino fields  $\nu_e$ ,  $\nu_\mu$ ,  $\nu_\tau$  and the left-handed components of the three charged leptons  $e$ ,  $\mu$ , and  $\tau$ . The  $SU(2)$  singlet  $R_A$  contains the right-handed components of the three charged leptons  $e$ ,  $\mu$ , and  $\tau$ . The derivative  $D_\mu$  is both spacetime and  $SU(3) \times SU(2) \times U(1)$  covariant. The quark fields are denoted  $U_A$ ,  $D_A$ , and  $Q_A$ , where  $A$  is the generation index. The right-handed components of the  $u$ ,  $c$ , and  $t$  quarks are the  $SU(2)$  singlets  $U_A$ , while the right-handed components of  $d$ ,  $s$ , and  $b$  are the  $SU(2)$  singlets  $D_A$ . The six left-handed quark fields are contained in the  $SU(2)$  doublet  $Q_A$ . The Yukawa sector involves terms coupling the Higgs doublet  $\phi$  to the leptons and to the quarks. The conventional Yukawa-coupling matrices are denoted  $(G_L)_{AB}$ ,  $(G_U)_{AB}$ , and  $(G_D)_{AB}$ . The hermitian conjugate of an operator is abbreviated h.c. in the table.

Table P45 presents information about the Higgs, gauge, and pure-gravity sectors for the Lagrange density of the minimal SME in Riemann-Cartan spacetime. The structure of the table is the same as that of Table P44. As before,  $D_\mu$  is both a spacetime and an  $SU(3) \times SU(2) \times U(1)$  covariant derivative. The complex Higgs field is denoted  $\phi$ , the  $SU(3)$  color gauge fields and the  $SU(2)$  gauge fields are the hermitian adjoint matrices  $G_\mu$  and  $W_\mu$ , respectively, while the  $U(1)$  hypercharge gauge field is the singlet  $B_\mu$ . Each gauge field has an associated field strength, denoted  $G_{\mu\nu}$  for the strong interactions,  $W_{\mu\nu}$  for the weak interactions, and  $B_{\mu\nu}$  for the hypercharge. The pure-gravity sector of Table P45 differs from that in Table P39 only in the addition of terms involving the torsion field  $T^\alpha_{\mu\nu}$ .

The minimal SME in Riemann-Cartan spacetime described in Tables P44 and P45 can be reduced to the minimal QED in Riemann spacetime described in Table P39 as follows. For the gauge sector, including the covariant derivatives, remove all the gauge fields except the charge  $U(1)$  field in the photon limit  $B_\mu \rightarrow A_\mu$ , and remove all the Higgs terms. For the gravity sector, remove all the torsion terms. For the fermion sector, restrict the lepton generation index to a single value, remove all quark and neutrino terms, and replace the Yukawa-coupling terms with the relevant fermion mass terms.

Table P46 concerns the piece of the neutrino sector in the SME that contains operators of renormalizable dimension, including both neutrino masses and Lorentz-violating terms. We restrict attention to three generations of active neutrinos and antineutrinos, allowing for possible violations of  $SU(3) \times SU(2) \times U(1)$  gauge sym-

metry and lepton number [29]. In the table, the first row involves the usual neutrino mass matrix  $(m_l m_l^\dagger)_{ab}$ , where the indices  $a, b$  take values  $e, \mu$ , and  $\tau$ . The next four rows concern coefficients for Lorentz violation that enter without mass in the relativistic limit of the effective hamiltonian for neutrino propagation and mixing, while the final five rows list coefficients that involve combinations of mass and Lorentz-violating effects [15]. The first column labels the coefficient types, the second column lists the coefficients, and the third gives the dimensions of the corresponding operators. The fourth column indicates generically the type of neutrino oscillations controlled by the coefficients. The final two columns list the properties of the operators under CPT and Lorentz transformations.

Table P47 provides the relationships between the spherical coefficients for neutrino operators of renormalizable dimension listed in Table P52 and the cartesian ones listed in Table P46. The spherical coefficients for Lorentz violation arise when a basis of spin-weighted spherical harmonics is chosen [15]. Table P47 is split into six sections according to the dimension and CPT properties of the operators, with each section having three columns. The first column lists the spherical coefficients, the second shows the corresponding combination of cartesian components, and the third shows the number of independent spherical coefficients appearing in each section. On each spherical coefficient, superscript flavor-space indices  $ab$  are understood. To keep the expressions compact, some of the combinations are expressed using cartesian components of the complex vector  $\hat{X}_\pm = \hat{X} \mp i\hat{Y}$ . The counting of independent spherical coefficients is obtained by taking into account hermiticity, symmetry, or antisymmetry conditions in flavor space. In total, 369 independent spherical coefficients exist for neutrino operators of renormalizable dimension.

### C. Nonminimal sectors

Tables P48 and P49 present information about the nonminimal quadratic sector for a single Dirac fermion in Minkowski spacetime [18]. Contributions to the Lagrange density at arbitrary  $d$  are provided in Table P48. The structure of this table is similar to that adopted for Tables P39, P44, and P45, with each row associated with a term in the Lagrange density. The entries in the first two columns display the coefficients for Lorentz violation in the cartesian basis and the number of independent degrees of freedom they contain. The third column shows the hermitian operator to which each coefficient corresponds, while the fourth gives the operator mass dimension  $d$ . The conventional multiplicative factor for the coefficient and the operator is given in the next column. The final two columns specify the CPT handedness and the Lorentz property of the operator, using the same

conventions as Table P39.

Table P48 is split into two pieces by a horizontal line. The entries above the line represent conventional Lorentz-invariant contributions of renormalizable dimension to the Lagrange density. The row containing  $m_5$  is included for generality, although in the absence of a chiral anomaly the corresponding term in the Lagrange density can be absorbed into the usual mass  $m$  via a chiral transformation. The entries below the horizontal line concern operators of arbitrary mass dimensions  $d$ . Most of these operators are Lorentz violating, but a few trace components are Lorentz invariant. The value of  $d$  for each coefficient is indicated by a superscript, and the coefficient itself has mass dimension  $4 - d$ . For  $d = 3$  and  $d = 4$ , the entries reduce to those of the minimal theory given in Table P39. Note that none of the operators associated with the coefficients  $m^{(d)\alpha_1 \dots \alpha_{d-3}}$  and  $m_5^{(d)\alpha_1 \dots \alpha_{d-3}}$  are of renormalizable dimension.

Table P49 contains properties of the spherical coefficients for Lorentz violation in the nonminimal quadratic sector for a single Dirac fermion in Minkowski spacetime. The spherical coefficients have comparatively simple transformation properties under rotations. The table has six sections separated by horizontal lines, each representing a specific theoretical scenario labeled in the first column. The second and third columns contain the spherical coefficients and the dimensions  $d$  at which they occur. The next two columns show the allowed ranges of the subscripts on the coefficients, while the final column lists the number of independent components appearing in each coefficient.

Allowing for operators of arbitrary mass dimension, the general effective hamiltonian describing the propagation of a Dirac fermion is formed from the eight sets of spherical coefficients that are listed in the first section of Table P49. The subset of these coefficients that is isotropic in a fixed inertial frame is given in the second section of this table. The superscript  $d$  denotes the mass dimension of the corresponding operator, while the subscripts  $n, j, m$  specify the energy dependence, the total angular momentum, and the azimuthal component of the angular momentum, respectively. The superscripts  $E$  and  $B$  fix the parity of the corresponding operator, while the numerals 0, 1, or 2 preceding  $E$  or  $B$  provide its spin weight. For a nonrelativistic Dirac fermion, only certain linear combinations of coefficients appear in the effective hamiltonian, and the physics is governed by the eight smaller sets of nonrelativistic coefficients listed in the third section of the table. The isotropic restriction of these coefficients is shown in the fourth section. Similarly, only a subset of the general spherical coefficients appears in the effective hamiltonian for a high-energy fermion, and these ultrarelativistic coefficients and their isotropic restriction are displayed in the final two sections of the table.

Table P50 and P51 provide information about the non-



minimal photon sector of the full SME in Minkowski spacetime. The relevant part of the Lagrange density includes operators of arbitrary dimension  $d$  that are both gauge invariant and quadratic in the photon field  $A_\mu$  [14], and these are listed in Table P50. This table is organized along the same lines as Table P48. The first column lists the coefficient for Lorentz violation, while the second column counts its independent components. The next three columns provide the corresponding operator appearing in the Lagrange density, its mass dimension, and the factor contracting the coefficient and the operator. The last two columns list the properties of the operator under CPT and Lorentz transformations, using the same conventions as Table P48.

Three sections appear in Table P50, separated by horizontal lines. The first section concerns the conventional Lorentz-preserving Maxwell term in the Lagrange density for the photon sector. The second and third sections concern coefficients associated with operators of odd and even dimensions  $d$ , respectively. Each of these sections has three rows for the lowest three values of  $d$ , along with a final row applicable to the case of general  $d$ . The notation for the coefficients incorporates a superscript specifying the dimension  $d$  of the corresponding operator. Note that the mass dimension of the coefficients is  $4 - d$ . In each section, the first row describes terms in the minimal SME, and the match is provided between the general notation for nonminimal coefficients and the standard notation used for the minimal SME in Table P39. In the case of mass dimension four, there are 19 independent Lorentz-violating operators. However, for this case the number in the second column is listed as  $19 + 1$  to allow for an additional Lorentz-preserving trace term, which maintains consistency with the expression for general  $d$  in the last row.

Table P51 summarizes properties of spherical coefficients for Lorentz violation in the nonminimal photon sector of the full SME in Minkowski spacetime [14]. The spherical coefficients are combinations of the coefficients listed in Table P50 that are of particular relevance for observation and experiment. They can be separated into two types. One set consists of vacuum coefficients that control leading-order effects on photon propagation in the vacuum, including birefringence and dispersion. The complementary set contains the vacuum-orthogonal coefficients, which leave photon propagation in the vacuum unaffected at leading order. The two parts of Table P51 reflect this separation, with the part above the horizontal line involving the vacuum coefficients and the part below involving the vacuum-orthogonal ones.

In Table P51, the first column of the table identifies the type of spherical coefficients, while the second column lists the specific coefficient. The coefficient notation reflects properties of the corresponding operator. Coefficients associated with operators leaving unchanged the leading-order photon propagation in the vacuum are dis-

tinguished by a negation diacritic  $\bar{\phantom{x}}$ . A symbol  $k$  denotes coefficients for birefringent operators, while  $c$  denotes nonbirefringent ones. The superscript  $d$  refers to the operator mass dimension, while the subscripts  $n, j, m$  determine the frequency or wavelength dependence, the total angular momentum, and the  $z$ -component of the angular momentum, respectively. The superscripts  $E$  and  $B$  refer to the parity of the operator, while the numerals 0, 1, or 2 preceding  $E$  or  $B$  refer to the spin weight. Note that the photon-sector coefficients in the minimal SME correspond to the vacuum coefficients with  $d = 3, 4$ . The third, fourth, and fifth columns of Table P51 provide the allowed ranges of the dimension  $d$  and of the indices  $n$  and  $j$ . The index  $m$  can take values ranging from  $-j$  to  $j$  in unit increments. The final column gives the number of independent coefficient components for each operator of dimension  $d$ .

Table P52 provides a guide to spherical coefficients for Lorentz violation in the neutrino sector of the full SME [15]. Use of these coefficients simplifies analyses in many circumstances because their rotation properties are comparatively simple. The table is separated by horizontal lines into seven sections, reflecting the various theoretical scenarios listed in the first column. The spherical coefficients relevant to each scenario are given in the second column. The third column shows the allowed values of the corresponding operator dimensions, while the fourth column provides the permissible range of the angular-momentum quantum number  $j$ . The last two columns display the number of independent coefficients involved and the CPT properties of the corresponding operators.

The effective hamiltonian governing the propagation and mixing of neutrinos in the presence of Lorentz-violating operators of arbitrary dimensions contains four sets of effective spherical coefficients, listed in the first section of Table P52. The observables in a given experiment lie among these effective coefficients. In the leading-order relativistic limit, they are formed from combinations of the ten sets of basic spherical coefficients displayed in the second section of the table. Six of these basic sets involve Dirac-type operators coupling neutrinos to neutrinos and antineutrinos to antineutrinos, while four involve Majorana-type operators coupling neutrinos to antineutrinos. The remainder of the sections in the table concern various limiting cases of the general formalism. The spherical coefficients associated with operators of renormalizable dimension are shown in the third section. Their relation to the cartesian coefficients listed in Table P46 is given in Table P47. The fourth section of Table P52 contains the coefficients appearing in the massless limit of the SME. The remaining three sections concern the special limits of flavor-blind and oscillation-free operators, of simultaneously diagonalizable operators, and of various types of isotropic operators. Constraints on many of the coefficients listed in these sections have been obtained and are given in the data tables.

## Acknowledgments

This work was supported in part by the U.S. Department of Energy under grant DE-SC0010120 and by the Indiana University Center for Spacetime Symmetries.

- 
- [1] V.A. Kostelecký and R. Potting, Phys. Rev. D **51**, 3923 (1995) [hep-ph/9501341].
- [2] D. Colladay and V.A. Kostelecký, Phys. Rev. D **55**, 6760 (1997) [hep-ph/9703464]; Phys. Rev. D **58**, 116002 (1998) [hep-ph/9809521].
- [3] V.A. Kostelecký, Phys. Rev. D **69**, 105009 (2004) [hep-th/0312310].
- [4] O.W. Greenberg, Phys. Rev. Lett. **89**, 231602 (2002) [hep-ph/0201258].
- [5] V.A. Kostelecký, ed., *CPT and Lorentz Symmetry I, II, III, IV, V, VI*, World Scientific, Singapore 1999, 2002, 2005, 2008, 2011, 2014.
- [6] R. Bluhm, Lect. Notes Phys. **702**, 191 (2006) [hep-ph/0506054].
- [7] J.D. Tasson, Rept. Prog. Phys. **77**, 062901 (2014) [arXiv:1403.7785].
- [8] T. Adam *et al.*, OPERA Collaboration, JHEP **1210**, 093 (2012) [arXiv:1109.4897].
- [9] N.Yu. Agafonova *et al.*, LVD and OPERA Collaborations, Eur. Phys. J. Plus **127**, 71 (2012) [arXiv:1206.2488].
- [10] R. Bluhm *et al.*, Phys. Rev. Lett. **88**, 090801 (2002) [hep-ph/0111141]; Phys. Rev. D **68**, 125008 (2003) [hep-ph/0306190].
- [11] V.A. Kostelecký and M. Mewes, Phys. Rev. D **66**, 056005 (2002) [hep-ph/0205211].
- [12] R. Bluhm *et al.*, Phys. Rev. Lett. **82**, 2254 (1999) [hep-ph/9810269].
- [13] V.A. Kostelecký and M. Mewes, Phys. Rev. Lett. **99**, 011601 (2007) [astro-ph/0702379].
- [14] V.A. Kostelecký and M. Mewes, Phys. Rev. D **80**, 015020 (2009) [arXiv:0905.0031].
- [15] V.A. Kostelecký and M. Mewes, Phys. Rev. D **85**, 096005 (2012) [arXiv:1112.6395].
- [16] V.A. Kostelecký and J.D. Tasson, Phys. Rev. Lett. **102**, 010402 (2009) [arXiv:0810.1459].
- [17] Q. Bailey, V.A. Kostelecký, and R. Xu, Phys. Rev. D **91**, 022006 (2015) [arXiv:1410.6162].
- [18] V.A. Kostelecký and M. Mewes, Phys. Rev. D **88**, 096006 (2013) [arXiv:1308.4973].
- [19] V.A. Kostelecký and M. Mewes, Astrophys. J. Lett. **689**, L1 (2008) [arXiv:0809.2846].
- [20] A.H. Gomes, V.A. Kostelecký, and A.J. Vargas, Phys. Rev. D **90**, 076009 (2014) [arXiv:1407.7748].
- [21] R. Utiyama, Phys. Rev. **101**, 1597 (1956); T.W.B. Kibble, J. Math. Phys. **2**, 212 (1961).
- [22] V.A. Kostelecký, N. Russell, and J.D. Tasson, Phys. Rev. Lett. **100**, 111102 (2008) [arXiv:0712.4393].
- [23] R. Lehnert, W.M. Snow, and H. Yan, Phys. Lett. B **730**, 353 (2014) [arXiv:1311.0467].
- [24] B. Altschul, J. Phys. A **39**, 13757 (2006) [hep-th/0602235].
- [25] V.A. Kostelecký and N. Russell, Phys. Lett. B **693**, 443 (2010) [arXiv:1008.5062].
- [26] V.A. Kostelecký and J. Tasson, Phys. Rev. D **83**, 016013 (2011) [arXiv:1006.4106].
- [27] V.A. Kostelecký, C.D. Lane, and A.G.M. Pickering, Phys. Rev. D **65**, 056006 (2002) [hep-th/0111123].
- [28] A. Fittante and N. Russell, J. Phys. G **39**, 125004 (2012) [arXiv:1210.2003].
- [29] V.A. Kostelecký and M. Mewes, Phys. Rev. D **69**, 016005 (2004) [hep-ph/0309025].
- [30] V.A. Kostelecký and A.J. Vargas, Phys. Rev. D **92**, 056002 (2015) [arXiv:1506.01706].
- [31] B.M. Roberts *et al.*, Phys. Rev. Lett. **113**, 081601 (2014) [arXiv:1404.2723].
- [32] B.M. Roberts *et al.*, Phys. Rev. D **90**, 096005 (2014) [arXiv:1409.2564].
- [33] B.R. Heckel *et al.*, Phys. Rev. D **78**, 092006 (2008) [arXiv:0808.2673].
- [34] T.W. Kornack, G. Vasilakis, and M.V. Romalis, in Ref. [5], vol. *IV*.
- [35] B.R. Heckel *et al.*, Phys. Rev. Lett. **97**, 021603 (2006) [hep-ph/0606218].
- [36] L.-S. Hou, W.-T. Ni, and Y.-C.M. Li, Phys. Rev. Lett. **90**, 201101 (2003) [physics/0009012].
- [37] L. Hunter *et al.*, in Ref. [5], vol. *I*.
- [38] H. Dehmelt *et al.*, Phys. Rev. Lett. **83**, 4694 (1999) [hep-ph/9906262].
- [39] R.K. Mittleman *et al.*, Phys. Rev. Lett. **83**, 2116 (1999).
- [40] C.J. Berglund *et al.*, Phys. Rev. Lett. **75**, 1879 (1995).
- [41] V.A. Kostelecký and C.D. Lane, Phys. Rev. D **60**, 116010 (1999) [hep-ph/9908504].
- [42] S.R. Coleman and S.L. Glashow, Phys. Lett. B **405**, 249 (1997) [hep-ph/9703240].
- [43] S.R. Coleman and S.L. Glashow, Phys. Rev. D **59**, 116008 (1999) [hep-ph/9812418].
- [44] F.W. Stecker and S.L. Glashow, Astropart. Phys. **16**, 97 (2001) [astro-ph/0102226].
- [45] T. Jacobson, S. Liberati and D. Mattingly, Phys. Rev. D **67**, 124011 (2003) [hep-ph/0211466].
- [46] F.W. Stecker, Astropart. Phys. **56**, 16 (2014) [arXiv:1306.6095].
- [47] T. Pruttivarasin *et al.*, Nature **517**, 592 (2015) [arXiv:1412.2194].
- [48] B. Botermann *et al.*, Phys. Rev. Lett. **113**, 120405 (2014) [arXiv:1409.7951]; Phys. Rev. Lett. **114**, 239902 (2015).
- [49] A. Matveev *et al.*, Phys. Rev. Lett. **110**, 230801 (2013).
- [50] M.A. Hohensee *et al.*, Phys. Rev. Lett. **111**, 050401 (2013) [arXiv:1303.2747].
- [51] M.A. Hohensee, H. Müller, and R.B. Wiringa, Phys. Rev. Lett. **111**, 151102 (2013) [arXiv:1308.2936].
- [52] M.A. Hohensee *et al.*, Phys. Rev. Lett. **106**, 151102 (2011) [arXiv:1102.4362].
- [53] B. Altschul, Phys. Rev. D **82**, 016002 (2010) [arXiv:1005.2994].
- [54] B. Altschul, Phys. Rev. D **81**, 041701 (2010) [arXiv:0912.0530].
- [55] H. Müller *et al.*, Phys. Rev. Lett. **99**, 050401 (2007) [arXiv:0706.2031].
- [56] H. Müller, Phys. Rev. D **71**, 045004 (2005) [hep-ph/0412385].
- [57] H. Müller *et al.*, Phys. Rev. D **68**, 116006 (2003) [hep-ph/0401016].
- [58] C.D. Lane, Phys. Rev. D **72**, 016005 (2005) [hep-ph/0505130].

- [59] B. Altschul, Phys. Rev. D **74**, 083003 (2006) [hep-ph/0608332].
- [60] B. Altschul, Astropart. Phys. **28**, 380 (2007) [hep-ph/0610324].
- [61] B. Altschul, Phys. Rev. D **75**, 041301(R) (2007) [hep-ph/0612288].
- [62] M. Schreck, arXiv:1512.04299.
- [63] T.J. Konopka and S.A. Major, New J. Phys. **4**, 57 (2002) [hep-ph/0201184].
- [64] T. Jacobson, S. Liberati and D. Mattingly, Nature **424**, 1019 (2003) [astro-ph/0212190].
- [65] O. Gagnon and G.D. Moore, Phys. Rev. D **70**, 065002 (2004) [hep-ph/0404196].
- [66] L. Maccione, S. Liberati, A. Celotti and J.G. Kirk, JCAP **0710**, 013 (2007). [arXiv:0707.2673].
- [67] B. Altschul, Phys. Rev. D **83**, 056012 (2011) [arXiv:1010.2779].
- [68] Y.V. Stadnik and V.V. Flambaum, Eur. Phys. J. C **75**, 110 (2015) [arXiv:1408.2184].
- [69] S.K. Peck *et al.*, Phys. Rev. A **86**, 012109 (2012) [arXiv:1205.5022].
- [70] J.M. Brown *et al.*, Phys. Rev. Lett. **105**, 151604 (2010) [arXiv:1006.5425]; in Ref. [5], vol. V.
- [71] M.A. Humphrey *et al.*, Phys. Rev. A **68**, 063807 (2003) [physics/0103068].
- [72] D.F. Phillips *et al.*, Phys. Rev. D **63**, 111101 (2001) [physics/0008230].
- [73] F.W. Stecker and S.T. Scully, New J. Phys. **11**, 085003 (2009) [arXiv:0906.1735].
- [74] J. Alfaro and G.A. Palma, hep-th/0501116.
- [75] P. Wolf *et al.*, Phys. Rev. Lett. **96**, 060801 (2006) [hep-ph/0601024].
- [76] S. Ulmer *et al.*, Nature **524**, 196 (2015).
- [77] G. Gabrielse *et al.*, Phys. Rev. Lett. **82**, 3198 (1999).
- [78] L. Maccione, A.M. Taylor, D.M. Mattingly and S. Liberati, JCAP **0904**, 022 (2009) [arXiv:0902.1756].
- [79] F. Allmendinger *et al.*, Phys. Rev. Lett. **112**, 110801 (2014). [arXiv:1312.3225]; in Ref. [5], vol. VI [arXiv:1307.5604].
- [80] C. Gemmel *et al.*, Phys. Rev. D **82**, 111901(R) (2010) [arXiv:1011.2143].
- [81] K. Tullney *et al.*, in Ref. [5], vol. V [arXiv:1008.0579].
- [82] I. Altarev *et al.*, Phys. Rev. Lett. **103**, 081602 (2009) [arXiv:0905.3221].
- [83] V. Flambaum, S. Lambert, and M. Pospelov, Phys. Rev. D **80**, 105021 (2009) [arXiv:0902.3217].
- [84] B. Altschul, Phys. Rev. D **79**, 061702(R) (2009) [arXiv:0901.1870].
- [85] F. Canè *et al.*, Phys. Rev. Lett. **93**, 230801 (2004) [physics/0309070].
- [86] D. Bear *et al.*, Phys. Rev. Lett. **85**, 5038 (2000) [physics/0007049]; **89**, 209902 (2002).
- [87] A. Lo *et al.*, arXiv:1412.2142.
- [88] M. Smiciklas *et al.*, Phys. Rev. Lett. **107**, 171604 (2011) [arXiv:1106.0738].
- [89] B. Altschul, Phys. Rev. D **78**, 085018 (2008) [arXiv:0805.0781].
- [90] B. Altschul, Phys. Rev. D **75**, 023001 (2007) [hep-ph/0608094].
- [91] J.D. Prestage *et al.*, Phys. Rev. Lett. **54**, 2387 (1985).
- [92] S.K. Lamoreaux *et al.*, Phys. Rev. Lett. **57**, 3125 (1986).
- [93] T.E. Chupp *et al.*, Phys. Rev. Lett. **63**, 1541 (1989).
- [94] M. Mewes, Phys. Rev. D **78**, 096008 (2008) [arXiv:0809.4801].
- [95] S.M. Carroll and G.B. Field, Phys. Rev. Lett. **79**, 2394 (1997) [arXiv:astro-ph/9704263].
- [96] B. Nodland and J.P. Ralston, Phys. Rev. Lett. **78**, 3043 (1997) [arXiv:astro-ph/9704196].
- [97] S.M. Carroll, G.B. Field, and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [98] E. Komatsu *et al.*, Astrophys. J. Suppl. **192**, 18 (2011). [arXiv:1001.4538].
- [99] J.-Q. Xia, H. Li, and X. Zhang, Phys. Lett. B **687**, 129 (2010) [arXiv:0908.1876].
- [100] M.L. Brown *et al.*, QUaD Collaboration, Astrophys. J. **705**, 978 (2009) [arXiv:0906.1003].
- [101] L. Pagano *et al.*, Phys. Rev. D **80**, 043522 (2009) [arXiv:0905.1651].
- [102] E.Y.S. Wu *et al.*, QUaD Collaboration, Phys. Rev. Lett. **102**, 161302 (2009) [arXiv:0811.0618].
- [103] T. Kahniashvili, R. Durrer, and Y. Maravin, Phys. Rev. D **78**, 123006 (2008) [arXiv:0807.2593].
- [104] E. Komatsu *et al.*, WMAP Collaboration, Astrophys. J. Suppl. **180**, 330 (2009) [arXiv:0803.0547].
- [105] J.-Q. Xia *et al.*, Astron. Astrophys. **483**, 715 (2008) [arXiv:0710.3325].
- [106] P. Cabella, P. Natoli, and J. Silk, Phys. Rev. D **76**, 123014 (2007) [arXiv:0705.0810].
- [107] B. Feng *et al.*, Phys. Rev. Lett. **96**, 221302 (2006) [astro-ph/0601095].
- [108] Q. Chen, E. Magoulakis, and S. Schiller, Phys. Rev. D **93**, 022003 (2016).
- [109] M. Nagel *et al.*, Nature Commun. **6**, 8174 (2015) [arXiv:1412.6954].
- [110] M.A. Hohensee *et al.*, Phys. Rev. D **82**, 076001 (2010) [arXiv:1006.1376].
- [111] S. Herrmann *et al.*, Phys. Rev. D **80**, 105011 (2009) [arXiv:1002.1284].
- [112] Ch. Eisele, A.Yu. Nevsky, and S. Schiller, Phys. Rev. Lett. **103**, 090401 (2009).
- [113] S. Herrmann *et al.*, in Ref. [5], vol. IV.
- [114] P.L. Stanwix *et al.*, Phys. Rev. D **74**, 081101(R) (2006) [gr-qc/0609072].
- [115] S. Herrmann *et al.*, Phys. Rev. Lett. **95**, 150401 (2005) [physics/0508097].
- [116] P.L. Stanwix *et al.*, Phys. Rev. Lett. **95**, 040404 (2005) [hep-ph/0506074].
- [117] P. Wolf *et al.*, Phys. Rev. D **70**, 051902 (2004) [hep-ph/0407232].
- [118] H. Müller *et al.*, Phys. Rev. Lett. **91** 020401 (2003) [physics/0305117].
- [119] J.A. Lipa *et al.*, Phys. Rev. Lett. **90**, 060403 (2003) [physics/0302093].
- [120] P. Antonini *et al.*, Phys. Rev. A **72**, 066102 (2005) [physics/0602115].
- [121] F.R. Klinkhamer and M. Risse, Phys. Rev. D **77**, 117901 (2008) [arXiv:0806.4351].
- [122] Y. Michimura *et al.*, Phys. Rev. Lett. **110**, 200401 (2013) [arXiv:1303.6709].
- [123] F.N. Baynes, M.E. Tobar, and A.N. Luiten, Phys. Rev. Lett. **108**, 260801 (2012).
- [124] F. Baynes, A. Luiten, and M. Tobar, Phys. Rev. D **84**, 081101 (2011) [arXiv:1108.5414].
- [125] P. Mohanmurthy, D. Dutta, and A. Narayan, arXiv:1509.08297.
- [126] J.-P. Bocquet *et al.*, Phys. Rev. Lett. **104**, 241601 (2010) [arXiv:1005.5230].
- [127] M. Schreck, in Ref. [5], vol. VI [arXiv:1310.5159].

- [128] M.E. Tobar *et al.*, Phys. Rev. D **80**, 125024 (2009) [arXiv:0909.2076].
- [129] B. Altschul, Phys. Rev. D **80** 091901(R) (2009) [arXiv:0905.4346].
- [130] M.A. Hohensee *et al.*, Phys. Rev. D **80** 036010 (2009) [arXiv:0809.3442]; Phys. Rev. Lett. **102**, 170402 (2009) [arXiv:0904.2031].
- [131] F.R. Klinkhamer and M. Schreck, Phys. Rev. D **78**, 085026 (2008) [arXiv:0809.3217].
- [132] S. Reinhardt *et al.*, Nature Physics **3**, 861 (2007).
- [133] M. Hohensee *et al.*, Phys. Rev. D **75**, 049902 (2007) [hep-ph/0701252]; M. Tobar *et al.*, Phys. Rev. D **71**, 025004 (2005) [hep-ph/0408006].
- [134] C.D. Carone, M. Sher, and M. Vanderhaeghen, Phys. Rev. D **74**, 077901 (2006) [hep-ph/0609150].
- [135] J.P. Cotter and B.T.H. Varcoe, physics/0603111.
- [136] S. Parker *et al.*, Phys. Rev. Lett. **106**, 180401 (2011) [arXiv:1102.0081].
- [137] V.A. Kostelecký and M. Mewes, Phys. Rev. Lett. **110**, 201601 (2013) [arXiv:1301.5367].
- [138] Q. Exirifard, Phys. Lett. B **699**, 1 (2011) [arXiv:1010.2054].
- [139] V.A. Kostelecký and M. Mewes, Phys. Rev. Lett. **97**, 140401 (2006) [hep-ph/0607084].
- [140] K. Toma *et al.*, Phys. Rev. Lett. **109**, 241104 (2012) [arXiv:1208.5288].
- [141] P. Laurent *et al.*, Phys. Rev. D **83**, 121301(R) (2011) [arXiv:1106.1068].
- [142] F.W. Stecker, Astropart. Phys. **35**, 95 (2011) [arXiv:1102.2784].
- [143] G. Gubitosi *et al.*, JCAP **08**, 021 (2009) [arXiv:0904.3201].
- [144] S.R. Parker, M. Mewes, F.N. Baynes, and M.E. Tobar, Phys. Lett. A **379**, 2681 (2015) [arXiv:1508.02490].
- [145] Y. Michimura *et al.*, Phys. Rev. D **88**, 111101(R) (2013) [arXiv:1310.1952].
- [146] F. Kislak and H. Krawczynski, Phys. Rev. D **92**, 045016 (2015) [arXiv:1505.02669].
- [147] V. Vasileiou *et al.*, Phys. Rev. D **87**, 122001 (2013) [arXiv:1305.3463].
- [148] V. Vasileiou, Fermi GBM and LAT Collaborations, in Ref. [5], vol. V [arXiv:1008.2913].
- [149] A.A. Abdo *et al.*, Fermi LAT and GBM Collaborations, Science **323**, 1688 (2009).
- [150] F. Aharonian *et al.*, H.E.S.S. Collaboration, Phys. Rev. Lett. **101**, 170402 (2008) [arXiv:0810.3475].
- [151] J. Albert *et al.*, MAGIC Collaboration, Phys. Lett. B **668**, 253 (2008) [arXiv:0708.2889].
- [152] S.E. Boggs *et al.*, Astrophys. J. Lett. **611**, 77 (2004) [astro-ph/0310307].
- [153] J.P. Noordmans, C.J.G. Onderwater, H.W. Wilschut, and R.G.E. Timmermans, arXiv:1412.3257.
- [154] G.W. Bennett *et al.*, Muon  $g-2$  Collaboration, Phys. Rev. Lett. **100**, 091602 (2008) [arXiv:0709.4670].
- [155] V.W. Hughes *et al.*, Phys. Rev. Lett. **87**, 111804 (2001) [hep-ex/0106103].
- [156] M. Deile *et al.*, Muon  $g-2$  Collaboration, in Ref. [5], vol. II [hep-ex/0110044].
- [157] R. Bluhm, V.A. Kostelecký, and C.D. Lane, Phys. Rev. Lett. **84**, 1098 (2000) [hep-ph/9912451].
- [158] J.S. Díaz, V.A. Kostelecký, and R. Lehnert, Phys. Rev. D **88**, 071902(R) (2013) [arXiv:1305.4636].
- [159] J.B. Albert *et al.*, EXO-200 Collaboration, arXiv:1601.07266.
- [160] K. Abe *et al.*, Super-Kamiokande Collaboration, Phys. Rev. D **91**, 052003 (2015) [arXiv:1410.4267]; R. Wendell, Super-Kamiokande Collaboration, AIP Conf. Proc. **1666**, 100001 (2015) [arXiv:1412.5234].
- [161] T. Katori, MiniBooNE Collaboration, Mod. Phys. Lett. A **27**, 1230024 (2012) [arXiv:1206.6915].
- [162] T. Katori, MiniBooNE Collaboration, in Ref. [5], vol. V [arXiv:1008.0906].
- [163] T. Katori and J. Spitz, in Ref. [5], vol. VI [arXiv:1307.5805].
- [164] P. Adamson *et al.*, MINOS Collaboration, Phys. Rev. D **85**, 031101 (2012) [arXiv:1201.2631].
- [165] Y. Abe *et al.*, Double Chooz Collaboration, Phys. Rev. D **86**, 112009 (2012) [arXiv:1209.5810].
- [166] A.A. Aguilar-Arevalo *et al.*, MiniBooNE Collaboration, Phys. Lett. B **718**, 1303 (2013) [arXiv:1109.3480].
- [167] L.B. Auerbach *et al.*, LSND Collaboration, Phys. Rev. D **72**, 076004 (2005) [hep-ex/0506067].
- [168] J.S. Díaz, arXiv:1506.01936.
- [169] T. Akiri, in Ref. [5], vol. VI [arXiv:1308.2210].
- [170] R. Abbasi *et al.*, IceCube Collaboration, Phys. Rev. D **82**, 112003 (2010) [arXiv:1010.4096].
- [171] P. Adamson *et al.*, MINOS Collaboration, Phys. Rev. Lett. **105**, 151601 (2010) [arXiv:1007.2791].
- [172] P. Adamson *et al.*, MINOS Collaboration, Phys. Rev. Lett. **101**, 151601 (2008) [arXiv:0806.4945].
- [173] V. Barger, D. Marfatia, and K. Whisnant, Phys. Lett. B **653**, 267 (2007) [arXiv:0706.1085].
- [174] M.D. Messier, in Ref. [5], vol. III.
- [175] B. Rebel and S. Mufson, Astropart. Phys. **48**, 78 (2013) [arXiv:1301.4684].
- [176] J.S. Díaz, T. Katori, J. Spitz, and J. Conrad, Phys. Lett. B **727**, 412 (2013) [arXiv:1307.5789].
- [177] J.S. Díaz, V.A. Kostelecký, and M. Mewes, Phys. Rev. D **89**, 043005 (2014) [arXiv:1308.6344].
- [178] F.W. Stecker, S.T. Scully, S. Liberati and D. Mattingly, Phys. Rev. D **91**, 045009 (2015) [arXiv:1411.5889]; F.W. Stecker and S.T. Scully, Phys. Rev. D **90**, 043012 (2014) [arXiv:1404.7025].
- [179] F.W. Stecker, in Ref. [5], vol. VI [arXiv:1307.5212].
- [180] R. Cowsik, S. Nussinov, and U. Sarkar, Phys. Rev. Lett. **107**, 251801 (2011) [arXiv:1110.0241].
- [181] X.-J. Bi, P.-F. Yin, Z.-H. Yu, and Q. Yuan, Phys. Rev. Lett. **107**, 241802 (2011) [arXiv:1109.6667].
- [182] A.G. Cohen and S.L. Glashow, Phys. Rev. Lett. **107**, 181803 (2011) [arXiv:1109.6562].
- [183] M. Antonello *et al.*, ICARUS Collaboration, Phys. Lett. B **711**, 270 (2012) [arXiv:1110.3763].
- [184] B. Altschul, Phys. Rev. D **87**, 096004 (2013) [arXiv:1302.2598].
- [185] B. Altschul, J. Phys. Conf. Ser. **173** 012003 (2009).
- [186] J.S. Díaz, Phys. Rev. D **89**, 036002 (2014) [arXiv:1311.0930].
- [187] D. Babusci *et al.*, KLOE Collaboration, Phys. Lett. B **730**, 89 (2014) [arXiv:1312.6818]; A. De Santis, KLOE Collaboration, in Ref. [5], vol. VI.
- [188] A. Di Domenico, KLOE Collaboration, Found. Phys. **40**, 852 (2010); J. Phys. Conf. Ser. **171**, 012008 (2009); F. Bossi *et al.*, KLOE Collaboration, Riv. Nuov. Cim. **031**, 531 (2008) [arXiv:0811.1929]; M. Testa, KLOE Collaboration, arXiv:0805.1969.
- [189] A. Di Domenico, KLOE Collaboration, in Ref. [5], vol. IV.
- [190] H. Nguyen, KTeV Collaboration, in Ref. [5], vol. II [hep-

- ex/0112046].
- [191] V.A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (1998) [hep-ph/9809572].
- [192] V.A. Kostelecký and R.J. Van Kooten, Phys. Rev. D **82**, 101702(R) (2010) [arXiv:1007.5312].
- [193] J. Link *et al.*, FOCUS Collaboration, Phys. Lett. B **556**, 7 (2003) [hep-ex/0208034].
- [194] J. van Tilburg and M. van Veghel, Phys. Lett. B **742**, 236 (2015) [arXiv:1407.1269].
- [195] B. Aubert *et al.*, BaBar Collaboration, Phys. Rev. Lett. **100**, 131802 (2008) [arXiv:0711.2713].
- [196] B. Aubert *et al.*, BaBar Collaboration, hep-ex/0607103.
- [197] V.M. Abazov *et al.*, Phys. Rev. Lett. **115**, 161601 (2015) [arXiv:1506.04123].
- [198] B. Altschul, Phys. Rev. D **77**, 105018 (2008) [arXiv:0712.1579].
- [199] V.M. Abazov *et al.*, D0 Collaboration, Phys. Rev. Lett. **108**, 261603 (2012) [arXiv:1203.6106].
- [200] D.L. Anderson, M. Sher, and I. Turan, Phys. Rev. D **70**, 016001 (2004) [hep-ph/0403116].
- [201] K.K. Vos *et al.*, Phys. Lett. B **729**, 112 (2014) [arXiv:1401.2245].
- [202] S.E. Müller *et al.*, Phys. Rev. D **88**, 071901(R) (2013) [arXiv:1310.3206].
- [203] J.P. Noordmans, H.W. Wilschut, and R.G.E. Timmermans, Phys. Rev. Lett. **111**, 171601 (2013) [arXiv:1308.5570].
- [204] A. Hees *et al.*, Phys. Rev. D **92**, 064049 (2015) [arXiv:1508.03478].
- [205] J.D. Tasson, Phys. Rev. D **86**, 124021 (2012) [arXiv:1211.4850].
- [206] H. Panjwani, L. Carbone, and C.C. Speake, in Ref. [5], vol. V.
- [207] V.A. Kostelecký and J.D. Tasson, Phys. Lett. B **749**, 551 (2015) [arXiv:1508.07007].
- [208] L. Shao, Phys. Rev. D **90**, 122009 (2014) [arXiv:1412.2320].
- [209] L. Shao, Phys. Rev. Lett. **112**, 111103 (2014) [arXiv:1402.6452].
- [210] Q.G. Bailey, R.D. Everett, and J.M. Overduin, Phys. Rev. D **88**, 102001 (2013) [arXiv:1309.6399].
- [211] J.M. Overduin, R.D. Everett, and Q.G. Bailey, in Ref. [5], vol. VI [arXiv:1307.4826].
- [212] K.-Y. Chung *et al.*, Phys. Rev. D **80**, 016002 (2009) [arXiv:0905.1929].
- [213] H. Müller *et al.*, Phys. Rev. Lett. **100**, 031101 (2008) [arXiv:0710.3768].
- [214] J.B.R. Battat, J.F. Chandler, and C.W. Stubbs, Phys. Rev. Lett. **99**, 241103 (2007) [arXiv:0710.0702].
- [215] D. Bennett, V. Skavysh, and J. Long, in Ref. [5], vol. V [arXiv:1008.3670].
- [216] L. Iorio, Class. Quant. Grav. **29**, 175007 (2012) [arXiv:1203.1859].
- [217] Q.G. Bailey and V.A. Kostelecký, Phys. Rev. D **74**, 045001 (2006) [gr-qc/0603030].
- [218] C.G. Shao *et al.*, Phys. Rev. D **91**, 102007 (2015) [arXiv:1504.03280].
- [219] J.C. Long and V.A. Kostelecký, Phys. Rev. D **91**, 092003 (2015) [arXiv:1412.8362].

Table 1. List of tables

Type	Table	Content	
<b>Summary</b>	S2	Maximal sensitivities for the matter sector	
	S3	Maximal sensitivities for the photon sector	
	S4	Maximal sensitivities for the neutrino sector	
	S5	Maximal sensitivities for the gravity sector	
	<b>Data</b>	D6	Electron sector, $d = 3, 4$
D7		Nonminimal electron sector, $d = 5, 6$	
D8		Nonminimal electron sector, $d \geq 7$	
D9		Proton sector, $d = 3, 4$	
D10		Nonminimal proton sector, $d \geq 5$	
D11		Neutron sector, $d = 3, 4$	
D12		Photon sector, $d = 3$	
D13		Photon sector, $d = 4$	
D14		Nonminimal photon sector, $d = 5$	
D15		Nonminimal photon sector, $d = 6$	
D16		Nonminimal photon sector, $d = 7$	
D17		Nonminimal photon sector, $d = 8$	
D18		Nonminimal photon sector, $d \geq 9$	
D19		Muon sector, $d = 3$	
D20		Muon sector, $d = 4$	
D21		Nonminimal muon sector, $d \geq 5$	
D22		Tau sector, $d \geq 3$	
D23		Neutrino sector, $d = 2$	
D24		Neutrino sector, $d = 3$	
D25		Neutrino sector, $d = 4$	
D26		Nonminimal neutrino sector, $d = 5$	
D27		Nonminimal neutrino sector, $d = 6$	
D28		Nonminimal neutrino sector, $d = 7$	
D29		Nonminimal neutrino sector, $d = 8$	
D30		Nonminimal neutrino sector, $d \geq 9$	
D31		Quark sector, $d = 3$	
D32		Quark sector, $d \geq 4$	
D33		Electroweak sector, $d = 3, 4$	
D34		Gluon sector, $d = 4$	
D35		Gravity sector, $d = 3$	
D36		Gravity sector, $d = 4$	
D37		Nonminimal gravity sector, $d = 5, 6$	
D38		Nonminimal gravity sector, $d \geq 7$	
<b>Properties</b>		P39	Lagrange density for the minimal QED extension in Riemann spacetime
		P40	C, P, T properties for operators for Lorentz violation in QED
		P41	Definitions for the fermion sector of the minimal QED extension
		P42	Inverse combinations for the fermion sector of the minimal QED extension
		P43	Definitions for the photon sector of the minimal QED extension
	P44	Lagrange density for the fermion sector of the minimal SME in Riemann-Cartan spacetime	
	P45	Lagrange density for the boson sector of the minimal SME in Riemann-Cartan spacetime	
	P46	Coefficients in the renormalizable neutrino sector	
	P47	Definitions for the minimal neutrino sector	
	P48	Quadratic Lagrange density for a nonminimal Dirac fermion in Minkowski spacetime	
	P49	Spherical coefficients for a nonminimal Dirac fermion in Minkowski spacetime	
	P50	Quadratic Lagrange density for the nonminimal photon sector in Minkowski spacetime	
	P51	Spherical coefficients for the nonminimal photon sector in Minkowski spacetime	
	P52	Spherical coefficients for the nonminimal neutrino sector in Minkowski spacetime	

## SUMMARY TABLES S2–S5

Table S2. Maximal sensitivities for the matter sector

Coefficient	Electron	Proton	Neutron
$\tilde{b}_X$	$10^{-31}$ GeV	$10^{-33}$ GeV	$10^{-33}$ GeV
$\tilde{b}_Y$	$10^{-31}$ GeV	$10^{-33}$ GeV	$10^{-33}$ GeV
$\tilde{b}_Z$	$10^{-29}$ GeV	$10^{-28}$ GeV	$10^{-29}$ GeV
$\tilde{b}_T$	$10^{-26}$ GeV	$10^{-7}$ GeV	$10^{-26}$ GeV
$\tilde{b}_J^*$ , ( $J = X, Y, Z$ )	$10^{-22}$ GeV	–	–
$\tilde{c}_-$	$10^{-20}$ GeV	$10^{-24}$ GeV	$10^{-28}$ GeV
$\tilde{c}_Q$	$10^{-17}$ GeV	$10^{-21}$ GeV	$10^{-10}$ GeV
$\tilde{c}_X$	$10^{-21}$ GeV	$10^{-25}$ GeV	$10^{-28}$ GeV
$\tilde{c}_Y$	$10^{-21}$ GeV	$10^{-25}$ GeV	$10^{-28}$ GeV
$\tilde{c}_Z$	$10^{-20}$ GeV	$10^{-24}$ GeV	$10^{-29}$ GeV
$\tilde{c}_{TX}$	$10^{-18}$ GeV	$10^{-20}$ GeV	$10^{-5}$ GeV
$\tilde{c}_{TY}$	$10^{-18}$ GeV	$10^{-20}$ GeV	$10^{-5}$ GeV
$\tilde{c}_{TZ}$	$10^{-20}$ GeV	$10^{-20}$ GeV	$10^{-5}$ GeV
$\tilde{c}_{TT}$	$10^{-18}$ GeV	$10^{-11}$ GeV	$10^{-11}$ GeV
$\tilde{d}_+$	$10^{-27}$ GeV	$10^{-7}$ GeV	$10^{-27}$ GeV
$\tilde{d}_-$	$10^{-26}$ GeV	–	$10^{-26}$ GeV
$\tilde{d}_Q$	$10^{-26}$ GeV	$10^{-7}$ GeV	$10^{-26}$ GeV
$\tilde{d}_{XY}$	$10^{-26}$ GeV	–	$10^{-27}$ GeV
$\tilde{d}_{YZ}$	$10^{-26}$ GeV	–	$10^{-26}$ GeV
$\tilde{d}_{ZX}$	$10^{-26}$ GeV	–	–
$\tilde{d}_X$	$10^{-22}$ GeV	$10^{-27}$ GeV	$10^{-28}$ GeV
$\tilde{d}_Y$	$10^{-22}$ GeV	$10^{-27}$ GeV	$10^{-28}$ GeV
$\tilde{d}_Z$	$10^{-19}$ GeV	–	–
$\tilde{H}_{XT}$	$10^{-26}$ GeV	–	$10^{-26}$ GeV
$\tilde{H}_{YT}$	$10^{-26}$ GeV	–	$10^{-26}$ GeV
$\tilde{H}_{ZT}$	$10^{-26}$ GeV	–	$10^{-27}$ GeV
$\tilde{g}_T$	$10^{-27}$ GeV	$10^{-7}$ GeV	$10^{-27}$ GeV
$\tilde{g}_c$	$10^{-26}$ GeV	–	$10^{-27}$ GeV
$\tilde{g}_Q$	–	–	–
$\tilde{g}_-$	–	–	–
$\tilde{g}_{TJ}$ , ( $J = X, Y, Z$ )	–	–	–
$\tilde{g}_{XY}$	$10^{-17}$ GeV	–	–
$\tilde{g}_{YX}$	$10^{-17}$ GeV	–	–
$\tilde{g}_{ZX}$	$10^{-18}$ GeV	–	–
$\tilde{g}_{XZ}$	$10^{-17}$ GeV	–	–
$\tilde{g}_{YZ}$	$10^{-17}$ GeV	–	–
$\tilde{g}_{ZY}$	$10^{-18}$ GeV	–	–
$\tilde{g}_{DX}$	$10^{-22}$ GeV	$10^{-27}$ GeV	$10^{-28}$ GeV
$\tilde{g}_{DY}$	$10^{-22}$ GeV	$10^{-27}$ GeV	$10^{-28}$ GeV
$\tilde{g}_{DZ}$	$10^{-22}$ GeV	–	–



Table S3. Maximal sensitivities for the photon sector

$d = 3$	Coefficient	Sensitivity		
	$k_{(V)00}^{(3)}$	$10^{-43}$ GeV		
	$k_{(V)10}^{(3)}$	$10^{-42}$ GeV		
	$\text{Re } k_{(V)11}^{(3)}$	$10^{-42}$ GeV		
	$\text{Im } k_{(V)11}^{(3)}$	$10^{-42}$ GeV		
$d = 4$	Coefficient	Sensitivity	Coefficient	Sensitivity
	$(\tilde{\kappa}_{e+})^{XY}$	$10^{-32}$	$(\tilde{\kappa}_{e-})^{XY}$	$10^{-17}$
	$(\tilde{\kappa}_{e+})^{XZ}$	$10^{-32}$	$(\tilde{\kappa}_{e-})^{XZ}$	$10^{-17}$
	$(\tilde{\kappa}_{e+})^{YZ}$	$10^{-32}$	$(\tilde{\kappa}_{e-})^{YZ}$	$10^{-17}$
	$(\tilde{\kappa}_{e+})^{XX} - (\tilde{\kappa}_{e+})^{YY}$	$10^{-32}$	$(\tilde{\kappa}_{e-})^{XX} - (\tilde{\kappa}_{e-})^{YY}$	$10^{-17}$
	$(\tilde{\kappa}_{e+})^{ZZ}$	$10^{-32}$	$(\tilde{\kappa}_{e-})^{ZZ}$	$10^{-16}$
	$(\tilde{\kappa}_{o-})^{XY}$	$10^{-32}$	$(\tilde{\kappa}_{o+})^{XY}$	$10^{-13}$
	$(\tilde{\kappa}_{o-})^{XZ}$	$10^{-32}$	$(\tilde{\kappa}_{o+})^{XZ}$	$10^{-14}$
	$(\tilde{\kappa}_{o-})^{YZ}$	$10^{-32}$	$(\tilde{\kappa}_{o+})^{YZ}$	$10^{-14}$
	$(\tilde{\kappa}_{o-})^{XX} - (\tilde{\kappa}_{o-})^{YY}$	$10^{-32}$		
	$(\tilde{\kappa}_{o-})^{ZZ}$	$10^{-32}$	$\tilde{\kappa}_{\text{tr}}$	$10^{-14}$
Isotropic	Coefficient	Sensitivity		
	$k_{(V)00}^{(3)}$	$10^{-43}$ GeV		
	$c_{(I)00}^{(4)} = \sqrt{4\pi}\tilde{\kappa}_{\text{tr}}$	$10^{-14}$		
	$k_{(V)00}^{(5)}$	$10^{-34}$ GeV $^{-1}$		
	$c_{(I)00}^{(6)}$	$10^{-21}$ GeV $^{-2}$		
	$k_{(V)00}^{(7)}$	$10^{-27}$ GeV $^{-3}$		
	$c_{(I)00}^{(8)}$	$10^{-24}$ GeV $^{-4}$		
	$k_{(V)00}^{(9)}$	$10^{-21}$ GeV $^{-5}$		

Table S4. Maximal sensitivities for the neutrino sector

$d = 3$	<b>Coefficient</b>	$e\mu$	$e\tau$	$\mu\tau$	<b>Coefficient</b>	$e\mu$	$e\tau$	$\mu\tau$
	$\text{Re}(a_L)^T$	$10^{-20}$ GeV	$10^{-19}$ GeV	$10^{-24}$ GeV	$\text{Im}(a_L)^T$	–	$10^{-19}$ GeV	–
	$\text{Re}(a_L)^X$	$10^{-20}$ GeV	$10^{-19}$ GeV	$10^{-23}$ GeV	$\text{Im}(a_L)^X$	$10^{-20}$ GeV	$10^{-19}$ GeV	$10^{-20}$ GeV
	$\text{Re}(a_L)^Y$	$10^{-21}$ GeV	$10^{-19}$ GeV	$10^{-23}$ GeV	$\text{Im}(a_L)^Y$	$10^{-21}$ GeV	$10^{-19}$ GeV	$10^{-20}$ GeV
	$\text{Re}(a_L)^Z$	$10^{-19}$ GeV	$10^{-19}$ GeV	–	$\text{Im}(a_L)^Z$	$10^{-19}$ GeV	$10^{-19}$ GeV	–
$d = 4$	<b>Coefficient</b>	$e\mu$	$e\tau$	$\mu\tau$	<b>Coefficient</b>	$e\mu$	$e\tau$	$\mu\tau$
	$\text{Re}(c_L)^{XY}$	$10^{-21}$	$10^{-17}$	$10^{-23}$	$\text{Im}(c_L)^{XY}$	$10^{-21}$	$10^{-17}$	$10^{-21}$
	$\text{Re}(c_L)^{XZ}$	$10^{-21}$	$10^{-17}$	$10^{-23}$	$\text{Im}(c_L)^{XZ}$	$10^{-21}$	$10^{-17}$	$10^{-21}$
	$\text{Re}(c_L)^{YZ}$	$10^{-21}$	$10^{-16}$	$10^{-23}$	$\text{Im}(c_L)^{YZ}$	$10^{-21}$	$10^{-16}$	$10^{-21}$
	$\text{Re}(c_L)^{XX}$	$10^{-21}$	$10^{-16}$	$10^{-23}$	$\text{Im}(c_L)^{XX}$	$10^{-21}$	$10^{-16}$	$10^{-21}$
	$\text{Re}(c_L)^{YY}$	$10^{-21}$	$10^{-16}$	$10^{-23}$	$\text{Im}(c_L)^{YY}$	$10^{-21}$	$10^{-16}$	$10^{-21}$
	$\text{Re}(c_L)^{ZZ}$	$10^{-19}$	$10^{-16}$	–	$\text{Im}(c_L)^{ZZ}$	–	$10^{-16}$	–
	$\text{Re}(c_L)^{TT}$	$10^{-19}$	$10^{-17}$	–	$\text{Im}(c_L)^{TT}$	–	$10^{-17}$	–
	$\text{Re}(c_L)^{TX}$	$10^{-22}$	$10^{-17}$	$10^{-27}$	$\text{Im}(c_L)^{TX}$	$10^{-22}$	$10^{-17}$	$10^{-22}$
	$\text{Re}(c_L)^{TY}$	$10^{-22}$	$10^{-17}$	$10^{-27}$	$\text{Im}(c_L)^{TY}$	$10^{-22}$	$10^{-17}$	$10^{-22}$
	$\text{Re}(c_L)^{TZ}$	$10^{-20}$	$10^{-16}$	–	$\text{Im}(c_L)^{TZ}$	–	$10^{-16}$	–
Isotropic	<b>Coefficient</b>	<b>Sensitivity</b>			<b>Coefficient</b>	<b>Sensitivity</b>		
	$\hat{a}^{(3)}$	$10^{-7}$ GeV			$\hat{a}_{e\mu}^{(3)}$	$10^{-20}$ GeV		
	$\hat{c}^{(4)}$	$10^{-9}$			$\hat{c}_{e\mu}^{(4)}$	$10^{-19}$		
	$\hat{a}^{(5)}$	$10^{-18}$ GeV $^{-1}$			$\hat{a}_{e\mu}^{(5)}$	$10^{-19}$ GeV $^{-1}$		
	$\hat{c}^{(6)}$	$10^{-9}$ GeV $^{-2}$			$\hat{c}_{e\mu}^{(6)}$	$10^{-19}$ GeV $^{-2}$		
	$\hat{a}^{(7)}$	$10^{-29}$ GeV $^{-3}$			$\hat{a}_{e\mu}^{(7)}$	$10^{-19}$ GeV $^{-3}$		
	$\hat{c}^{(8)}$	$10^{-11}$ GeV $^{-4}$			$\hat{c}_{e\mu}^{(8)}$	$10^{-18}$ GeV $^{-4}$		
	$\hat{a}^{(9)}$	$10^{-40}$ GeV $^{-5}$			$\hat{a}_{e\mu}^{(9)}$	$10^{-18}$ GeV $^{-5}$		
	$\hat{c}^{(10)}$	$10^{-14}$ GeV $^{-6}$			$\hat{c}_{e\mu}^{(10)}$	$10^{-18}$ GeV $^{-6}$		

Table S5. Maximal sensitivities for the gravity sector

$d = 3$	Coefficient	Electron	Proton	Neutron
	$\alpha\bar{a}_T$	$10^{-11}$ GeV	$10^{-11}$ GeV	$10^{-11}$ GeV
	$\alpha\bar{a}_X$	$10^{-6}$ GeV	$10^{-6}$ GeV	$10^{-5}$ GeV
	$\alpha\bar{a}_Y$	$10^{-5}$ GeV	$10^{-5}$ GeV	$10^{-4}$ GeV
	$\alpha\bar{a}_Z$	$10^{-5}$ GeV	$10^{-5}$ GeV	$10^{-4}$ GeV
$d = 4$	Coefficient	Electron	Proton	Neutron
	$\alpha\bar{e}_T$	$10^{-8}$	$10^{-11}$	$10^{-11}$
	$\alpha\bar{e}_X$	$10^{-3}$	$10^{-6}$	$10^{-5}$
	$\alpha\bar{e}_Y$	$10^{-2}$	$10^{-5}$	$10^{-4}$
	$\alpha\bar{e}_Z$	$10^{-2}$	$10^{-5}$	$10^{-4}$
	Coefficient	Sensitivity		
	$\bar{s}^{XY}$	$10^{-10}$		
	$\bar{s}^{XZ}$	$10^{-11}$		
	$\bar{s}^{YZ}$	$10^{-11}$		
	$\bar{s}^{XX} - \bar{s}^{YY}$	$10^{-10}$		
	$\bar{s}^{XX} + \bar{s}^{YY} - 2\bar{s}^{ZZ}$	$10^{-10}$		
	$\bar{s}^{TT}$	$10^{-5}$		
	$\bar{s}^{TX}$	$10^{-8}$		
	$\bar{s}^{TY}$	$10^{-8}$		
	$\bar{s}^{TZ}$	$10^{-8}$		

**DATA TABLES D6–D38**

Table D6. Electron sector,  $d = 3, 4$  (part 1 of 3)

Combination	Result	System	Ref.
$ \operatorname{Re} H_{011}^{\text{NR}(0B)} ,  \operatorname{Im} H_{011}^{\text{NR}(0B)} ,  \operatorname{Re} g_{011}^{\text{NR}(0B)} ,  \operatorname{Im} g_{011}^{\text{NR}(0B)} $	$< 9 \times 10^{-27}$ GeV	H maser	[30]*
$ \operatorname{Re} H_{011}^{\text{NR}(1B)} ,  \operatorname{Im} H_{011}^{\text{NR}(1B)} ,  \operatorname{Re} g_{011}^{\text{NR}(1B)} ,  \operatorname{Im} g_{011}^{\text{NR}(1B)} $	$< 5 \times 10^{-27}$ GeV	"	[30]*
$ b_0 $	$< 2 \times 10^{-14}$ GeV	Cs spectroscopy	[31]*, [32]*
"	$< 2 \times 10^{-12}$ GeV	Tl spectroscopy	[31]*, [32]*
"	$< 7 \times 10^{-15}$ GeV	Dy spectroscopy	[31]*, [32]*
"	$< 2 \times 10^{-12}$ GeV	Yb spectroscopy	[32]*
$\tilde{b}_X$	$(-0.9 \pm 1.4) \times 10^{-31}$ GeV	Torsion pendulum	[33]
$\tilde{b}_Y$	$(-0.9 \pm 1.4) \times 10^{-31}$ GeV	"	[33]
$\tilde{b}_Z$	$(-0.3 \pm 4.4) \times 10^{-30}$ GeV	"	[33]
$\frac{1}{2}(\tilde{b}_T + \tilde{d}_- - 2\tilde{g}_c - 3\tilde{g}_T + 4\tilde{d}_+ - \tilde{d}_Q)$	$(0.9 \pm 2.2) \times 10^{-27}$ GeV	"	[33]
$\frac{1}{2}(2\tilde{g}_c - \tilde{g}_T - \tilde{b}_T + 4\tilde{d}_+ - \tilde{d}_- - \tilde{d}_Q)$ $+ \tan \eta(\tilde{d}_{YZ} - \tilde{H}_{XT})$	$(-0.8 \pm 2.0) \times 10^{-27}$ GeV	"	[33]
$\tilde{b}_X$	$(2.8 \pm 6.1) \times 10^{-29}$ GeV	K/He magnetometer	[34]
$\tilde{b}_Y$	$(6.8 \pm 6.1) \times 10^{-29}$ GeV	"	[34]
$\tilde{b}_X$	$(0.1 \pm 2.4) \times 10^{-31}$ GeV	Torsion pendulum	[35]
$\tilde{b}_Y$	$(-1.7 \pm 2.5) \times 10^{-31}$ GeV	"	[35]
$\tilde{b}_Z$	$(-29 \pm 39) \times 10^{-31}$ GeV	"	[35]
$\tilde{b}_\perp$	$< 3.1 \times 10^{-29}$ GeV	"	[36]
$ \tilde{b}_Z $	$< 7.1 \times 10^{-28}$ GeV	"	[36]
$r_e$	$< 3.2 \times 10^{-24}$	Hg/Cs comparison	[37]
$ \vec{b} $	$< 50$ radians/s	Penning trap	[38]
$r_{\omega_a^-, \text{diurnal}}$	$< 1.6 \times 10^{-21}$	"	[39]
$ \tilde{b}_J , (J = X, Y)$	$< 10^{-27}$ GeV	Hg/Cs comparison	[40], [41]*
$ \tilde{b}_J^* , (J = X, Y, Z)$	$< 10^{-22}$ GeV	Penning trap	[28]*

Table D6. Electron sector,  $d = 3, 4$  (part 2 of 3)

Combination	Result	System	Ref.
$\dot{c}^{\text{UR}(4)}$	$< 1.5 \times 10^{-15}$	Astrophysics	[42]*, [18]*
"	$> -5 \times 10^{-13}$	"	[43]*, [18]*
"	$(-1.3 \text{ to } 0.2) \times 10^{-15}$	"	[44]*, [18]*
"	$> -1.2 \times 10^{-16}$	"	[45]*, [18]*
"	$> -6 \times 10^{-20}$	"	[46]*, [18]*
$-0.16c'_{X-Y} + 0.33c'_{XY} - 0.92c'_{XZ} - 0.16c'_{YZ}$	$(0.1 \pm 1.0) \times 10^{-18}$	Trapped Calcium ions	[47]
$-0.04c'_{X-Y} - 0.32c'_{XY} - 0.35c'_{XZ} + 0.88c'_{YZ}$	$(2.4 \pm 7.4) \times 10^{-19}$	"	[47]
$0.29c'_{X-Y} - 0.38c'_{XY} - 0.73c'_{XZ} - 0.48c'_{YZ}$	$(5.9 \pm 9.5) \times 10^{-19}$	"	[47]
$-0.31c'_{X-Y} - 0.65c'_{XY} + 0.07c'_{XZ} - 0.69c'_{YZ}$	$(0.7 \pm 1.2) \times 10^{-18}$	"	[47]
$ \bar{c}_Q $	$< 2 \times 10^{-8}$ GeV	Relativistic Lithium ions	[48]
$c_{(TX)}$	$(3.1 \pm 1.9) \times 10^{-11}$	1S-2S transition	[49]
$0.92c_{(TY)} + 0.40c_{(TZ)}$	$(2.6 \pm 5.3) \times 10^{-11}$	"	[49]
$c_{TT}$	$> -5 \times 10^{-21}$	Astrophysics	[46]*
$0.10c_{X-Y} - 0.99c_{XZ}$	$(-9.0 \pm 11) \times 10^{-17}$	Dysprosium spectroscopy	[50]
$0.99c_{X-Y} + 0.10c_{XZ}$	$(3.8 \pm 5.6) \times 10^{-17}$	"	[50]
$0.94c_{XY} - 0.35c_{YZ}$	$(-0.4 \pm 2.8) \times 10^{-17}$	"	[50]
$0.35c_{XY} + 0.94c_{YZ}$	$(3.2 \pm 7.0) \times 10^{-17}$	"	[50]
$0.18c_{TX} - 0.98c_{T(Y+Z)}$	$(0.95 \pm 18) \times 10^{-15}$	"	[50]
$0.98c_{TX} + 0.18c_{T(Y+Z)}$	$(5.6 \pm 7.7) \times 10^{-15}$	"	[50]
$c_{T(Y-Z)}$	$(-21 \pm 19) \times 10^{-13}$	"	[50]
$c_{TT}$	$(-8.8 \pm 5.1) \times 10^{-9}$	"	[50]
"	$(-14 \pm 28) \times 10^{-9}$	"	[50]
$\bar{c}_{TT}$	$(-0.014 \pm 0.028) \times 10^{-6}$	Nuclear binding energy	[51]
"	$(4.6 \pm 4.6) \times 10^{-6}$	Cs interferometer	[52]
$c_{TT}$	$(-4 \text{ to } 2) \times 10^{-15}$	Collider physics	[53]*
$c_{(TX)}$	$(-30 \text{ to } 1) \times 10^{-14}$	"	[53]*
$c_{(TY)}$	$(-80 \text{ to } 6) \times 10^{-15}$	"	[53]*
$c_{(TZ)}$	$(-11 \text{ to } 1.3) \times 10^{-13}$	"	[53]*
$0.83c_{(TX)} + 0.51c_{(TY)} + 0.22c_{(TZ)}$	$(4 \pm 8) \times 10^{-11}$	1S-2S transition	[54]*
$c_{XX} - c_{YY}$	$(-2.9 \pm 6.3) \times 10^{-16}$	Optical, microwave resonators	[55]*
$\frac{1}{2}c_{(XY)}$	$(2.1 \pm 0.9) \times 10^{-16}$	"	[55]*
$\frac{1}{2}c_{(XZ)}$	$(-1.5 \pm 0.9) \times 10^{-16}$	"	[55]*
$\frac{1}{2}c_{(YZ)}$	$(-0.5 \pm 1.2) \times 10^{-16}$	"	[55]*
$c_{XX} + c_{YY} - 2c_{ZZ}$	$(-106 \pm 147) \times 10^{-16}$	"	[55]*
$\lambda^{ZZ}$	$(13.3 \pm 9.8) \times 10^{-16}$	"	[55]*

Table D6. Electron sector,  $d = 3, 4$  (part 3 of 3)

Combination	Result	System	Ref.
$\frac{1}{2}c_{(YZ)}$	$(2.1 \pm 4.6) \times 10^{-16}$	Optical, microwave resonators	[56]*
$\frac{1}{2}c_{(XZ)}$	$(-1.6 \pm 6.3) \times 10^{-16}$	"	[56]*
$\frac{1}{2}c_{(XY)}$	$(7.6 \pm 3.5) \times 10^{-16}$	"	[56]*
$c_{XX} - c_{YY}$	$(1.15 \pm 0.64) \times 10^{-15}$	"	[56]*
$ c_{XX} + c_{YY} - 2c_{ZZ} - 0.25(\tilde{\kappa}_{e-})^{ZZ} $	$< 10^{-12}$	"	[56]*
$ \frac{1}{2}c_{(XY)} $	$< 8 \times 10^{-15}$	Optical resonators	[57]*
$ c_{XX} - c_{YY} $	$< 1.6 \times 10^{-14}$	"	[57]*
$ c_{XX} + c_{YY} - 2c_{ZZ} $	$< 10^{-5}$	Doppler shift	[58]*
$ c_{TJ} + c_{JT} , (J = X, Y, Z)$	$< 10^{-2}$	"	[58]*
$c_{XX}$	$(-3 \text{ to } 5) \times 10^{-15}$	Astrophysics	[59]*
$c_{YY}$	$(-0.7 \text{ to } 2.5) \times 10^{-15}$	"	[59]*
$c_{ZZ}$	$(-1.6 \text{ to } 2.5) \times 10^{-15}$	"	[59]*
$c_{(YZ)}$	$(-2.5 \text{ to } 1.8) \times 10^{-15}$	"	[59]*
$c_{0X}$	$(-7 \text{ to } 4) \times 10^{-15}$	"	[59]*
$c_{0Y}$	$(-0.5 \text{ to } 1.5) \times 10^{-15}$	"	[59]*
$c_{0Z}$	$(-4 \text{ to } 2) \times 10^{-17}$	"	[59]*
$ 0.05c_{XX} + 0.55c_{YY} + 0.41c_{ZZ}$ $+0.16c_{(XY)} - 0.14c_{(XZ)} - 0.47c_{(YZ)}$ $+0.22c_{(0X)} + 0.74c_{(0Y)} - 0.64c_{(0Z)} + c_{00} $	$< 1.3 \times 10^{-15}$	"	[60]*
$ 0.58c_{XX} + 0.04c_{YY} + 0.38c_{ZZ}$ $-0.14c_{(XY)} - 0.47c_{(XZ)} + 0.12c_{(YZ)}$ $+0.76c_{(0X)} - 0.19c_{(0Y)} - 0.62c_{(0Z)} + c_{00} $	$< 2.5 \times 10^{-15}$	"	[60]*
$\tilde{d}_{XY} - \tilde{H}_{ZT} + \tan \eta \tilde{H}_{YT}$	$(0.1 \pm 1.8) \times 10^{-27}$ GeV	Torsion pendulum	[33]
$\tilde{H}_{ZT}$	$(-4.1 \pm 2.4) \times 10^{-27}$ GeV	"	[33]
$\tilde{H}_{YT} - \tilde{d}_{ZX}$	$(-4.9 \pm 8.9) \times 10^{-27}$ GeV	"	[33]
$-\tilde{H}_{XT} + \tan \eta (\tilde{g}_T - 2\tilde{d}_+ + \tilde{d}_Q)$	$(1.1 \pm 9.2) \times 10^{-27}$ GeV	"	[33]
$ d_{XX} $	$< 2 \times 10^{-14}$	Astrophysics	[61]*
$ d_{YY} ,  d_{ZZ} $	$< 3 \times 10^{-15}$	"	[61]*
$ d_{(XY)} $	$< 2 \times 10^{-15}$	"	[61]*
$ d_{(XZ)} $	$< 2 \times 10^{-14}$	"	[61]*
$ d_{(YZ)} $	$< 7 \times 10^{-15}$	"	[61]*
$ d_{TX} $	$< 5 \times 10^{-14}$	"	[61]*
$ d_{TY} $	$< 5 \times 10^{-15}$	"	[61]*
$ d_{TZ} $	$< 8 \times 10^{-17}$	"	[61]*
$ \tilde{d}_J , (J = X, Y)$	$< 10^{-22}$ GeV	Hg/Cs comparison	[40], [41]*
$ \tilde{d}_Z $	$< 10^{-19}$ GeV	Astrophysics	[28]*
$ \tilde{g}_{XY} ,  \tilde{g}_{XZ} $	$< 10^{-17}$ GeV	Astrophysics	[28]*
$ \tilde{g}_{YZ} ,  \tilde{g}_{YX} $	$< 10^{-17}$ GeV	"	[28]*
$ \tilde{g}_{ZX} ,  \tilde{g}_{ZY} $	$< 10^{-18}$ GeV	"	[28]*
$ \tilde{g}_{D,J} , (J = X, Y, Z)$	$< 10^{-22}$ GeV	Penning trap	[28]*
$ \tilde{g}_{D,J} , (J = X, Y)$	$< 10^{-22}$ GeV	Hg/Cs comparison	[40], [41]*

Table D7. Nonminimal electron sector,  $d = 5, 6$ 

Combination	Result	System	Ref.
$\hat{c}_{e,2}^{\text{NR}} + \frac{67}{12}(\alpha m_r)^2 \hat{c}_{e,4}^{\text{NR}}$	$(4.5 \pm 2.5) \times 10^{-6} \text{ GeV}^{-1}$	Positronium 1S-2S	[30]*
$ \text{Re } H_{211}^{\text{NR}(0B)} ,  \text{Im } H_{211}^{\text{NR}(0B)} ,  \text{Re } g_{211}^{\text{NR}(0B)} ,  \text{Im } g_{211}^{\text{NR}(0B)} $	$< 7 \times 10^{-16} \text{ GeV}^{-1}$	H maser	[30]*
$ \text{Re } H_{211}^{\text{NR}(1B)} ,  \text{Im } H_{211}^{\text{NR}(1B)} ,  \text{Re } g_{211}^{\text{NR}(1B)} ,  \text{Im } g_{211}^{\text{NR}(1B)} $	$< 4 \times 10^{-16} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)TTX} $	$< 3.4 \times 10^{-8} \text{ GeV}^{-1}$	1S-2S transition	[30]*
$ a_{\text{eff}}^{(5)TTY} $	$< 5.6 \times 10^{-8} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)TTZ} $	$< 1.3 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKX} $	$< 6.7 \times 10^{-8} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKY} $	$< 1.1 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKZ} $	$< 2.5 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$\hat{a}_0^{(5)} + \hat{a}_2^{(5)}$	$(-9.4 \text{ to } 11) \times 10^{-2} \text{ GeV}^{-1}$	Electron kinematics	[62]*
"	$> 8.0 \times 10^{-7} \text{ GeV}^{-1}$	Time-of-flight	[62]*
$\hat{a}^{\text{UR}(5)}$	$< 6.5 \times 10^{-27} \text{ GeV}^{-1}$	Astrophysics	[63]*, [18]*
"	$> -3.5 \times 10^{-27} \text{ GeV}^{-1}$	"	[64]*, [18]*
$\hat{a}^{\text{UR}(5)} - m\hat{g}^{\text{UR}(6)}$	$(-1 \text{ to } 1) \times 10^{-34} \text{ GeV}^{-1}$	"	[65]*, [18]*
$\hat{a}^{\text{UR}(5)} \pm m\hat{g}^{\text{UR}(6)}$	$(-4 \text{ to } 4) \times 10^{-25} \text{ GeV}^{-1}$	"	[66]*, [18]*
$\hat{a}^{\text{UR}(5)}$	$(-0.001 \text{ to } 2.8) \times 10^{-17} \text{ GeV}^{-1}$	"	[67]*, [18]*
$ c_{\text{eff}}^{(6)TTTX} $	$< 3.3 \times 10^{-5} \text{ GeV}^{-2}$	1S-2S transition	[30]*
$ c_{\text{eff}}^{(6)TTY} $	$< 5.5 \times 10^{-5} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TTZ} $	$< 1.3 \times 10^{-4} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKX} $	$< 3.3 \times 10^{-5} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKY} $	$< 5.5 \times 10^{-5} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKZ} $	$< 1.3 \times 10^{-4} \text{ GeV}^{-2}$	"	[30]*
$\hat{c}_0^{(6)} + \hat{c}_2^{(6)} + \hat{c}_4^{(6)}$	$-17 \text{ to } 19 \text{ GeV}^{-2}$	Electron kinematics	[62]*
"	$< 4.5 \times 10^{-7} \text{ GeV}^{-2}$	Time-of-flight	[62]*
$\hat{e}_0^{(6)} + \hat{e}_2^{(6)}$	$(-1.8 \text{ to } 2.1) \times 10^2 \text{ GeV}^{-2}$	Electron kinematics	[62]*
"	$< 1.6 \times 10^{-3} \text{ GeV}^{-2}$	Time-of-flight	[62]*
$\hat{f}_0^{(6)} + \hat{f}_2^{(6)}$	$(-1.3 \text{ to } 1.3) \times 10^3 \text{ GeV}^{-2}$	Electron kinematics	[62]*
$\hat{c}^{\text{UR}(6)}$	$(-8.5 \text{ to } 0.0025) \times 10^{-20} \text{ GeV}^{-2}$	Astrophysics	[65]*, [18]*
$\hat{g}^{\text{UR}(6)}$	$(-5.4 \text{ to } 5.4) \times 10^{-14} \text{ GeV}^{-2}$	"	[67]*, [18]*



Table D8. Nonminimal electron sector,  $d \geq 7$ 

Combination	Result	System	Ref.
$ \text{Re } H_{411}^{\text{NR}(0B)} ,  \text{Im } H_{411}^{\text{NR}(0B)} ,  \text{Re } g_{411}^{\text{NR}(0B)} ,  \text{Im } g_{411}^{\text{NR}(0B)} $	$< 9 \times 10^{-6} \text{ GeV}^{-3}$	H maser	[30]*
$ \text{Re } H_{411}^{\text{NR}(1B)} ,  \text{Im } H_{411}^{\text{NR}(1B)} ,  \text{Re } g_{411}^{\text{NR}(1B)} ,  \text{Im } g_{411}^{\text{NR}(1B)} $	$< 5 \times 10^{-6} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTTTX} $	$< 3.9 \times 10^{-2} \text{ GeV}^{-3}$	1S-2S transition	[30]*
$ a_{\text{eff}}^{(7)TTTTY} $	$< 6.5 \times 10^{-2} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTTTZ} $	$< 0.15 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKX} $	$< 2.6 \times 10^{-2} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKY} $	$< 4.3 \times 10^{-2} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKZ} $	$< 0.1 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLX} $	$< 8.7 \times 10^2 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLY} $	$< 1.5 \times 10^3 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLZ} $	$< 3.5 \times 10^3 \text{ GeV}^{-3}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTX} $	$< 51 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTY} $	$< 85 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTZ} $	$< 2.0 \times 10^2 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKX} $	$< 25 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKY} $	$< 43 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKZ} $	$< 98 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLX} $	$< 2.8 \times 10^5 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLY} $	$< 4.7 \times 10^5 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLZ} $	$< 1.1 \times 10^6 \text{ GeV}^{-4}$	"	[30]*

Table D9. Proton sector,  $d = 3, 4$ 

Combination	Result	System	Ref.
$ \text{Re } H_{011}^{\text{NR}(0B)} ,  \text{Im } H_{011}^{\text{NR}(0B)} ,  \text{Re } g_{011}^{\text{NR}(0B)} ,  \text{Im } g_{011}^{\text{NR}(0B)} $	$< 9 \times 10^{-27}$ GeV	H maser	[30]*
$ \text{Re } H_{011}^{\text{NR}(1B)} ,  \text{Im } H_{011}^{\text{NR}(1B)} ,  \text{Re } g_{011}^{\text{NR}(1B)} ,  \text{Im } g_{011}^{\text{NR}(1B)} $	$< 5 \times 10^{-27}$ GeV	"	[30]*
$ \tilde{b}_\perp $	$< 7.6 \times 10^{-33}$ GeV	He/Xe magnetometer	[68]*
$ b_0 $	$< 3 \times 10^{-8}$ GeV	Cs spectroscopy	[31]*
"	$< 7 \times 10^{-8}$ GeV	"	[68]*
"	$< 4 \times 10^{-8}$ GeV	"	[32]*
"	$< 8 \times 10^{-8}$ GeV	Tl spectroscopy	[31]*, [32]*
$\tilde{b}_Z$	$< 7 \times 10^{-29}$ GeV	Hg/Cs comparison	[69]
$\tilde{b}_\perp$	$< 4 \times 10^{-30}$ GeV	"	[69]
$\tilde{b}_\perp$	$< 6 \times 10^{-32}$ GeV	K/He magnetometer	[70]
$\tilde{b}_X$	$(6.0 \pm 1.3) \times 10^{-31}$ GeV	"	[34]
$\tilde{b}_Y$	$(1.5 \pm 1.2) \times 10^{-31}$ GeV	"	[34]
$\sqrt{(\tilde{b}_X^e + \tilde{b}_X^p)^2 + (\tilde{b}_Y^e + \tilde{b}_Y^p)^2}$	$(3 \pm 2) \times 10^{-27}$ GeV	H maser	[71]
$ \tilde{b}_J , (J = X, Y)$	$< 2 \times 10^{-27}$ GeV	"	[72]
"	$< 10^{-27}$ GeV	Hg/Cs comparison	[40], [41]*
$\tilde{c}^{\text{UR}(4)}$	$> -5 \times 10^{-23}$	Astrophysics	[42]*, [18]*
"	$< 5 \times 10^{-24}$	"	[43]*, [18]*
"	$> -2 \times 10^{-22}$	"	[45]*, [18]*
"	$< 4.5 \times 10^{-23}$	"	[73]*, [18]*
$\tilde{c}_p^{\text{UR}(4)} - \tilde{c}_e^{\text{UR}(4)}$	$(-9.8 \text{ to } 9.8) \times 10^{-22}$	"	[74]*, [18]*
$ \tilde{c}_Q $	$< 2 \times 10^{-11}$ GeV	Relativistic Lithium ions	[48]
$\tilde{c}_{TT}$	$(0.24 \pm 0.30) \times 10^{-6}$	Nuclear binding energy	[51]
"	$(-3.3 \pm 3.5) \times 10^{-6}$	Cs interferometer	[52]
$\tilde{c}_Q$	$(-0.3 \pm 2.2) \times 10^{-22}$ GeV	Cs fountain	[75]
$\tilde{c}_-$	$(-1.8 \pm 2.8) \times 10^{-25}$ GeV	"	[75]
$\tilde{c}_X$	$(0.6 \pm 1.2) \times 10^{-25}$ GeV	"	[75]
$\tilde{c}_Y$	$(-1.9 \pm 1.2) \times 10^{-25}$ GeV	"	[75]
$\tilde{c}_Z$	$(-1.4 \pm 2.8) \times 10^{-25}$ GeV	"	[75]
$\tilde{c}_{TX}$	$(-2.7 \pm 3.0) \times 10^{-21}$ GeV	"	[75]
$\tilde{c}_{TY}$	$(-0.2 \pm 3.0) \times 10^{-21}$ GeV	"	[75]
$\tilde{c}_{TZ}$	$(-0.4 \pm 2.0) \times 10^{-21}$ GeV	"	[75]
$ c_{XX} + c_{YY} - 2c_{ZZ} $	$< 10^{-11}$	Doppler shift	[58]*
$ c_{TJ} + c_{JT} , (J = X, Y, Z)$	$< 10^{-8}$	"	[58]*
$r^{H^-}$	$< 9 \times 10^{-27}$	Penning trap	[76]
$r_{\omega_c}^{H^-}$	$< 4 \times 10^{-26}$	"	[77]
$ d_{00} $	$< 8 \times 10^{-8}$	Cs spectroscopy	[68]*
"	$< 5 \times 10^{-8}$	"	[32]*
"	$< 3 \times 10^{-8}$	"	[68]*
"	$< 9 \times 10^{-8}$	Tl spectroscopy	[32]*, [68]*
$\tilde{d}_J, \tilde{g}_{DJ}, (J = X, Y)$	$< 3 \times 10^{-28}$ GeV	Hg/Cs comparison	[69]
$ \tilde{d}_J ,  \tilde{g}_{D,J} , (J = X, Y)$	$< 10^{-25}$ GeV	"	[40], [41]*

Table D10. Nonminimal proton sector,  $d \geq 5$ 

Combination	Result	System	Ref.
$ 3\hat{a}_{p,2}^{\text{NR}} - \hat{a}_{n,2}^{\text{NR}} - 4(\mathbf{p}_{pd}^2)(\hat{a}_{p,4}^{\text{NR}} + \hat{a}_{n,4}^{\text{NR}}) - 3\hat{c}_{p,2}^{\text{NR}} + \hat{c}_{n,2}^{\text{NR}} - 4(\mathbf{p}_{pd}^2)(\hat{c}_{p,4}^{\text{NR}} + \hat{c}_{n,4}^{\text{NR}}) $	$< 2 \times 10^{-7} \text{ GeV}^{-1}$	Deuterium 1S-2S	[30]*
$ \text{Re } H_{211}^{\text{NR}(0B)} ,  \text{Im } H_{211}^{\text{NR}(0B)} ,  \text{Re } g_{211}^{\text{NR}(0B)} ,  \text{Im } g_{211}^{\text{NR}(0B)} $	$< 7 \times 10^{-16} \text{ GeV}^{-1}$	H maser	[30]*
$ \text{Re } H_{211}^{\text{NR}(1B)} ,  \text{Im } H_{211}^{\text{NR}(1B)} ,  \text{Re } g_{211}^{\text{NR}(1B)} ,  \text{Im } g_{211}^{\text{NR}(1B)} $	$< 4 \times 10^{-16} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)TTX} $	$< 3.4 \times 10^{-8} \text{ GeV}^{-1}$	1S-2S transition	[30]*
$ a_{\text{eff}}^{(5)TTY} $	$< 5.6 \times 10^{-8} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)TTZ} $	$< 1.3 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKX} $	$< 6.7 \times 10^{-8} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKY} $	$< 1.1 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$ a_{\text{eff}}^{(5)KKZ} $	$< 2.5 \times 10^{-7} \text{ GeV}^{-1}$	"	[30]*
$ c_{\text{eff}}^{(6)TTTTX} $	$< 1.8 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TTTTY} $	$< 3.0 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TTTTZ} $	$< 6.9 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKKX} $	$< 1.8 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKKY} $	$< 3.0 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$ c_{\text{eff}}^{(6)TKKKZ} $	$< 6.9 \times 10^{-8} \text{ GeV}^{-2}$	"	[30]*
$\hat{c}^{\text{UR}(6)}$	$(-0.0034 \text{ to } 3.4) \times 10^{-42} \text{ GeV}^{-2}$	Astrophysics	[78]*, [18]*
$ \text{Re } H_{411}^{\text{NR}(0B)} ,  \text{Im } H_{411}^{\text{NR}(0B)} ,  \text{Re } g_{411}^{\text{NR}(0B)} ,  \text{Im } g_{411}^{\text{NR}(0B)} $	$< 9 \times 10^{-6} \text{ GeV}^{-3}$	H maser	[30]*
$ \text{Re } H_{411}^{\text{NR}(1B)} ,  \text{Im } H_{411}^{\text{NR}(1B)} ,  \text{Re } g_{411}^{\text{NR}(1B)} ,  \text{Im } g_{411}^{\text{NR}(1B)} $	$< 5 \times 10^{-6} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTTTX} $	$< 1.1 \times 10^{-8} \text{ GeV}^{-3}$	1S-2S transition	[30]*
$ a_{\text{eff}}^{(7)TTTTY} $	$< 1.9 \times 10^{-8} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTTTZ} $	$< 4.4 \times 10^{-8} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKX} $	$< 7.6 \times 10^{-9} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKY} $	$< 1.3 \times 10^{-8} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)TTKKZ} $	$< 2.9 \times 10^{-8} \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLX} $	$< 8.7 \times 10^2 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLY} $	$< 1.5 \times 10^3 \text{ GeV}^{-3}$	"	[30]*
$ a_{\text{eff}}^{(7)KKLLZ} $	$< 3.4 \times 10^3 \text{ GeV}^{-3}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTTX} $	$< 8.1 \times 10^{-9} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTTY} $	$< 1.4 \times 10^{-8} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTTTZ} $	$< 3.1 \times 10^{-8} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKX} $	$< 4.1 \times 10^{-9} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKY} $	$< 6.8 \times 10^{-9} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TTTKKZ} $	$< 1.6 \times 10^{-8} \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLX} $	$< 1.5 \times 10^2 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLY} $	$< 2.6 \times 10^2 \text{ GeV}^{-4}$	"	[30]*
$ c_{\text{eff}}^{(8)TKKLLZ} $	$< 6.0 \times 10^2 \text{ GeV}^{-4}$	"	[30]*

Table D11. Neutron sector,  $d = 3, 4$  (part 1 of 2)

Combination	Result	System	Ref.
$ \tilde{b}_\perp $	$< 1.5 \times 10^{-33}$ GeV	He/Xe magnetometer	[68]*
$ b_0 $	$< 3 \times 10^{-7}$ GeV	Cs spectroscopy	[68]*
"	$< 2 \times 10^{-7}$ GeV	"	[32]*
$\tilde{b}_x$	$(4.1 \pm 4.7) \times 10^{-34}$ GeV	He/Xe magnetometer	[79]
$\tilde{b}_y$	$(2.9 \pm 6.2) \times 10^{-34}$ GeV	"	[79]
$\tilde{b}_\perp$	$< 1.3 \times 10^{-33}$ GeV	"	[79]
$\tilde{b}_Z$	$< 7 \times 10^{-30}$ GeV	Hg/Cs magnetometer	[69]
$\tilde{b}_\perp$	$< 3.7 \times 10^{-31}$ GeV	"	[69]
$\tilde{b}_X$	$(3.36 \pm 1.72) \times 10^{-32}$ GeV	He/Xe magnetometer	[80]
$\tilde{b}_Y$	$(1.43 \pm 1.33) \times 10^{-32}$ GeV	"	[80]
$ \tilde{b}_\perp $	$< 3.72 \times 10^{-32}$ GeV	"	[80], [81]
$\tilde{b}_X$	$(0.1 \pm 1.6) \times 10^{-33}$ GeV	K/He magnetometer	[70]
$\tilde{b}_Y$	$(2.5 \pm 1.6) \times 10^{-33}$ GeV	"	[70]
$ \tilde{b}_\perp $	$< 3.7 \times 10^{-33}$ GeV	"	[70]
$b_\perp$	$< 2 \times 10^{-29}$ GeV	Ultra-cold neutrons	[82]
$-4.2b_i^{(n)} + 0.7b_i^{(p)}$	$2\pi(53 \pm 45$ nHz)	Xe/He maser	[83]*
$ b_J - \frac{1}{2}\epsilon_{JKL}H_{KL} $ , ( $J = X, Y$ )	$< 10^{-28}$ GeV	Maser/magnetometer	[84]*
$\tilde{b}_X$	$(-3.7 \pm 8.1) \times 10^{-32}$ GeV	K/He magnetometer	[34]
$\tilde{b}_Y$	$(-9.0 \pm 7.5) \times 10^{-32}$ GeV	"	[34]
$\tilde{b}_Y - 0.0034\tilde{d}_Y + 0.0034\tilde{g}_{DY}$	$(8.0 \pm 9.5) \times 10^{-32}$ GeV	Xe/He maser	[85]
$-\tilde{b}_X - 0.0034\tilde{d}_X - 0.0034\tilde{g}_{DX}$	$(2.2 \pm 7.9) \times 10^{-32}$ GeV	"	[85]
$-\cos\eta(\frac{1}{2}\tilde{b}_T + \frac{1}{2}\tilde{d}_- - \tilde{g}_c - \frac{1}{2}\tilde{g}_T)$ $-\cos\eta(\tilde{g}_T - 2\tilde{d}_+ + \frac{1}{2}\tilde{d}_Q)$ $+\sin\eta(\tilde{d}_{YZ} - \tilde{H}_{XT})$ $-\tilde{H}_{ZT}$	$(-1.1 \pm 1.0) \times 10^{-27}$ GeV	"	[85]
$(\frac{1}{2}\tilde{b}_T + \frac{1}{2}\tilde{d}_- - \tilde{g}_c - \frac{1}{2}\tilde{g}_T)$ $-(\tilde{g}_T - 2\tilde{d}_+ + \frac{1}{2}\tilde{d}_Q)$	$(0.2 \pm 1.8) \times 10^{-27}$ GeV	"	[85]
$(\frac{1}{2}\tilde{b}_T + \frac{1}{2}\tilde{d}_- - \tilde{g}_c - \frac{1}{2}\tilde{g}_T)$ $-(\tilde{g}_T - 2\tilde{d}_+ + \frac{1}{2}\tilde{d}_Q)$	$(-1.8 \pm 1.9) \times 10^{-27}$ GeV	"	[85]
$\cos\eta(\tilde{H}_{ZT} - \tilde{d}_{XY}) - \sin\eta\tilde{H}_{YT}$	$(-1.1 \pm 0.8) \times 10^{-27}$ GeV	"	[85]
$\sqrt{(\tilde{b}_X)^2 + (\tilde{b}_Y)^2}$	$(6.4 \pm 5.4) \times 10^{-32}$ GeV	"	[86]
$r_n$	$< 1.5 \times 10^{-30}$	Hg/Cs comparison	[37]
$ \tilde{b}_J $ , ( $J = X, Y$ )	$< 10^{-30}$ GeV	"	[40], [41]*

Table D11. Neutron sector,  $d = 3, 4$  (part 2 of 2)

Combination	Result	System	Ref.
$\tilde{c}_Q$	$(-1.8 \pm 2.2) \times 10^{-14}$ GeV	Quartz oscillators	[87]
$\bar{c}_{TT}$	$(1.1 \pm 1.4) \times 10^{-6}$	Nuclear binding energy	[51]
"	$(7.6 \pm 6.7) \times 10^{-6}$	Cs interferometer	[52]
$c_{YZ} + c_{ZY}$	$(4.8 \pm 4.4) \times 10^{-29}$	Ne/Rb/K magnetometer	[88]
$c_{XZ} + c_{ZX}$	$(-2.8 \pm 3.4) \times 10^{-29}$	"	[88]
$c_{XY} + c_{YX}$	$(-1.2 \pm 1.4) \times 10^{-29}$	"	[88]
$c_{XX} - c_{YY}$	$(1.4 \pm 1.7) \times 10^{-29}$	"	[88]
$ \bar{c}_Q $	$< 10^{-8}$	Combined WEP	[26]*
$ \bar{c}_\oplus $	$< 10^{-8}$	Solar system	[26]*
$ \bar{c}_\ominus $	$< 10^{-7}$	"	[26]*
$\frac{1}{4} c_Q ,  c_{(TJ)} , (J = X, Y, Z)$	$< 5 \times 10^{-14}$	Astrophysics	[89]*
$\min( c_{11} - c_{22} ,  c_{11} - c_{33} ,  c_{22} - c_{33} )$	$< 1.7 \times 10^{-8}$	Pulsar timing	[90]*
$ \bar{c}_J , (J = X, Y)$	$< 10^{-25}$ GeV	Be/H comparison	[91], [41]*
$ \tilde{c}_- ,  \tilde{c}_Z $	$< 10^{-27}$ GeV	Hg/Hg & Ne/He comparison	[92], [93], [41]*
$ d_{00} $	$< 3 \times 10^{-7}$	Cs spectroscopy	[68]*
"	$< 2 \times 10^{-7}$	"	[32]*
$\tilde{d}_J, \tilde{g}_{DJ}, (J = X, Y)$	$< 3 \times 10^{-29}$ GeV	Hg/Cs comparison	[69]
$ md_{JT} - \frac{1}{2}\epsilon_{JKLM}g_{KLT} , (J = X, Y)$	$< 10^{-28}$ GeV	Maser/magnetometer	[84]*
$\frac{1}{2} d_{(XZ)} ,  d_{(TZ)} $	$< 5 \times 10^{-14}$	Astrophysics	[89]*
$ \tilde{d}_J ,  \tilde{g}_{D,J} , (J = X, Y)$	$< 10^{-28}$ GeV	Hg/Cs comparison	[40], [41]*

Table D12. Photon sector,  $d = 3$ 

Combination	Result	System	Ref.
$ k_{(V)10}^{(3)} $	$< 16 \times 10^{-21}$ GeV	Schumann resonances	[94]*
$ k_{(V)11}^{(3)} $	$< 12 \times 10^{-21}$ GeV	"	[94]*
$ \mathbf{k}_{\mathbf{AF}}^{(3)}  \equiv \left(6 k_{(V)11}^{(3)} ^2 + 3 k_{(V)10}^{(3)} ^2\right)^{1/2} / \sqrt{4\pi}$	$(10_{-8}^{+4}) \times 10^{-43}$ GeV	CMB polarization	[19]*
$ \mathbf{k}_{\mathbf{AF}}^{(3)} $	$(15 \pm 6) \times 10^{-43}$ GeV	"	[13]*, [14]*
$k_{(V)10}^{(3)}$	$\pm(3 \pm 1) \times 10^{-42}$ GeV	"	[13]*
$\text{Re } k_{(V)11}^{(3)}$	$\pm(21_{-9}^{+7}) \times 10^{-43}$ GeV	"	[13]*
$ \mathbf{k}_{\mathbf{AF}}^{(3)} $	$(0.57 \pm 0.70)H_0$	Astrophysical birefringence	[95]*
$ 2\mathbf{k}_{\mathbf{AF}}^{(3)} $	$10^{-41}$ GeV	"	[96]*
$ \sum_{jm} Y_{jm} k_{(V)jm}^{(3)} $	$< 6 \times 10^{-43}$ GeV	"	[97]*, [14]*
$ k_{(V)00}^{(3)} $	$< 14 \times 10^{-21}$ GeV	Schumann resonances	[94]*
$k_{(V)00}^{(3)}$	$(1.1 \pm 1.3 \pm 1.5) \times 10^{-43}$ GeV	CMB polarization	[98]
"	$(0.04 \pm 0.35) \times 10^{-43}$ GeV	"	[99]*
"	$(-0.64 \pm 0.50 \pm 0.50) \times 10^{-43}$ GeV	"	[100]
"	$(4.3 \pm 4.1) \times 10^{-43}$ GeV	"	[101]
"	$(-1.4 \pm 0.9 \pm 0.5) \times 10^{-43}$ GeV	"	[102]
"	$(2.3 \pm 5.4) \times 10^{-43}$ GeV	"	[19]*
"	$< 2.5 \times 10^{-43}$ GeV	"	[103]*, [19]*
"	$(1.2 \pm 2.2) \times 10^{-43}$ GeV	"	[104], [19]*
"	$(12 \pm 7) \times 10^{-43}$ GeV	"	[13]*
"	$(2.6 \pm 1.9) \times 10^{-43}$ GeV	"	[105]*, [19]*
"	$(2.5 \pm 3.0) \times 10^{-43}$ GeV	"	[106]*, [19]*
"	$(6.0 \pm 4.0) \times 10^{-43}$ GeV	"	[107]*, [13]*
"	$(1.1 \pm 1.4)H_0$	Astrophysical birefringence	[95]*
"	$< 2 \times 10^{-42}$ GeV	"	[97]*, [14]*

Table D13. Photon sector,  $d = 4$  (part 1 of 4)

Combination	Result	System	Ref.
$(\tilde{\kappa}_{e-})^{XY}$	$(0.8 \pm 0.4) \times 10^{-17}$	Rotating optical resonators	[108]
"	$(-0.7 \pm 1.6) \times 10^{-18}$	Sapphire cavity oscillators	[109]
"	$(0.8 \pm 0.6) \times 10^{-16}$	Rotating microwave resonators	[110]
"	$(-0.31 \pm 0.73) \times 10^{-17}$	Rotating optical resonators	[111]
"	$(0.0 \pm 1.0 \pm 0.3) \times 10^{-17}$	"	[112]
"	$(-0.1 \pm 0.6) \times 10^{-17}$	"	[113]
"	$(-7.7 \pm 4.0) \times 10^{-16}$	Optical, microwave resonators	[55]*
"	$(2.9 \pm 2.3) \times 10^{-16}$	Rotating microwave resonators	[114]
"	$(-3.1 \pm 2.5) \times 10^{-16}$	Rotating optical resonators	[115]
"	$(-0.63 \pm 0.43) \times 10^{-15}$	Rotating microwave resonators	[116]
"	$(-1.7 \pm 1.6) \times 10^{-15}$	Optical, microwave resonators	[56]*
"	$(-5.7 \pm 2.3) \times 10^{-15}$	Microwave resonator, maser	[117]
"	$(1.7 \pm 2.6) \times 10^{-15}$	Optical resonators	[118]
"	$(1.4 \pm 1.4) \times 10^{-13}$	Microwave resonators	[119]
$(\tilde{\kappa}_{e-})^{XZ}$	$(-0.3 \pm 0.6) \times 10^{-17}$	Rotating optical resonators	[108]
"	$(-5.5 \pm 4.0) \times 10^{-18}$	Sapphire cavity oscillators	[109]
"	$(1.5 \pm 1.3) \times 10^{-16}$	Rotating microwave resonators	[110]
"	$(0.54 \pm 0.70) \times 10^{-17}$	Rotating optical resonators	[111]
"	$(0.4 \pm 1.5 \pm 0.1) \times 10^{-17}$	"	[112]
"	$(-2.0 \pm 0.9) \times 10^{-17}$	"	[113]
"	$(-10.3 \pm 3.9) \times 10^{-16}$	Optical, microwave resonators	[55]*
"	$(-6.9 \pm 2.2) \times 10^{-16}$	Rotating microwave resonators	[114]
"	$(5.7 \pm 4.9) \times 10^{-16}$	Rotating optical resonators	[115]
"	$(0.19 \pm 0.37) \times 10^{-15}$	Rotating microwave resonators	[116]
"	$(-4.0 \pm 3.3) \times 10^{-15}$	Optical, microwave resonators	[56]*
"	$(-3.2 \pm 1.3) \times 10^{-15}$	Microwave resonator, maser	[117]
"	$(-6.3 \pm 12.4) \times 10^{-15}$	Optical resonators	[118]
"	$(-3.5 \pm 4.3) \times 10^{-13}$	Microwave resonators	[119]
$(\tilde{\kappa}_{e-})^{YZ}$	$(0.0 \pm 0.6) \times 10^{-17}$	Rotating optical resonators	[108]
"	$(-1.9 \pm 3.2) \times 10^{-18}$	Sapphire cavity oscillators	[109]
"	$(1.7 \pm 1.3) \times 10^{-16}$	Rotating microwave resonators	[110]
"	$(-0.97 \pm 0.74) \times 10^{-17}$	Rotating optical resonators	[111]
"	$(-0.6 \pm 1.4 \pm 0.5) \times 10^{-17}$	"	[112]
"	$(-0.3 \pm 1.4) \times 10^{-17}$	"	[113]
"	$(0.9 \pm 4.2) \times 10^{-16}$	Optical, microwave resonators	[55]*
"	$(2.1 \pm 2.1) \times 10^{-16}$	Rotating microwave resonators	[114]
"	$(-1.5 \pm 4.4) \times 10^{-16}$	Rotating optical resonators	[115]
"	$(-0.45 \pm 0.37) \times 10^{-15}$	Rotating microwave resonators	[116]
"	$(0.52 \pm 2.52) \times 10^{-15}$	Optical, microwave resonators	[56]*
"	$(-0.5 \pm 1.3) \times 10^{-15}$	Microwave resonator, maser	[117]
"	$(3.6 \pm 9.0) \times 10^{-15}$	Optical resonators	[118]
"	$(1.7 \pm 3.6) \times 10^{-13}$	Microwave resonators	[119]

Table D13. Photon sector,  $d = 4$  (part 2 of 4)

Combination	Result	System	Ref.
$(\tilde{\kappa}_{e-})^{XX} - (\tilde{\kappa}_{e-})^{YY}$	$(0.8 \pm 0.8) \times 10^{-17}$	Rotating optical resonators	[108]
”	$(-1.5 \pm 3.4) \times 10^{-18}$	Sapphire cavity oscillators	[109]
”	$(0.2 \pm 1.0) \times 10^{-16}$	Rotating microwave resonators	[110]
”	$(0.80 \pm 1.27) \times 10^{-17}$	Rotating optical resonators	[111]
”	$(0.8 \pm 2.0 \pm 0.3) \times 10^{-17}$	”	[112]
”	$(-2.0 \pm 1.7) \times 10^{-17}$	”	[113]
”	$(-12 \pm 16) \times 10^{-16}$	Optical, microwave resonators	[55]*
”	$(-5.0 \pm 4.7) \times 10^{-16}$	Rotating microwave resonators	[114]
”	$(5.4 \pm 4.8) \times 10^{-16}$	Rotating optical resonators	[115]
”	$(-1.3 \pm 0.9) \times 10^{-15}$	Rotating microwave resonators	[116]
”	$(2.8 \pm 3.3) \times 10^{-15}$	Optical, microwave resonators	[56]*
”	$(-3.2 \pm 4.6) \times 10^{-15}$	Microwave resonator, maser	[117]
”	$(8.9 \pm 4.9) \times 10^{-15}$	Optical resonators	[118]
”	$(-1.0 \pm 2.1) \times 10^{-13}$	Microwave resonators	[119]
$(\tilde{\kappa}_{e-})^{ZZ}$	$(4.8 \pm 1.1) \times 10^{-17}$	Rotating optical resonators	[108]
”	$(-286 \pm 279) \times 10^{-18}$	Sapphire cavity oscillators	[109]
”	$(143 \pm 179) \times 10^{-16}$	Rotating microwave resonators	[110]
”	$(-0.04 \pm 1.73) \times 10^{-17}$	Rotating optical resonators	[111]
”	$(1.6 \pm 2.4 \pm 1.1) \times 10^{-17}$	”	[112]
”	$(-0.2 \pm 3.1) \times 10^{-17}$	”	[113]
”	$(223 \pm 290) \times 10^{-16}$	Optical, microwave resonators	[55]*
”	$(143 \pm 179) \times 10^{-16}$	Rotating microwave resonators	[114]
”	$(-1.9 \pm 5.2) \times 10^{-15}$	Rotating optical resonators	[115]
”	$(21 \pm 57) \times 10^{-15}$	Rotating microwave resonators	[116]
”	$(-2.9 \pm 2.2) \times 10^{-14}$	Optical resonators	[120]
$ (\tilde{\kappa}_{e-})^{(kl)} $	$< 4 \times 10^{-18}$	Astrophysics	[121]*



Table D13. Photon sector,  $d = 4$  (part 3 of 4)

Combination	Result	System	Ref.
$(\tilde{\kappa}_{o+})^{XY}$	$(-3.0 \pm 3.4) \times 10^{-14}$	Sapphire cavity oscillators	[109]
$\tilde{\kappa}_{o+}^{XY}$	$(0.7 \pm 1.0) \times 10^{-14}$	Optical ring cavity	[122]
$(\tilde{\kappa}_{o+})^{XY}$	$(-1.5 \pm 1.2) \times 10^{-12}$	Rotating microwave resonators	[110]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{XY}$	$(-0.14 \pm 0.78) \times 10^{-17}$	Rotating optical resonators	[111]
$(\tilde{\kappa}_{o+})^{XY}$	$(1.5 \pm 1.5 \pm 0.2) \times 10^{-13}$	"	[112]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{XY}$	$(-2.5 \pm 2.5) \times 10^{-17}$	"	[113]
$(\tilde{\kappa}_{o+})^{XY}$	$(1.7 \pm 2.0) \times 10^{-12}$	Optical, microwave resonators	[55]*
"	$(-0.9 \pm 2.6) \times 10^{-12}$	Rotating microwave resonators	[114]
"	$(-2.5 \pm 5.1) \times 10^{-12}$	Rotating optical resonators	[115]
"	$(0.20 \pm 0.21) \times 10^{-11}$	Rotating microwave resonators	[116]
"	$(-1.8 \pm 1.5) \times 10^{-11}$	Microwave resonator, maser	[117]
"	$(14 \pm 14) \times 10^{-11}$	Optical resonators	[118]
$(\tilde{\kappa}_{o+})^{XZ}$	$(0.21 \pm 1.7) \times 10^{-14}$	Sapphire cavity oscillators	[109]
$\tilde{\kappa}_{o+}^{XZ}$	$(-0.6 \pm 1.2) \times 10^{-14}$	Optical ring cavity	[122]
$(\tilde{\kappa}_{o+})^{XZ}$	$(1.6 \pm 2.2) \times 10^{-13}$	Asymmetric optical resonator	[123]
"	$(0.7 \pm 1.4) \times 10^{-12}$	"	[124]
"	$(1.7 \pm 0.7) \times 10^{-12}$	Rotating microwave resonators	[110]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{XZ}$	$(-0.45 \pm 0.62) \times 10^{-17}$	Rotating optical resonators	[111]
$(\tilde{\kappa}_{o+})^{XZ}$	$(-0.1 \pm 1.0 \pm 0.2) \times 10^{-13}$	"	[112]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{XZ}$	$(1.5 \pm 1.7) \times 10^{-17}$	"	[113]
$(\tilde{\kappa}_{o+})^{XZ}$	$(-3.1 \pm 2.3) \times 10^{-12}$	Optical, microwave resonators	[55]*
"	$(-4.4 \pm 2.5) \times 10^{-12}$	Rotating microwave resonators	[114]
"	$(-3.6 \pm 2.7) \times 10^{-12}$	Rotating optical resonators	[115]
"	$(-0.91 \pm 0.46) \times 10^{-11}$	Rotating microwave resonators	[116]
"	$(-1.4 \pm 2.3) \times 10^{-11}$	Microwave resonator, maser	[117]
"	$(-1.2 \pm 2.6) \times 10^{-11}$	Optical resonators	[118]
$(\tilde{\kappa}_{o+})^{YZ}$	$(-2.0 \pm 1.6) \times 10^{-14}$	Sapphire cavity oscillators	[109]
$\tilde{\kappa}_{o+}^{YZ}$	$(0.5 \pm 1.0) \times 10^{-14}$	Optical ring cavity	[122]
$(\tilde{\kappa}_{o+})^{YZ}$	$(0.2 \pm 0.7) \times 10^{-12}$	Rotating microwave resonators	[110]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{YZ}$	$(-0.34 \pm 0.61) \times 10^{-17}$	Rotating optical resonators	[111]
$(\tilde{\kappa}_{o+})^{YZ}$	$(-0.1 \pm 1.0 \pm 0.4) \times 10^{-13}$	"	[112]
$\beta_{\oplus}(\tilde{\kappa}_{o+})^{YZ}$	$(-1.0 \pm 1.5) \times 10^{-17}$	"	[113]
$(\tilde{\kappa}_{o+})^{YZ}$	$(-2.8 \pm 2.2) \times 10^{-12}$	Optical, microwave resonators	[55]*
"	$(-3.2 \pm 2.3) \times 10^{-12}$	Rotating microwave resonators	[114]
"	$(2.9 \pm 2.8) \times 10^{-12}$	Rotating optical resonators	[115]
"	$(0.44 \pm 0.46) \times 10^{-11}$	Rotating microwave resonators	[116]
"	$(2.7 \pm 2.2) \times 10^{-11}$	Microwave resonator, maser	[117]
"	$(0.1 \pm 2.7) \times 10^{-11}$	Optical resonators	[118]
$\sqrt{(2c_{TX} - \tilde{\kappa}_{o+}^{YZ})^2 + (2c_{TY} - \tilde{\kappa}_{o+}^{ZX})^2}$	$< 4.6 \times 10^{-10}$	Compton scattering	[125]*
"	$< 1.6 \times 10^{-14}$	"	[126]
$(\tilde{\kappa}_{o+})^{YX} - 0.432(\tilde{\kappa}_{o+})^{ZX}$	$(4.0 \pm 8.4) \times 10^{-9}$	Microwave resonators	[119]
$(\tilde{\kappa}_{o+})^{XY} - 0.209(\tilde{\kappa}_{o+})^{YZ}$	$(4.0 \pm 4.9) \times 10^{-9}$	"	[119]
$(\tilde{\kappa}_{o+})^{XZ} - 0.484(\tilde{\kappa}_{o+})^{YZ}$	$(1.6 \pm 1.7) \times 10^{-9}$	"	[119]
$(\tilde{\kappa}_{o+})^{YZ} + 0.484(\tilde{\kappa}_{o+})^{XZ}$	$(0.6 \pm 1.9) \times 10^{-9}$	"	[119]
$ (\tilde{\kappa}_{o+})^{(ij)} $	$< 2 \times 10^{-18}$	Astrophysics	[121]*

Table D13. Photon sector,  $d = 4$  (part 4 of 4)

Combination	Result	System	Ref.
$\tilde{\kappa}_{\text{tr}}$	$(-6.0 \pm 4.0) \times 10^{-10}$	Sapphire cavity oscillators	[109]
$ \tilde{\kappa}_{\text{tr}} $	$< 2 \times 10^{-8}$	Relativistic Lithium ions	[48]
$\tilde{\kappa}_{\text{tr}}$	$(-2 \text{ to } 0.0006) \times 10^{-16}$	Astrophysics	[127]*
"	$(-0.4 \pm 0.9) \times 10^{-10}$	Optical ring cavity	[122]
"	$(3 \pm 11) \times 10^{-10}$	Asymmetric optical resonator	[123]
"	$(3.4 \pm 6.2) \times 10^{-9}$	"	[124]
"	$(-1.5 \pm 0.74) \times 10^{-8}$	Rotating microwave resonators	[110]
"	$(-0.3 \pm 3) \times 10^{-7}$	Microwave interferometer	[128]
$ \tilde{\kappa}_{\text{tr}} - \frac{4}{3}c_{00}^e $	$< 5 \times 10^{-15}$	Collider physics	[129]*
$\tilde{\kappa}_{\text{tr}} - \frac{4}{3}c_{00}^e$	$(-5.8 \text{ to } 12) \times 10^{-12}$	"	[130]*
$\tilde{\kappa}_{\text{tr}} - \frac{4}{3}c_{00}^p$	$< 6 \times 10^{-20}$	Astrophysics	[131]*
$-\left[\tilde{\kappa}_{\text{tr}} - \frac{4}{3}c_{00}^e\right]$	$< 9 \times 10^{-16}$	"	[131]*
$\tilde{\kappa}_{\text{tr}}$	$< 1.4 \times 10^{-19}$	"	[121]*
$ \tilde{\kappa}_{\text{tr}} $	$< 8.4 \times 10^{-8}$	Optical atomic clocks	[132]
"	$< 2.2 \times 10^{-7}$	Heavy-ion storage ring	[133]*
"	$< 2 \times 10^{-14}$	Astrophysics	[134]*
"	$< 3 \times 10^{-8}$	$g_e - 2$	[134]*
"	$< 1.6 \times 10^{-5}$	Sagnac interferometer	[135]
$c_{(I)20}^{(4)}$	$(3 \pm 16) \times 10^{-15}$	Rotating microwave resonators	[136]
$\text{Re}(c_{(I)21}^{(4)})$	$(20 \pm 23) \times 10^{-17}$	"	[136]
$\text{Im}(c_{(I)21}^{(4)})$	$(137 \pm 71) \times 10^{-18}$	"	[136]
$\text{Re}(c_{(I)22}^{(4)})$	$(-4 \pm 23) \times 10^{-17}$	"	[136]
$\text{Im}(c_{(I)22}^{(4)})$	$(20 \pm 22) \times 10^{-18}$	"	[136]
$ \sum_{jm} 2Y_{jm}(27^\circ, 6^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-37}$	Astrophysical birefringence	[137]*
$ \sum_{jm} 2Y_{jm}(112^\circ, 279^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-38}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(61^\circ, 229^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-38}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(129^\circ, 333^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-38}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(98.2^\circ, 182.1^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-37}$	"	[14]*
$ \sum_{jm} 2Y_{jm}(87.3^\circ, 37.3^\circ)(k_{(E)jm}^{(4)} + ik_{(B)jm}^{(4)}) $	$\lesssim 10^{-37}$	"	[14]*
$k_{(E)20}^{(4)}$	$\pm(17_{-9}^{+7}) \times 10^{-31}$	CMB polarization	[13]*
$k_{(B)20}^{(4)}$	$\pm(17_{-9}^{+7}) \times 10^{-31}$	"	[13]*
$\sqrt{(\sum_{i=1}^3 k_2^i)^2}$	$< 1.4 \times 10^{-16}$	Astrophysical birefringence	[138]*
$\sqrt{\sum_m ( k_{(E)2m}^{(4)} ^2 +  k_{(B)2m}^{(4)} ^2)}$	$< 5 \times 10^{-32}$	"	[11]*, [14]*
$ k^a $ for some $a$	$< 2 \times 10^{-37}$	"	[139]*
$ k^a $ for $a = 1, \dots, 10$	$< 2 \times 10^{-32}$	"	[11]*

Table D14. Nonminimal photon sector,  $d = 5$ 

Combination	Result	System	Ref.
$ \sum_{jm} Y_{jm}(27^\circ, 6^\circ)k_{(V)jm}^{(5)} $	$< 2 \times 10^{-34} \text{ GeV}^{-1}$	Astrophysical birefringence	[137]*
$ k_{(V)00}^{(5)} $	$< 8 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ \sum_{jm} Y_{jm}(112^\circ, 279^\circ)k_{(V)jm}^{(5)} $	$< 7 \times 10^{-35} \text{ GeV}^{-1}$	"	[137]*
$ k_{(V)00}^{(5)} $	$< 2 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ \sum_{jm} Y_{jm}(61^\circ, 229^\circ)k_{(V)jm}^{(5)} $	$< 2 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ k_{(V)00}^{(5)} $	$< 9 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ \sum_{jm} Y_{jm}(129^\circ, 333^\circ)k_{(V)jm}^{(5)} $	$< 1 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ k_{(V)00}^{(5)} $	$< 4 \times 10^{-34} \text{ GeV}^{-1}$	"	[137]*
$ k_{(V)00}^{(5)} $	$< 6 \times 10^{-34} \text{ GeV}^{-1}$	"	[140], [137]*
$ k_{(V)00}^{(5)} $	$< 3 \times 10^{-33} \text{ GeV}^{-1}$	"	[141]*, [137]*
$ \sum_{jm} Y_{jm}(37^\circ, 0^\circ)k_{(V)jm}^{(5)} $	$< 1.2 \times 10^{-34} \text{ GeV}^{-1}$	"	[142]*
$ k_{(V)00}^{(5)} $	$< 4.2 \times 10^{-34} \text{ GeV}^{-1}$	"	[142]*
$ \sum_{jm} Y_{jm}(98.2^\circ, 182.1^\circ)k_{(V)jm}^{(5)} $	$< 7 \times 10^{-33} \text{ GeV}^{-1}$	"	[14]*
$ k_{(V)00}^{(5)} $	$< 2 \times 10^{-32} \text{ GeV}^{-1}$	"	[14]*
$ \sum_{jm} Y_{jm}(87.3^\circ, 37.3^\circ)k_{(V)jm}^{(5)} $	$< 4 \times 10^{-33} \text{ GeV}^{-1}$	"	[14]*
$ k_{(V)00}^{(5)} $	$< 1 \times 10^{-32} \text{ GeV}^{-1}$	"	[14]*
$k_{(V)00}^{(5)}$	$(3.2 \pm 2.1) \times 10^{-20} \text{ GeV}^{-1}$	CMB polarization	[143]*
"	$(3 \pm 2) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*
$k_{(V)10}^{(5)}$	$(8_{-3}^{+2}) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*
	$-(8_{-4}^{+3}) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*
$k_{(V)20}^{(5)}$	$-(10 \pm 3) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*
$k_{(V)30}^{(5)}$	$(8_{-4}^{+3}) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*
	$-(8 \pm 3) \times 10^{-20} \text{ GeV}^{-1}$	"	[13]*

Table D15. Nonminimal photon sector,  $d = 6$  (part 1 of 2)

Combination	Result	System	Ref.
$\text{Re}(\bar{c}_F^6)_{111}$	$(-1.55 \pm 1.08) \times 10^6 \text{ GeV}^{-2}$	Asymmetric optical resonator	[144]
$\text{Im}(\bar{c}_F^6)_{111}$	$(-1.51 \pm 1.08) \times 10^6 \text{ GeV}^{-2}$	"	[144]
$(\bar{c}_F^{(6)})_{110}^{(0E)}$	$(-0.1 \pm 1.5) \times 10^3 \text{ GeV}^{-2}$	Optical ring cavity	[145]
$\text{Re}[(\bar{c}_F^{(6)})_{111}^{(0E)}]$	$(0.8 \pm 1.1) \times 10^3 \text{ GeV}^{-2}$	"	[145]
$\text{Im}[(\bar{c}_F^{(6)})_{111}^{(0E)}]$	$(-0.6 \pm 1.0) \times 10^3 \text{ GeV}^{-2}$	"	[145]
$(\bar{c}_F^{(6)})_{220}^{(0E)}$	$(3 \pm 13) \times 10^{10} \text{ GeV}^{-2}$	Rotating microwave resonators	[136]
$\text{Re}((\bar{c}_F^{(6)})_{221}^{(0E)})$	$(17 \pm 19) \times 10^8 \text{ GeV}^{-2}$	"	[136]
$\text{Im}((\bar{c}_F^{(6)})_{221}^{(0E)})$	$(114 \pm 59) \times 10^7 \text{ GeV}^{-2}$	"	[136]
$\text{Re}((\bar{c}_F^{(6)})_{222}^{(0E)})$	$(-3 \pm 19) \times 10^8 \text{ GeV}^{-2}$	"	[136]
$\text{Im}((\bar{c}_F^{(6)})_{222}^{(0E)})$	$(37 \pm 40) \times 10^7 \text{ GeV}^{-2}$	"	[136]
$c_{(I)00}^{(6)}$	$(-2.705 \text{ to } 3.925) \times 10^{-14} \text{ GeV}^{-2}$	Astrophysical dispersion	[146]
$c_{(I)10}^{(6)}$	$(-3.753 \text{ to } 2.889) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)11}^{(6)})$	$(-2.816 \text{ to } 3.574) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)11}^{(6)})$	$(-3.299 \text{ to } 5.984) \times 10^{-15} \text{ GeV}^{-2}$	"	[146]
$c_{(I)20}^{(6)}$	$(-4.232 \text{ to } 3.032) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)21}^{(6)})$	$(-1.590 \text{ to } 1.043) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)21}^{(6)})$	$(-4.412 \text{ to } 3.288) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)22}^{(6)})$	$(-2.353 \text{ to } 3.113) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)22}^{(6)})$	$(-5.144 \text{ to } 6.634) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$c_{(I)30}^{(6)}$	$(-4.823 \text{ to } 6.435) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)31}^{(6)})$	$(-2.439 \text{ to } 1.798) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)31}^{(6)})$	$(-2.822 \text{ to } 2.078) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)32}^{(6)})$	$(-3.125 \text{ to } 3.855) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)32}^{(6)})$	$(-2.171 \text{ to } 1.624) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)33}^{(6)})$	$(-3.693 \text{ to } 2.943) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)33}^{(6)})$	$(-4.216 \text{ to } 5.656) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$c_{(I)40}^{(6)}$	$(-2.313 \text{ to } 2.739) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)41}^{(6)})$	$(-9.021 \text{ to } 11.31) \times 10^{-15} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)41}^{(6)})$	$(-2.953 \text{ to } 3.904) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)42}^{(6)})$	$(-4.650 \text{ to } 6.846) \times 10^{-15} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)42}^{(6)})$	$(-2.489 \text{ to } 1.961) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)43}^{(6)})$	$(-7.276 \text{ to } 10.14) \times 10^{-15} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)43}^{(6)})$	$(-1.246 \text{ to } 1.343) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Re}(c_{(I)44}^{(6)})$	$(-3.919 \text{ to } 2.923) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]
$\text{Im}(c_{(I)44}^{(6)})$	$(-1.801 \text{ to } 1.427) \times 10^{-14} \text{ GeV}^{-2}$	"	[146]

Table D15. Nonminimal photon sector,  $d = 6$  (part 2 of 2)

Combination	Result	System	Ref.
$\sum_{jm} {}_0Y_{jm}(145^\circ, 120^\circ)c_{(I)jm}^{(6)}$	$(-8.7 \text{ to } 20) \times 10^{-20} \text{ GeV}^{-2}$	Astrophysical dispersion	[147]
$c_{(I)00}^{(6)}$	$(-31 \text{ to } 70) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$\sum_{jm} {}_0Y_{jm}(117^\circ, 334^\circ)c_{(I)jm}^{(6)}$	$(-0.31 \text{ to } 0.16) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$c_{(I)00}^{(6)}$	$(-1.1 \text{ to } 0.57) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$\sum_{jm} {}_0Y_{jm}(63^\circ, 265^\circ)c_{(I)jm}^{(6)}$	$(-3.4 \text{ to } 5.2) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$c_{(I)00}^{(6)}$	$(-12 \text{ to } 18) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$\sum_{jm} {}_0Y_{jm}(156^\circ, 353^\circ)c_{(I)jm}^{(6)}$	$(-11 \text{ to } 5.2) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$c_{(I)00}^{(6)}$	$(-37 \text{ to } 19) \times 10^{-20} \text{ GeV}^{-2}$	"	[147]
$\sum_{jm} Y_{jm}(116^\circ, 334^\circ)c_{(I)jm}^{(6)}$	$< 3.9 \times 10^{-22} \text{ GeV}^{-2}$	Astrophysical dispersion	[148]
$c_{(I)00}^{(6)}$	$< 1.4 \times 10^{-21} \text{ GeV}^{-2}$	"	[148]
$\sum_{jm} Y_{jm}(147^\circ, 120^\circ)c_{(I)jm}^{(6)}$	$< 3.2 \times 10^{-20} \text{ GeV}^{-2}$	"	[149], [14]*
$c_{(I)00}^{(6)}$	$< 1.1 \times 10^{-19} \text{ GeV}^{-2}$	"	[149], [14]*
$ \sum_{jm} Y_{jm}(120^\circ, 330^\circ)c_{(I)jm}^{(6)} $	$< 7.4 \times 10^{-22} \text{ GeV}^{-2}$	"	[150], [14]*
$ c_{(I)00}^{(6)} $	$< 2.6 \times 10^{-21} \text{ GeV}^{-2}$	"	[150], [14]*
$\sum_{jm} Y_{jm}(50.2^\circ, 253^\circ)c_{(I)jm}^{(6)}$	$3_{-2}^{+1} \times 10^{-22} \text{ GeV}^{-2}$	"	[151], [14]*
$c_{(I)00}^{(6)}$	$10_{-7}^{+4} \times 10^{-22} \text{ GeV}^{-2}$	"	[151], [14]*
$ \sum_{jm} Y_{jm}(99.7^\circ, 240^\circ)c_{(I)jm}^{(6)} $	$< 1 \times 10^{-16} \text{ GeV}^{-2}$	"	[152], [14]*
$ c_{(I)00}^{(6)} $	$< 4 \times 10^{-16} \text{ GeV}^{-2}$	"	[152], [14]*
$ \sum_{jm} {}_2Y_{jm}(27^\circ, 6^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-31} \text{ GeV}^{-2}$	Astrophysical birefringence	[137]*
$ \sum_{jm} {}_2Y_{jm}(112^\circ, 279^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-32} \text{ GeV}^{-2}$	"	[137]*
$ \sum_{jm} {}_2Y_{jm}(61^\circ, 229^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-31} \text{ GeV}^{-2}$	"	[137]*
$ \sum_{jm} {}_2Y_{jm}(129^\circ, 333^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-31} \text{ GeV}^{-2}$	"	[137]*
$ \sum_{jm} {}_2Y_{jm}(98.2^\circ, 182.1^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-29} \text{ GeV}^{-2}$	"	[14]*
$ \sum_{jm} {}_2Y_{jm}(87.3^\circ, 37.3^\circ)(k_{(E)jm}^{(6)} + ik_{(B)jm}^{(6)}) $	$\lesssim 10^{-29} \text{ GeV}^{-2}$	"	[14]*
$k_{(E)20}^{(6)}$	$\pm(11_{-5}^{+4}) \times 10^{-10} \text{ GeV}^{-2}$	CMB polarization	[13]*
$k_{(E)30}^{(6)}$	$\pm(11_{-6}^{+5}) \times 10^{-10} \text{ GeV}^{-2}$	"	[13]*
$k_{(E)40}^{(6)}$	$\pm(11_{-6}^{+5}) \times 10^{-10} \text{ GeV}^{-2}$	"	[13]*

Table D16. Nonminimal photon sector,  $d = 7$ 

Combination	Result	System	Ref.
$ \sum_{jm} Y_{jm}(27^\circ, 6^\circ)k_{(V)jm}^{(7)} $	$< 2 \times 10^{-28} \text{ GeV}^{-3}$	Astrophysical birefringence	[137]*
$ k_{(V)00}^{(7)} $	$< 8 \times 10^{-28} \text{ GeV}^{-3}$	"	[137]*
$ \sum_{jm} Y_{jm}(112^\circ, 279^\circ)k_{(V)jm}^{(7)} $	$< 4 \times 10^{-28} \text{ GeV}^{-3}$	"	[137]*
$ k_{(V)00}^{(7)} $	$< 1 \times 10^{-27} \text{ GeV}^{-3}$	"	[137]*
$ \sum_{jm} Y_{jm}(61^\circ, 229^\circ)k_{(V)jm}^{(7)} $	$< 2 \times 10^{-27} \text{ GeV}^{-3}$	"	[137]*
$ k_{(V)00}^{(7)} $	$< 8 \times 10^{-27} \text{ GeV}^{-3}$	"	[137]*
$ \sum_{jm} Y_{jm}(129^\circ, 333^\circ)k_{(V)jm}^{(7)} $	$< 8 \times 10^{-28} \text{ GeV}^{-3}$	"	[137]*
$ k_{(V)00}^{(7)} $	$< 3 \times 10^{-27} \text{ GeV}^{-3}$	"	[137]*
$ \sum_{jm} Y_{jm}(98.2^\circ, 182.1^\circ)k_{(V)jm}^{(7)} $	$< 2 \times 10^{-24} \text{ GeV}^{-3}$	"	[14]*
$ k_{(V)00}^{(7)} $	$< 7 \times 10^{-24} \text{ GeV}^{-3}$	"	[14]*
$ \sum_{jm} Y_{jm}(87.3^\circ, 37.3^\circ)k_{(V)jm}^{(7)} $	$< 5 \times 10^{-25} \text{ GeV}^{-3}$	"	[14]*
$ k_{(V)00}^{(7)} $	$< 2 \times 10^{-24} \text{ GeV}^{-3}$	"	[14]*

Table D17. Nonminimal photon sector,  $d = 8$  (part 1 of 2)

Combination	Result	System	Ref.
$\text{Re}(\bar{c}_F^{(8)})_{311} - 0.093 \text{Re}(\bar{c}_F^{(8)})_{111}$	$(-5.33 \pm 3.70) \times 10^{21} \text{ GeV}^{-4}$	Asymmetric optical resonator	[144]
$\text{Im}(\bar{c}_F^{(8)})_{311} - 0.093 \text{Im}(\bar{c}_F^{(8)})_{111}$	$(-5.21 \pm 3.68) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Re}(\bar{c}_F^{(8)})_{111}$	$(7.32 \pm 5.04) \times 10^{22} \text{ GeV}^{-4}$	"	[144]
$\text{Im}(\bar{c}_F^{(8)})_{111}$	$(7.14 \pm 5.00) \times 10^{22} \text{ GeV}^{-4}$	"	[144]
$\text{Re}(\bar{c}_F^{(8)})_{311}$	$(1.48 \pm 0.98) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Im}(\bar{c}_F^{(8)})_{311}$	$(1.43 \pm 0.98) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Re}(\bar{c}_F^{(8)})_{332}$	$(20.19 \pm 12.24) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Im}(\bar{c}_F^{(8)})_{332}$	$(-3.60 \pm 12.24) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Re}(\bar{c}_F^{(8)})_{333}$	$(-2.78 \pm 3.80) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$\text{Im}(\bar{c}_F^{(8)})_{333}$	$(-4.22 \pm 3.82) \times 10^{21} \text{ GeV}^{-4}$	"	[144]
$(\bar{c}_F^{(8)})_{310}^{(0E)} - 0.020(\bar{c}_F^{(8)})_{110}^{(0E)}$	$(-0.2 \pm 1.9) \times 10^{19} \text{ GeV}^{-4}$	Optical ring cavity	[145]
$\text{Re}[(\bar{c}_F^{(8)})_{311}^{(0E)} - 0.020(\bar{c}_F^{(8)})_{111}^{(0E)}]$	$(1.4 \pm 1.3) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Im}[(\bar{c}_F^{(8)})_{311}^{(0E)} - 0.020(\bar{c}_F^{(8)})_{111}^{(0E)}]$	$(0.1 \pm 1.3) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$(\bar{c}_F^{(8)})_{330}^{(0E)}$	$(-0.8 \pm 3.3) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Re}[(\bar{c}_F^{(8)})_{331}^{(0E)}]$	$(-0.3 \pm 1.9) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Im}[(\bar{c}_F^{(8)})_{331}^{(0E)}]$	$(-2.8 \pm 1.9) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Re}[(\bar{c}_F^{(8)})_{332}^{(0E)}]$	$(2.2 \pm 1.3) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Im}[(\bar{c}_F^{(8)})_{332}^{(0E)}]$	$(0.2 \pm 1.3) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Re}[(\bar{c}_F^{(8)})_{333}^{(0E)}]$	$(-0.1 \pm 1.6) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$\text{Im}[(\bar{c}_F^{(8)})_{333}^{(0E)}]$	$(-0.1 \pm 1.6) \times 10^{19} \text{ GeV}^{-4}$	"	[145]
$(\bar{c}_F^{(8)})_{220}^{(0E)} - 94(\bar{c}_F^{(8)})_{420}^{(0E)} - 20(\bar{c}_F^{(8)})_{440}^{(0E)}$	$(5 \pm 23) \times 10^{37} \text{ GeV}^{-4}$	Rotating microwave resonators	[136]
$\text{Re}((\bar{c}_F^{(8)})_{221}^{(0E)} - 98 \text{Re}((\bar{c}_F^{(8)})_{421}^{(0E)}) + 48 \text{Re}((\bar{c}_F^{(8)})_{441}^{(0E)}))$	$(11 \pm 15) \times 10^{36} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{221}^{(0E)} - 98 \text{Im}((\bar{c}_F^{(8)})_{421}^{(0E)}) + 48 \text{Im}((\bar{c}_F^{(8)})_{441}^{(0E)}))$	$(39 \pm 15) \times 10^{36} \text{ GeV}^{-4}$	"	[136]
$\text{Re}((\bar{c}_F^{(8)})_{221}^{(0E)} - 92 \text{Re}((\bar{c}_F^{(8)})_{421}^{(0E)}) - 11 \text{Re}((\bar{c}_F^{(8)})_{441}^{(0E)}))$	$(58 \pm 55) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{221}^{(0E)} - 92 \text{Im}((\bar{c}_F^{(8)})_{421}^{(0E)}) - 11 \text{Im}((\bar{c}_F^{(8)})_{441}^{(0E)}))$	$(7 \pm 55) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Re}((\bar{c}_F^{(8)})_{222}^{(0E)} - 96 \text{Re}((\bar{c}_F^{(8)})_{422}^{(0E)}) + 3 \text{Re}((\bar{c}_F^{(8)})_{442}^{(0E)}))$	$(66 \pm 29) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{222}^{(0E)} - 96 \text{Im}((\bar{c}_F^{(8)})_{422}^{(0E)}) + 3 \text{Im}((\bar{c}_F^{(8)})_{442}^{(0E)}))$	$(-26 \pm 28) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Re}((\bar{c}_F^{(8)})_{222}^{(0E)} - 94 \text{Re}((\bar{c}_F^{(8)})_{422}^{(0E)}) + 49 \text{Re}((\bar{c}_F^{(8)})_{442}^{(0E)}))$	$(-8 \pm 24) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{222}^{(0E)} - 94 \text{Im}((\bar{c}_F^{(8)})_{422}^{(0E)}) + 49 \text{Im}((\bar{c}_F^{(8)})_{442}^{(0E)}))$	$(-25 \pm 25) \times 10^{35} \text{ GeV}^{-4}$	"	[136]
$\text{Re}((\bar{c}_F^{(8)})_{443}^{(0E)})$	$(-11 \pm 15) \times 10^{33} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{443}^{(0E)})$	$(-89 \pm 53) \times 10^{32} \text{ GeV}^{-4}$	"	[136]
$\text{Re}((\bar{c}_F^{(8)})_{444}^{(0E)})$	$(-5 \pm 29) \times 10^{33} \text{ GeV}^{-4}$	"	[136]
$\text{Im}((\bar{c}_F^{(8)})_{444}^{(0E)})$	$(-10 \pm 28) \times 10^{32} \text{ GeV}^{-4}$	"	[136]

Table D17. Nonminimal photon sector,  $d = 8$  (part 2 of 2)

Combination	Result	System	Ref.
$\sum_{jm} Y_{jm}(116^\circ, 334^\circ) c_{(I)jm}^{(8)}$	$< 2.1 \times 10^{-25} \text{ GeV}^{-4}$	Astrophysical dispersion	[148]
$c_{(I)00}^{(8)}$	$< 7.6 \times 10^{-25} \text{ GeV}^{-4}$	"	[148]
$\sum_{jm} Y_{jm}(147^\circ, 120^\circ) c_{(I)jm}^{(8)}$	$< 2.6 \times 10^{-23} \text{ GeV}^{-4}$	"	[149], [14]*
$c_{(I)00}^{(8)}$	$< 9.2 \times 10^{-23} \text{ GeV}^{-4}$	"	[149], [14]*
$ \sum_{jm} Y_{jm}(99.7^\circ, 240^\circ) c_{(I)jm}^{(8)} $	$< 3 \times 10^{-13} \text{ GeV}^{-4}$	"	[152], [14]*
$ c_{(I)00}^{(8)} $	$< 9 \times 10^{-13} \text{ GeV}^{-4}$	"	[152], [14]*
$ \sum_{jm} 2Y_{jm}(27^\circ, 6^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-25} \text{ GeV}^{-4}$	Astrophysical birefringence	[137]*
$ \sum_{jm} 2Y_{jm}(112^\circ, 279^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-25} \text{ GeV}^{-4}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(61^\circ, 229^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-24} \text{ GeV}^{-4}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(129^\circ, 333^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-24} \text{ GeV}^{-4}$	"	[137]*
$ \sum_{jm} 2Y_{jm}(98.2^\circ, 182.1^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-20} \text{ GeV}^{-4}$	"	[14]*
$ \sum_{jm} 2Y_{jm}(87.3^\circ, 37.3^\circ)(k_{(E)jm}^{(8)} + ik_{(B)jm}^{(8)}) $	$\lesssim 10^{-20} \text{ GeV}^{-4}$	"	[14]*

Table D18. Nonminimal photon sector,  $d \geq 9$ 

Combination	Result	System	Ref.
$ \sum_{jm} Y_{jm}(27^\circ, 6^\circ) k_{(V)jm}^{(9)} $	$< 2 \times 10^{-22} \text{ GeV}^{-5}$	Astrophysical birefringence	[137]*
$ k_{(V)00}^{(9)} $	$< 8 \times 10^{-22} \text{ GeV}^{-5}$	"	[137]*
$ \sum_{jm} Y_{jm}(112^\circ, 279^\circ) k_{(V)jm}^{(9)} $	$< 2 \times 10^{-21} \text{ GeV}^{-5}$	"	[137]*
$ k_{(V)00}^{(9)} $	$< 7 \times 10^{-21} \text{ GeV}^{-5}$	"	[137]*
$ \sum_{jm} Y_{jm}(61^\circ, 229^\circ) k_{(V)jm}^{(9)} $	$< 2 \times 10^{-20} \text{ GeV}^{-5}$	"	[137]*
$ k_{(V)00}^{(9)} $	$< 7 \times 10^{-20} \text{ GeV}^{-5}$	"	[137]*
$ \sum_{jm} Y_{jm}(129^\circ, 333^\circ) k_{(V)jm}^{(9)} $	$< 5 \times 10^{-21} \text{ GeV}^{-5}$	"	[137]*
$ k_{(V)00}^{(9)} $	$< 2 \times 10^{-20} \text{ GeV}^{-5}$	"	[137]*
$ \sum_{jm} Y_{jm}(98.2^\circ, 182.1^\circ) k_{(V)jm}^{(9)} $	$< 6 \times 10^{-16} \text{ GeV}^{-5}$	"	[14]*
$ k_{(V)00}^{(9)} $	$< 2 \times 10^{-15} \text{ GeV}^{-5}$	"	[14]*
$ \sum_{jm} Y_{jm}(87.3^\circ, 37.3^\circ) k_{(V)jm}^{(9)} $	$< 1 \times 10^{-16} \text{ GeV}^{-5}$	"	[14]*
$ k_{(V)00}^{(9)} $	$< 4 \times 10^{-16} \text{ GeV}^{-5}$	"	[14]*

Table D19. Muon sector,  $d = 3$ 

Combination	Result	System	Ref.
$ \text{Re } H_{011}^{\text{NR}(0B)} ,  \text{Im } H_{011}^{\text{NR}(0B)} ,  \text{Re } g_{011}^{\text{NR}(0B)} ,  \text{Im } g_{011}^{\text{NR}(0B)} $	$< 2 \times 10^{-22}$ GeV	Muonium spectroscopy	[20]*
$ \text{Re } H_{011}^{\text{NR}(1B)} ,  \text{Im } H_{011}^{\text{NR}(1B)} ,  \text{Re } g_{011}^{\text{NR}(1B)} ,  \text{Im } g_{011}^{\text{NR}(1B)} $	$< 7 \times 10^{-23}$ GeV	"	[20]*
$b^T/m_\mu$	$(7.3 \pm 5.0) \times 10^{-7}$	Muon decay	[153]*
$b_Z$	$-(1.0 \pm 1.1) \times 10^{-23}$ GeV	BNL $g_\mu - 2$	[154]
$\sqrt{(\check{b}_X^{\mu+})^2 + (\check{b}_Y^{\mu+})^2}$	$< 1.4 \times 10^{-24}$ GeV	"	[154]
$\sqrt{(\check{b}_X^{\mu-})^2 + (\check{b}_Y^{\mu-})^2}$	$< 2.6 \times 10^{-24}$ GeV	"	[154]
$\sqrt{(\check{b}_X)^2 + (\check{b}_Y)^2}$	$< 2 \times 10^{-23}$ GeV	Muonium spectroscopy	[155]
$b_Z - 1.19(m_\mu d_{Z0} + H_{XY})$	$(-1.4 \pm 1.0) \times 10^{-22}$ GeV	BNL, CERN $g_\mu - 2$ data	[156]
$b_Z$	$(-2.3 \pm 1.4) \times 10^{-22}$ GeV	CERN $g_\mu - 2$ data	[156], [157]*
$ \text{Re } H_{011}^{(3)(0B)} ,  \text{Im } H_{011}^{(3)(0B)} $	$< 5 \times 10^{-23}$ GeV	"	[20]*
$\check{H}_{010}^{(3)}$	$(-1.6 \pm 1.7) \times 10^{-22}$ GeV	BNL, CERN $g_\mu - 2$ data	[20]*
$ \text{Re } \check{H}_{011}^{(3)} ,  \text{Im } \check{H}_{011}^{(3)} $	$< 2.0 \times 10^{-24}$ GeV	BNL $g_\mu - 2$	[20]*
$m_\mu d_{Z0} + H_{XY}$	$(1.8 \pm 6.0) \times 10^{-23}$ GeV	"	[154]

Table D20. Muon sector,  $d = 4$ 

Combination	Result	System	Ref.
$c^{TT} - 0.05c^{ZZ}$	$(4.9 \pm 1.1) \times 10^{-8}$	Muon decay	[153]*
$ c $	$< 10^{-11}$	Astrophysics	[60]*
$ \text{Re } g_{011}^{(4)(0B)} ,  \text{Im } g_{011}^{(4)(0B)} $	$< 5 \times 10^{-22}$	Muonium spectroscopy	[20]*
$ \text{Re } \check{g}_{011}^{(4)} ,  \text{Im } \check{g}_{011}^{(4)} $	$< 6.6 \times 10^{-25}$	BNL $g_\mu - 2$	[20]*
$\check{g}_{010}^{(4)}$	$(-2.3 \pm 2.4) \times 10^{-25}$	"	[20]*
$m_\mu g_{XYT}^{(M)}$	$-(7.8 \pm 8.5) \times 10^{-27}$ GeV	"	[20]*
$m_\mu \sqrt{(g_{XZT}^{(M)})^2 + (g_{YZT}^{(M)})^2}$	$< 1.1 \times 10^{-27}$ GeV	"	[20]*



Table D21. Nonminimal muon sector,  $d \geq 5$ 

Combination	Result	System	Ref.
$ \check{a}_2^{\text{NR}} $	$< 8 \times 10^{-6} \text{ GeV}^{-1}$	Muonium spectroscopy	[20]*
$ \check{c}_2^{\text{NR}} $	$< 8 \times 10^{-6} \text{ GeV}^{-1}$	"	[20]*
$ \text{Re } H_{211}^{\text{NR}(0B)} ,  \text{Im } H_{211}^{\text{NR}(0B)} ,  \text{Re } g_{211}^{\text{NR}(0B)} ,  \text{Im } g_{211}^{\text{NR}(0B)} $	$< 1 \times 10^{-11} \text{ GeV}^{-1}$	"	[20]*
$ \text{Re } H_{211}^{\text{NR}(1B)} ,  \text{Im } H_{211}^{\text{NR}(1B)} ,  \text{Re } g_{211}^{\text{NR}(1B)} ,  \text{Im } g_{211}^{\text{NR}(1B)} $	$< 6 \times 10^{-12} \text{ GeV}^{-1}$	"	[20]*
$\check{a}^{\text{UR}(5)} - m_\mu \check{g}^{\text{UR}(6)}$	$(-1 \text{ to } 1) \times 10^{-34} \text{ GeV}^{-1}$	Astrophysics	[65]*, [18]*
$ \text{Re } H_{011}^{(5)(0B)} ,  \text{Im } H_{011}^{(5)(0B)} $	$< 5 \times 10^{-21} \text{ GeV}^{-1}$	Muonium spectroscopy	[20]*
$ \text{Re } \check{H}_{011}^{(5)} ,  \text{Im } \check{H}_{011}^{(5)} ,  \text{Re } \check{H}_{211}^{(5)} ,  \text{Im } \check{H}_{211}^{(5)} $	$< 2.1 \times 10^{-25} \text{ GeV}^{-1}$	BNL $g_\mu - 2$	[20]*
$ \text{Re } \check{H}_{231}^{(5)} ,  \text{Im } \check{H}_{231}^{(5)} $	$< 1.3 \times 10^{-25} \text{ GeV}^{-1}$	"	[20]*
$\check{H}_{010}^{(5)}, \check{H}_{210}^{(5)}$	$(-1.7 \pm 1.7) \times 10^{-23} \text{ GeV}^{-1}$	BNL, CERN $g_\mu - 2$ data	[20]*
$\check{H}_{230}^{(5)}$	$(2.9 \pm 3.0) \times 10^{-24} \text{ GeV}^{-1}$	"	[20]*
$\check{c}^{\text{UR}(6)}$	$(-8.5 \text{ to } 0.0025) \times 10^{-20} \text{ GeV}^{-2}$	Astrophysics	[65]*, [18]*
$ \text{Re } g_{011}^{(6)(0B)} ,  \text{Im } g_{011}^{(6)(0B)} $	$< 5 \times 10^{-20} \text{ GeV}^{-2}$	Muonium spectroscopy	[20]*
$ \text{Re } \check{g}_{011}^{(6)} ,  \text{Im } \check{g}_{011}^{(6)} ,  \text{Re } \check{g}_{211}^{(6)} ,  \text{Im } \check{g}_{211}^{(6)} $	$< 6.8 \times 10^{-26} \text{ GeV}^{-2}$	BNL $g_\mu - 2$	[20]*
$ \text{Re } \check{g}_{231}^{(6)} ,  \text{Im } \check{g}_{231}^{(6)} $	$< 4.3 \times 10^{-26} \text{ GeV}^{-2}$	"	[20]*
$\check{g}_{010}^{(6)}, \check{g}_{210}^{(6)}$	$(-2.4 \pm 2.5) \times 10^{-26} \text{ GeV}^{-2}$	"	[20]*
$\check{g}_{230}^{(6)}$	$(-2.5 \pm 2.5) \times 10^{-26} \text{ GeV}^{-2}$	"	[20]*
$ \check{a}_4^{\text{NR}} $	$< 1 \times 10^5 \text{ GeV}^{-3}$	Muonium spectroscopy	[20]*
$ \check{a}_4^{\text{NR}} $	$< 1 \times 10^6 \text{ GeV}^{-3}$	"	[20]*
$ \check{c}_4^{\text{NR}} $	$< 1 \times 10^5 \text{ GeV}^{-3}$	"	[20]*
$ \check{c}_4^{\text{NR}} $	$< 1 \times 10^6 \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } H_{411}^{\text{NR}(0B)} ,  \text{Im } H_{411}^{\text{NR}(0B)} ,  \text{Re } g_{411}^{\text{NR}(0B)} ,  \text{Im } g_{411}^{\text{NR}(0B)} $	$< 2 \times 10^{-1} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } H_{411}^{\text{NR}(1B)} ,  \text{Im } H_{411}^{\text{NR}(1B)} ,  \text{Re } g_{411}^{\text{NR}(1B)} ,  \text{Im } g_{411}^{\text{NR}(1B)} $	$< 8 \times 10^{-2} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } H_{011}^{(7)(0B)} ,  \text{Im } H_{011}^{(7)(0B)} $	$< 4 \times 10^{-19} \text{ GeV}^{-3}$	Muonium spectroscopy	[20]*
$\check{H}_{010}^{(7)}, \check{H}_{210}^{(7)}, \check{H}_{410}^{(7)}$	$(-1.7 \pm 1.8) \times 10^{-24} \text{ GeV}^{-3}$	BNL, CERN $g_\mu - 2$ data	[20]*
$\check{H}_{230}^{(7)}, \check{H}_{430}^{(7)}$	$(3.0 \pm 3.1) \times 10^{-25} \text{ GeV}^{-3}$	"	[20]*
$\check{H}_{450}^{(7)}$	$(2.6 \pm 2.6) \times 10^{-25} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } \check{H}_{011}^{(7)} ,  \text{Im } \check{H}_{011}^{(7)} ,  \text{Re } \check{H}_{211}^{(7)} ,  \text{Im } \check{H}_{211}^{(7)} $	$< 2.2 \times 10^{-26} \text{ GeV}^{-3}$	BNL $g_\mu - 2$	[20]*
$ \text{Re } \check{H}_{411}^{(7)} ,  \text{Im } \check{H}_{411}^{(7)} $	$< 2.2 \times 10^{-26} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } \check{H}_{231}^{(7)} ,  \text{Im } \check{H}_{231}^{(7)} ,  \text{Re } \check{H}_{431}^{(7)} ,  \text{Im } \check{H}_{431}^{(7)} $	$< 1.4 \times 10^{-26} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } \check{H}_{451}^{(7)} ,  \text{Im } \check{H}_{451}^{(7)} $	$< 1.1 \times 10^{-26} \text{ GeV}^{-3}$	"	[20]*
$ \text{Re } g_{011}^{(8)(0B)} ,  \text{Im } g_{011}^{(8)(0B)} $	$< 4 \times 10^{-18} \text{ GeV}^{-4}$	Muonium spectroscopy	[20]*
$\check{g}_{010}^{(8)}, \check{g}_{210}^{(8)}, \check{g}_{410}^{(8)}$	$(-2.5 \pm 2.6) \times 10^{-27} \text{ GeV}^{-4}$	BNL $g_\mu - 2$	[20]*
$\check{g}_{230}^{(8)}, \check{g}_{430}^{(8)}$	$(-2.6 \pm 2.6) \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$\check{g}_{450}^{(8)}$	$(1.6 \pm 1.7) \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$ \text{Re } \check{g}_{011}^{(8)} ,  \text{Im } \check{g}_{011}^{(8)} ,  \text{Re } \check{g}_{211}^{(8)} ,  \text{Im } \check{g}_{211}^{(8)} $	$< 7.1 \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$ \text{Re } \check{g}_{411}^{(8)} ,  \text{Im } \check{g}_{411}^{(8)} $	$< 7.1 \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$ \text{Re } \check{g}_{231}^{(8)} ,  \text{Im } \check{g}_{231}^{(8)} ,  \text{Re } \check{g}_{431}^{(8)} ,  \text{Im } \check{g}_{431}^{(8)} $	$< 4.5 \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$ \text{Re } \check{g}_{451}^{(8)} ,  \text{Im } \check{g}_{451}^{(8)} $	$< 3.6 \times 10^{-27} \text{ GeV}^{-4}$	"	[20]*
$\check{H}_{010}^{(9)}, \check{H}_{210}^{(9)}, \check{H}_{410}^{(9)}, \check{H}_{610}^{(9)}$	$(-1.8 \pm 1.9) \times 10^{-25} \text{ GeV}^{-5}$	BNL, CERN $g_\mu - 2$ data	[20]*
$\check{H}_{230}^{(9)}, \check{H}_{430}^{(9)}, \check{H}_{630}^{(9)}$	$(3.2 \pm 3.3) \times 10^{-26} \text{ GeV}^{-5}$	"	[20]*
$\check{H}_{450}^{(9)}, \check{H}_{650}^{(9)}$	$(2.7 \pm 2.7) \times 10^{-26} \text{ GeV}^{-5}$	"	[20]*
$\check{H}_{670}^{(9)}$	$(-1.1 \pm 1.1) \times 10^{-26} \text{ GeV}^{-5}$	"	[20]*
$\check{g}_{010}^{(10)}, \check{g}_{210}^{(10)}, \check{g}_{410}^{(10)}, \check{g}_{610}^{(10)}$	$(-2.6 \pm 2.7) \times 10^{-28} \text{ GeV}^{-6}$	BNL $g_\mu - 2$	[20]*
$\check{g}_{230}^{(10)}, \check{g}_{430}^{(10)}, \check{g}_{630}^{(10)}$	$(-2.7 \pm 2.7) \times 10^{-28} \text{ GeV}^{-6}$	"	[20]*
$\check{g}_{450}^{(10)}, \check{g}_{650}^{(10)}$	$(1.7 \pm 1.7) \times 10^{-28} \text{ GeV}^{-6}$	"	[20]*
$\check{g}_{670}^{(10)}$	$(1.3 \pm 1.4) \times 10^{-28} \text{ GeV}^{-6}$	"	[20]*

Table D22. Tau sector,  $d \geq 3$ 

Combination	Result	System	Ref.
$\tilde{a}^{\text{UR}(5)} - m_\tau \tilde{g}^{\text{UR}(6)}$	$(-2 \text{ to } 2) \times 10^{-33} \text{ GeV}^{-1}$	Astrophysics	[65]*, [18]*
$ c $	$< 10^{-8}$	"	[60]*

Table D23. Neutrino sector,  $d = 2$ 

Combination	Result	System	Ref.
$(c_{\text{eff}}^{(2)})^{ee}_{10}$	$> -1 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$> -2 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*
$(c_{\text{eff}}^{(2)})^{\mu\mu}_{10}$	$> -2 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$> -4 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*
$(c_{\text{eff}}^{(2)})^{\tau\tau}_{10}$	$> -2 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$> -5 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*
$\text{Re}(c_{\text{eff}}^{(2)})^{e\mu}_{10}$	$< 1 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$< 3 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*
$ \text{Re}(c_{\text{eff}}^{(2)})^{e\mu}_{10} $	$< 1.5 \times 10^{-19} \text{ GeV}^2$	LSND	[15]*
"	$< 4.1 \times 10^{-20} \text{ GeV}^2$	MiniBooNE	[15]*
"	$< 4.3 \times 10^{-20} \text{ GeV}^2$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(2)})^{e\mu}_{11} $	$< 3.0 \times 10^{-20} \text{ GeV}^2$	LSND	[15]*
"	$< 3.1 \times 10^{-20} \text{ GeV}^2$	MiniBooNE	[15]*
"	$< 4.8 \times 10^{-20} \text{ GeV}^2$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(2)})^{e\mu}_{11} $	$< 3.0 \times 10^{-20} \text{ GeV}^2$	LSND	[15]*
"	$< 2.6 \times 10^{-20} \text{ GeV}^2$	MiniBooNE	[15]*
"	$< 5.1 \times 10^{-20} \text{ GeV}^2$	MiniBooNE $\bar{\nu}$	[15]*
$\text{Re}(c_{\text{eff}}^{(2)})^{e\tau}_{10}$	$> -3 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$> -8 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*
$\text{Re}(c_{\text{eff}}^{(2)})^{\mu\tau}_{10}$	$< 1 \times 10^{-17} \text{ GeV}^2$	Troitsk tritium decay	[158]*
"	$< 3 \times 10^{-17} \text{ GeV}^2$	Mainz tritium decay	[158]*

Table D24. Neutrino sector,  $d = 3$  (part 1 of 3)

Combination	Result	System	Ref.
$\hat{a}_{\text{of}}^{(3)}$	$(-26.5 \text{ to } 7.60) \times 10^{-6} \text{ GeV}$	Double beta decay	[159]
$ (a_{\text{of}}^{(3)})_{00} $	$< 3 \times 10^{-8} \text{ GeV}$	Tritium decay	[158]*
$ (a_{\text{of}}^{(3)})_{10} $	$< 3 \times 10^{-8} \text{ GeV}$	"	[158]*
$ \hat{a}^{(3)} $	$< 1.9 \times 10^{-7} \text{ GeV}$	IceCube meson threshold	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{00}^{e\mu} $	$< 1.3 \times 10^{-18} \text{ GeV}$	LSND	[15]*
"	$< 1.5 \times 10^{-19} \text{ GeV}$	MiniBooNE	[15]*
"	$< 9.2 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{10}^{e\mu} $	$< 3.8 \times 10^{-18} \text{ GeV}$	LSND	[15]*
"	$< 1.1 \times 10^{-19} \text{ GeV}$	MiniBooNE	[15]*
"	$< 7.1 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{11}^{e\mu} $	$< 7.6 \times 10^{-19} \text{ GeV}$	LSND	[15]*
"	$< 8.7 \times 10^{-20} \text{ GeV}$	MiniBooNE	[15]*
"	$< 8.1 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(3)})_{11}^{e\mu} $	$< 7.6 \times 10^{-19} \text{ GeV}$	LSND	[15]*
"	$< 7.2 \times 10^{-20} \text{ GeV}$	MiniBooNE	[15]*
"	$< 8.5 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{20}^{e\mu} $	$< 1.3 \times 10^{-18} \text{ GeV}$	LSND	[15]*
"	$< 2.0 \times 10^{-19} \text{ GeV}$	MiniBooNE	[15]*
"	$< 1.2 \times 10^{-19} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{21}^{e\mu} $	$< 1.7 \times 10^{-18} \text{ GeV}$	LSND	[15]*
"	$< 5.2 \times 10^{-20} \text{ GeV}$	MiniBooNE	[15]*
"	$< 4.8 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(3)})_{21}^{e\mu} $	$< 1.7 \times 10^{-18} \text{ GeV}$	LSND	[15]*
"	$< 4.3 \times 10^{-20} \text{ GeV}$	MiniBooNE	[15]*
"	$< 5.1 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(3)})_{22}^{e\mu} $	$< 6.9 \times 10^{-19} \text{ GeV}$	LSND	[15]*
$ \text{Im}(a_{\text{eff}}^{(3)})_{22}^{e\mu} $	$< 6.9 \times 10^{-19} \text{ GeV}$	"	[15]*
$ \hat{a}_{e\mu}^{(3)} $	$< 3.6 \times 10^{-19} \text{ GeV}$	"	[15]*
"	$< 4.2 \times 10^{-20} \text{ GeV}$	MiniBooNE	[15]*
"	$< 2.6 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[15]*
$\text{Re}(a_{e\mu}^T), \text{Im}(a_{e\mu}^T)$	$< 1.8 \times 10^{-23} \text{ GeV}$	Super-Kamiokande	[160]
$ \text{Re}(a_L)_{e\mu}^T ,  \text{Im}(a_L)_{e\mu}^T $	$< 4.2 \times 10^{-20} \text{ GeV}$	MiniBooNE	[161]
"	$< 2.6 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[161]
$(a_L)_{e\mu}^T$	$(-3.1 \pm 0.9) \times 10^{-20} \text{ GeV}$	MiniBooNE	[162]
$(a_L)_{e\bar{\mu}}^T$	$(0.2 \pm 1.0) \times 10^{-19} \text{ GeV}$	LSND	[162]
$ \text{Re}(a_L)_{e\mu}^X ,  \text{Im}(a_L)_{e\mu}^X $	$< 1.6 \times 10^{-21} \text{ GeV}$	Double Chooz	[163]
"	$< 6.0 \times 10^{-20} \text{ GeV}$	MiniBooNE	[161]
"	$< 5.6 \times 10^{-20} \text{ GeV}$	MiniBooNE $\bar{\nu}$	[161]
"	$< 2.2 \times 10^{-20} \text{ GeV}$	MINOS	[164]
$(a_L)_{e\mu}^X$	$(0.6 \pm 1.9) \times 10^{-20} \text{ GeV}$	MiniBooNE	[162]
$(a_L)_{e\bar{\mu}}^X$	$(4.2 \pm 1.5) \times 10^{-19} \text{ GeV}$	LSND	[162]

Table D24. Neutrino sector,  $d = 3$  (part 2 of 3)

Combination	Result	System	Ref.
$ \text{Re}(a_L)_{e\mu}^Y ,  \text{Im}(a_L)_{e\mu}^Y $	$< 6.1 \times 10^{-20}$ GeV	Double Chooz	[163]
”	$< 5.0 \times 10^{-20}$ GeV	MiniBooNE	[161]
”	$< 5.9 \times 10^{-20}$ GeV	MiniBooNE $\bar{\nu}$	[161]
”	$< 2.2 \times 10^{-20}$ GeV	MINOS	[164]
$(a_L)_{e\mu}^Y$	$(-0.9 \pm 1.8) \times 10^{-20}$ GeV	MiniBooNE	[162]
$(a_L)_{\bar{e}\bar{\mu}}^Y$	$(-1.7 \pm 1.8) \times 10^{-19}$ GeV	LSND	[162]
$ \text{Re}(a_L)_{e\mu}^Z ,  \text{Im}(a_L)_{e\mu}^Z $	$< 5.6 \times 10^{-20}$ GeV	MiniBooNE	[161]
”	$< 3.5 \times 10^{-20}$ GeV	MiniBooNE $\bar{\nu}$	[161]
$(a_L)_{e\mu}^Z$	$(-4.2 \pm 1.2) \times 10^{-20}$ GeV	MiniBooNE	[162]
$(a_L)_{\bar{e}\bar{\mu}}^Z$	$(1.0 \pm 5.4) \times 10^{-19}$ GeV	LSND	[162]
$-(a_L)_{e\mu}^T - 0.29(a_L)_{e\mu}^Z$	$(5.8 \pm 1.7) \times 10^{-20}$ GeV	Double Chooz	[165]
$-0.91(a_L)_{e\mu}^X + 0.29(a_L)_{e\mu}^Y$	$(-0.4 \pm 0.7) \times 10^{-20}$ GeV	”	[165]
$0.29(a_L)_{e\mu}^X + 0.91(a_L)_{e\mu}^Y$	$(0.5 \pm 0.8) \times 10^{-20}$ GeV	”	[165]
$ (a_L)_{e\mu}^T + 0.75(a_L)_{e\mu}^Z$ $-0.36[1.22(c_L)_{e\mu}^{TT} + 1.50(c_L)_{e\mu}^{TZ} + 0.34(c_L)_{e\mu}^{ZZ}]$ GeV	$(3.1 \pm 0.6 \pm 0.9) \times 10^{-20}$ GeV	MiniBooNE	[166]
$ -(a_L)_{e\mu}^T - 0.75(a_L)_{e\mu}^Z$ $-0.60[1.22(c_L)_{e\mu}^{TT} + 1.50(c_L)_{e\mu}^{TZ} + 0.34(c_L)_{e\mu}^{ZZ}]$ GeV	$(0.1 \pm 0.8 \pm 0.1) \times 10^{-20}$ GeV	”	[166]
$ 0.66(a_L)_{e\mu}^Y$ $-0.36[1.33(c_L)_{e\mu}^{TY} + 0.99(c_L)_{e\mu}^{YZ}]$ GeV	$(0.6 \pm 0.9 \pm 0.3) \times 10^{-20}$ GeV	”	[166]
$  -0.66(a_L)_{e\mu}^Y$ $-0.60[1.33(c_L)_{e\mu}^{TY} + 0.99(c_L)_{e\mu}^{YZ}]$ GeV	$(2.4 \pm 1.3 \pm 0.5) \times 10^{-20}$ GeV	”	[166]
$ 0.66(a_L)_{e\mu}^X$ $-0.36[1.33(c_L)_{e\mu}^{TX} + 0.99(c_L)_{e\mu}^{XZ}]$ GeV	$(0.4 \pm 0.9 \pm 0.4) \times 10^{-20}$ GeV	”	[166]
$  -0.66(a_L)_{e\mu}^X$ $-0.60[1.33(c_L)_{e\mu}^{TX} + 0.99(c_L)_{e\mu}^{XZ}]$ GeV	$(2.1 \pm 1.2 \pm 0.4) \times 10^{-20}$ GeV	”	[166]
$ (C)_{\bar{e}\bar{\mu}} ^2$	$(10.7 \pm 2.6 \pm 1.3) \times (10^{-19} \text{ GeV})^2$	LSND	[167]
$ (C)_{\bar{e}\bar{\mu}} ^2 + \frac{1}{2} (\mathcal{A}_s)_{\bar{e}\bar{\mu}} ^2 + \frac{1}{2} (\mathcal{A}_c)_{\bar{e}\bar{\mu}} ^2$	$(9.9 \pm 2.3 \pm 1.4) \times (10^{-19} \text{ GeV})^2$	”	[167]
$ (C)_{\bar{e}\bar{\mu}} ^2 + \frac{1}{2} (\mathcal{A}_s)_{\bar{e}\bar{\mu}} ^2 + \frac{1}{2} (\mathcal{A}_c)_{\bar{e}\bar{\mu}} ^2$ $+ \frac{1}{2} (\mathcal{B}_s)_{\bar{e}\bar{\mu}} ^2 + \frac{1}{2} (\mathcal{B}_c)_{\bar{e}\bar{\mu}} ^2$	$(10.5 \pm 2.4 \pm 1.4) \times (10^{-19} \text{ GeV})^2$	”	[167]
$\text{Re}(a_{e\tau}^T)$	$< 4.1 \times 10^{-23}$ GeV	Super-Kamiokande	[160]
$\text{Im}(a_{e\tau}^T)$	$< 2.8 \times 10^{-23}$ GeV	”	[160]
$ \text{Re}(a_L)_{e\tau}^T ,  \text{Im}(a_L)_{e\tau}^T $	$< 7.8 \times 10^{-20}$ GeV	Double Chooz	[163]
$ \text{Re}(a_L)_{e\tau}^X ,  \text{Im}(a_L)_{e\tau}^X $	$< 4.4 \times 10^{-20}$ GeV	”	[163]
$ \text{Re}(a_L)_{e\tau}^Y ,  \text{Im}(a_L)_{e\tau}^Y $	$< 9.0 \times 10^{-20}$ GeV	”	[163]
$ \text{Re}(a_L)_{e\tau}^Z ,  \text{Im}(a_L)_{e\tau}^Z $	$< 2.7 \times 10^{-19}$ GeV	”	[163]
$ -(a_L)_{e\tau}^T - 0.29(a_L)_{e\tau}^Z + 0.0042[-1.46(c_L)_{e\tau}^{TT}$ $-0.57(c_L)_{e\tau}^{TZ} + 0.38(c_L)_{e\tau}^{ZZ}]$ GeV	$< 7.8 \times 10^{-20}$ GeV	”	[165]
$  -0.91(a_L)_{e\tau}^X + 0.29(a_L)_{e\tau}^Y + 0.0042[-1.83(c_L)_{e\tau}^{TX}$ $+0.58(c_L)_{e\tau}^{TY} - 0.52(c_L)_{e\tau}^{XZ} + 0.16(c_L)_{e\tau}^{YZ}]$ GeV	$< 6.6 \times 10^{-20}$ GeV	”	[165]
$ 0.29(a_L)_{e\tau}^X + 0.91(a_L)_{e\tau}^Y + 0.0042[0.58(c_L)_{e\tau}^{TX}$ $+1.83(c_L)_{e\tau}^{TY} + 0.16(c_L)_{e\tau}^{XZ} + 0.52(c_L)_{e\tau}^{YZ}]$ GeV	$< 7.0 \times 10^{-20}$ GeV	”	[165]
$ 0.0042[0.26((c_L)_{e\tau}^{XX} - (c_L)_{e\tau}^{YY}) + 0.75(c_L)_{e\tau}^{XY}]$ GeV	$< 5.4 \times 10^{-20}$ GeV	”	[165]
$ 0.0042[0.38((c_L)_{e\tau}^{XX} - (c_L)_{e\tau}^{YY}) - 0.53(c_L)_{e\tau}^{XY}]$ GeV	$< 5.4 \times 10^{-20}$ GeV	”	[165]

Table D24. Neutrino sector,  $d = 3$  (part 3 of 3)

Combination	Result	System	Ref.
$ (a_L)_{\mu\mu}^0 - (a_L)_{\tau\tau}^0 $	$< 6.3 \times 10^{-24}$ GeV	Super-Kamiokande	[168]*
$ \text{Re}(a_L)_{\mu\tau}^0 $	$< 1.4 \times 10^{-24}$ GeV	"	[168]*
$\text{Re}(a_L)_{\mu\tau}^0$	$(-1.95 \text{ to } 0.683) \times 10^{-23}$ GeV	MINOS	[168]*
$\text{Re}(a_{\mu\tau}^T)$	$< 6.5 \times 10^{-24}$ GeV	Super-Kamiokande	[160]
$\text{Im}(a_{\mu\tau}^T)$	$< 5.1 \times 10^{-24}$ GeV	"	[160]
$\text{Re}(a_{\mu\tau})$	$< 4.1 \times 10^{-24}$ GeV	"	[169]
$\text{Im}(a_{\mu\tau})$	$< 5.1 \times 10^{-24}$ GeV	"	[169]
$ \text{Re}(a_L)_{\mu\tau}^X $	$< 5.9 \times 10^{-23}$ GeV	MINOS	[164]
$ \text{Im}(a_L)_{\mu\tau}^X $	$< 2.2 \times 10^{-20}$ GeV	"	[164]
$ (a_L)_{\mu\tau}^X $	$< 1.8 \times 10^{-23}$ GeV	IceCube	[170]
"	$< 5.9 \times 10^{-23}$ GeV	MINOS FD	[171]
$ \text{Re}(a_L)_{\mu\tau}^Y $	$< 6.1 \times 10^{-23}$ GeV	MINOS	[164]
$ \text{Im}(a_L)_{\mu\tau}^Y $	$< 2.2 \times 10^{-20}$ GeV	"	[164]
$ (a_L)_{\mu\tau}^Y $	$< 1.8 \times 10^{-23}$ GeV	IceCube	[170]
"	$< 6.1 \times 10^{-23}$ GeV	MINOS FD	[171]
$ a_L^X ,  a_L^Y $	$< 3.0 \times 10^{-20}$ GeV	MINOS ND	[172]
$a \cos \rho$	excluded	Multiple	[173]*
$a \sin \rho \hat{n}$	excluded	"	[173]*
$b$	$< 1.6 \times 10^{-23}$ GeV	Atmospheric	[174]
$\check{a}/\check{c}$	$< 5$ GeV	"	[174]
$ \text{Re} \tilde{H}_{e\bar{\mu}}^X ,  \text{Im} \tilde{H}_{e\bar{\mu}}^X $	$< 3.3 \times 10^{-21}$ GeV	MINOS FD	[175]
$ \text{Re} \tilde{H}_{e\bar{\mu}}^Y ,  \text{Im} \tilde{H}_{e\bar{\mu}}^Y $	$< 3.3 \times 10^{-21}$ GeV	"	[175]
$ \text{Re} \tilde{H}_{e\bar{\mu}}^Z ,  \text{Im} \tilde{H}_{e\bar{\mu}}^Z $	$< 1.4 \times 10^{-19}$ GeV	Double Chooz	[176]
$ \text{Re} \tilde{H}_{e\bar{\tau}}^X ,  \text{Im} \tilde{H}_{e\bar{\tau}}^X $	$< 3.4 \times 10^{-21}$ GeV	MINOS FD	[175]
$ \text{Re} \tilde{H}_{e\bar{\tau}}^Y ,  \text{Im} \tilde{H}_{e\bar{\tau}}^Y $	$< 3.6 \times 10^{-21}$ GeV	"	[175]
$ \text{Re} \tilde{H}_{e\bar{\tau}}^Z ,  \text{Im} \tilde{H}_{e\bar{\tau}}^Z $	$< 1.4 \times 10^{-19}$ GeV	Double Chooz	[176]
$ \text{Re} \tilde{H}_{\mu\bar{\tau}}^X ,  \text{Im} \tilde{H}_{\mu\bar{\tau}}^X $	$< 1.1 \times 10^{-22}$ GeV	MINOS FD	[175]
$ \text{Re} \tilde{H}_{\mu\bar{\tau}}^Y ,  \text{Im} \tilde{H}_{\mu\bar{\tau}}^Y $	$< 1.0 \times 10^{-22}$ GeV	"	[175]
$ \text{Re} \tilde{H}_{\mu\bar{\tau}}^Z ,  \text{Im} \tilde{H}_{\mu\bar{\tau}}^Z $	$< 1.7 \times 10^{-18}$ GeV	Double Chooz	[176]

Table D25. Neutrino sector,  $d = 4$  (part 1 of 8)

Combination	Result	System	Ref.
$(c_{\text{of}}^{(4)})_{00}$	$> -4 \times 10^{-19}$	IceCube	[177]*
$ (c_{\text{of}}^{(4)})_{00} $	$< 7.1 \times 10^{-9}$	SN1987A time of flight	[15]*
"	$< 1.4 \times 10^{-4}$	Fermilab time of flight	[15]*
$(c_{\text{of}}^{(4)})_{00}$	$-8.4 \pm 1.1_{-0.9}^{+1.2} \times 10^{-5}$	OPERA time of flight	[15]*
"	$-1.8 \pm 1.0 \times 10^{-4}$	MINOS time of flight	[15]*
$(c_{\text{of}}^{(4)})_{10}$	$(-1 \text{ to } 4) \times 10^{-17}$	IceCube	[177]*
$ (c_{\text{of}}^{(4)})_{10} $	$< 4.4 \times 10^{-9}$	SN1987A time of flight	[15]*
"	$< 1.6 \times 10^{-4}$	Fermilab time of flight	[15]*
$(c_{\text{of}}^{(4)})_{10}$	$11.8 \pm 1.6_{-1.2}^{+1.7} \times 10^{-5}$	OPERA time of flight	[15]*
"	$-1.6 \pm 0.9 \times 10^{-4}$	MINOS time of flight	[15]*
$\text{Re}(c_{\text{of}}^{(4)})_{11}$	$(-3 \text{ to } 2) \times 10^{-17}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(4)})_{11} $	$< 7.7 \times 10^{-8}$	SN1987A time of flight	[15]*
$\text{Im}(c_{\text{of}}^{(4)})_{11}$	$(-2 \text{ to } 2) \times 10^{-17}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(4)})_{11} $	$< 8.2 \times 10^{-9}$	SN1987A time of flight	[15]*
$(c_{\text{of}}^{(4)})_{20}$	$(-1 \text{ to } 7) \times 10^{-17}$	IceCube	[177]*
$ (c_{\text{of}}^{(4)})_{20} $	$< 3.9 \times 10^{-9}$	SN1987A time of flight	[15]*
"	$< 6.2 \times 10^{-4}$	Fermilab time of flight	[15]*
$(c_{\text{of}}^{(4)})_{20}$	$15.2 \pm 2.1_{-1.5}^{+2.2} \times 10^{-5}$	OPERA time of flight	[15]*
"	$-5.6 \pm 3.2 \times 10^{-4}$	MINOS time of flight	[15]*
$\text{Re}(c_{\text{of}}^{(4)})_{21}$	$(-2 \text{ to } 3) \times 10^{-17}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(4)})_{21} $	$< 3.7 \times 10^{-8}$	SN1987A time of flight	[15]*
$\text{Im}(c_{\text{of}}^{(4)})_{21}$	$(-2 \text{ to } 5) \times 10^{-17}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(4)})_{21} $	$< 3.9 \times 10^{-9}$	SN1987A time of flight	[15]*
$\text{Re}(c_{\text{of}}^{(4)})_{22}$	$(-5 \text{ to } 2) \times 10^{-17}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(4)})_{22} $	$< 2.1 \times 10^{-8}$	SN1987A time of flight	[15]*
$\text{Im}(c_{\text{of}}^{(4)})_{22}$	$(-3 \text{ to } 4) \times 10^{-17}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(4)})_{22} $	$< 9.8 \times 10^{-8}$	SN1987A time of flight	[15]*

Table D25. Neutrino sector,  $d = 4$  (part 2 of 8)

Combination	Result	System	Ref.
$\hat{c}^{(4)}$	$> -5.2 \times 10^{-21}$	IceCube	[178]*
"	$> -2 \times 10^{-12}$	Atmospheric threshold	[177]*
"	$> -3 \times 10^{-13}$	Atmospheric Čerenkov	[177]*
"	$> -5 \times 10^{-19}$	Astrophysical Čerenkov	[177]*
"	$> -3 \times 10^{-19}$	IceCube	[179]*
$ \hat{c}^{(4)} $	$< 2.0 \times 10^{-9}$	SN1987A time of flight	[15]*
"	$< 4.0 \times 10^{-5}$	Fermilab time of flight	[15]*
$\hat{c}^{(4)}$	$-23.7 \pm 3.2_{-2.4}^{+3.4} \times 10^{-6}$	OPERA time of flight	[15]*
"	$-5.1 \pm 2.9 \times 10^{-5}$	MINOS time of flight	[15]*
"	$> -4.7 \times 10^{-13}$	IceCube meson threshold	[15]*
"	$> -10^{-12}$	"	[180]*, [15]*
"	$> -4 \times 10^{-15}$	"	[181]*, [15]*
"	$> -3 \times 10^{-5}$	17 GeV, Čerenkov emission	[15]*
"	$> -6 \times 10^{-6}$	43 GeV, Čerenkov emission	[15]*
"	$> -3 \times 10^{-11}$	100 TeV, Čerenkov emission	[15]*
"	$> -8.5 \times 10^{-12}$	"	[182]*, [15]*
"	$> -2 \times 10^{-8}$	ICARUS, Čerenkov emission	[183], [15]*
$ (c_L)_{(TX)} $	$< 2.3 \times 10^{-5}$	Pion decay	[184]*
$ (c_L)_{(TY)} $	$< 2.3 \times 10^{-5}$	"	[184]*
$ (c_L)_{-} $	$< 5.4 \times 10^{-5}$	"	[184]*
$ (c_L)_{(XY)} $	$< 5.4 \times 10^{-5}$	"	[184]*
$ (c_L)_{(XZ)} $	$< 4.0 \times 10^{-5}$	"	[184]*
$ (c_L)_{(YZ)} $	$< 4.0 \times 10^{-5}$	"	[184]*
$(c_L^{\nu e})_{00}$	$< 2 \times 10^{-11}$	Cosmic rays	[185]*

Table D25. Neutrino sector,  $d = 4$  (part 3 of 8)

Combination	Result	System	Ref.
$ \text{Re}(c_{\text{eff}}^{(4)})_{00}^{e\mu} $	$< 3.2 \times 10^{-17}$	LSND	[15]*
”	$< 4.1 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 1.5 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{10}^{e\mu} $	$< 9.6 \times 10^{-17}$	LSND	[15]*
”	$< 3.2 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 1.2 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{11}^{e\mu} $	$< 1.9 \times 10^{-17}$	LSND	[15]*
”	$< 2.4 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 1.3 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(4)})_{11}^{e\mu} $	$< 1.9 \times 10^{-17}$	LSND	[15]*
”	$< 2.0 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 1.4 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{20}^{e\mu} $	$< 3.3 \times 10^{-17}$	LSND	[15]*
”	$< 5.5 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 2.0 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{21}^{e\mu} $	$< 4.4 \times 10^{-17}$	LSND	[15]*
”	$< 1.4 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 8.0 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(4)})_{21}^{e\mu} $	$< 4.4 \times 10^{-17}$	LSND	[15]*
”	$< 1.2 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 8.5 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{22}^{e\mu} $	$< 1.7 \times 10^{-17}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(4)})_{22}^{e\mu} $	$< 1.7 \times 10^{-17}$	”	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{30}^{e\mu} $	$< 4.5 \times 10^{-17}$	LSND	[15]*
”	$< 2.0 \times 10^{-18}$	MiniBooNE	[15]*
”	$< 7.6 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{31}^{e\mu} $	$< 2.5 \times 10^{-17}$	LSND	[15]*
”	$< 1.4 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 8.0 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(4)})_{31}^{e\mu} $	$< 2.5 \times 10^{-17}$	LSND	[15]*
”	$< 1.2 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 8.4 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(4)})_{32}^{e\mu} $	$< 3.4 \times 10^{-17}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(4)})_{32}^{e\mu} $	$< 3.4 \times 10^{-17}$	LSND	[15]*
$ \hat{c}_{e\mu}^{(4)} $	$< 9.1 \times 10^{-18}$	LSND	[15]*
”	$< 1.2 \times 10^{-19}$	MiniBooNE	[15]*
”	$< 4.3 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[15]*



Table D25. Neutrino sector,  $d = 4$  (part 4 of 8)

Combination	Result	System	Ref.
$\text{Re}(c_{e\mu}^{TT}), \text{Im}(c_{e\mu}^{TT})$	$< 1.1 \times 10^{-26}$	Super-Kamiokande	[160]
$ \text{Re}(c_L)_{e\mu}^{XY} ,  \text{Im}(c_L)_{e\mu}^{XY} $	$< 2.2 \times 10^{-21}$	MINOS	[164]
$ \text{Re}(c_L)_{e\mu}^{XZ} ,  \text{Im}(c_L)_{e\mu}^{XZ} $	$< 1.1 \times 10^{-19}$	MiniBooNE	[161]
”	$< 6.2 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
”	$< 1.1 \times 10^{-21}$	MINOS	[164]
$(c_L)_{e\mu}^{XZ}$	$(-1.1 \pm 3.7) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{XZ}$	$(-2.7 \pm 1.0) \times 10^{-17}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{YZ} ,  \text{Im}(c_L)_{e\mu}^{YZ} $	$< 9.2 \times 10^{-20}$	MiniBooNE	[161]
”	$< 6.5 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
”	$< 1.1 \times 10^{-21}$	MINOS	[164]
$(c_L)_{e\mu}^{YZ}$	$(1.7 \pm 3.4) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{YZ}$	$(1.1 \pm 1.2) \times 10^{-17}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{XX} ,  \text{Im}(c_L)_{e\mu}^{XX} $	$< 4.6 \times 10^{-21}$	MINOS	[164]
$ \text{Re}(c_L)_{e\mu}^{YY} ,  \text{Im}(c_L)_{e\mu}^{YY} $	$< 4.5 \times 10^{-21}$	MINOS	[164]
$ \text{Re}(c_L)_{e\mu}^{ZZ} ,  \text{Im}(c_L)_{e\mu}^{ZZ} $	$< 3.4 \times 10^{-19}$	MiniBooNE	[161]
”	$< 1.3 \times 10^{-19}$	MiniBooNE $\bar{\nu}$	[161]
$(c_L)_{e\mu}^{ZZ}$	$(2.6 \pm 0.8) \times 10^{-19}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{ZZ}$	$(-1.1 \pm 5.9) \times 10^{-18}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{TT} ,  \text{Im}(c_L)_{e\mu}^{TT} $	$< 9.6 \times 10^{-20}$	MiniBooNE	[161]
”	$< 3.6 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
$(c_L)_{e\mu}^{TT}$	$(7.2 \pm 2.1) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{TT}$	$(0.3 \pm 1.8) \times 10^{-18}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{TX} ,  \text{Im}(c_L)_{e\mu}^{TX} $	$< 8.4 \times 10^{-20}$	MiniBooNE	[161]
”	$< 4.6 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
”	$< 9.0 \times 10^{-23}$	MINOS	[164]
$(c_L)_{e\mu}^{TX}$	$(-0.9 \pm 2.8) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{TX}$	$(-5.2 \pm 1.9) \times 10^{-18}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{TY} ,  \text{Im}(c_L)_{e\mu}^{TY} $	$< 6.9 \times 10^{-20}$	MiniBooNE	[161]
”	$< 4.9 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
”	$< 9.0 \times 10^{-23}$	MINOS	[164]
$(c_L)_{e\mu}^{TY}$	$(1.3 \pm 2.6) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{TY}$	$(2.1 \pm 2.2) \times 10^{-18}$	LSND	[162]
$ \text{Re}(c_L)_{e\mu}^{TZ} ,  \text{Im}(c_L)_{e\mu}^{TZ} $	$< 7.8 \times 10^{-20}$	MiniBooNE	[161]
”	$< 2.9 \times 10^{-20}$	MiniBooNE $\bar{\nu}$	[161]
$(c_L)_{e\mu}^{TZ}$	$(5.9 \pm 1.7) \times 10^{-20}$	MiniBooNE	[162]
$(c_L)_{\bar{e}\bar{\mu}}^{TZ}$	$(1.3 \pm 6.7) \times 10^{-18}$	LSND	[162]
$c$	excluded	Multiple	[173]*
”	$< 1.4 \times 10^{-26}$	Atmospheric	[174]

Table D25. Neutrino sector,  $d = 4$  (part 5 of 8)

Combination	Result	System	Ref.
$\text{Re}(c_{e\tau}^{TT})$	$< 1.2 \times 10^{-24}$	Super-Kamiokande	[160]
$\text{Im}(c_{e\tau}^{TT})$	$< 1.4 \times 10^{-24}$	"	[160]
$ \text{Re}(c_L)_{e\tau}^{XY} ,  \text{Im}(c_L)_{e\tau}^{XY} $	$< 3.4 \times 10^{-18}$	Double Chooz	[163]
$ \text{Re}(c_L)_{e\tau}^{XZ} ,  \text{Im}(c_L)_{e\tau}^{XZ} $	$< 1.8 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{YZ} ,  \text{Im}(c_L)_{e\tau}^{YZ} $	$< 3.8 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{XX} ,  \text{Im}(c_L)_{e\tau}^{XX} $	$< 3.9 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{YY} ,  \text{Im}(c_L)_{e\tau}^{YY} $	$< 3.9 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{ZZ} ,  \text{Im}(c_L)_{e\tau}^{ZZ} $	$< 4.9 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{TT} ,  \text{Im}(c_L)_{e\tau}^{TT} $	$< 1.3 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{TX} ,  \text{Im}(c_L)_{e\tau}^{TX} $	$< 5.2 \times 10^{-18}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{TY} ,  \text{Im}(c_L)_{e\tau}^{TY} $	$< 1.1 \times 10^{-17}$	"	[163]
$ \text{Re}(c_L)_{e\tau}^{TZ} ,  \text{Im}(c_L)_{e\tau}^{TZ} $	$< 3.2 \times 10^{-17}$	"	[163]

Table D25. Neutrino sector,  $d = 4$  (part 6 of 8)

Combination	Result	System	Ref.
$\text{Re}(c_{\mu\tau}^{TT})$	$< 5.8 \times 10^{-27}$	Super-Kamiokande	[160]
$\text{Im}(c_{\mu\tau}^{TT})$	$< 5.6 \times 10^{-27}$	"	[160]
$\text{Re}(c_{\mu\tau})$	$< 1.7 \times 10^{-27}$	"	[169]
$\text{Im}(c_{\mu\tau})$	$< 1.7 \times 10^{-27}$	"	[169]
$ \text{Re}(c_L)_{\mu\tau}^{XY} $	$< 1.2 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{XY} $	$< 2.2 \times 10^{-21}$	"	[164]
$ (c_L)_{\mu\tau}^{XY} $	$< 1.2 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{XZ} $	$< 0.7 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{XZ} $	$< 1.1 \times 10^{-21}$	"	[164]
$ (c_L)_{\mu\tau}^{XZ} $	$< 0.7 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{YZ} $	$< 0.7 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{YZ} $	$< 1.1 \times 10^{-21}$	"	[164]
$ (c_L)_{\mu\tau}^{YZ} $	$< 0.7 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{XX} $	$< 2.5 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{XX} $	$< 4.6 \times 10^{-21}$	"	[164]
$ (c_L)_{\mu\tau}^{XX} $	$< 2.5 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{YY} $	$< 2.4 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{YY} $	$< 4.5 \times 10^{-21}$	"	[164]
$ (c_L)_{\mu\tau}^{YY} $	$< 2.4 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{TX} $	$< 0.5 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{TX} $	$< 9.0 \times 10^{-23}$	"	[164]
$ (c_L)_{\mu\tau}^{TX} $	$< 3.7 \times 10^{-27}$	IceCube	[170]
"	$< 0.5 \times 10^{-23}$	MINOS FD	[171]
$ \text{Re}(c_L)_{\mu\tau}^{TY} $	$< 0.5 \times 10^{-23}$	MINOS	[164]
$ \text{Im}(c_L)_{\mu\tau}^{TY} $	$< 9.0 \times 10^{-23}$	"	[164]
$ (c_L)_{\mu\tau}^{TY} $	$< 3.7 \times 10^{-27}$	IceCube	[170]
"	$< 0.5 \times 10^{-23}$	MINOS FD	[171]
$ c_L^{XY} $	$< 2.7 \times 10^{-21}$	MINOS ND	[172]
$ c_L^{XZ} $	$< 1.3 \times 10^{-21}$	"	[172]
$ c_L^{YZ} $	$< 1.2 \times 10^{-21}$	"	[172]
$ c_L^{XX} $	$< 5.6 \times 10^{-21}$	"	[172]
$ c_L^{YY} $	$< 5.5 \times 10^{-21}$	"	[172]
$ c_L^{TX} ,  c_L^{TY} $	$< 9 \times 10^{-23}$	"	[172]

Table D25. Neutrino sector,  $d = 4$  (part 7 of 8)

Combination	Result	System	Ref.
$ \text{Re } \tilde{g}_{e\bar{e}}^{JK} ,  \text{Im } \tilde{g}_{e\bar{e}}^{JK} , (J, K = X, Y, Z)$	$< 1 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{e\bar{e}}^{ZT} ,  \text{Im } \tilde{g}_{e\bar{e}}^{ZT} $	$< 9.7 \times 10^{-18}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{e\bar{e}}^{ZZ} ,  \text{Im } \tilde{g}_{e\bar{e}}^{ZZ} $	$< 3.3 \times 10^{-17}$	"	[176]
$ \text{Re } \tilde{g}_{e\bar{e}}^{XT} ,  \text{Im } \tilde{g}_{e\bar{e}}^{XT} $	$< 7.6 \times 10^{-22}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{YT} ,  \text{Im } \tilde{g}_{e\bar{e}}^{YT} $	$< 7.6 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{XZ} ,  \text{Im } \tilde{g}_{e\bar{e}}^{XZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{YZ} ,  \text{Im } \tilde{g}_{e\bar{e}}^{YZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{ZX} ,  \text{Im } \tilde{g}_{e\bar{e}}^{ZX} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{ZY} ,  \text{Im } \tilde{g}_{e\bar{e}}^{ZY} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{XX} ,  \text{Im } \tilde{g}_{e\bar{e}}^{XX} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{YY} ,  \text{Im } \tilde{g}_{e\bar{e}}^{YY} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{XY} ,  \text{Im } \tilde{g}_{e\bar{e}}^{XY} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{e}}^{YX} ,  \text{Im } \tilde{g}_{e\bar{e}}^{YX} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{JK} , (J, K = X, Y, Z)$	$< 1 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{ZT} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{ZT} $	$< 2.7 \times 10^{-17}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{ZZ} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{ZZ} $	$< 9.3 \times 10^{-17}$	"	[176]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{XT} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{XT} $	$< 7.6 \times 10^{-22}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{YT} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{YT} $	$< 7.6 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{XZ} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{XZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{YZ} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{YZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{ZX} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{ZX} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{ZY} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{ZY} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{XX} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{XX} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{YY} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{YY} $	$< 2.1 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{XY} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{XY} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\mu}}^{YX} ,  \text{Im } \tilde{g}_{e\bar{\mu}}^{YX} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{JK} , (J, K = X, Y, Z)$	$< 2 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{ZT} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{ZT} $	$< 2.7 \times 10^{-17}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{ZZ} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{ZZ} $	$< 9.3 \times 10^{-17}$	"	[176]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{XT} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{XT} $	$< 8.2 \times 10^{-22}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{YT} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{YT} $	$< 7.6 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{XZ} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{XZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{YZ} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{YZ} $	$< 1.2 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{ZX} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{ZX} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{ZY} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{ZY} $	$< 1.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{XX} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{XX} $	$< 2.0 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{YY} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{YY} $	$< 2.1 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{XY} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{XY} $	$< 2.1 \times 10^{-21}$	"	[175]
$ \text{Re } \tilde{g}_{e\bar{\tau}}^{YX} ,  \text{Im } \tilde{g}_{e\bar{\tau}}^{YX} $	$< 2.0 \times 10^{-21}$	"	[175]

Table D25. Neutrino sector,  $d = 4$  (part 8 of 8)

Combination	Result	System	Ref.
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{JK} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{JK} , (J, K = X, Y, Z)$	$< 3 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{ZT} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{ZT} $	$< 2.3 \times 10^{-16}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{ZZ} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{ZZ} $	$< 8.1 \times 10^{-16}$	"	[176]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{XT} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{XT} $	$< 8.9 \times 10^{-24}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{YT} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{YT} $	$< 8.6 \times 10^{-24}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{XZ} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{XZ} $	$< 1.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{YZ} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{YZ} $	$< 1.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{ZX} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{ZX} $	$< 1.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{ZY} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{ZY} $	$< 1.1 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{XX} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{XX} $	$< 2.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{YY} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{YY} $	$< 2.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{XY} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{XY} $	$< 2.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\mu}}^{YX} ,  \text{Im } \tilde{g}_{\mu\bar{\mu}}^{YX} $	$< 2.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{JK} , (J, K = X, Y, Z)$	$< 3 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{ZT} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{ZT} $	$< 4.4 \times 10^{-16}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{ZZ} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{ZZ} $	$< 1.5 \times 10^{-15}$	"	[176]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{XT} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{XT} $	$< 8.6 \times 10^{-24}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{YT} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{YT} $	$< 8.4 \times 10^{-24}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{XZ} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{XZ} $	$< 1.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{YZ} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{YZ} $	$< 1.4 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{ZX} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{ZX} $	$< 1.1 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{ZY} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{ZY} $	$< 1.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{XX} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{XX} $	$< 2.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{YY} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{YY} $	$< 2.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{XY} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{XY} $	$< 2.3 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\mu\bar{\tau}}^{YX} ,  \text{Im } \tilde{g}_{\mu\bar{\tau}}^{YX} $	$< 2.2 \times 10^{-23}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{JK} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{JK} , (J, K = X, Y, Z)$	$< 9 \times 10^{-8}$	Double beta decay	[186]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{ZT} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{ZT} $	$< 2.3 \times 10^{-16}$	Double Chooz	[176]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{ZZ} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{ZZ} $	$< 8.1 \times 10^{-16}$	"	[176]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{XT} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{XT} $	$< 1.8 \times 10^{-22}$	MINOS FD	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{YT} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{YT} $	$< 1.8 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{XZ} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{XZ} $	$< 2.9 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{YZ} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{YZ} $	$< 2.9 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{ZX} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{ZX} $	$< 2.4 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{ZY} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{ZY} $	$< 2.4 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{XX} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{XX} $	$< 4.8 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{YY} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{YY} $	$< 4.8 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{XY} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{XY} $	$< 4.8 \times 10^{-22}$	"	[175]
$ \text{Re } \tilde{g}_{\tau\bar{\tau}}^{YX} ,  \text{Im } \tilde{g}_{\tau\bar{\tau}}^{YX} $	$< 4.8 \times 10^{-22}$	"	[175]

Table D26. Nonminimal neutrino sector,  $d = 5$  (part 1 of 3)

Combination	Result	System	Ref.
$(a_{\text{of}}^{(5)})_{00}$	$25.7 \pm 3.3_{-2.5}^{+3.5} \times 10^{-7} \text{ GeV}^{-1}$	OPERA time of flight	[15]*
"	$3.0 \pm 1.7 \times 10^{-5} \text{ GeV}^{-1}$	MINOS time of flight	[15]*
$ (a_{\text{of}}^{(5)})_{00} $	$< 2.4 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab time of flight	[15]*
"	$< 2.1 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 3.5 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 3.5 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$(a_{\text{of}}^{(5)})_{10}$	$-34.7 \pm 4.7_{-3.5}^{+5.0} \times 10^{-7} \text{ GeV}^{-1}$	OPERA time of flight	[15]*
"	$2.7 \pm 1.5 \times 10^{-5} \text{ GeV}^{-1}$	MINOS time of flight	[15]*
$ (a_{\text{of}}^{(5)})_{10} $	$< 2.7 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab time of flight	[15]*
"	$< 2.3 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 2.2 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.2 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{11} $	$< 3.8 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 3.8 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{11} $	$< 4.1 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 4.1 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$(a_{\text{of}}^{(5)})_{20}$	$-44.8 \pm 6.1_{-4.5}^{+6.4} \times 10^{-7} \text{ GeV}^{-1}$	OPERA time of flight	[15]*
"	$9.4 \pm 5.3 \times 10^{-5} \text{ GeV}^{-1}$	MINOS time of flight	[15]*
$ (a_{\text{of}}^{(5)})_{20} $	$< 1.0 \times 10^{-5} \text{ GeV}^{-1}$	Fermilab time of flight	[15]*
"	$< 9.0 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 2.0 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.0 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{21} $	$< 1.8 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 1.8 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{21} $	$< 2.0 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.0 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{22} $	$< 1.1 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 1.1 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{22} $	$< 4.9 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 4.9 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*

Table D26. Nonminimal neutrino sector,  $d = 5$  (part 2 of 3)

Combination	Result	System	Ref.
$(a_{\text{of}}^{(5)})_{30}$	$21.1 \pm 2.9_{-2.1}^{+3.0} \times 10^{-7} \text{ GeV}^{-1}$	OPERA time of flight	[15]*
"	$-4.1 \pm 2.3 \times 10^{-5} \text{ GeV}^{-1}$	MINOS time of flight	[15]*
$ (a_{\text{of}}^{(5)})_{30} $	$< 2.1 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab time of flight	[15]*
"	$< 1.8 \times 10^{-6} \text{ GeV}^{-1}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 2.1 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.1 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{31} $	$< 1.2 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 1.2 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{31} $	$< 1.3 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 1.3 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{32} $	$< 4.3 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 4.3 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{32} $	$< 2.0 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.0 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{of}}^{(5)})_{33} $	$< 8.6 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 8.6 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Im}(a_{\text{of}}^{(5)})_{33} $	$< 2.8 \times 10^{-6} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 2.8 \times 10^{-9} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$\sum_j 0\mathcal{N}_{j0}(86.7^\circ, 52.4^\circ, 47.5^\circ) (a_{\text{of}}^{(5)})_{j0}$	$10 \pm 19 \times 10^{-8} \text{ GeV}^{-1}$	OPERA time of flight	[15]*
$\hat{a}^{(5)}$	$69.7 \pm 9.4_{-7.1}^{+10.0} \times 10^{-8} \text{ GeV}^{-1}$	"	[15]*
"	$10 \pm 19 \times 10^{-8} \text{ GeV}^{-1}$	"	[15]*
"	$8.5 \pm 4.8 \times 10^{-6} \text{ GeV}^{-1}$	MINOS time of flight	[15]*
"	$< 1 \times 10^{-6} \text{ GeV}^{-1}$	17 GeV, Čerenkov emission	[15]*
"	$< 1 \times 10^{-7} \text{ GeV}^{-1}$	43 GeV, Čerenkov emission	[15]*
$ \hat{a}^{(5)} $	$< 6.7 \times 10^{-7} \text{ GeV}^{-1}$	Fermilab time of flight	[15]*
"	$< 5.8 \times 10^{-7} \text{ GeV}^{-1}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 1.2 \times 10^{-18} \text{ GeV}^{-1}$	IceCube meson threshold	[15]*
"	$< 4 \times 10^{-16} \text{ GeV}^{-1}$	100 TeV, Čerenkov emission	[15]*
"	$< 1.0 \times 10^{-7} \text{ GeV}^{-1}$	SN1987A time of flight	[15]*
"	$< 1.0 \times 10^{-10} \text{ GeV}^{-1}$	SN1987A dispersion	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)e\mu})_{00} $	$< 8.1 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 1.1 \times 10^{-18} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.6 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)e\mu})_{10} $	$< 2.4 \times 10^{-15} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 8.9 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.0 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)e\mu})_{11} $	$< 4.8 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 6.7 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.2 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)e\mu})_{11} $	$< 4.8 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 5.6 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*

Table D26. Nonminimal neutrino sector,  $d = 5$  (part 3 of 3)

Combination	Result	System	Ref.
$ \text{Re}(a_{\text{eff}}^{(5)})_{20}^{e\mu} $	$< 8.1 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 1.5 \times 10^{-18} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 3.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{21}^{e\mu} $	$< 1.1 \times 10^{-15} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 4.0 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 1.3 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{21}^{e\mu} $	$< 1.1 \times 10^{-15} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 3.3 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 1.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{22}^{e\mu} $	$< 4.3 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{22}^{e\mu} $	$< 4.3 \times 10^{-16} \text{ GeV}^{-1}$	"	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{30}^{e\mu} $	$< 1.1 \times 10^{-15} \text{ GeV}^{-1}$	"	[15]*
"	$< 5.7 \times 10^{-18} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 1.3 \times 10^{-18} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{31}^{e\mu} $	$< 6.3 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 4.0 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 1.3 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{31}^{e\mu} $	$< 6.3 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 3.3 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 1.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{32}^{e\mu} $	$< 8.4 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{32}^{e\mu} $	$< 8.4 \times 10^{-16} \text{ GeV}^{-1}$	"	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{40}^{e\mu} $	$< 1.1 \times 10^{-15} \text{ GeV}^{-1}$	"	[15]*
"	$< 1.1 \times 10^{-18} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{41}^{e\mu} $	$< 6.5 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 7.2 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.4 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{41}^{e\mu} $	$< 6.5 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
"	$< 5.9 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 2.5 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(a_{\text{eff}}^{(5)})_{42}^{e\mu} $	$< 6.8 \times 10^{-16} \text{ GeV}^{-1}$	LSND	[15]*
$ \text{Im}(a_{\text{eff}}^{(5)})_{42}^{e\mu} $	$< 6.8 \times 10^{-16} \text{ GeV}^{-1}$	"	[15]*
$ \hat{a}_{e\mu}^{(5)} $	$< 2.3 \times 10^{-16} \text{ GeV}^{-1}$	"	[15]*
"	$< 3.2 \times 10^{-19} \text{ GeV}^{-1}$	MiniBooNE	[15]*
"	$< 7.2 \times 10^{-20} \text{ GeV}^{-1}$	MiniBooNE $\bar{\nu}$	[15]*



Table D27. Nonminimal neutrino sector,  $d = 6$  (part 1 of 4)

Combination	Result	System	Ref.
$(c_{\text{of}}^{(6)})_{00}$	$> -3 \times 10^{-31} \text{ GeV}^{-2}$	IceCube	[177]*
"	$-9.7 \pm 1.3_{-1.0}^{+1.4} \times 10^{-8} \text{ GeV}^{-2}$	OPERA time of flight	[15]*
"	$-6.7 \pm 3.8 \times 10^{-6} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
$ (c_{\text{of}}^{(6)})_{00} $	$< 5.3 \times 10^{-8} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 2.4 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 7.9 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$(c_{\text{of}}^{(6)})_{10}$	$(-2 \text{ to } 9) \times 10^{-28} \text{ GeV}^{-2}$	IceCube	[177]*
"	$13.6 \pm 1.8_{-1.3}^{+2.0} \times 10^{-8} \text{ GeV}^{-2}$	OPERA time of flight	[15]*
"	$-5.9 \pm 3.4 \times 10^{-6} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
$ (c_{\text{of}}^{(6)})_{10} $	$< 5.9 \times 10^{-8} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 1.5 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.9 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{11}$	$(-6 \text{ to } 5) \times 10^{-28} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{11} $	$< 2.6 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 8.6 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{11}$	$(-3 \text{ to } 3) \times 10^{-28} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{11} $	$< 2.7 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 9.1 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$(c_{\text{of}}^{(6)})_{20}$	$(-0.4 \text{ to } 7) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
"	$17.6 \pm 2.4_{-1.8}^{+2.5} \times 10^{-8} \text{ GeV}^{-2}$	OPERA time of flight	[15]*
"	$2.1 \pm 1.2 \times 10^{-5} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
$ (c_{\text{of}}^{(6)})_{20} $	$< 2.3 \times 10^{-7} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 1.3 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.3 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{21}$	$(-1 \text{ to } 2) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{21} $	$< 1.2 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.1 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{21}$	$(-1 \text{ to } 3) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{21} $	$< 1.3 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.4 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{22}$	$(-5 \text{ to } 0.6) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{22} $	$< 7.0 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.3 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{22}$	$(-1 \text{ to } 4) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{22} $	$< 3.3 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 1.1 \times 10^{-7} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*

Table D27. Nonminimal neutrino sector,  $d = 6$  (part 2 of 4)

Combination	Result	System	Ref.
$(c_{\text{of}}^{(6)})_{30}$	$(-1 \text{ to } 0.4) \times 10^{-26} \text{ GeV}^{-2}$	IceCube	[177]*
"	$-8.3 \pm 1.1_{-0.8}^{+1.2} \times 10^{-8}$	OPERA time of flight	[15]*
"	$9.1 \pm 5.4 \times 10^{-6} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
$ (c_{\text{of}}^{(6)})_{30} $	$< 4.6 \times 10^{-8} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 1.4 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.6 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{31}$	$(-0.2 \text{ to } 1) \times 10^{-26} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{31} $	$< 8.1 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.7 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{31}$	$(-5 \text{ to } 3) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{31} $	$< 8.7 \times 10^{-6} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.9 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{32}$	$(-0.2 \text{ to } 1) \times 10^{-26} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{32} $	$< 2.8 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 9.5 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{32}$	$(-4 \text{ to } 6) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{32} $	$< 1.3 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 4.4 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{33}$	$(-5 \text{ to } 6) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{33} $	$< 5.7 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 1.9 \times 10^{-7} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{33}$	$(-1 \text{ to } 0.07) \times 10^{-26} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{33} $	$< 1.9 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 6.3 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$(c_{\text{of}}^{(6)})_{40}$	$(-5 \text{ to } 2) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
"	$24.2 \pm 3.3_{-2.5}^{+3.5} \times 10^{-8} \text{ GeV}^{-2}$	OPERA time of flight	[15]*
"	$5.2 \pm 3.0 \times 10^{-6} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
$ (c_{\text{of}}^{(6)})_{40} $	$< 5.6 \times 10^{-8} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 1.8 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 5.9 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{41}$	$(-1 \text{ to } 3) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{41} $	$< 6.4 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.1 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{41}$	$(-1 \text{ to } 0.6) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{41} $	$< 6.9 \times 10^{-6} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.3 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{42}$	$(-1 \text{ to } 3) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{42} $	$< 1.6 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 5.3 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{42}$	$(-2 \text{ to } 1) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{42} $	$< 7.3 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.4 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*

Table D27. Nonminimal neutrino sector,  $d = 6$  (part 3 of 4)

Combination	Result	System	Ref.
$\text{Re}(c_{\text{of}}^{(6)})_{43}$	$(-1 \text{ to } 0.9) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{43} $	$< 2.0 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 6.8 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{43}$	$(-2 \text{ to } 1) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{43} $	$< 6.8 \times 10^{-5} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.3 \times 10^{-8} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Re}(c_{\text{of}}^{(6)})_{44}$	$(-2 \text{ to } 1) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Re}(c_{\text{of}}^{(6)})_{44} $	$< 5.3 \times 10^{-4} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 1.8 \times 10^{-7} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\text{Im}(c_{\text{of}}^{(6)})_{44}$	$(-0.5 \text{ to } 1) \times 10^{-27} \text{ GeV}^{-2}$	IceCube	[177]*
$ \text{Im}(c_{\text{of}}^{(6)})_{44} $	$< 1.2 \times 10^{-3} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 3.9 \times 10^{-7} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$\check{c}^{(6)}$	$> -5.23 \times 10^{-35} \text{ GeV}^{-2}$	IceCube	[178]*
"	$> -2 \times 10^{-24} \text{ GeV}^{-2}$	Atmospheric threshold	[177]*
"	$> -3 \times 10^{-25} \text{ GeV}^{-2}$	Atmospheric Čerenkov	[177]*
"	$> -5 \times 10^{-31} \text{ GeV}^{-2}$	Astrophysical Čerenkov	[177]*
"	$-27.3 \pm 3.7_{-2.8}^{+3.9} \times 10^{-9} \text{ GeV}^{-2}$	OPERA time of flight	[15]*
"	$-1.9 \pm 1.1 \times 10^{-6} \text{ GeV}^{-2}$	MINOS time of flight	[15]*
"	$> -2.9 \times 10^{-24} \text{ GeV}^{-2}$	IceCube meson threshold	[15]*
"	$> -8 \times 10^{-8} \text{ GeV}^{-2}$	17 GeV, Čerenkov emission	[15]*
"	$> -3 \times 10^{-9} \text{ GeV}^{-2}$	43 GeV, Čerenkov emission	[15]*
"	$> -3 \times 10^{-21} \text{ GeV}^{-2}$	100 TeV, Čerenkov emission	[15]*
$ \check{c}^{(6)} $	$< 1.5 \times 10^{-8} \text{ GeV}^{-2}$	Fermilab time of flight	[15]*
"	$< 6.7 \times 10^{-6} \text{ GeV}^{-2}$	SN1987A time of flight	[15]*
"	$< 2.2 \times 10^{-9} \text{ GeV}^{-2}$	SN1987A dispersion	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{00}^{e\mu} $	$< 2.0 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
"	$< 3.2 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
"	$< 4.3 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{10}^{e\mu} $	$< 6.0 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
"	$< 2.5 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
"	$< 3.3 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{11}^{e\mu} $	$< 1.2 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
"	$< 1.9 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
"	$< 3.7 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{11}^{e\mu} $	$< 1.2 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
"	$< 1.5 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
"	$< 3.9 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*

Table D27. Nonminimal neutrino sector,  $d = 6$  (part 4 of 4)

Combination	Result	System	Ref.
$ \text{Re}(c_{\text{eff}}^{(6)})_{20}^{e\mu} $	$< 2.0 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 4.2 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 5.6 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{21}^{e\mu} $	$< 2.7 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 1.1 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 2.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{21}^{e\mu} $	$< 2.7 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 9.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 2.4 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{22}^{e\mu} $	$< 1.1 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{22}^{e\mu} $	$< 1.1 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{30}^{e\mu} $	$< 2.8 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
”	$< 1.6 \times 10^{-17} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 2.1 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{31}^{e\mu} $	$< 1.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 1.1 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 2.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{31}^{e\mu} $	$< 1.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 9.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 2.3 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{32}^{e\mu} $	$< 2.1 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{32}^{e\mu} $	$< 2.1 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{40}^{e\mu} $	$< 2.8 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
”	$< 3.0 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 4.0 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{41}^{e\mu} $	$< 1.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 2.0 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 4.0 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{41}^{e\mu} $	$< 1.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 1.6 \times 10^{-17} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 4.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{42}^{e\mu} $	$< 1.7 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{42}^{e\mu} $	$< 1.7 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{50}^{e\mu} $	$< 2.0 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 2.3 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 3.1 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{51}^{e\mu} $	$< 2.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 7.7 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 1.5 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{51}^{e\mu} $	$< 2.6 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 6.3 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 1.6 \times 10^{-18} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*
$ \text{Re}(c_{\text{eff}}^{(6)})_{52}^{e\mu} $	$< 1.4 \times 10^{-14} \text{ GeV}^{-2}$	LSND	[15]*
$ \text{Im}(c_{\text{eff}}^{(6)})_{52}^{e\mu} $	$< 1.4 \times 10^{-14} \text{ GeV}^{-2}$	”	[15]*
$ \tilde{c}_{e\mu}^{(6)} $	$< 5.7 \times 10^{-15} \text{ GeV}^{-2}$	LSND	[15]*
”	$< 9.0 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE	[15]*
”	$< 1.2 \times 10^{-19} \text{ GeV}^{-2}$	MiniBooNE $\bar{\nu}$	[15]*

Table D28. Nonminimal neutrino sector,  $d = 7$ 

Combination	Result	System	Ref.
$\overset{\circ}{a}^{(7)}$	$12.1 \pm 1.6_{-1.2}^{+1.7} \times 10^{-10} \text{ GeV}^{-3}$	OPERA time of flight	[15]*
"	$4.7 \pm 2.7 \times 10^{-7} \text{ GeV}^{-3}$	MINOS time of flight	[15]*
"	$< 5 \times 10^{-9} \text{ GeV}^{-3}$	17 GeV, Čerenkov emission	[15]*
"	$< 6 \times 10^{-11} \text{ GeV}^{-3}$	43 GeV, Čerenkov emission	[15]*
$ \overset{\circ}{a}^{(7)} $	$< 3.7 \times 10^{-10} \text{ GeV}^{-3}$	Fermilab time of flight	[15]*
"	$< 3.2 \times 10^{-10} \text{ GeV}^{-3}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 5.0 \times 10^{-4} \text{ GeV}^{-3}$	SN1987A time of flight	[15]*
"	$< 7.1 \times 10^{-8} \text{ GeV}^{-3}$	SN1987A dispersion	[15]*
"	$< 7.3 \times 10^{-30} \text{ GeV}^{-3}$	IceCube meson threshold	[15]*
"	$< 2 \times 10^{-26} \text{ GeV}^{-3}$	100 TeV, Čerenkov emission	[15]*
$ \overset{\circ}{a}_{e\mu}^{(7)} $	$< 1.4 \times 10^{-13} \text{ GeV}^{-3}$	LSND	[15]*
"	$< 2.5 \times 10^{-18} \text{ GeV}^{-3}$	MiniBooNE	[15]*
"	$< 2.0 \times 10^{-19} \text{ GeV}^{-3}$	MiniBooNE $\bar{\nu}$	[15]*

Table D29. Nonminimal neutrino sector,  $d = 8$ 

Combination	Result	System	Ref.
$\overset{\circ}{c}^{(8)}$	$> -2 \times 10^{-36} \text{ GeV}^{-4}$	Atmospheric threshold	[177]*
"	$> -2 \times 10^{-37} \text{ GeV}^{-4}$	Atmospheric Čerenkov	[177]*
"	$> -5 \times 10^{-43} \text{ GeV}^{-4}$	Astrophysical Čerenkov	[177]*
"	$-56.8 \pm 7.7_{-5.8}^{+8.1} \times 10^{-12} \text{ GeV}^{-4}$	OPERA time of flight	[15]*
"	$-12.6 \pm 7.2 \times 10^{-8} \text{ GeV}^{-4}$	MINOS time of flight	[15]*
"	$> -1.8 \times 10^{-35} \text{ GeV}^{-4}$	IceCube meson threshold	[15]*
"	$> -3 \times 10^{-10} \text{ GeV}^{-4}$	17 GeV, Čerenkov emission	[15]*
"	$> -1 \times 10^{-12} \text{ GeV}^{-4}$	43 GeV, Čerenkov emission	[15]*
"	$> -2 \times 10^{-31} \text{ GeV}^{-4}$	100 TeV, Čerenkov emission	[15]*
$ \overset{\circ}{c}^{(8)} $	$< 9.9 \times 10^{-12} \text{ GeV}^{-4}$	Fermilab time of flight	[15]*
"	$< 4.0 \times 10^{-2} \text{ GeV}^{-4}$	SN1987A time of flight	[15]*
"	$< 2.7 \times 10^{-6} \text{ GeV}^{-4}$	SN1987A dispersion	[15]*
$ \overset{\circ}{c}_{e\mu}^{(8)} $	$< 3.6 \times 10^{-12} \text{ GeV}^{-4}$	LSND	[15]*
"	$< 6.9 \times 10^{-18} \text{ GeV}^{-4}$	MiniBooNE	[15]*
"	$< 3.3 \times 10^{-19} \text{ GeV}^{-4}$	MiniBooNE $\bar{\nu}$	[15]*

Table D30. Nonminimal neutrino sector,  $d \geq 9$ 

Combination	Result	System	Ref.
$\hat{a}^{(9)}$	$27.8 \pm 3.8_{-2.8}^{+4.0} \times 10^{-13} \text{ GeV}^{-5}$	OPERA time of flight	[15]*
"	$3.5 \pm 2.0 \times 10^{-8} \text{ GeV}^{-5}$	MINOS time of flight	[15]*
"	$< 2 \times 10^{-11} \text{ GeV}^{-5}$	17 GeV, Čerenkov emission	[15]*
"	$< 3 \times 10^{-14} \text{ GeV}^{-5}$	43 GeV, Čerenkov emission	[15]*
$ \hat{a}^{(9)} $	$< 2.7 \times 10^{-13} \text{ GeV}^{-5}$	Fermilab time of flight	[15]*
"	$< 2.4 \times 10^{-13} \text{ GeV}^{-5}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
"	$< 3.3 \text{ GeV}^{-5}$	SN1987A time of flight	[15]*
"	$< 1.1 \times 10^{-4} \text{ GeV}^{-5}$	SN1987A dispersion	[15]*
"	$< 4.6 \times 10^{-41} \text{ GeV}^{-5}$	IceCube meson threshold	[15]*
"	$< 2 \times 10^{-36} \text{ GeV}^{-5}$	100 TeV, Čerenkov emission	[15]*
$ \hat{a}_{e\mu}^{(9)} $	$< 8.9 \times 10^{-11} \text{ GeV}^{-5}$	LSND	[15]*
"	$< 1.9 \times 10^{-17} \text{ GeV}^{-5}$	MiniBooNE	[15]*
"	$< 5.6 \times 10^{-19} \text{ GeV}^{-5}$	MiniBooNE $\bar{\nu}$	[15]*
$\hat{c}^{(10)}$	$> -2 \times 10^{-48} \text{ GeV}^{-6}$	Atmospheric threshold	[177]*
"	$> -2 \times 10^{-49} \text{ GeV}^{-6}$	Atmospheric Čerenkov	[177]*
"	$> -5 \times 10^{-55} \text{ GeV}^{-6}$	Astrophysical Čerenkov	[177]*
"	$-14.0 \pm 1.9_{-1.4}^{+2.0} \times 10^{-14} \text{ GeV}^{-6}$	OPERA time of flight	[15]*
"	$-10.0 \pm 5.7 \times 10^{-9} \text{ GeV}^{-6}$	MINOS time of flight	[15]*
"	$> -1.1 \times 10^{-46} \text{ GeV}^{-6}$	IceCube meson threshold	[15]*
"	$> -9 \times 10^{-13} \text{ GeV}^{-6}$	17 GeV, Čerenkov emission	[15]*
"	$> -8 \times 10^{-16} \text{ GeV}^{-6}$	43 GeV, Čerenkov emission	[15]*
"	$> -2 \times 10^{-41} \text{ GeV}^{-6}$	100 TeV, Čerenkov emission	[15]*
$ \hat{c}^{(10)} $	$< 7.8 \times 10^{-15} \text{ GeV}^{-6}$	Fermilab time of flight	[15]*
"	$< 2.9 \times 10^2 \text{ GeV}^{-6}$	SN1987A time of flight	[15]*
"	$< 4.5 \times 10^{-3} \text{ GeV}^{-6}$	SN1987A dispersion	[15]*
$ \hat{c}_{e\mu}^{(10)} $	$< 2.2 \times 10^{-9} \text{ GeV}^{-6}$	LSND	[15]*
"	$< 5.4 \times 10^{-17} \text{ GeV}^{-6}$	MiniBooNE	[15]*
"	$< 9.3 \times 10^{-19} \text{ GeV}^{-6}$	MiniBooNE $\bar{\nu}$	[15]*
$\sum_{dj} (d-3)(17 \text{ GeV})^{d-4} \times_0 \mathcal{N}_{j0}(86.7^\circ, 52.4^\circ, 47.5^\circ) [(a_{\text{of}}^{(d)})_{j0} - (c_{\text{of}}^{(d)})_{j0}]$	$2.37 \pm 0.32_{-0.24}^{+0.34} \times 10^{-5}$	OPERA time of flight	[15]*
$\sum_{dj} (d-3)(3 \text{ GeV})^{d-4} \times_0 \mathcal{N}_{j0}(86.7^\circ, 203.9^\circ, 42.2^\circ) [(a_{\text{of}}^{(d)})_{j0} - (c_{\text{of}}^{(d)})_{j0}]$	$5.1 \pm 2.9 \times 10^{-5}$	MINOS time of flight	[15]*
$ \sum_{dj} (d-3)(30 \text{ GeV})^{d-4} \times_0 \mathcal{N}_{j0}(90^\circ, 140^\circ, 48.2^\circ) [\pm (a_{\text{of}}^{(d)})_{j0} - (c_{\text{of}}^{(d)})_{j0}] $	$< 4 \times 10^{-5}$	Fermilab time of flight	[15]*
$ \sum_{dj} (d-3)(30 \text{ GeV})^{d-4} \times_0 \mathcal{N}_{j0}(90^\circ, 140^\circ, 48.2^\circ) (a_{\text{of}}^{(d)})_{j0} $	$< 3.5 \times 10^{-5}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
$ \sum_{djm} (d-3)(10 \text{ MeV})^{d-4} \times Y_{jm}(20.7^\circ, 185.6^\circ) [(a_{\text{of}}^{(d)})_{jm} + (c_{\text{of}}^{(d)})_{jm}] $	$< 2 \times 10^{-9}$	SN1987A time of flight	[15]*
$\sum_d (d-3)(17 \text{ GeV})^{d-4} (\hat{a}^{(d)} - \hat{c}^{(d)})$	$2.37 \pm 0.32_{-0.24}^{+0.34} \times 10^{-5}$	OPERA time of flight	[15]*
$\sum_d (d-3)(3 \text{ GeV})^{d-4} (\hat{a}^{(d)} - \hat{c}^{(d)})$	$5.1 \pm 2.9 \times 10^{-5}$	MINOS time of flight	[15]*
$ \sum_d (d-3)(30 \text{ GeV})^{d-4} (\pm \hat{a}^{(d)} - \hat{c}^{(d)}) $	$< 4 \times 10^{-5}$	Fermilab time of flight	[15]*
$ \sum_d (d-3)(30 \text{ GeV})^{d-4} \hat{a}^{(d)} $	$< 3.5 \times 10^{-5}$	Fermilab $\nu\bar{\nu}$ comparison	[15]*
$ \sum_d (d-3)(10 \text{ MeV})^{d-4} (\hat{a}^{(d)} + \hat{c}^{(d)}) $	$< 2 \times 10^{-9}$	SN1987A time of flight	[15]*

Table D31. Quark sector,  $d = 3$ 

Combination	Result	System	Ref.
$\Delta a_0^K$	$(-6.0 \pm 7.7_{stat} \pm 3.1_{syst}) \times 10^{-18}$ GeV	$K$ oscillations	[187]
$\Delta a_X^K$	$(0.9 \pm 1.5_{stat} \pm 0.6_{syst}) \times 10^{-18}$ GeV	"	[187]
$\Delta a_Y^K$	$(-2.0 \pm 1.5_{stat} \pm 0.5_{syst}) \times 10^{-18}$ GeV	"	[187]
$\Delta a_Z^K$	$(3.1 \pm 1.7_{stat} \pm 0.5_{syst}) \times 10^{-18}$ GeV	"	[187]
$\Delta a_0^K$	$(0.4 \pm 1.8) \times 10^{-17}$ GeV	"	[188], [189]
$\Delta a_X^K$	$(-6.3 \pm 6.0) \times 10^{-18}$ GeV	"	[188]
$\Delta a_Y^K$	$(2.8 \pm 5.9) \times 10^{-18}$ GeV	"	[188]
$\Delta a_Z^K$	$(2.4 \pm 9.7) \times 10^{-18}$ GeV	"	[188]
$\Delta a_Z^K$	$(-1 \pm 4) \times 10^{-17}$ GeV	"	[189]
$ \Delta a_1^K $	$< 9.2 \times 10^{-22}$ GeV	"	[190]
$ \Delta a_2^K $	$< 9.2 \times 10^{-22}$ GeV	"	[190]
$ (\Delta a^K)_T - 0.60(\Delta a^K)_Z $	$< 5 \times 10^{-21}$ GeV	"	[191]*, [192]*
$N^D(\Delta a_0^D - 0.6\Delta a_Z^D)$	$(-2.8 \text{ to } 4.8) \times 10^{-16}$ GeV	$D$ oscillations	[193]
$N^D \Delta a_X^D$	$(-7 \text{ to } 3.8) \times 10^{-16}$ GeV	"	[193]
$N^D \Delta a_Y^D$	$(-7 \text{ to } 3.8) \times 10^{-16}$ GeV	"	[193]
$\Delta a_0^B - 0.38\Delta a_Z^B$	$(0.9 \pm 2.8) \times 10^{-15}$ GeV	$B_d$ oscillations	[194]*
$N^B(\Delta a_0^B - 0.30\Delta a_Z^B)$	$(-3.0 \pm 2.4) \times 10^{-15}$ GeV	"	[195]
$N^B \Delta a_X$	$(-22 \pm 7) \times 10^{-15}$ GeV	"	[195]
$N^B \Delta a_Y$	$(-27 \text{ to } -4) \times 10^{-15}$ GeV	"	[195]
$N^B(\Delta a_0^B - 0.3\Delta a_Z^B)$	$(-5.2 \pm 4.0) \times 10^{-15}$ GeV	"	[196]
$N^B \sqrt{(\Delta a_X^B)^2 + (\Delta a_Y^B)^2}$	$(37 \pm 16) \times 10^{-15}$ GeV	"	[196]
$\Delta a_{\perp}^{B_s}$	$< 1.2 \times 10^{-12}$ GeV	$B_s$ oscillations	[197]
$\Delta a_T^{B_s} - 0.396\Delta a_Z^{B_s}$	$(-0.8 \text{ to } 3.9) \times 10^{-13}$ GeV	"	[197]
$\Delta a_0^{B_s} - 0.38\Delta a_Z^{B_s}$	$(5 \pm 3) \times 10^{-14}$ GeV	"	[194]*
$(\Delta a^{B_s})_T$	$(3.7 \pm 3.8) \times 10^{-12}$ GeV	"	[192]*

Table D32. Quark sector,  $d \geq 4$ 

Combination	Result	System	Ref.
$\hat{c}_q^{\text{UR}(4)}$	$(-0.01 \text{ to } 1.8) \times 10^{-21}$	Astrophysics	[65]*, [18]*
$\hat{c}_q^{\text{UR}(4)} - 2\hat{c}_e^{\text{UR}(4)}$	$(-0.001 \text{ to } 2) \times 10^{-20}$	"	[65]*, [18]*
$\delta\pi$	$(-1.5 \text{ to } 200) \times 10^{-11}$	"	[198]*
$ c^\pi $	$< 10^{-10}$	"	[60]*
$ c^K $	$< 10^{-9}$	"	[60]*
$ c^D $	$< 10^{-8}$	"	[60]*
$ c^{B_d} ,  c^{B_s} $	$< 10^{-7}$	"	[60]*
$(c_Q)_{XX33}$	$-0.12 \pm 0.11 \pm 0.02$	$t\bar{t}$ production	[199]
$(c_Q)_{YY33}$	$0.12 \pm 0.11 \pm 0.02$	"	[199]
$(c_Q)_{XY33}$	$-0.04 \pm 0.11 \pm 0.01$	"	[199]
$(c_Q)_{XZ33}$	$0.15 \pm 0.08 \pm 0.02$	"	[199]
$(c_Q)_{YZ33}$	$-0.03 \pm 0.08 \pm 0.01$	"	[199]
$(c_U)_{XX33}$	$0.1 \pm 0.09 \pm 0.02$	"	[199]
$(c_U)_{YY33}$	$-0.1 \pm 0.09 \pm 0.02$	"	[199]
$(c_U)_{XY33}$	$0.04 \pm 0.09 \pm 0.01$	"	[199]
$(c_U)_{XZ33}$	$-0.14 \pm 0.07 \pm 0.02$	"	[199]
$(c_U)_{YZ33}$	$0.01 \pm 0.07 \pm < 0.01$	"	[199]
$d_{XX}$	$-0.11 \pm 0.1 \pm 0.02$	"	[199]
$d_{YY}$	$0.11 \pm 0.1 \pm 0.02$	"	[199]
$d_{XY}$	$-0.04 \pm 0.1 \pm 0.01$	"	[199]
$d_{XZ}$	$0.14 \pm 0.07 \pm 0.02$	"	[199]
$d_{YZ}$	$-0.02 \pm 0.07 \pm < 0.01$	"	[199]
$\hat{c}_q^{\text{UR}(6)}$	$(-0.63 \text{ to } 1.7) \times 10^{-22} \text{ GeV}^{-2}$	Astrophysics	[65]*, [18]*



Table D33. Electroweak sector,  $d = 3, 4$ 

Combination	Result	System	Ref.
$ (k_\phi)_X ,  (k_\phi)_Y $	$< 10^{-31}$ GeV	Xe-He maser	[200]*
$ (k_\phi)_Z ,  (k_\phi)_T $	$< 2.8 \times 10^{-27}$ GeV	"	[200]*
$ (k_{\phi\phi})_S^{XT} $	$< 3.3 \times 10^{-3}$	Kaon decay	[201]*
$ (k_{\phi\phi})_S^{YT} $	$< 6.3 \times 10^{-3}$	"	[201]*
$ (k_{\phi\phi})_S^{ZT} $	$< 6.0 \times 10^{-3}$	"	[201]*
$2(k_{\phi\phi}^A)^{32} + \frac{1}{g}(k_{\phi W})^{32}$	$(-9 \text{ to } 2) \times 10^{-3}$	Nuclear beta decay	[202]
$2(k_{\phi\phi}^A)^{13} + \frac{1}{g}(k_{\phi W})^{13}$	$(-6 \text{ to } 4) \times 10^{-3}$	"	[202]
$(k_{\phi\phi}^S)^{ZT}, (k_{\phi\phi}^A)^{YX}, (k_{\phi W})^{YX}$	$(-0.5 \text{ to } 1) \times 10^{-8}$	"	[203]*
$(k_{\phi\phi}^S)^{ZZ}$	$(-1 \text{ to } 0.4) \times 10^{-6}$	"	[203]*
$(k_{\phi\phi}^S)^{TT}$	$(-1 \text{ to } 3) \times 10^{-6}$	"	[203]*
$ (k_{\phi\phi}^S)^{XX} ,  (k_{\phi\phi}^S)^{YY} $	$< 1 \times 10^{-6}$	"	[203]*
$ (k_{\phi\phi}^S)^{XT} ,  (k_{\phi\phi}^S)^{YT} ,  (k_{\phi\phi}^A)^{XZ} ,  (k_{\phi\phi}^A)^{YZ} ,  (k_{\phi W})^{XZ} ,  (k_{\phi W})^{YZ} $	$< 2 \times 10^{-8}$	"	[203]*
$ (k_{\phi\phi}^S)^{XY} ,  (k_{\phi\phi}^S)^{XZ} ,  (k_{\phi\phi}^S)^{YZ} $	$< 5 \times 10^{-7}$	"	[203]*
$ (k_{\phi\phi}^A)_{\mu\nu} $	$< 3 \times 10^{-16}$	Cosmological birefringence	[200]*
$ (k_{\phi B})_{\mu\nu} $	$< 0.9 \times 10^{-16}$	"	[200]*
$ (k_{\phi W})_{\mu\nu} $	$< 1.7 \times 10^{-16}$	"	[200]*
$ (k_{\phi\phi}^S)_{XX} ,  (k_{\phi\phi}^S)_{YY} ,  (k_{\phi\phi}^S)_{ZZ} $	$< 10^{-27}$	Clock comparisons	[200]*
$ (k_{\phi\phi}^S)_{XY} $	$< 10^{-27}$	"	[200]*
$ (k_{\phi\phi}^S)_{XZ} ,  (k_{\phi\phi}^S)_{YZ} $	$< 10^{-25}$	"	[200]*
$ (k_{\phi\phi}^S)_{TT} $	$< 4 \times 10^{-13}$	$H^-$ ion, $\bar{p}$ comparison	[200]*
$ (k_W)^\alpha_{\mu\alpha\nu} $	$< 10^{-5}$	Astrophysics	[60]*

Table D34. Gluon sector,  $d = 4$ 

Combination	Result	System	Ref.
$ \tilde{\kappa}_{\text{tr}}^{\text{QCD}} $	$< 2 \times 10^{-13}$	Astrophysics [134]*	

Table D35. Gravity sector,  $d = 3$ 

Combination	Result	System	Ref.
$\alpha(\bar{a}^{e+p})_T$	$(0.22 \pm 0.28) \times 10^{-6}$ GeV	Nuclear binding energy	[51]
$\alpha(\bar{a}^n)_T$	$(0.51 \pm 0.64) \times 10^{-6}$ GeV	"	[51]
$\alpha(\bar{a}_{\text{eff}}^{e+p})_T$	$(0.8 \pm 1.0) \times 10^{-6}$ GeV	Cs interferometer	[52]
$\alpha(\bar{a}_{\text{eff}}^n)_T$	$(4.3 \pm 3.7) \times 10^{-6}$ GeV	"	[52]
$ \alpha(\bar{a}_{\text{eff}}^{e+p})_T $	$< 10^{-7}$ GeV	Free-fall WEP	[26]*
"	$< 10^{-10}$ GeV	Force-comparison WEP	[26]*
$ \alpha(\bar{a}_{\text{eff}}^n)_T $	$< 10^{-7}$ GeV	Free-fall WEP	[26]*
"	$< 10^{-10}$ GeV	Force-comparison WEP	[26]*
$ \alpha(\bar{a}_{\text{eff}}^{e+p-n})_T - \frac{1}{3}m^p(\bar{c}^{e+p-n})_{TT} $	$< 10^{-8}$ GeV	Combined WEP	[26]*
$ \alpha(\bar{a}_{\text{eff}}^{e+p-n})_T - \frac{1}{3}m^p(\bar{c}^{e+p-n})_{TT} + (\frac{1}{2}\cos^2\chi - \frac{1}{6})m^n(\bar{c}^n)_Q $	$< 10^{-8}$ GeV	Free-fall WEP	[26]*
$ \alpha(\bar{a}_{\text{eff}}^{e+p-n})_T - \frac{1}{3}m^p(\bar{c}^{e+p-n})_{TT} - \frac{1}{6}m^n(\bar{c}^n)_Q $	$< 10^{-11}$ GeV	Force-comparison WEP	[26]*
$ \alpha\bar{a}_T^e + \alpha\bar{a}_T^p - 0.8\alpha\bar{a}_T^n $	$< 1 \times 10^{-11}$ GeV	Torsion pendulum	[16]*
$\alpha(\bar{a}_{\text{eff}}^e)^X + \alpha(\bar{a}_{\text{eff}}^p)^X$	$(-7.6 \pm 9.0) \times 10^{-6}$ GeV	Combined	[204]*
$\alpha(\bar{a}_{\text{eff}}^n)^X$	$(-5.4 \pm 6.3) \times 10^{-6}$ GeV	"	[204]*
$\bar{s}^{TX} - \frac{\alpha}{m_{\oplus}}(\bar{a}_{\text{eff}}^{\oplus})^X$	$0.24 \pm 0.15$	K/He magnetometer	[205]*
$ \alpha[\bar{a}_X^n + 0.83(\bar{a}_X^p + \bar{a}_X^e)] $	$\leq 2 \times 10^{-1}$ GeV	Torsion-strip balance	[206]
$\alpha(\bar{a}_{\text{eff}}^e)^Y + \alpha(\bar{a}_{\text{eff}}^p)^Y$	$(-6.2 \pm 9.5) \times 10^{-5}$ GeV	Combined	[204]*
$\alpha(\bar{a}_{\text{eff}}^n)^Y$	$(4.8 \pm 8.2) \times 10^{-4}$ GeV	"	[204]*
$\bar{s}^{TY} - \frac{\alpha}{m_{\oplus}}(\bar{a}_{\text{eff}}^{\oplus})^Y$	$0.02 \pm 0.13$	K/He magnetometer	[205]*
$ \alpha[\bar{a}_Y^n + 0.83(\bar{a}_Y^p + \bar{a}_Y^e)] $	$\leq 4 \times 10^{-1}$ GeV	Torsion-strip balance	[206]
$\alpha(\bar{a}_{\text{eff}}^e)^Z + \alpha(\bar{a}_{\text{eff}}^p)^Z$	$(1.3 \pm 2.2) \times 10^{-4}$ GeV	Combined	[204]*
$\alpha(\bar{a}_{\text{eff}}^n)^Z$	$(-1.1 \pm 1.9) \times 10^{-3}$ GeV	"	[204]*
$ (\bar{a}_{\text{eff}}^e)_{\text{}}^{\text{}} + (\bar{a}_{\text{eff}}^p)_{\text{}}^{\text{}} + 0.1(\bar{a}_{\text{eff}}^n)_{\text{}}^{\text{}} $	$< 10^{-6}$ GeV	Solar system	[26]*
$ (\bar{a}_{\text{eff}}^e)_{\oplus} + (\bar{a}_{\text{eff}}^p)_{\oplus} + 0.1(\bar{a}_{\text{eff}}^n)_{\oplus} $	$< 10^{-6}$ GeV	"	[26]*

Table D36. Gravity sector,  $d = 4$  (part 1 of 2)

Combination	Result	System	Ref.
$\bar{s}_{00}^{(4)}$	$> -3 \times 10^{-14}$	Cosmic rays	[207]*
$\bar{s}_{10}^{(4)}$	$(-10 \text{ to } 7) \times 10^{-14}$	"	[207]*
$\text{Re } \bar{s}_{11}^{(4)}$	$(-8 \text{ to } 8) \times 10^{-14}$	"	[207]*
$\text{Im } \bar{s}_{11}^{(4)}$	$(-7 \text{ to } 9) \times 10^{-14}$	"	[207]*
$\bar{s}_{20}^{(4)}$	$(-7 \text{ to } 10) \times 10^{-14}$	"	[207]*
$\text{Re } \bar{s}_{21}^{(4)}$	$(-7 \text{ to } 7) \times 10^{-14}$	"	[207]*
$\text{Im } \bar{s}_{21}^{(4)}$	$(-5 \text{ to } 8) \times 10^{-14}$	"	[207]*
$\text{Re } \bar{s}_{22}^{(4)}$	$(-6 \text{ to } 8) \times 10^{-14}$	"	[207]*
$\text{Im } \bar{s}_{22}^{(4)}$	$(-7 \text{ to } 7) \times 10^{-14}$	"	[207]*
$\bar{s}^{XY}$	$(6.5 \pm 3.2) \times 10^{-11}$	Combined	[204]*
$\bar{s}^{XZ}$	$(2.0 \pm 1.0) \times 10^{-11}$	"	[204]*
$\bar{s}^{YZ}$	$(4.1 \pm 5.0) \times 10^{-12}$	"	[204]*
$\bar{s}^{XX} - \bar{s}^{YY}$	$(9.6 \pm 5.6) \times 10^{-11}$	"	[204]*
$\bar{s}^{XX} + \bar{s}^{YY} - 2\bar{s}^{ZZ}$	$(1.6 \pm 0.78) \times 10^{-10}$	"	[204]*
$\bar{s}^{TX}$	$(-7.4 \pm 8.7) \times 10^{-6}$	"	[204]*
$\bar{s}^{TY}$	$(-0.8 \pm 2.5) \times 10^{-5}$	"	[204]*
$\bar{s}^{TZ}$	$(0.8 \pm 5.8) \times 10^{-5}$	"	[204]*
$ \bar{s}^{TT} $	$< 1.6 \times 10^{-5}$	Binary pulsars	[208]*
"	$< 2.8 \times 10^{-4}$	"	[208]*
$\bar{s}^{TX}$	$(-5.2 \text{ to } 5.3) \times 10^{-9}$	"	[209]*
$\bar{s}^{TY}$	$(-7.5 \text{ to } 8.5) \times 10^{-9}$	"	[209]*
$\bar{s}^{TZ}$	$(-5.9 \text{ to } 5.8) \times 10^{-9}$	"	[209]*
$\bar{s}^{XY}$	$(-3.5 \text{ to } 3.6) \times 10^{-11}$	"	[209]*
$\bar{s}^{XZ}$	$(-2.0 \text{ to } 2.0) \times 10^{-11}$	"	[209]*
$\bar{s}^{YZ}$	$(-3.3 \text{ to } 3.3) \times 10^{-11}$	"	[209]*
$\bar{s}^{XX} - \bar{s}^{YY}$	$(-9.7 \text{ to } 10.1) \times 10^{-11}$	"	[209]*
$\bar{s}^{XX} + \bar{s}^{YY} - 2\bar{s}^{ZZ}$	$(-12.3 \text{ to } 12.2) \times 10^{-11}$	"	[209]*
$ \bar{s}^{TT} $	$< 3.8 \times 10^{-3}$	Combined	[210]*
$ \bar{s}^{XX} $	$< 1.3 \times 10^{-3}$	"	[210]*
$ \bar{s}^{YY} $	$< 1.3 \times 10^{-3}$	"	[210]*
$ \bar{s}^{ZZ} $	$< 1.3 \times 10^{-3}$	"	[210]*
$ \bar{s}^{TT} + 0.14(\bar{s}^{XX} - \bar{s}^{YY}) - 0.054(\bar{s}^{XX} + \bar{s}^{YY} - 2\bar{s}^{ZZ}) - 0.18\bar{s}^{XY} $	$< 3.8 \times 10^{-3}$	Geodetic effect	[210]*
$ (\bar{s}^{XX} - \bar{s}^{YY}) + 3.0\bar{s}^{XY} + 0.32\bar{s}^{XZ} + 1.0\bar{s}^{YZ} $	$< 4.9 \times 10^{-7}$	Frame dragging	[210]*
$\bar{s}^{TT}$	$< 4.4 \times 10^{-3}$	Gravity Probe B	[211]
$\bar{s}^{XX}, \bar{s}^{YY}, \bar{s}^{ZZ}$	$< 1.5 \times 10^{-3}$	"	[211]

Table D36. Gravity sector,  $d = 4$  (part 2 of 2)

Combination	Result	System	Ref.
$\sigma^{XX} - \sigma^{YY}$	$(4.4 \pm 11) \times 10^{-9}$	Atom interferometry	[212]
$\sigma^{XY}$	$(0.2 \pm 3.9) \times 10^{-9}$	"	[212]
$\sigma^{XZ}$	$(-2.6 \pm 4.4) \times 10^{-9}$	"	[212]
$\sigma^{YZ}$	$(-0.3 \pm 4.5) \times 10^{-9}$	"	[212]
$\sigma^{TX}$	$(-3.1 \pm 5.1) \times 10^{-5}$	"	[212]
$\sigma^{TY}$	$(0.1 \pm 5.4) \times 10^{-5}$	"	[212]
$\sigma^{TZ}$	$(1.4 \pm 6.6) \times 10^{-5}$	"	[212]
$\sigma^{XX} - \sigma^{YY}$	$(-5.6 \pm 2.1) \times 10^{-9}$	"	[213]
$\sigma^{XY}$	$(-0.09 \pm 79) \times 10^{-9}$	"	[213]
$\sigma^{XZ}$	$(-13 \pm 37) \times 10^{-9}$	"	[213]
$\sigma^{YZ}$	$(-61 \pm 38) \times 10^{-9}$	"	[213]
$\sigma^{TX}$	$(5.4 \pm 4.5) \times 10^{-5}$	"	[213]
$\sigma^{TY}$	$(-2.0 \pm 4.4) \times 10^{-5}$	"	[213]
$\sigma^{TZ}$	$(1.1 \pm 26) \times 10^{-5}$	"	[213]
$\bar{s}^{XX} - \bar{s}^{YY}$	$(-1.2 \pm 1.6) \times 10^{-9}$	LLR & Atom interferometry	[214]*, [212]*
$\bar{s}^{XX} + \bar{s}^{YY} - 2\bar{s}^{ZZ}$	$(1.8 \pm 38) \times 10^{-9}$	"	[214]*, [212]*
$\bar{s}^{XY}$	$(-0.6 \pm 1.5) \times 10^{-9}$	"	[214]*, [212]*
$\bar{s}^{XZ}$	$(-2.7 \pm 1.4) \times 10^{-9}$	"	[214]*, [212]*
$\bar{s}^{YZ}$	$(0.6 \pm 1.4) \times 10^{-9}$	"	[214]*, [212]*
$\bar{s}^{TX}$	$(0.5 \pm 6.2) \times 10^{-7}$	"	[214]*, [212]*
$\bar{s}^{TY}$	$(0.1 \pm 1.3) \times 10^{-6}$	"	[214]*, [212]*
$\bar{s}^{TZ}$	$(-0.4 \pm 3.8) \times 10^{-6}$	"	[214]*, [212]*
$\bar{s}^{11} - \bar{s}^{22}$	$(1.3 \pm 0.9) \times 10^{-10}$	Lunar laser ranging (LLR)	[214]*
$\bar{s}^{12}$	$(6.9 \pm 4.5) \times 10^{-11}$	"	[214]*
$\bar{s}^{01}$	$(-0.8 \pm 1.1) \times 10^{-6}$	"	[214]*
$\bar{s}^{02}$	$(-5.2 \pm 4.8) \times 10^{-7}$	"	[214]*
$\bar{s}_{\oplus c}$	$(0.2 \pm 3.9) \times 10^{-7}$	"	[214]*
$\bar{s}_{\oplus s}$	$(-1.3 \pm 4.1) \times 10^{-7}$	"	[214]*
$\bar{s}^{XX}$	$(-0.04 \pm 4.90) \times 10^4$	Short-range gravity	[215]
$\bar{s}^{XY}$	$(-0.07 \pm 6.12) \times 10^4$	"	[215]
$\bar{s}^{XZ}$	$(-0.01 \pm 2.56) \times 10^3$	"	[215]
$\bar{s}^{YZ}$	$(-0.06 \pm 5.83) \times 10^4$	"	[215]
$\bar{s}^{ZZ}$	$(-0.08 \pm 6.68) \times 10^4$	"	[215]
$\bar{s}^{01}$	$(-0.9 \pm 1.5) \times 10^{-8}$	Perihelion precession	[216]*
$\bar{s}^{02}$	$(4 \pm 6) \times 10^{-9}$	"	[216]*
$\bar{s}^{03}$	$(-0.3 \pm 1) \times 10^{-9}$	"	[216]*
$ \bar{s}^{01} $	$< 1.4 \times 10^{-3}$	LAGEOS satellite	[216]*
$ \bar{s}^{02} $	$< 2 \times 10^{-4}$	"	[216]*
$ \bar{s}^{03} $	$< 1.3 \times 10^{-3}$	"	[216]*
$ \bar{s}_{\ddagger} $	$\leq 10^{-9}$	Perihelion precession	[217]*
$ \bar{s}_{\oplus} $	$\leq 10^{-8}$	"	[217]*
$ \bar{s}_{\text{SSP}} $	$\leq 10^{-13}$	Solar-spin precession	[217]*

Table D37. Nonminimal gravity sector,  $d = 5, 6$  (part 1 of 2)

Combination	Result	System	Ref.
$\bar{s}_{00}^{(6)}$	$< 2 \times 10^{-31} \text{ GeV}^{-2}$	Cosmic rays	[207]*
$\bar{s}_{10}^{(6)}$	$(-6 \text{ to } 10) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{11}^{(6)}$	$(-6 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{11}^{(6)}$	$(-8 \text{ to } 5) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\bar{s}_{20}^{(6)}$	$(-1 \text{ to } 1) \times 10^{-29} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{21}^{(6)}$	$(-7 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{21}^{(6)}$	$(-9 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{22}^{(6)}$	$(-9 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{22}^{(6)}$	$(-8 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\bar{s}_{30}^{(6)}$	$(-10 \text{ to } 8) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{31}^{(6)}$	$(-8 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{31}^{(6)}$	$(-6 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{32}^{(6)}$	$(-6 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{32}^{(6)}$	$(-7 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{33}^{(6)}$	$(-7 \text{ to } 5) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{33}^{(6)}$	$(-7 \text{ to } 8) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\bar{s}_{40}^{(6)}$	$(-10 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{41}^{(6)}$	$(-8 \text{ to } 5) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{41}^{(6)}$	$(-8 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{42}^{(6)}$	$(-7 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{42}^{(6)}$	$(-8 \text{ to } 6) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{43}^{(6)}$	$(-9 \text{ to } 4) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{43}^{(6)}$	$(-6 \text{ to } 7) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Re } \bar{s}_{44}^{(6)}$	$(-8 \text{ to } 9) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$\text{Im } \bar{s}_{44}^{(6)}$	$(-4 \text{ to } 8) \times 10^{-30} \text{ GeV}^{-2}$	"	[207]*
$(\bar{k}_{\text{eff}})_{XXXX}$	$(-0.2 \pm 2.8) \times 10^{-8} \text{ m}^2$	Tungsten torsion pendulum	[218]
$(\bar{k}_{\text{eff}})_{YYYY}$	$(0.4 \pm 2.8) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{ZZZZ}$	$(-0.9 \pm 7.7) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{XXXY}$	$(0.4 \pm 1.3) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{XXZX}$	$(-0.1 \pm 0.5) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{YY YX}$	$(0.6 \pm 1.3) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{YY YZ}$	$(0.4 \pm 0.5) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{ZZZX}$	$(-1.3 \pm 1.4) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{ZZZY}$	$(-0.2 \pm 1.3) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{XXYY}$	$(-0.1 \pm 1.7) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{XXZZ}$	$(-0.2 \pm 1.0) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{YYZZ}$	$(0.2 \pm 1.0) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{XXYZ}$	$(0.5 \pm 0.5) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{YYXZ}$	$(-0.2 \pm 0.5) \times 10^{-8} \text{ m}^2$	"	[218]
$(\bar{k}_{\text{eff}})_{ZZXY}$	$(-0.2 \pm 0.5) \times 10^{-8} \text{ m}^2$	"	[218]

Table D37. Nonminimal gravity sector,  $d = 5, 6$  (part 2 of 2)

Combination	Result	System	Ref.
$(\bar{k}_{\text{eff}})_{XXXX}$	$(1.1 \pm 3.1) \times 10^{-7} \text{ m}^2$	Tungsten oscillators	[219]
$(\bar{k}_{\text{eff}})_{YYYY}$	$(0.4 \pm 3.1) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{ZZZZ}$	$(0.6 \pm 2.5) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{XXXY}$	$(-3.4 \pm 15.8) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{XXZX}$	$(-8.1 \pm 10.7) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{YYYX}$	$(-4.4 \pm 15.8) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{YYYZ}$	$(4.6 \pm 9.6) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{ZZZX}$	$(-5.4 \pm 16.6) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{ZZZY}$	$(3.4 \pm 17.8) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{XXYY}$	$(0.4 \pm 1.5) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{XXZZ}$	$(0.2 \pm 1.6) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{YYZZ}$	$(0.5 \pm 1.6) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{XXYZ}$	$(-2.7 \pm 5.5) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{YYXZ}$	$(5.0 \pm 6.6) \times 10^{-7} \text{ m}^2$	"	[219]
$(\bar{k}_{\text{eff}})_{ZZXY}$	$(-0.7 \pm 1.9) \times 10^{-7} \text{ m}^2$	"	[219]

Table D38. Nonminimal gravity sector,  $d \geq 7$  (part 1 of 2)

Combination	Result	System	Ref.
$\bar{s}_{00}^{(8)}$	$> -7 \times 10^{-49} \text{ GeV}^{-4}$	Cosmic rays	[207]*
$\bar{s}_{10}^{(8)}$	$(-1 \text{ to } 1) \times 10^{-45} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{11}^{(8)}$	$(-9 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{11}^{(8)}$	$(-9 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\bar{s}_{20}^{(8)}$	$(-9 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{21}^{(8)}$	$(-10 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{21}^{(8)}$	$(-8 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{22}^{(8)}$	$(-10 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{22}^{(8)}$	$(-10 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\bar{s}_{30}^{(8)}$	$(-1 \text{ to } 1) \times 10^{-45} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{31}^{(8)}$	$(-10 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{31}^{(8)}$	$(-9 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{32}^{(8)}$	$(-8 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{32}^{(8)}$	$(-9 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{33}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{33}^{(8)}$	$(-1 \text{ to } 1) \times 10^{-45} \text{ GeV}^{-4}$	"	[207]*
$\bar{s}_{40}^{(8)}$	$(-1 \text{ to } 1) \times 10^{-45} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{41}^{(8)}$	$(-6 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{41}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{42}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{42}^{(8)}$	$(-6 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{43}^{(8)}$	$(-7 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{43}^{(8)}$	$(-8 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{44}^{(8)}$	$(-10 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{44}^{(8)}$	$(-9 \text{ to } 6) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\bar{s}_{50}^{(8)}$	$(-1 \text{ to } 1) \times 10^{-45} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{51}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{51}^{(8)}$	$(-8 \text{ to } 7) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{52}^{(8)}$	$(-9 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{52}^{(8)}$	$(-8 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{53}^{(8)}$	$(-10 \text{ to } 7) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{53}^{(8)}$	$(-6 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{54}^{(8)}$	$(-9 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{54}^{(8)}$	$(-8 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{55}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{55}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*

Table D38. Nonminimal gravity sector,  $d \geq 7$  (part 2 of 2)

Combination	Result	System	Ref.
$\bar{s}_{60}^{(8)}$	$(-1 \text{ to } 2) \times 10^{-45} \text{ GeV}^{-4}$	Cosmic rays	[207]*
$\text{Re } \bar{s}_{61}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{61}^{(8)}$	$(-7 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{62}^{(8)}$	$(-10 \text{ to } 6) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{62}^{(8)}$	$(-6 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{63}^{(8)}$	$(-7 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{63}^{(8)}$	$(-7 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{64}^{(8)}$	$(-8 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{64}^{(8)}$	$(-9 \text{ to } 8) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{65}^{(8)}$	$(-8 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{65}^{(8)}$	$(-8 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Re } \bar{s}_{66}^{(8)}$	$(-10 \text{ to } 9) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*
$\text{Im } \bar{s}_{66}^{(8)}$	$(-7 \text{ to } 10) \times 10^{-46} \text{ GeV}^{-4}$	"	[207]*



## **PROPERTIES TABLES P39–P52**

Table P39. Lagrange density for the minimal QED extension in Riemann spacetime

Sector	Coeff.	#	Operator	Dim.	Factor	CPT	L.V.
Fermion	$m$		$\bar{\psi}\psi$	3	$-e$	+	
	$m_5$		$\bar{\psi}\gamma_5\psi$	3	$-ie$	+	
			$\bar{\psi}\gamma^a\overleftrightarrow{D}_\mu\psi$	4	$\frac{1}{2}iee^\mu_a$	+	
	$a_\mu$	4	$\bar{\psi}\gamma^a\psi$	3	$-ee^\mu_a$	-	✓
	$b_\mu$	4	$\bar{\psi}\gamma_5\gamma^a\psi$	3	$-ee^\mu_a$	-	✓
	$H_{\mu\nu}$	6	$\bar{\psi}\sigma^{ab}\psi$	3	$-ee^\mu_ae^\nu_b$	+	✓
	$c_{\lambda\nu}$	16	$\bar{\psi}\gamma^b\overleftrightarrow{D}_\mu\psi$	4	$-\frac{1}{2}iee^\mu_ae^{\nu a}e^\lambda_b$	+	✓
	$d_{\lambda\nu}$	16	$\bar{\psi}\gamma_5\gamma^b\overleftrightarrow{D}_\mu\psi$	4	$-\frac{1}{2}iee^\mu_ae^{\nu a}e^\lambda_b$	+	✓
	$e_\lambda$	4	$\bar{\psi}\overleftrightarrow{D}_\mu\psi$	4	$-\frac{1}{2}iee^\mu_ae^{\lambda a}$	-	✓
	$f_\lambda$	4	$\bar{\psi}\gamma_5\overleftrightarrow{D}_\mu\psi$	4	$-\frac{1}{2}iee^\mu_ae^{\lambda a}$	-	✓
	$g_{\lambda\kappa\nu}$	24	$\bar{\psi}\sigma^{bc}\overleftrightarrow{D}_\mu\psi$	4	$-\frac{1}{4}iee^\mu_ae^{\nu a}e^\lambda_be^\kappa_c$	-	✓
Photon			$F_{\mu\nu}F^{\mu\nu}$	4	$-\frac{1}{4}e$	+	
	$(k_{AF})^\kappa$	4	$A^\lambda F^{\mu\nu}$	3	$\frac{1}{2}e\epsilon_{\kappa\lambda\mu\nu}$	-	✓
	$(k_F)_{\kappa\lambda\mu\nu}$	19	$F^{\kappa\lambda}F^{\mu\nu}$	4	$-\frac{1}{4}e$	+	✓
Gravity			$R$	2	$e/2\kappa$	+	
	$\Lambda$		1	0	$-e/\kappa$	+	
	$u$	1	$R$	2	$-e/2\kappa$	+	
	$s^{\mu\nu}$	9	$R_{\mu\nu}$	2	$e/2\kappa$	+	✓
	$t^{\kappa\lambda\mu\nu}$	10	$R_{\kappa\lambda\mu\nu}$	2	$e/2\kappa$	+	✓

Table P40. C, P, T properties of operators for Lorentz violation in QED

Coefficient	C	P	T	CP	CT	PT	CPT
$c_{TT}, c_{JK}, (k_F)_{TJTK}, (k_F)_{JKLM}$	+	+	+	+	+	+	+
$b_J, g_{JTL}, g_{JKT}, (k_{AF})_J$	+	+	-	+	-	-	-
$b_T, g_{JTT}, g_{JKL}, (k_{AF})_T$	+	-	+	-	+	-	-
$c_{TJ}, c_{JT}, (k_F)_{TJKL}$	+	-	-	-	-	+	+
$a_T, e_T, f_J$	-	+	+	-	-	+	-
$H_{JK}, d_{TJ}, d_{JT}$	-	+	-	-	+	-	+
$H_{TJ}, d_{TT}, d_{JK}$	-	-	+	+	-	-	+
$a_J, e_J, f_T$	-	-	-	+	+	+	-

Table P41. Definitions for the fermion sector of the minimal QED extension

Symbol	Combination	Components
$\tilde{b}_J$	$b_J - \frac{1}{2}\varepsilon_{JKL}H_{KL} - m(d_{JT} - \frac{1}{2}\varepsilon_{JKL}g_{KLT})$	3
$\tilde{b}_J^*$	$b_J + \frac{1}{2}\varepsilon_{JKL}H_{KL} + m(d_{JT} + \frac{1}{2}\varepsilon_{JKL}g_{KLT}),$	3
$\tilde{b}_T$	$b_T + mg_{XYZ}$	1
$\tilde{g}_T$	$b_T - m(g_{XYZ} - g_{YZX} - g_{ZXY})$	1
$\tilde{H}_{XT}$	$H_{XT} + m(d_{ZY} - g_{XTT} - g_{XY})$	1
$\tilde{H}_{YT}$	$H_{YT} + m(d_{XZ} - g_{YTT} - g_{YZ})$	1
$\tilde{H}_{ZT}$	$H_{ZT} + m(d_{YX} - g_{ZTT} - g_{ZX})$	1
$\tilde{d}_\pm$	$m(d_{XX} \pm d_{YY})$	2
$\tilde{d}_Q$	$m(d_{XX} + d_{YY} - 2d_{ZZ} - g_{YZX} - g_{ZXY} + 2g_{XYZ})$	1
$\tilde{d}_J$	$m(d_{TJ} + \frac{1}{2}d_{JT}) - \frac{1}{4}\varepsilon_{JKL}H_{KL}$	3
$\tilde{d}_{YZ}$	$m(d_{YZ} + d_{ZY} - g_{XY} + g_{XZ})$	1
$\tilde{d}_{ZX}$	$m(d_{ZX} + d_{XZ} - g_{YZ} + g_{YX})$	1
$\tilde{d}_{XY}$	$m(d_{XY} + d_{YX} - g_{ZX} + g_{ZY})$	1
$\tilde{g}_e$	$m(g_{XYZ} - g_{ZXY})$	1
$\tilde{g}_-$	$m(g_{XTX} - g_{YTY})$	1
$\tilde{g}_Q$	$m(g_{XTX} + g_{YTY} - 2g_{ZTZ})$	1
$\tilde{g}_{TJ}$	$m \varepsilon_{JKL} g_{KTL}$	3
$\tilde{g}_{DJ}$	$-b_J + m\varepsilon_{JKL}(g_{KTL} + \frac{1}{2}g_{KLT})$	3
$\tilde{g}_{JK}$	$m(g_{JTT} + g_{JKK}),$ (no $K$ sum, $J \neq K$ )	6
$\tilde{c}_Q$	$m(c_{XX} + c_{YY} - 2c_{ZZ})$	1
$\tilde{c}_-$	$m(c_{XX} - c_{YY})$	1
$\tilde{c}_J$	$m \varepsilon_{JKL} c_{KL}$	3
$\tilde{c}_{TJ}$	$m(c_{TJ} + c_{JT})$	3
$\tilde{c}_{TT}$	$m c_{TT}$	1 <b>total: 44</b>

Table P42. Inverse combinations for the fermion sector of the minimal QED extension

Coefficient	Combination	Components
$b_T - mg_T^{(A)}$	$\frac{1}{3}(2\tilde{b}_T + \tilde{g}_T)$	
$md_{XX}$	$\frac{1}{2}(\tilde{d}_+ + \tilde{d}_-)$	
$md_{YY}$	$\frac{1}{2}(\tilde{d}_+ - \tilde{d}_-)$	
$md_{ZZ}$	$\frac{1}{2}(\tilde{b}_T - \tilde{g}_T + \tilde{d}_+ - \tilde{d}_Q)$	
$mg_{XYZ}^{(M)}$	$\frac{1}{3}(\tilde{b}_T - \tilde{g}_T)$	
$mg_{YZX}^{(M)}$	$-\frac{2}{3}(\tilde{b}_T - \tilde{g}_T) + \tilde{g}_c$	6
$b_J - mg_J^{(A)}$	$\frac{1}{3}(\tilde{b}_J + \tilde{b}_J^* - \tilde{g}_{DJ})$	
$H_{KL} - md_{TJ}^-$	$-\frac{3}{8}(\tilde{b}_J - \tilde{b}_J^*) - \frac{1}{2}\tilde{d}_J$	
$md_{TJ}^+$	$-\frac{1}{8}(\tilde{b}_J - \tilde{b}_J^*) + \frac{1}{2}\tilde{d}_J$	
$mg_{KLT}^{(M)}$	$\frac{1}{6}(\tilde{b}_J + \tilde{b}_J^* + 2\tilde{g}_{DJ})$	
$mg_{TKL}^{(M)}$	$-\frac{1}{12}(\tilde{b}_J + \tilde{b}_J^* + 2\tilde{g}_{DJ} + 6\tilde{g}_{TJ})$	
	$(J, KL) = (X, YZ), (Y, ZX), (Z, XY)$	5+5+5
$H_{TJ} + md_{KL}^-$	$\tilde{H}_{TJ} + \frac{1}{2}(\tilde{d}_{KL} - \tilde{g}_{JK} - \tilde{g}_{JL})$	
$md_{KL}^+$	$\frac{1}{2}(\tilde{d}_{KL} + \tilde{g}_{JK} - \tilde{g}_{JL})$	
$mg_{TJT}^{(M)}$	$-\frac{1}{3}(\tilde{g}_{JK} + \tilde{g}_{JL})$	
$mg_{JKK}^{(M)}$	$\frac{1}{3}(2\tilde{g}_{JK} - \tilde{g}_{JL})$	
	$(J, K, L) = (X, Y, Z), (Y, Z, X), (Z, X, Y)$	4+4+4
$mg_{TXX}^{(M)}$	$-\frac{1}{6}(3\tilde{g}_- + \tilde{g}_Q)$	
$mg_{TY Y}^{(M)}$	$\frac{1}{6}(3\tilde{g}_- - \tilde{g}_Q)$	2
$mc_{XX}$	$\frac{1}{6}(\tilde{c}_Q + 3\tilde{c}_- + 2\tilde{c}_{TT})$	
$mc_{YY}$	$\frac{1}{6}(\tilde{c}_Q - 3\tilde{c}_- + 2\tilde{c}_{TT})$	
$mc_{ZZ}$	$\frac{1}{3}(-\tilde{c}_Q + \tilde{c}_{TT})$	3
$m \varepsilon_{JKL} c_{KL}$	$\tilde{c}_J$	
	$J = X, Y, Z, \text{ sum on } K, L$	1+1+1
$m(c_{TJ} + c_{JT})$	$\tilde{c}_{TJ}$	
	$J = X, Y, Z$	1+1+1 <b>Total: 44</b>

Table P43. Definitions for the photon sector of the minimal QED extension

Symbol	Combination	Components
$(\tilde{\kappa}_{e+})^{JK}$	$-(k_F)^{TJTK} + \frac{1}{4}\epsilon^{JPQ}\epsilon^{KRS}(k_F)^{PQRS}$	5
$(\tilde{\kappa}_{o-})^{JK}$	$\frac{1}{2}\epsilon^{KPQ}(k_F)^{TJPQ} + \frac{1}{2}\epsilon^{JPQ}(k_F)^{TKPQ}$	5
$(\tilde{\kappa}_{e-})^{JK}$	$-(k_F)^{TJTK} - \frac{1}{4}\epsilon^{JPQ}\epsilon^{KRS}(k_F)^{PQRS} + \frac{2}{3}(k_F)^{TLTL}\delta^{JK}$	5
$(\tilde{\kappa}_{o+})^{JK}$	$\frac{1}{2}\epsilon^{KPQ}(k_F)^{TJPQ} - \frac{1}{2}\epsilon^{JPQ}(k_F)^{TKPQ}$	3
$\tilde{\kappa}_{\text{tr}}$	$-\frac{2}{3}[(k_F)^{TXTX} + (k_F)^{TYTY} + (k_F)^{TZTZ}]$	1 <b>total: 19</b>
$k^1$	$(k_F)^{TYXZ}$	1
$k^2$	$(k_F)^{TXYZ}$	1
$k^3$	$(k_F)^{TYTY} - (k_F)^{XZZX}$	1
$k^4$	$(k_F)^{TZTZ} - (k_F)^{XYXY}$	1
$k^5$	$(k_F)^{TXTY} + (k_F)^{XZYZ}$	1
$k^6$	$(k_F)^{TXTZ} - (k_F)^{XYYZ}$	1
$k^7$	$(k_F)^{TYTZ} + (k_F)^{XYXZ}$	1
$k^8$	$(k_F)^{TXXY} + (k_F)^{TZYZ}$	1
$k^9$	$(k_F)^{TXXZ} - (k_F)^{TYYZ}$	1
$k^{10}$	$(k_F)^{TYXY} - (k_F)^{TZXX}$	1 <b>total: 10</b>
$k_{(V)00}^{(3)}$	$-\sqrt{4\pi}(k_{AF})^T$	1
$k_{(V)10}^{(3)}$	$-\sqrt{4\pi/3}(k_{AF})^Z$	1
$\text{Re } k_{(V)11}^{(3)}$	$\sqrt{2\pi/3}(k_{AF})^X$	1
$\text{Im } k_{(V)11}^{(3)}$	$-\sqrt{2\pi/3}(k_{AF})^Y$	1 <b>total: 4</b>

Table P44. Lagrange density for the fermion sector of the minimal SME in Riemann-Cartan spacetime

Sector	Coeff.	Operator	Dim.	Factor	CPT	L.V.
Lepton		$\bar{L}_A \gamma^a \overleftrightarrow{D}_\mu L_A$	4	$\frac{1}{2} i e e^\mu_a$	+	
		$\bar{R}_A \gamma^a \overleftrightarrow{D}_\mu R_A$	4	$\frac{1}{2} i e e^\mu_a$	+	
	$(a_L)_{\mu AB}$	$\bar{L}_A \gamma^a L_B$	3	$-e e^\mu_a$	-	✓
	$(a_R)_{\mu AB}$	$\bar{R}_A \gamma^a R_B$	3	$-e e^\mu_a$	-	✓
	$(c_L)_{\mu\nu AB}$	$\bar{L}_A \gamma^a \overleftrightarrow{D}^\nu L_B$	4	$-\frac{1}{2} i e e^\mu_a$	+	✓
	$(c_R)_{\mu\nu AB}$	$\bar{R}_A \gamma^a \overleftrightarrow{D}^\nu R_B$	4	$-\frac{1}{2} i e e^\mu_a$	+	✓
Quark		$\bar{Q}_A \gamma^a \overleftrightarrow{D}_\mu Q_A$	4	$\frac{1}{2} i e e^\mu_a$	+	
		$\bar{U}_A \gamma^a \overleftrightarrow{D}_\mu U_A$	4	$\frac{1}{2} i e e^\mu_a$	+	
		$\bar{D}_A \gamma^a \overleftrightarrow{D}_\mu D_A$	4	$\frac{1}{2} i e e^\mu_a$	+	
	$(a_Q)_{\mu AB}$	$\bar{Q}_A \gamma^a Q_B$	3	$-e e^\mu_a$	-	✓
	$(a_U)_{\mu AB}$	$\bar{U}_A \gamma^a U_B$	3	$-e e^\mu_a$	-	✓
	$(a_D)_{\mu AB}$	$\bar{D}_A \gamma^a D_B$	3	$-e e^\mu_a$	-	✓
	$(c_Q)_{\mu\nu AB}$	$\bar{Q}_A \gamma^a \overleftrightarrow{D}^\nu Q_B$	4	$-\frac{1}{2} i e e^\mu_a$	+	✓
	$(c_U)_{\mu\nu AB}$	$\bar{U}_A \gamma^a \overleftrightarrow{D}^\nu U_B$	4	$-\frac{1}{2} i e e^\mu_a$	+	✓
	$(c_D)_{\mu\nu AB}$	$\bar{D}_A \gamma^a \overleftrightarrow{D}^\nu D_B$	4	$-\frac{1}{2} i e e^\mu_a$	+	✓
Yukawa	$(G_L)_{AB}$	$\bar{L}_A \phi R_B + \text{h.c.}$	4	$-e$	+	
	$(G_U)_{AB}$	$\bar{Q}_A \phi^c U_B + \text{h.c.}$	4	$-e$	+	
	$(G_D)_{AB}$	$\bar{Q}_A \phi D_B + \text{h.c.}$	4	$-e$	+	
	$(H_L)_{\mu\nu AB}$	$\bar{L}_A \phi \sigma^{ab} R_B + \text{h.c.}$	4	$-\frac{1}{2} e e^\mu_a e^\nu_b$	+	✓
	$(H_U)_{\mu\nu AB}$	$\bar{Q}_A \phi^c \sigma^{ab} U_B + \text{h.c.}$	4	$-\frac{1}{2} e e^\mu_a e^\nu_b$	+	✓
	$(H_D)_{\mu\nu AB}$	$\bar{Q}_A \phi \sigma^{ab} D_B + \text{h.c.}$	4	$-\frac{1}{2} e e^\mu_a e^\nu_b$	+	✓

Table P45. Lagrange density for the boson sector of the minimal SME in Riemann-Cartan spacetime

Sector	Coeff.	Operator	Dim.	Factor	CPT	L.V.
Higgs	$\mu^2$	$\phi^\dagger \phi$	2	$e$	+	
	$\lambda$	$(\phi^\dagger \phi)^2$	4	$-\frac{1}{3!}e$	+	
		$(D_\mu \phi)^\dagger (D^\mu \phi)$	4	$-e$	+	
	$(k_\phi)^\mu$	$\phi^\dagger D_\mu \phi + \text{h.c.}$	3	$ie$	-	✓
	$(k_{\phi\phi})^{\mu\nu}$	$(D_\mu \phi)^\dagger (D_\nu \phi) + \text{h.c.}$	4	$\frac{1}{2}e$	+	✓
	$(k_{\phi W})^{\mu\nu}$	$\phi^\dagger W_{\mu\nu} \phi$	4	$-\frac{1}{2}e$	+	✓
	$(k_{\phi B})^{\mu\nu}$	$\phi^\dagger \phi B_{\mu\nu}$	4	$-\frac{1}{2}e$	+	✓
Gauge		$\text{Tr}(G_{\mu\nu} G^{\mu\nu})$	4	$-\frac{1}{2}e$	+	
		$\text{Tr}(W_{\mu\nu} W^{\mu\nu})$	4	$-\frac{1}{2}e$	+	
		$B_{\mu\nu} B^{\mu\nu}$	4	$-\frac{1}{4}e$	+	
	$(k_0)_\kappa$	$B^\kappa$	1	$e$	-	✓
	$(k_1)_\kappa$	$B_\lambda B_{\mu\nu}$	3	$e\epsilon^{\kappa\lambda\mu\nu}$	-	✓
	$(k_2)_\kappa$	$\text{Tr}(W_\lambda W_{\mu\nu} + \frac{2}{3}igW_\lambda W_\mu W_\nu)$	3	$e\epsilon^{\kappa\lambda\mu\nu}$	-	✓
	$(k_3)_\kappa$	$\text{Tr}(G_\lambda G_{\mu\nu} + \frac{2}{3}ig_3G_\lambda G_\mu G_\nu)$	3	$e\epsilon^{\kappa\lambda\mu\nu}$	-	✓
	$(k_G)_{\kappa\lambda\mu\nu}$	$\text{Tr}(G^{\kappa\lambda} G^{\mu\nu})$	4	$-\frac{1}{2}e$	+	✓
	$(k_W)_{\kappa\lambda\mu\nu}$	$\text{Tr}(W^{\kappa\lambda} W^{\mu\nu})$	4	$-\frac{1}{2}e$	+	✓
	$(k_B)_{\kappa\lambda\mu\nu}$	$B^{\kappa\lambda} B^{\mu\nu}$	4	$-\frac{1}{4}e$	+	✓
Gravity		$R$	2	$e/2\kappa$	+	
	$\Lambda$	1	0	$-e/\kappa$	+	
	$u$	$R$	2	$-e/2\kappa$	+	
	$(k_T)^{\lambda\mu\nu}$	$T_{\lambda\mu\nu}$	1	$e/2\kappa$	+	✓
	$s^{\mu\nu}$	$R_{\mu\nu}$	2	$e/2\kappa$	+	✓
	$t^{\kappa\lambda\mu\nu}$	$R_{\kappa\lambda\mu\nu}$	2	$e/2\kappa$	+	✓
	$(k_{TT})^{\alpha\beta\gamma\lambda\mu\nu}$	$T_{\alpha\beta\gamma} T_{\lambda\mu\nu}$	2	$e/2\kappa$	+	✓
	$(k_{DT})^{\kappa\lambda\mu\nu}$	$D_\kappa T_{\lambda\mu\nu}$	2	$e/2\kappa$	+	✓

Table P46. Coefficients in the renormalizable neutrino sector

Type	Coeff.	Dim.	Oscillation	CPT	L.V.
mass	$(m_l m_l^\dagger)_{ab}$	3	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	+	
massless	$(a_L)_{ab}^\mu \equiv [a_L^{(3)\mu}]_{ab}$	3	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	-	✓
	$H_{ab}^{\mu\nu} \equiv [H_{M+}^{(3)\mu\nu}]_{ab}$	3	$\nu \leftrightarrow \bar{\nu}$	+	✓
	$(c_L)_{ab}^{\mu\nu} \equiv [c_L^{(4)\mu\nu}]_{ab}$	4	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	+	✓
	$g_{ab}^{\mu\nu\sigma} \equiv [g_{M+}^{(4)\mu\nu\rho}]_{ab}$	4	$\nu \leftrightarrow \bar{\nu}$	-	✓
mass-induced	$[H_l^{(3)\mu\nu}]_{ab}$	3	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	+	✓
	$[a_l^{(3)\mu}]_{ab}$	3	$\nu \leftrightarrow \bar{\nu}$	-	✓
	$[e_l^{(4)\mu}]_{ab}$	4	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	-	✓
	$[g_l^{(4)\mu\nu\rho}]_{ab}$	4	$\nu \leftrightarrow \nu, \bar{\nu} \leftrightarrow \bar{\nu}$	-	✓
	$[c_l^{(4)\mu\nu}]_{ab}$	4	$\nu \leftrightarrow \bar{\nu}$	+	✓



Table P47. Definitions for the minimal neutrino sector

Symbol	Combination	Components
$(a_{\text{eff}}^{(3)})_{00}$	$\sqrt{4\pi}(a_L^{(3)T} - e_l^{(4)T} - 2\tilde{g}_l^{(4)TTT}) - (4\sqrt{\pi}/3)\tilde{g}_l^{(4)TJJ}$	
$(a_{\text{eff}}^{(3)})_{1-1}$	$-\sqrt{(2\pi/3)}\hat{X}_-^J(a_L^{(3)J} - e_l^{(4)J} - 2\tilde{g}_l^{(4)TJT})$	
$(a_{\text{eff}}^{(3)})_{10}$	$-\sqrt{(4\pi/3)}(a_L^{(3)Z} - e_l^{(4)Z} - 2\tilde{g}_l^{(4)TZT})$	
$(a_{\text{eff}}^{(3)})_{11}$	$\sqrt{(2\pi/3)}\hat{X}_+^J(a_L^{(3)J} - e_l^{(4)J} - 2\tilde{g}_l^{(4)TJT})$	
$(a_{\text{eff}}^{(3)})_{2-2}$	$-\sqrt{(8\pi/15)}\hat{X}_-\hat{X}_-^K\tilde{g}_l^{(4)TJK}$	
$(a_{\text{eff}}^{(3)})_{2-1}$	$-\sqrt{(8\pi/15)}\hat{X}_-^J(\tilde{g}_l^{(4)TZJ} + \tilde{g}_l^{(4)TJZ})$	
$(a_{\text{eff}}^{(3)})_{20}$	$-\sqrt{(16\pi/5)}(\tilde{g}_l^{(4)TZZ} - (1/3)\tilde{g}_l^{(4)TJJ})$	
$(a_{\text{eff}}^{(3)})_{21}$	$\sqrt{(8\pi/15)}\hat{X}_+^J(\tilde{g}_l^{(4)TZJ} + \tilde{g}_l^{(4)TJZ})$	
$(a_{\text{eff}}^{(3)})_{22}$	$-\sqrt{(8\pi/15)}\hat{X}_+\hat{X}_+^K\tilde{g}_l^{(4)TJK}$	$9 \times 9 = 81$
$(c_{\text{eff}}^{(2)})_{1-1}$	$\sqrt{(8\pi/3)}\hat{X}_-\tilde{H}_l^{(3)TJ}$	
$(c_{\text{eff}}^{(2)})_{10}$	$\sqrt{(16\pi/3)}\tilde{H}_l^{(3)TZ}$	
$(c_{\text{eff}}^{(2)})_{11}$	$-\sqrt{(8\pi/3)}\hat{X}_+\tilde{H}_l^{(3)TJ}$	$9 \times 3 = 27$
$(c_{\text{eff}}^{(4)})_{00}$	$\sqrt{4\pi}(c_L^{(4)TT} + (1/3)c_L^{(4)JJ})$	
$(c_{\text{eff}}^{(4)})_{1-1}$	$-\sqrt{(4\pi/6)}\hat{X}_-^J(c_L^{(4)TJ} + c_L^{(4)JT})$	
$(c_{\text{eff}}^{(4)})_{10}$	$-\sqrt{(4\pi/3)}(c_L^{(4)TZ} + c_L^{(4)ZT})$	
$(c_{\text{eff}}^{(4)})_{11}$	$\sqrt{(4\pi/6)}\hat{X}_+^J(c_L^{(4)TJ} + c_L^{(4)JT})$	
$(c_{\text{eff}}^{(4)})_{2-2}$	$\sqrt{(2\pi/15)}\hat{X}_-\hat{X}_-^Kc_L^{(4)JK}$	
$(c_{\text{eff}}^{(4)})_{2-1}$	$\sqrt{(2\pi/15)}\hat{X}_-^J(c_L^{(4)ZJ} + c_L^{(4)JZ})$	
$(c_{\text{eff}}^{(4)})_{20}$	$\sqrt{(4\pi/5)}(c_L^{(4)ZZ} - (1/3)c_L^{(4)JJ})$	
$(c_{\text{eff}}^{(4)})_{21}$	$-\sqrt{(2\pi/15)}\hat{X}_+^J(c_L^{(4)ZJ} + c_L^{(4)JZ})$	
$(c_{\text{eff}}^{(4)})_{22}$	$\sqrt{(2\pi/15)}\hat{X}_+\hat{X}_+^Kc_L^{(4)JK}$	$9 \times 9 = 81$
$(g_{\text{eff}}^{(2)})_{1-1}$	$\sqrt{(4\pi/3)}\hat{X}_-^J a_l^{(3)J}$	
$(g_{\text{eff}}^{(2)})_{10}$	$\sqrt{(8\pi/3)}a_l^{(3)Z}$	
$(g_{\text{eff}}^{(2)})_{11}$	$-\sqrt{(4\pi/3)}\hat{X}_+^J a_l^{(3)J}$	$12 \times 3 = 36$
$(g_{\text{eff}}^{(4)})_{1-1}$	$i\sqrt{(16\pi/3)}\hat{X}_-^J(g_{M+}^{(4)TJT} + (1/2)g_{M+}^{(4)TZJ} - (1/2)g_{M+}^{(4)TJZ})$	
$(g_{\text{eff}}^{(4)})_{10}$	$i\sqrt{(32\pi/3)}(g_{M+}^{(4)TZT} - (i/2)g_{M+}^{(4)TXY} + (i/2)g_{M+}^{(4)TYX})$	
$(g_{\text{eff}}^{(4)})_{11}$	$-i\sqrt{(16\pi/3)}\hat{X}_+^J(g_{M+}^{(4)TJT} - (1/2)g_{M+}^{(4)TZJ} + (1/2)g_{M+}^{(4)TJZ})$	
$(g_{\text{eff}}^{(4)})_{2-2}$	$-i\sqrt{(4\pi/5)}\hat{X}_-\hat{X}_-^K g_{M+}^{(4)TJK}$	
$(g_{\text{eff}}^{(4)})_{2-1}$	$-i\sqrt{(4\pi/5)}\hat{X}_-^J(g_{M+}^{(4)TZJ} + g_{M+}^{(4)TJZ})$	
$(g_{\text{eff}}^{(4)})_{20}$	$-i\sqrt{(24\pi/5)}(g_{M+}^{(4)TZZ} - (1/3)g_{M+}^{(4)TJJ})$	
$(g_{\text{eff}}^{(4)})_{21}$	$i\sqrt{(4\pi/5)}\hat{X}_+^J(g_{M+}^{(4)TZJ} + g_{M+}^{(4)TJZ})$	
$(g_{\text{eff}}^{(4)})_{22}$	$-i\sqrt{(4\pi/5)}\hat{X}_+\hat{X}_+^K g_{M+}^{(4)TJK}$	$12 \times 8 = 96$
$(H_{\text{eff}}^{(3)})_{1-1}$	$\sqrt{(4\pi/3)}\hat{X}_-^J(2iH_{M+}^{(3)TJ} + c_l^{(4)JT} + (1/2)c_l^{(4)ZJ} - (1/2)c_l^{(4)JZ})$	
$(H_{\text{eff}}^{(3)})_{10}$	$\sqrt{(8\pi/3)}(2iH_{M+}^{(3)TZ} + c_l^{(4)ZT} - (i/2)c_l^{(4)XY} + (i/2)c_l^{(4)YX})$	
$(H_{\text{eff}}^{(3)})_{11}$	$-\sqrt{(4\pi/3)}\hat{X}_+^J(2iH_{M+}^{(3)TJ} + c_l^{(4)JT} - (1/2)c_l^{(4)ZJ} + (1/2)c_l^{(4)JZ})$	
$(H_{\text{eff}}^{(3)})_{2-2}$	$-\sqrt{(\pi/5)}\hat{X}_-\hat{X}_-^Kc_l^{(4)JK}$	
$(H_{\text{eff}}^{(3)})_{2-1}$	$-\sqrt{(\pi/5)}\hat{X}_-^J(c_l^{(4)ZJ} + c_l^{(4)JZ})$	
$(H_{\text{eff}}^{(3)})_{20}$	$-\sqrt{(6\pi/5)}(c_l^{(4)ZZ} - (1/3)c_l^{(4)JJ})$	
$(H_{\text{eff}}^{(3)})_{21}$	$\sqrt{(\pi/5)}\hat{X}_+^J(c_l^{(4)ZJ} + c_l^{(4)JZ})$	
$(H_{\text{eff}}^{(3)})_{22}$	$-\sqrt{(\pi/5)}\hat{X}_+\hat{X}_+^Kc_l^{(4)JK}$	$6 \times 8 = 48$ <b>total: 369</b>

Table P48. Quadratic Lagrange density for a nonminimal Dirac fermion in Minkowski spacetime

Coeff.	#	Operator	Dim.	Factor	CPT	L.V.
$m$		$\bar{\psi}\psi$	3	-1	+	
$m_5$		$\bar{\psi}\gamma_5\psi$	3	$-i$	+	
		$\bar{\psi}\gamma^\mu\overleftrightarrow{\partial}_\mu\psi$	4	$\frac{1}{2}i$	+	
$m^{(d)\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)/6$	$\bar{\psi}\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	odd, $\geq 5$	$-\frac{1}{2}i^{d-3}$	+	✓
$m_5^{(d)\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)/6$	$\bar{\psi}\gamma_5\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	odd, $\geq 5$	$-\frac{1}{2}i^{d-2}$	+	✓
$a^{(d)\mu\alpha_1\dots\alpha_{d-3}}$	$2d(d-1)(d-2)/3$	$\bar{\psi}\gamma_\mu\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	odd, $\geq 3$	$-\frac{1}{2}i^{d-3}$	-	✓
$b^{(d)\mu\alpha_1\dots\alpha_{d-3}}$	$2d(d-1)(d-2)/3$	$\bar{\psi}\gamma_5\gamma_\mu\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	odd, $\geq 3$	$-\frac{1}{2}i^{d-3}$	-	✓
$c^{(d)\mu\alpha_1\dots\alpha_{d-3}}$	$2d(d-1)(d-2)/3$	$\bar{\psi}\gamma_\mu\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	even, $\geq 4$	$\frac{1}{2}i^{d-3}$	+	✓
$d^{(d)\mu\alpha_1\dots\alpha_{d-3}}$	$2d(d-1)(d-2)/3$	$\bar{\psi}\gamma_5\gamma_\mu\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	even, $\geq 4$	$\frac{1}{2}i^{d-3}$	+	✓
$e^{(d)\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)/6$	$\bar{\psi}\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	even, $\geq 4$	$\frac{1}{2}i^{d-3}$	-	✓
$f^{(d)\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)/6$	$\bar{\psi}\gamma_5\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	even, $\geq 4$	$-\frac{1}{2}i^{d-2}$	-	✓
$g^{(d)\mu\nu\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)$	$\bar{\psi}\sigma_{\mu\nu}\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	even, $\geq 4$	$\frac{1}{4}i^{d-3}$	-	✓
$H^{(d)\mu\nu\alpha_1\dots\alpha_{d-3}}$	$d(d-1)(d-2)$	$\bar{\psi}\sigma_{\mu\nu}\partial_{\alpha_1}\dots\partial_{\alpha_{d-3}}\psi + \text{h.c.}$	odd, $\geq 3$	$-\frac{1}{4}i^{d-3}$	+	✓

Table P49. Spherical coefficients for a nonminimal Dirac fermion in Minkowski spacetime

Type	Coeff.	Dim.	$n$	$j$	#	
spherical	$a_{njm}^{(d)}$	odd, $\geq 3$	$0, 1, \dots, d-2$	$n, n-2, n-4, \dots \geq 0$	$\frac{1}{6}(d+1)d(d-1)$	
	$c_{njm}^{(d)}$	even, $\geq 4$	$0, 1, \dots, d-2$	$n, n-2, n-4, \dots \geq 0$	$\frac{1}{6}(d+1)d(d-1)$	
	$g_{njm}^{(d)(0B)}$	even, $\geq 4$	$0, 1, \dots, d-3$	$n+1, n-1, n-3, \dots \geq 0$	$\frac{1}{6}(d+1)d(d-1) - 1$	
	$g_{njm}^{(d)(1B)}$	even, $\geq 4$	$2, 3, \dots, d-2$	$n-1, n-3, n-5, \dots \geq 1$	$\frac{1}{6}(d-2)(d^2 - d - 3)$	
	$g_{njm}^{(d)(1E)}$	even, $\geq 4$	$1, 2, \dots, d-2$	$n, n-2, n-4, \dots \geq 1$	$\frac{1}{6}(d+2)d(d-2)$	
	$H_{njm}^{(d)(0B)}$	odd, $\geq 3$	$0, 1, \dots, d-3$	$n+1, n-1, n-3, \dots \geq 0$	$\frac{1}{6}(d+1)d(d-1) - 1$	
	$H_{njm}^{(d)(1B)}$	odd, $\geq 5$	$2, 3, \dots, d-2$	$n-1, n-3, n-5, \dots \geq 1$	$\frac{1}{6}(d+1)(d-1)(d-3)$	
	$H_{njm}^{(d)(1E)}$	odd, $\geq 3$	$1, 2, \dots, d-2$	$n, n-2, n-4, \dots \geq 1$	$\frac{1}{6}(d-1)(d^2 + d - 3)$	
	isotropic spherical	$\hat{a}_n^{(d)}$	odd, $\geq 3$	$0, 2, 4, \dots, d-3$	0	$\frac{1}{2}(d-1)$
$\hat{c}_n^{(d)}$		even, $\geq 4$	$0, 2, 4, \dots, d-2$	0	$\frac{1}{2}d$	
$\hat{g}_n^{(d)}$		even, $\geq 4$	$1, 3, 5, \dots, d-3$	0	$\frac{1}{2}(d-2)$	
$\hat{H}_n^{(d)}$		odd, $\geq 5$	$1, 3, 5, \dots, d-4$	0	$\frac{1}{2}(d-3)$	
nonrelativistic		$a_{njm}^{\text{NR}}$	-	$\geq 0$	$n, n-2, n-4, \dots \geq 0$	$\frac{1}{2}(n+1)(n+2)$
	$c_{njm}^{\text{NR}}$	-	$\geq 0$	$n, n-2, n-4, \dots \geq 0$	$\frac{1}{2}(n+1)(n+2)$	
	$g_{njm}^{\text{NR}(0B)}$	-	$\geq 0$	$n+1, n-1, n-3, \dots \geq 0$	$\frac{1}{2}(n+2)(n+3)$	
	$g_{njm}^{\text{NR}(1B)}$	-	$\geq 0$	$n+1, n-1, n-3, \dots \geq 1$	$\frac{1}{2}(n+1+\iota_n)(n+4-\iota_n)$	
	$g_{njm}^{\text{NR}(1E)}$	-	$\geq 1$	$n, n-2, n-4, \dots \geq 1$	$\frac{1}{2}(n+1-\iota_n)(n+2+\iota_n)$	
	$H_{njm}^{\text{NR}(0B)}$	-	$\geq 0$	$n+1, n-1, n-3, \dots \geq 0$	$\frac{1}{2}(n+2)(n+3)$	
	$H_{njm}^{\text{NR}(1B)}$	-	$\geq 0$	$n+1, n-1, n-3, \dots \geq 1$	$\frac{1}{2}(n+1+\iota_n)(n+4-\iota_n)$	
	$H_{njm}^{\text{NR}(1E)}$	-	$\geq 1$	$n, n-2, n-4, \dots \geq 1$	$\frac{1}{2}(n+1-\iota_n)(n+2+\iota_n)$	
	isotropic nonrelativistic	$\hat{a}_n^{\text{NR}}$	-	even, $\geq 0$	0	1
$\hat{c}_n^{\text{NR}}$		-	even, $\geq 0$	0	1	
$\hat{g}_n^{\text{NR}}$		-	odd, $\geq 0$	0	1	
$\hat{H}_n^{\text{NR}}$		-	odd, $\geq 0$	0	1	
ultrarelativistic		$a_{jm}^{\text{UR}(d)}$	odd, $\geq 3$	-	$0 \leq j \leq d-2$	$(d-1)^2$
	$c_{jm}^{\text{UR}(d)}$	even, $\geq 4$	-	$0 \leq j \leq d-2$	$(d-1)^2$	
	$g_{jm}^{\text{UR}(d)(0B)}$	even, $\geq 4$	-	$0 \leq j \leq d-3$	$(d-2)^2$	
	$g_{jm}^{\text{UR}(d)(1B)}$	even, $\geq 4$	-	$1 \leq j \leq d-2$	$(d-2)d$	
	$g_{jm}^{\text{UR}(d)(1E)}$	even, $\geq 4$	-	$1 \leq j \leq d-2$	$(d-2)d$	
	$H_{jm}^{\text{UR}(d)(0B)}$	odd, $\geq 5$	-	$0 \leq j \leq d-3$	$(d-2)^2$	
	$H_{jm}^{\text{UR}(d)(1B)}$	odd, $\geq 3$	-	$1 \leq j \leq d-2$	$(d-2)d$	
	$H_{jm}^{\text{UR}(d)(1E)}$	odd, $\geq 3$	-	$1 \leq j \leq d-2$	$(d-2)d$	
	isotropic ultrarelativistic	$\hat{a}^{\text{UR}(d)}$	odd, $\geq 3$	-	0	1
		$\hat{c}^{\text{UR}(d)}$	even, $\geq 4$	-	0	1
$\hat{g}^{\text{UR}(d)}$		even, $\geq 4$	-	0	1	
$\hat{H}^{\text{UR}(d)}$		even, $\geq 5$	-	0	1	

Table P50. Quadratic Lagrange density for the nonminimal photon sector in Minkowski spacetime

Coeff.	#	Operator	Dim.	Factor	CPT	L.V.
		$F_{\mu\nu}F^{\mu\nu}$	4	$-\frac{1}{4}$	+	
$(k_{AF}^{(3)})_{\kappa} \equiv (k_{AF})_{\kappa}$	4	$A_{\lambda}F_{\mu\nu}$	3	$\frac{1}{2}\epsilon^{\kappa\lambda\mu\nu}$	-	✓
$(k_{AF}^{(5)})_{\kappa}^{\alpha_1\alpha_2}$	36	$A_{\lambda}\partial_{\alpha_1}\partial_{\alpha_2}F_{\mu\nu}$	5	$\frac{1}{2}\epsilon^{\kappa\lambda\mu\nu}$	-	✓
$(k_{AF}^{(7)})_{\kappa}^{\alpha_1\alpha_2\alpha_3\alpha_4}$	120	$A_{\lambda}\partial_{\alpha_1}\partial_{\alpha_2}\partial_{\alpha_3}\partial_{\alpha_4}F_{\mu\nu}$	7	$\frac{1}{2}\epsilon^{\kappa\lambda\mu\nu}$	-	✓
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$		
$(k_{AF}^{(d)})_{\kappa}^{\alpha_1\cdots\alpha_{(d-3)}}$	$\frac{1}{2}(d+1)(d-1)(d-2)$	$A_{\lambda}\partial_{\alpha_1}\cdots\partial_{\alpha_{(d-3)}}F_{\mu\nu}$	odd $d$	$\frac{1}{2}\epsilon^{\kappa\lambda\mu\nu}$	-	✓
$(k_F^{(4)})^{\kappa\lambda\mu\nu} \equiv (k_F)^{\kappa\lambda\mu\nu}$	19 + 1	$F_{\kappa\lambda}F_{\mu\nu}$	4	$-\frac{1}{4}$	+	✓
$(k_F^{(6)})^{\kappa\lambda\mu\nu\alpha_1\alpha_2}$	126	$F_{\kappa\lambda}\partial_{\alpha_1}\partial_{\alpha_2}F_{\mu\nu}$	6	$-\frac{1}{4}$	+	✓
$(k_F^{(8)})^{\kappa\lambda\mu\nu\alpha_1\alpha_2\alpha_3\alpha_4}$	360	$F_{\kappa\lambda}\partial_{\alpha_1}\partial_{\alpha_2}\partial_{\alpha_3}\partial_{\alpha_4}F_{\mu\nu}$	8	$-\frac{1}{4}$	+	✓
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$		
$(k_F^{(d)})^{\kappa\lambda\mu\nu\alpha_1\cdots\alpha_{(d-4)}}$	$(d+1)d(d-3)$	$F_{\kappa\lambda}\partial_{\alpha_1}\cdots\partial_{\alpha_{(d-4)}}F_{\mu\nu}$	even $d$	$-\frac{1}{4}$	+	✓

Table P51. Spherical coefficients for the nonminimal photon sector in Minkowski spacetime

Type	Coeff.	Dim.	$n$	$j$	#
vacuum	$c_{(I)jm}^{(d)}$	even, $\geq 4$	-	$0, 1, \dots, d-2$	$(d-1)^2$
	$k_{(E)jm}^{(d)}$	even, $\geq 4$	-	$2, 3, \dots, d-2$	$(d-1)^2 - 4$
	$k_{(B)jm}^{(d)}$	even, $\geq 4$	-	$2, 3, \dots, d-2$	$(d-1)^2 - 4$
	$k_{(V)jm}^{(d)}$	odd, $\geq 3$	-	$0, 1, \dots, d-2$	$(d-1)^2$
vacuum orthogonal	$(\bar{c}_F^{(d)})_{njm}^{(0E)}$	even, $\geq 4$	$0, 1, \dots, d-4$	$n, n-2, n-4, \dots, \geq 0$	$\frac{(d-1)(d-2)(d-3)}{6}$
	$(\bar{k}_F^{(d)})_{njm}^{(0E)}$	even, $\geq 6$	$1, 2, \dots, d-4$	$n, n-2, n-4, \dots, \geq 0$	$\frac{(d-1)(d-2)(d-3)}{6} - 1$
	$(\bar{k}_F^{(d)})_{njm}^{(1E)}$	even, $\geq 6$	$1, 2, \dots, d-4$	$n+1, n-1, n-3, \dots, \geq 1$	$\frac{(d-4)(d^2+d+3)}{6}$
	$(\bar{k}_F^{(d)})_{njm}^{(2E)}$	even, $\geq 6$	$2, 3, \dots, d-4$	$n, n-2, n-4, \dots, \geq 2$	$\frac{(d-4)(d^2-2d-9)}{6}$
	$(\bar{k}_F^{(d)})_{njm}^{(1B)}$	even, $\geq 6$	$1, 2, \dots, d-4$	$n, n-2, n-4, \dots, \geq 1$	$\frac{d(d-2)(d-4)}{6}$
	$(\bar{k}_F^{(d)})_{njm}^{(2B)}$	even, $\geq 6$	$1, 2, \dots, d-4$	$n+1, n-1, n-3, \dots, \geq 2$	$\frac{(d+3)(d-2)(d-4)}{6}$
	$(\bar{k}_{AF}^{(d)})_{njm}^{(0B)}$	odd, $\geq 5$	$0, 1, \dots, d-4$	$n, n-2, n-4, \dots, \geq 0$	$\frac{(d-1)(d-2)(d-3)}{6}$
	$(\bar{k}_{AF}^{(d)})_{njm}^{(1B)}$	odd, $\geq 5$	$0, 1, \dots, d-4$	$n+1, n-1, n-3, \dots, \geq 1$	$\frac{(d+1)(d-1)(d-3)}{6}$
	$(\bar{k}_{AF}^{(d)})_{njm}^{(1E)}$	odd, $\geq 5$	$1, 2, \dots, d-3$	$n, n-2, n-4, \dots, \geq 1$	$\frac{(d+1)(d-1)(d-3)}{6}$

Table P52. Spherical coefficients for the nonminimal neutrino sector in Minkowski spacetime

Type	Coeff.	Dim.	$j$	#	CPT
effective	$(a_{\text{eff}}^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 1 \geq j \geq 0$	$9d^2$	-
	$(c_{\text{eff}}^{(d)})_{jm}^{ab}$	$\begin{cases} d = 2 \\ \text{even, } \geq 4 \end{cases}$	$\begin{cases} j = 1 \\ d - 1 \geq j \geq 0 \end{cases}$	$\begin{cases} 27 \\ 9d^2 \end{cases}$	+
	$(g_{\text{eff}}^{(d)})_{jm}^{ab}$	even, $\geq 2$	$d - 1 \geq j \geq 1$	$12(d^2 - 1)$	-
	$(H_{\text{eff}}^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 1 \geq j \geq 1$	$6(d^2 - 1)$	+
Dirac	$(a_L^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 2 \geq j \geq 0$	$9(d - 1)^2$	-
	$(c_L^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$9(d - 1)^2$	+
	$(m_l^{(d)})_{jm}^{ab}$	odd, $\geq 5$	$d - 3 \geq j \geq 0$	$9(d - 2)^2$	+
	$(e_l^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 3 \geq j \geq 0$	$9(d - 2)^2$	-
	$(g_l^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$9(d - 1)^2$	-
	$(H_l^{(d)})_{jm}^{ab}$	$\begin{cases} d = 3 \\ \text{odd, } \geq 5 \end{cases}$	$\begin{cases} j = 1 \\ d - 2 \geq j \geq 0 \end{cases}$	$\begin{cases} 27 \\ 9(d - 1)^2 \end{cases}$	+
Majorana	$(g_{M+}^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 1$	$12d(d - 2)$	-
	$(H_{M+}^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 2 \geq j \geq 1$	$6d(d - 2)$	+
	$(a_l^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 2 \geq j \geq 1$	$12d(d - 2)$	-
	$(c_l^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 1$	$6d(d - 2)$	+
renormalizable	$(a_{\text{eff}}^{(3)})_{jm}$	3	0, 1, 2	81	-
	$(c_{\text{eff}}^{(2)})_{jm}$	2	1	27	+
	$(c_{\text{eff}}^{(4)})_{jm}$	4	0, 1, 2	81	+
	$(g_{\text{eff}}^{(2)})_{jm}$	2	1	36	-
	$(g_{\text{eff}}^{(4)})_{jm}$	4	1, 2	96	-
	$(H_{\text{eff}}^{(3)})_{jm}$	3	1, 2	48	+
massless	$(a_L^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 2 \geq j \geq 0$	$9(d - 1)^2$	-
	$(c_L^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$9(d - 1)^2$	+
	$(g_{M+}^{(d)})_{jm}^{ab}$	even, $\geq 4$	$d - 2 \geq j \geq 1$	$12d(d - 2)$	-
	$(H_{M+}^{(d)})_{jm}^{ab}$	odd, $\geq 3$	$d - 2 \geq j \geq 1$	$6d(d - 2)$	+
flavor-blind	$(a_{\text{fb}}^{(d)})_{jm}$	odd, $\geq 3$	$d - 1 \geq j \geq 0$	$d^2$	-
	$(c_{\text{fb}}^{(d)})_{jm}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$(d - 1)^2$	+
	$(g_{\text{fb}}^{(d)})_{jm}$	even, $\geq 2$	$d - 1 \geq j \geq 1$	$2(d^2 - 1)$	-
oscillation-free	$(a_{\text{of}}^{(d)})_{jm}$	odd, $\geq 3$	$d - 2 \geq j \geq 0$	$(d - 1)^2$	-
	$(c_{\text{of}}^{(d)})_{jm}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$(d - 1)^2$	+
diagonalizable	$(a_d^{(d)})_{jm}^{a'}$	odd, $\geq 3$	$d - 1 \geq j \geq 0$	$3d^2$	-
	$(c_d^{(d)})_{jm}^{a'}$	even, $\geq 4$	$d - 2 \geq j \geq 0$	$3(d - 1)^2$	+
	$(g_d^{(d)})_{jm}^{a'}$	even, $\geq 2$	$d - 1 \geq j \geq 1$	$6(d^2 - 1)$	-
generic isotropic	$\hat{a}_{ab}^{(d)}$	odd, $\geq 3$		9	-
	$\hat{c}_{ab}^{(d)}$	even, $\geq 4$		9	+
isotropic diag.	$\hat{a}_{a'}^{(d)}$	odd, $\geq 3$		3	-
	$\hat{c}_{a'}^{(d)}$	even, $\geq 4$		3	+
isotropic osc.-free	$\hat{a}^{(d)}$	odd, $\geq 3$		1	-
	$\hat{c}^{(d)}$	even, $\geq 4$		1	+