

# The Black Hole ?

Ying-Qiu Gu\*

School of Mathematical Science, Fudan University, Shanghai 200433, China

## Abstract

In this paper, we revisit the formation of black hole in a collapsing star via fully dynamical approach. We derive and simplify the complete dynamical equation system for a collapsing star with spherical symmetry, and establish some new constraints on the initial and boundary conditions. From the analysis we find that, the black hole might not exist if the initial distributions of mass density and velocity are reasonable, and the dynamical system can even automatically remove the weak singularity in the initial data.

**Key Words:** *black hole, singularity, gravitational collapse, dynamical process*

**PACS numbers:** 04.20.Dw, 04.70.-s, 97.60.Lf, 98.35.Jk

## 1 Dynamical Equations for a Collapsing Star

The black hole come forth in physical theory when Schwarzschild got his solution to the Einstein equation in 1916[1]. After a series singularity theorems were proved, the existence of the practical black hole has been naturally accepted by most scientists. There are only a very few skeptics, who proposed some special matter models which lead to global solutions or the violation of the conditions for singularity theorems(see [2]-[6] and references therein). However the evolution of a collapsing star and the formation of a black hole was not clearly checked with dynamical model. In [7], the stellar models are all singularity-free, and the analysis hint that the singularity of the space-time may generally not exist. Here we give a fully dynamical approach to this problem. The analysis clearly shows that, the singularity can be avoidable if the initial distributions

---

\*email: yqgu@fudan.edu.cn

of the mass density and velocity are reasonable, and the dynamical system even can automatically remove the weak singularity in the initial data.

Since the asymmetry such as rotation always alleviate the singularity of solutions to the Einstein equation, it is enough to check the case of spherical symmetry, which is the worst case. The line element for a spherical symmetric collapsing star is generally given by

$$ds^2 = u^2 d\tilde{t}^2 - (v d\tilde{t} - w dr)^2 - r^2(d\theta^2 + \sin^2 \theta d\varphi^2). \quad (1.1)$$

In this paper, we set light velocity  $c = 1$  as unit of speed. Making transformation

$$\tilde{t} = t + \int \frac{w}{u+v} dr, \quad (1.2)$$

we get the equivalent canonical form for (1.1) as[8, 9]

$$ds^2 = abdt^2 + 2\sqrt{a} dt dr - r^2(d\theta^2 + \sin^2 \theta d\varphi^2), \quad (1.3)$$

where  $a, b$  are continuous functions of  $(t, r)$  before the star becomes singular. For the external Schwarzschild solution, we have

$$a = 1, \quad b = 1 - \frac{R_s}{r}, \quad (\text{for } r \geq R > R_s), \quad (1.4)$$

where  $R = R(t)$  and  $R_s$  are respectively radius and Schwarzschild radius of the star,

$$R_s \equiv 8\pi G \int_0^{R(t)} \rho_{\text{tot}} r^2 dr, \quad (1.5)$$

where  $\rho_{\text{tot}}$  is the total gravitational mass density defined below, which includes the influence of pressure or momentum of the fluid, and is different from the usual  $\rho$ .

Denote the 4-vector speed of the fluid by

$$U^\mu = \{U, V, 0, 0\}, \quad U_\mu = \{abU + \sqrt{a}V, \sqrt{a}U, 0, 0\}, \quad (1.6)$$

which satisfying the line element equation

$$1 = g_{\mu\nu} U^\mu U^\nu = (abU + 2\sqrt{a}V) U. \quad (1.7)$$

So between  $(U, V)$  only  $V$  is independent variable, we get

$$U = \frac{1}{\sqrt{a}(\sqrt{b+V^2} + V)} = \frac{1}{\sqrt{a}b}(\sqrt{b+V^2} - V). \quad (1.8)$$

The physical meaning of  $V$  is approximately the usual speed when  $|V| \ll 1$ .

For the energy momentum tensor of perfect fluid  $T_{\mu\nu} = (\rho + P)U_\mu U_\nu - Pg_{\mu\nu}$ , we have the nonzero components as follow

$$T_{tt} = a(\rho + P) \left( \sqrt{ab}U + V \right)^2 - abP, \quad (1.9)$$

$$T_{tr} = a(\rho + P) \left( \sqrt{ab}U + V \right) U - \sqrt{a}P = T_{rt}, \quad (1.10)$$

$$T_{rr} = a(\rho + P) U^2, \quad T_{\theta\theta} = Pr^2, \quad T_{\varphi\varphi} = Pr^2 \sin^2 \theta, \quad (1.11)$$

where  $P = P(\rho)$  is a given equation of state. For the classical particles, we have  $(\rho, P)$  in comoving coordinate system as[10]-[13]

$$\rho \equiv \sum_k \frac{m_k}{\sqrt{1 - v_k^2}} \delta^3(\vec{x} - \vec{X}_k), \quad P \equiv \frac{1}{3} \sum_k \frac{m_k v_k^2}{\sqrt{1 - v_k^2}} \delta^3(\vec{x} - \vec{X}_k), \quad (1.12)$$

Obviously, we have  $P \rightarrow \frac{1}{3}\rho$  for high temperature due to  $\forall v_k \rightarrow 1$ . So for reasonable matter model we should have

$$\frac{dP}{d\rho} \geq 0, \quad C_s = \sqrt{\frac{dP}{d\rho}} \leq \frac{\sqrt{3}}{3}, \quad (1.13)$$

where  $C_s$  stands for the velocity of sound, which is a increasing function of  $\rho$ . Even for the dark spinors with negative pressure, calculation also shows  $\frac{dP}{d\rho} > 0$  for adequate parameters[14]. However, for the following discussion, we can weaken the requirements for  $C_s$  as the following requirement of causality

$$C_s \rightarrow C_0 < 1, \quad (\rho \rightarrow \infty). \quad (1.14)$$

The nonzero components of the Einstein tensor reads

$$G_{tt} = -\frac{1}{r} \left( \sqrt{a}\partial_t b - ab\partial_r b \right) - \frac{1}{r^2} ab(1 - b), \quad (1.15)$$

$$G_{tr} = \frac{1}{r} \sqrt{a}\partial_r b - \frac{1}{r^2} \sqrt{a}(1 - b) = G_{rt}, \quad G_{rr} = -\frac{\partial_r a}{ra}, \quad (1.16)$$

$$G_{\theta\theta} = \left( \frac{b}{r} \left( \frac{\partial_r a}{2a} + \frac{\partial_r b}{b} \right) - \frac{1 - b}{r^2} + \frac{\mathcal{R}}{2} \right) r^2, \quad G_{\varphi\varphi} = G_{\theta\theta} \sin^2 \theta, \quad (1.17)$$

where the scalar curvature  $\mathcal{R}$  depends on the second order derivatives of  $(a, b)$ , which has no use in the following calculation.

For the Einstein equation  $G_{\mu\nu} = -8\pi GT_{\mu\nu}$ , detailed calculation shows only the following 3 equations are independent,

$$\partial_r a = 8\pi Gr(\rho + P)a^2 U^2, \quad (1.18)$$

$$\partial_t b = 8\pi G(\rho + P)rV\sqrt{a(b + V^2)}, \quad (1.19)$$

$$\partial_r b = -4\pi Gr \left( (\rho - P) + (\rho + P)abU^2 \right) + \frac{1 - b}{r}. \quad (1.20)$$

By (1.4), (1.5) and (1.19), we find  $R_s$  is a conserved constant for an evolutionary star.

For the energy momentum conservation law  $T^{\mu\nu}_{;\nu} = 0$ , calculation also shows only one independent equation due to (1.8). We choose the continuity equation  $U_\mu T^{\mu\nu}_{;\nu} = 0$ , which leads to

$$U^\mu \partial_\mu \rho + (\rho + P) U^\mu_{;\mu} = 0. \quad (1.21)$$

Simplifying (1.21) and the consistent equation of (1.20) and (1.19), we get a hyperbolic system for  $(\rho, V)$  as follows

$$\begin{aligned} (1 - C_s^2) \sqrt{a} U^2 \partial_t \rho + \frac{1}{2} [2 - (1 - C_s^2)(1 + abU^2)] \partial_r \rho + \frac{2(\rho + P) \sqrt{a} U}{1 + abU^2} \partial_r V \\ = -\frac{1}{r} \left( \frac{aU^2}{1 + abU^2} (1 - b + 8\pi G r^2 P) + 2\sqrt{a} UV \right) (\rho + P), \end{aligned} \quad (1.22)$$

$$\begin{aligned} (1 - C_s^2) \sqrt{a} U^2 \partial_t V + \frac{1}{2} [2 - (1 - C_s^2)(1 + abU^2)] \partial_r V + \frac{(1 + abU^2) C_s^2}{2(\rho + P) \sqrt{a} U} \partial_r \rho \\ = -\frac{1}{2r} \left( \sqrt{a} U (1 - b + 8\pi G r^2 P) + 2C_s^2 (1 + abU^2) V \right). \end{aligned} \quad (1.23)$$

The characteristic velocity for the above system is given by

$$V_1 = \frac{V}{U} + \frac{C_s}{(1 - C_s) \sqrt{a} U^2} = \frac{V + C_s \sqrt{b + V^2}}{U(1 - C_s)}, \quad (1.24)$$

$$V_2 = \frac{V}{U} - \frac{C_s}{(1 + C_s) \sqrt{a} U^2} = \frac{V - C_s \sqrt{b + V^2}}{U(1 + C_s)}. \quad (1.25)$$

Four equations among (1.18)-(1.19) and (1.22)-(1.23) form a closed system, and the redundant equation can be treated as identity.

As done in [7], we can get the dimensionless form of the equation system by setting  $(8\pi G \varrho)^{-\frac{1}{2}}$  as length unit, where  $\varrho$  stands for a characteristic mass density. A theoretical analysis on this scale invariance and its implications see [15]. A direct result of this property is that, even if the black hole exists, its formation is irrelevant with the total mass of a star, but relevant with the distribution of mass density and velocity.

For a regular star, we have the following ranges for the variables

$$0 < a \leq 1, \quad 0 < b \leq 1, \quad 0 \leq \rho < \infty, \quad |V| < \infty. \quad (1.26)$$

By (1.18) and (1.20) we find that,  $a(\cdot, r) \in C^0([0, R])$  is a monotone increasing function with respect to  $r$ , and  $b(\cdot, r) \in C^1([0, \infty))$  has a positive minimum. The reasonable initial and boundary conditions are as follow:

$$\text{at } r = 0: \quad b = 1, \quad \partial_r b = 0, \quad V = 0; \quad (1.27)$$

$$\text{at } r = R: \quad a = 1, \quad b = 1 - \frac{R_s}{R}, \quad \rho = 0. \quad (1.28)$$

The initial data of  $(\rho, V)$  is quite arbitrary, but they should satisfy (1.18), (1.20) and the above boundary conditions, as well as a constraint similar to

$$\frac{8\pi G}{r} \int_0^r \rho(0, r) r^2 dr < 1, \quad (1.29)$$

or in differential form

$$8\pi G \rho(0, r) r^2 < 1. \quad (1.30)$$

(1.29) or (1.30) means that, we can not set a small black hole inside the star in the initial data, because a proof of singularity from singular conditions is meaningless. The concrete form for such condition will be derived in the next section.

## 2 Analysis of the Singularity

Define an auxiliary energy function by

$$F \equiv (\rho + P)abU^2 = (\rho + P) \frac{\sqrt{b + V^2} - V}{\sqrt{b + V^2} + V}. \quad (2.1)$$

For a static star, we have  $F = \rho + P$ . Substituting (2.1) into equation (1.20), we get

$$1 - b - r\partial_r b = 4\pi G r^2 (\rho - P + F). \quad (2.2)$$

Instead of  $V$ , if we take  $F$  as independent variable in some analysis, then solving (2.2) we get

$$b = 1 - \frac{4\pi G}{r} \int_0^r (\rho - P + F) r^2 dr. \quad (2.3)$$

In contrast (2.3) with (1.4), we get the gravitational mass density defined in (1.5) as

$$\rho_{\text{tot}} = \frac{1}{2}(\rho - P + F) = \frac{\rho\sqrt{b + V^2} - PV}{\sqrt{b + V^2} + V}. \quad (2.4)$$

(2.4) shows how the gravitational mass density, and then the ADM mass, are related to the pressure and velocity of the fluid  $(P, V)$ . By conditions (1.27) and (1.28), we find  $F$  satisfies

$$F(t, 0) = \rho(t, 0) + P(t, 0), \quad F(t, R) = 0. \quad (2.5)$$

By (1.18) and (1.19), we get

$$4\sqrt{a}\partial_t b + b^2\partial_r a = 8\pi G r a (\rho + P) \left( \sqrt{b + V^2} + V \right)^2 \geq 0. \quad (2.6)$$

The solution is given by

$$b(t, r) \geq b_0 \left( 1 + \frac{b_0}{2} \int_{t_0}^t (\partial_r \sqrt{a}) dt \right)^{-1} > 0, \quad b_0 = b(t_0, r), \quad (2.7)$$

which means  $b(t, r)$  is regular until  $a(t, r)$  becomes singular.

Substituting (1.8) into (1.18), we get

$$\partial_r a = \frac{8\pi G(\rho + P)ra}{(\sqrt{b + V^2} + V)^2}. \quad (2.8)$$

Since  $b > 0$ , solving (2.8) with  $a(t, R) = 1$ , we get

$$a = \exp \left( - \int_r^R \frac{8\pi G(\rho + P)rdr}{(\sqrt{b + V^2} + V)^2} \right). \quad (2.9)$$

Obviously,  $0 < a \leq 1$  is an increasing continuous function of  $r$ . The asymptotic singularity  $a(t, 0) \rightarrow 0$  only occurs as  $\rho(t, r) \rightarrow \rho_0 r^{-n}$ , ( $n \geq 2$ ). Noticing (2.3) and (2.9), even if  $\rho(t, r) \rightarrow \infty$  with  $0 < n < 2$ , the space-time still keep regular, and in what follows we show that the dynamical equation (1.22) and (1.23) can automatically remove such weak singularity soon.

We examine the asymptotic behavior of  $(\rho, V)$  near the center. Noticing the finite propagating speed (1.24) and (1.25), for a given time range  $0 \leq t \leq T_0$ ,  $\rho(t, 0)$  only depends on the initial data near the center. For a collapsing star, the worst case is the highly compact star with large  $(\rho, P)$  and  $V < 0$  near the center. Since the large  $P$  leads large temperature[7], for such extremely hot gas, the equation of state becomes simple, we have

$$P \doteq C_0^2 \rho, \quad C_s \doteq C_0 < 1. \quad (2.10)$$

Noticing  $V(t, 0) = 0$ , then we have the following approximations near the center by omitting  $O(r^2)$

$$|V| \ll 1, \quad U \doteq \frac{\sqrt{b} - V}{\sqrt{a} b}, \quad (2.11)$$

$$a \doteq a_0(t), \quad b \doteq 1. \quad (2.12)$$

Substituting (2.10)-(2.12) into (1.22) and (1.23), we get the following equations

$$\frac{1 - C_0^2}{1 + C_0^2} \frac{\partial_r \rho}{\rho} + \frac{C_0^2}{1 + C_0^2} \frac{\partial_r \rho}{\rho} + \partial_r V + \frac{2V}{r} + 4\pi G C_0^2 \rho r = 0, \quad (2.13)$$

$$\frac{1 - C_0^2}{C_0^2} \partial_r V + \frac{1}{1 + C_0^2} \frac{\partial_r \rho}{\rho} + \partial_r V + \frac{2V}{r} + 4\pi G \rho r = 0, \quad (2.14)$$

where  $d\tau \equiv \sqrt{a_0} dt$  is something similar to proper time. By (2.13) and (2.14), we get a linear equation for  $\rho^{-1}$

$$\partial_\tau \rho^{-1} - \partial_r \rho^{-1} + (1 + C_0^{-2}) \partial_\tau V \rho^{-1} + 4\pi G(1 + C_0^2)r = 0. \quad (2.15)$$

The solution is given by

$$\rho(\tau, r) = \frac{\rho(0, \tau + r)}{f(\tau, r)} \exp\left((1 + C_0^{-2})[V(\tau, r) - V(0, \tau + r)]\right), \quad (2.16)$$

where

$$f(\tau, r) \equiv 1 - 4\pi G(1 + C_0^2)\rho(0, \tau + r) \times \int_0^\tau (\tau + r - x) \exp\left((1 + C_0^{-2})[V(x, \tau + r - x) - V(0, \tau + r)]\right) dx. \quad (2.17)$$

Obviously,  $\rho(\tau, 0) \rightarrow \infty$  if and only if  $f(\tau, 0) \rightarrow 0$ .

Noticing the condition  $V(x, r) \leq 0$ , we have the following estimation

$$\exp\left((1 + C_0^{-2})[V(x, \tau - x) - V(0, \tau)]\right) \leq \exp\left(-(1 + C_0^{-2})V(0, \tau)\right). \quad (2.18)$$

Substituting it into (2.17), we get

$$f(\tau, 0) \geq 1 - 2\pi G(1 + C_0^2)\rho(0, \tau)\tau^2 \exp\left(-(1 + C_0^{-2})V(0, \tau)\right). \quad (2.19)$$

Then the black hole can not form if

$$2\pi G(1 + C_0^2)\rho(0, r)r^2 \exp\left(-(1 + C_0^{-2})V(0, r)\right) < 1. \quad (2.20)$$

(2.20) is a more definite but weaker constraint on initial data than (1.30).

By (2.16), we find that, even if the initial mass density has weak singularity  $\rho(0, r) \rightarrow \rho_0 r^{-n}$ , ( $0 < n < 2$ ), it will soon become regular. This is a negative feedback process, so it is really hard to form a black hole.

### 3 Discussion and Conclusion

The above analysis shows that, the black hole may not form under reasonable and regular initial and boundary conditions. The dynamical system can automatically remove the weak singularity in the initial data. Even if the high order true singularity occurs, it is an asymptotic singularity irrelative to the total mass of a star, but relative to the distribution of mass density and velocity, and it will spend infinity time  $t$  to reach the asymptotic singularity due to  $a_0 \rightarrow 0$  in (2.9) and  $t = \int_0^r a_0^{-1} d\tau$  for  $f(\tau, 0) \rightarrow 0$  in (2.19).

The physical phenomenon is a subtle collaboration of all dynamical variables, but the conventional research of black hole seems to use few values of dynamical variables.

The dynamical equations usually include second order derivatives of variables, and Jacobian matrix of coordinate transformation is constituted by the first order derivatives of coordinates, so a valid coordinate transformation should have bounded second order derivatives in its effective domain. However, in the research of black hole, some coordinate transformations without continuous derivatives may not be globally valid, which unconsciously add source terms like  $\delta(r - 2m)$  and  $\delta'(r - 2m)$  to the singular surface.

In the above discussion, we only used ( $\rho \geq 0, 0 < \frac{dP}{d\rho} < 1$ ) for matter state, which are quite usual conditions. What is most important to exclude the singularity is the distribution of  $\rho$  and  $V$ . In cosmology,  $\rho(0, r)$  is a constant, then the ADM mass

$$M(r) \equiv 4\pi \int_0^r \rho_{\text{tot}} r^2 dr \propto r^3, \quad (3.1)$$

which inevitably leads to  $b = 1 - \frac{2M}{r} \leq 0$  for large enough  $r$ . So we need negative pressure to get a singularity-free and evolutionary universe, and this dark energy should play an important role in the formation of large scale structure of galaxies. But in a star, the main particles can be treated as ideal gases[7]. The singularity theorems might include ambiguous concepts and unrealistic conditions similar to the singular coordinate transformation.

The conserved mass of a star should be ADM mass  $M$ , rather than  $4\pi \int_0^R \rho \sqrt{A} r^2 dr$  as used in some textbooks. This constraints are important for numerical simulation. In addition, the equation system (1.18)-(1.20) and (1.22)-(1.23) is strongly stiff when the star heavily shrinks with large  $R_s : R$ .

## Acknowledgments

The author is grateful to his supervisor Prof. Ta-Tsien Li and Prof. Han-Ji Shang for their encouragement. The work is directed by Prof. Li, but he does not enjoy the non-exact results and requires to cancel his name in the authors list. The work is supported by the Postdoctoral Station of Fudan University.

## References

- [1] S. W. Hawking, G. F. R. Ellis, *The Large Scale Structure of the Space-time*, Cambridge Univ. Press 1999



- [2] R. G. Cai, A. Z. Wang, *Black hole formation from collapsing dust fluid in a background of dark energy*, astro-ph/0505136
- [3] J. M. M. Senovilla, *Singularity Theorems in General Relativity: Achievements and Open Questions*, physics/0605007
- [4] H. Andréasson, M. K. Fachbereich, G. Rein, *Global existence for the spherically symmetric Einstein-Vlasov system with outgoing matter*, gr-qc/0611115
- [5] M. R. Mbonye, D. Kazanas, *Can gravitational collapse sustain singularity-free trapped surfaces?*, arXiv:0708.2360
- [6] C. Barceló, S. Liberati, S. Sonego, M. Visser, *Fate of gravitational collapse in semiclassical gravity*, arXiv:0712.1130
- [7] Y. Q. Gu, *Structure of the Star with Ideal Gases*, arXiv:0712.0219
- [8] Y. Q. Gu, *A Canonical Form of the Metric in General Relativity*, arXiv:0710.5792
- [9] Y. Q. Gu, *Exact Vacuum Solutions to the Einstein Equation*, Chin. Ann. Math. 28B(5), (2007)499-506, arXiv:0706.0318
- [10] S. L. Weinberg, *Gravitation and Cosmology*, Wiley, New York, 1972
- [11] Y. Q. Gu, *New Approach to N-body Relativistic Quantum Mechanics*, IJMPA Vol.22(No.11) 2007-2019(2007), hep-th/0610153
- [12] Y. Q. Gu, *Mass-Energy Relation of the Nonlinear Spinor*, hep-th/0701030
- [13] Y. Q. Gu, *Potentials in Energy Momentum Tensor and the Equation of State*, arXiv:0711.1243
- [14] Y. Q. Gu, *Accelerating Expansion of the Universe with Nonlinear Spinors*, gr-qc/0612176
- [15] J. Navarro, J. B. Sancho, *On the naturalness of Einsteins equation*, arXiv:0709.1928