

Analytic Continuation of q -Euler numbers and polynomials

Taekyun Kim¹

¹ School of Electrical Engineering and Computer Science ,
Kyungpook National University, Taegu 702-701, S. Korea
e-mail: tkim@knu.ac.kr

Abstract In this paper we study that the q -Euler numbers and polynomials are analytically continued to $E_q(s)$. A new formula for the Euler's q -Zeta function $\zeta_{E,q}(s)$ in terms of nested series of $\zeta_{E,q}(n)$ is derived. Finally we introduce the new concept of the dynamics of analytically continued q -Euler numbers and polynomials.

2000 Mathematics Subject Classification - 11B68, 11S40

Key words- q -Bernoulli polynomial, q -Riemann Zeta function

1 Introduction

Throughout this paper, \mathbb{Z}, \mathbb{R} and \mathbb{C} will denote the ring of integers, the field of real numbers and the complex numbers, respectively.

When one talks of q -extension, q is variously considered as an indeterminate, a complex numbers or p -adic numbers. Throughout this paper, we will assume that $q \in \mathbb{C}$ with $|q| < 1$. The q -symbol $[x]_q$ denotes $[x]_q = \frac{1-q^x}{1-q}$, (see [1-16]).

In this paper we study that the q -Euler numbers and polynomials are analytically continued to $E_q(s)$. A new formula for the Euler's q -Zeta function $\zeta_{E,q}(s)$ in terms of nested series of $\zeta_{E,q}(n)$ is derived. Finally we introduce the new concept of the dynamics of analytically continued q -Euler numbers and polynomials.

2 Generating q -Euler polynomials and numbers

For $h \in \mathbb{Z}$, the q -Euler polynomials were defined as

$$\sum_{n=0}^{\infty} \frac{E_n(x, h|q)}{n!} t^n = [2]_q \sum_{n=0}^{\infty} (-1)^n q^{hn} e^{[n+x]_q t}, \quad (2.1)$$

for $x, q \in \mathbb{C}$, cf. [1,7]. In the special case $x = 0$, $E_n(0, h|q) = E_n(h|q)$ are called q -Euler numbers, cf. [1,2,3,4]. By (2.1), we easily see that

$$E_n(x, h|q) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^{l+h}} q^{lx}, \quad \text{cf. [7,8],} \quad (2.2)$$

where $\binom{n}{j}$ is binomial coefficient. From (2.1), we derive

$$E_{n,q}(x, h|q) = (q^x E(h|q) + [x]_q)^n$$

with the usual convention of replacing $E^n(h|q)$ by $E_n(h|q)$. In the case $h = 0$, $E_n(x, 0|q)$ will be symbolically written as $E_{n,q}(x)$. Let $G_q(x, t)$ be generating function of q -Euler polynomials as follows:

$$G_q(x, t) = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}. \quad (2.3)$$

Then we easily see that

$$G_q(x, t) = [2]_q \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t}. \quad (2.4)$$

For $x = 0$, $E_{n,q} = E_{n,q}(0)$ will be called q -Euler numbers.

From (2.3), (2.4), we easily derive the following: For $k(= \text{even})$ and $n \in \mathbb{Z}_+$, we have

$$E_{n,q}(k) - E_{n,q} = [2]_q \sum_{l=0}^{k-1} (-1)^l [l]_q^n. \quad (2.5)$$

For $k(= \text{odd})$ and $n \in \mathbb{Z}_+$, we have

$$E_{n,q}(k) + E_{n,q} = [2]_q \sum_{l=0}^{k-1} (-1)^l [l]_q^n. \quad (2.6)$$

By (2.4), we easily see that

$$E_{m,q}(x) = \sum_{l=0}^m \binom{m}{l} q^{xl} E_{l,q}[x]_q^{m-l}. \quad (2.7)$$

From (2.5), (2.6), and (2.7), we derive

$$[2]_q \sum_{l=0}^{k-1} (-1)^{l-1} [l]_q^n = (q^{kn} - 1) E_{n,q} + \sum_{l=0}^{k-1} \binom{n}{l} q^{kl} E_{l,q}[k]_q^{n-l}, \quad (2.8)$$

where $k(= \text{even}) \in \mathbb{N}$. For $k(= \text{odd})$ and $n \in \mathbb{Z}_+$, we have

$$[2]_q \sum_{l=0}^{k-1} (-1)^l [l]_q^n = (q^{kn} + 1) E_{n,q} + \sum_{l=0}^{k-1} \binom{n}{l} q^{kl} E_{l,q}[k]_q^{n-l}. \quad (2.9)$$

3 q -Euler zeta function

It was known that the Euler polynomials are defined as

$$\frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} \frac{E_n(x)}{n!} t^n, \quad |t| < \pi, \text{ cf. [1-16]}. \quad (3.1)$$

For $s \in \mathbb{C}, x \in \mathbb{R}$ with $0 \leq x < 1$, define

$$\zeta_E(s, x) = 2 \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+x)^s}, \text{ and } \zeta_E(s) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}. \quad (3.2)$$

By (3.1) and (3.2) we see that Euler numbers are related to the Euler zeta function as

$$\zeta_E(-n) = E_n, \quad \zeta_E(-n, x) = E_n(x).$$

For $s, q, h \in \mathbb{C}$ with $|q| < 1$, we define q -Euler zeta function as follows:

$$\zeta_{E,q}(s, x|h) = [2]_q \sum_{n=0}^{\infty} \frac{(-1)^n q^{nh}}{[n+x]_q^s}, \text{ and } \zeta_{E,q}(s|h) = [2]_q \sum_{n=1}^{\infty} \frac{(-1)^n q^{nh}}{[n]_q^s}. \quad (3.3)$$

For $k \in \mathbb{N}, h \in \mathbb{Z}$, we have

$$\zeta_{E,q}(-n|h) = E_n(h|q).$$

In the special case $h = 0$, $\zeta_{E,q}(s|0)$ will be written as $\zeta_{E,q}(s)$. For $s \in \mathbb{C}$, we note that

$$\zeta_{E,q}(s) = [2]_q \sum_{n=1}^{\infty} \frac{(-1)^n}{[n]_q^s}.$$

We now consider the function $E_q(s)$ as the analytic continuation of Euler numbers. All the q -Euler numbers $E_{n,q}$ agree with $E_q(n)$, the analytic continuation of Euler numbers evaluated at n ,

$$E_q(n) = E_{n,q} \text{ for } n \geq 0.$$

Ordinary Euler numbers are defined by

$$\frac{2}{e^t + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \quad |t| < \pi. \quad (3.4)$$

By (3.4), it is easy to see that

$$E_0 = 1, \text{ and } E_n = -\frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l, \quad n = 0, 1, 2, \dots$$

From (2.9) and (3.3), we can consider the q -extension of Euler numbers E_n as follows:

$$E_{0,q} = \frac{[2]_q}{2}, \text{ and } E_{n,q} = -\frac{1}{[2]_q^n} \sum_{l=0}^{n-1} \binom{n}{l} q^l E_{l,q}, \quad n = 1, 2, 3, \dots, \quad (3.5)$$

In fact, we can express $E'_q(s)$ in terms of $\zeta'_{E,q}(s)$, the derivative of $\zeta_{E,q}(s)$.

$$E_q(s) = \zeta_{E,q}(-s), E'_q(s) = \zeta'_{E,q}(-s), E'_q(2n+1) = \zeta'_{E,q}(-2n-1), \quad (3.6)$$

for $n \in \mathbb{N} \cup \{0\}$. This is just the differential of the functional equation and so verifies the consistency of $E_q(s)$ and $E'_q(s)$ with $E_{n,q}$ and $\zeta(s)$.

From the above analytic continuation of q -Euler numbers, we derive

$$\begin{aligned} E_q(s) &= \zeta_{E,q}(-s), E_q(-s) = \zeta_{E,q}(s) \\ \Rightarrow E_{-n,q} &= E_q(-n) = \zeta_{E,q}(n), n \in \mathbb{Z}_+. \end{aligned} \quad (3.7)$$

The curve $E_q(s)$ runs through the points $E_{-n,q}$ and grows $\sim n$ asymptotically as $(-n) \rightarrow -\infty$. The curve $E_q(s)$ runs through the point $E_q(-n)$ and $\lim_{n \rightarrow \infty} E_q(-n) = \lim_{n \rightarrow \infty} \zeta_{E,q}(n) = -2$. From (3.5), (3.6) and (3.7), we note that

$$\zeta_{E,q}(-n) = E_q(n) \mapsto \zeta_{E,q}(-s) = E_q(s).$$

4 Analytic continuation of q -Euler polynomials

For consistency with the redefinition of $E_{n,q} = E_q(n)$ in (4.5) and (4.6), we have

$$E_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} E_{k,q} q^{kx} [x]_q^{n-k}.$$

Let $\Gamma(s)$ be the gamma function. Then the analytic continuation can be obtained as

$$\begin{aligned} n &\mapsto s \in \mathbb{R}, x \mapsto w \in \mathbb{C}, \\ E_{k,q} &\mapsto E_q(k+s-[s]) = \zeta_{E,q}(-(k+(s-[s]))) , \\ \binom{n}{k} &\mapsto \frac{\Gamma(1+s)}{\Gamma(1+k+(s-[s]))\Gamma(1+[s]-k)} \\ \Rightarrow E_{n,q}(s) &\mapsto E_q(s, w) = \sum_{k=-1}^{[s]} \frac{\Gamma(1+s)E_q(k+s-[s])q^{(k+s-[s])w} [w]_q^{[s]-k}}{\Gamma(1+k+(s-[s]))\Gamma(1+[s]-k)} \\ &= \sum_{k=0}^{[s]+1} \frac{\Gamma(1+s)E_q((k-1)+s-[s])q^{((k-1)+s-[s])w} [w]_q^{[s]+1-k}}{\Gamma(k+(s-[s]))\Gamma(2+[s]-k)}, \end{aligned}$$

where $[s]$ gives the integer part of s , and so $s - [s]$ gives the fractional part.

Deformation of the curve $E_q(2, w)$ into the curve of $E_q(3, w)$ via the real analytic continuation $E_q(s, w)$, $2 \leq s \leq 3$, $-0.5 \leq w \leq 0.5$.

ACKNOWLEDGEMENTS. This paper is supported by Jangjeon Mathematical Society and Jangjeon Research Institute for Mathematical Science and Physics(2007-001-JRIMS 1234567)

References

- [1] M. Cenkci, M. Can, *Some results on q -analogue of the Lerch zeta function*, Advan. Stud. Contemp. Math., Vol 12(2006), 213-223.
- [2] M. Cenkci, *The p -adic generalized twisted (h, q) -Euler- l -function and its applications*, Advan. Stud. Contemp. Math., Vol 15(2007), 37-47
- [3] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integrals*, J. Nonlinear Math. Phys., Vol 14(2007), 15-27.
- [4] T. Kim, *On p -adic interpolating function for q -Euler numbers and its derivatives*, J. Math. Anal. Appl., Vol 339(2008), 598-608.
- [5] T. Kim, *A Note on p -Adic q -integral on Z_p Associated with q -Euler Numbers*, Advan. Stud. Contemp. Math., Vol 15(2007), 133-137.
- [6] T. Kim, *On p -adic q - l -functions and sums of powers*, J. Math. Anal. Appl., Vol 329(2007), 1472-1481.
- [7] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys., Vol 9 (2002), 288-299.
- [8] T. Kim, *A note on some formulas for the q -Euler numbers and polynomials*, Proc. Jangjeon Math. Soc., Vol 9(2006), 227-232.
- [9] T. Kim, *Multiple p -adic L -function*, Russ. J. Math. Phys., Vol 13(2006), 151-157.
- [10] T. Kim, *On explicit formulas of p -adic q - L -functions*, Kyushu J. Math., Vol 43(1994), 73-86.
- [11] A. Kudo, *A congruence of generalized Bernoulli number for the character of the first kind*, Advan. Stud. Contemp. Math., Vol 2(2000), 1-8.
- [12] Q.-M. Luo, F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Advan. Stud. Contemp. Math., Vol 7(2003), 11-18.
- [13] Q.-M. Luo, *Some recursion formulae and relations for Bernoulli numbers and Euler numbers of higher order*, Advan. Stud. Contemp. Math., Vol 10 (2005), 63-70.
- [14] H. Ozden, Y. Simsek, S.H. Rim, I. Cangul, *A note on p -adic q -Euler measure*, Advan. Stud. Contemp. Math., Vol 14(2007), 233-239.
- [15] Y. Simsek, *Theorem on twisted L -function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math., Vol 12(2006), 237-246.
- [16] C. S. Ryoo, T. Kim, R. P. Agarwal, *Exploring the multiple Changhee q -Bernoulli polynomials*, Inter. J. Comput. Math., Vol 82(2005), 483-493.