

# Borg type uniqueness Theorems for periodic Jacobi operators with matrix valued coefficients

Evgeny Korotyaev <sup>\*</sup>      Anton Kutsenko <sup>†</sup>

November 21, 2018

## Abstract

We give a simple proof of Borg type uniqueness Theorems for periodic Jacobi operators with matrix valued coefficients.

## 1 Introduction

Consider a self-adjoint matrix-valued Jacobi operator  $J$  acting on  $\ell^2(\mathbb{Z})^m$  and given by

$$(Jy)_n = a_n y_{n+1} + b_n y_n + a_{n-1} y_{n-1}, \quad n \in \mathbb{Z}, \quad y_n \in \mathbb{C}^m, y = (y_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z})^m, m \geq 1, \quad (1.1)$$

where  $a_n > 0, b_n = b_n^*, n \in \mathbb{Z}$  are  $p$ -periodic sequences of the complex  $m \times m$  matrices. It is well known that the spectrum  $\sigma(J)$  of  $J$  is absolutely continuous and consists of non-degenerated intervals  $[\lambda_{n-1}^+, \lambda_n^-], \lambda_{n-1}^+ < \lambda_n^- \leq \lambda_n^+, n = 1, \dots, N < \infty$ . These intervals are separated by the gaps  $\gamma_n = (\lambda_n^-, \lambda_n^+), n = 1, \dots, N - 1$  with the length  $> 0$ . Introduce the fundamental  $m \times m$  matrix-valued solutions  $\varphi = (\varphi_n(z))_{n \in \mathbb{Z}}, \vartheta = (\vartheta_n(z))_{n \in \mathbb{Z}}$  of the equation

$$a_n y_{n+1} + b_n y_n + a_{n-1}^* y_{n-1} = z y_n, \quad \varphi_0 \equiv \vartheta_1 \equiv 0, \quad \varphi_1 \equiv \vartheta_0 \equiv I_m, \quad (z, n) \in \mathbb{C} \times \mathbb{Z}, \quad (1.2)$$

where  $I_m$  is the identity  $m \times m$  matrix. We define the monodromy  $2m \times 2m$  matrix  $\mathcal{M}_p$  by

$$\mathcal{M}_p(z) = \begin{pmatrix} \vartheta_p(z) & \varphi_p(z) \\ \vartheta_{p+1}(z) & \varphi_{p+1}(z) \end{pmatrix}, \quad z \in \mathbb{C}. \quad (1.3)$$

Let  $\tau_1(z), \dots, \tau_{2m}(z)$  be eigenvalues of  $\mathcal{M}_p(z)$ . Recall that  $\sigma(J) = \bigcup_{j=1}^{2m} \{z \in \mathbb{C} : |\tau_j(z)| = 1\}$ , see [KKu]. Let  $J^0$  be the unperturbed Jacobi matrix with  $a_n^0 = I_m, b_n^0 = 0$ . Note, that  $\sigma(J^0) = \sigma_{ac}(J^0) = [-2, 2]$ . Let  $\mathbb{N}_s = \{1, \dots, s\}$  and let  $c = \det \prod_{n=1}^p a_n$ . We formulate first Borg type uniqueness theorem

---

<sup>\*</sup>Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

<sup>†</sup>Department of Mathematics of Sankt-Petersburg State University, Russia e-mail: kucenko@rambler.u

**Theorem 1.1.** *Let  $c = 1$  and let  $\sigma(J) = [\lambda_0^+, \lambda_N^-] = \{\text{each } |\tau_j(z)| = 1, (z, j) \in [\lambda_0^+, \lambda_N^-] \times \mathbb{N}_{2m}\}$ , where  $\lambda_0^+ = -\lambda_N^-$ . Then  $J = J^0$ .*

This theorem was proved in [CGR]. We give a simple proof based on the trace formula and the properties of the Chebyshev polynomials proved in Lemma 2.3. There is an enormous literature on inverse spectral problems for scalar (i.e.,  $m = 1$ ) periodic Jacobi matrices (see [BGGK], [K],[KKu1], [KKu2], [vM], [T], book [T] and references therein), but very little for matrix-valued periodic Jacobi operators (see [CGR] and references therein). Note that the complete solution of inverse problem for finite matrix-valued Jacobi operators was obtained recently [BCK].

The operator  $J$  is unitarily equivalent to the operator  $\mathcal{J} = \int_{[0,2\pi]}^{\oplus} K_p(e^{ix}) \frac{dx}{2\pi}$  acting in  $\int_{[0,2\pi]}^{\oplus} \mathcal{H} \frac{dx}{2\pi}$ , where  $\mathcal{H} = \mathbb{C}^p$  and  $pm \times mp$  matrix  $K_p(\tau)$  is given by

$$K_p(\tau) = \begin{pmatrix} b_1 & a_1 & 0 & \dots & \tau^{-1}a_p \\ a_1 & b_2 & a_2 & \dots & 0 \\ 0 & a_2 & b_3 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ \tau a_p & 0 & \dots & a_{p-1} & b_p \end{pmatrix}, \quad \tau \in \mathbb{S}^1 = \{\tau : |\tau| = 1\}, \quad (1.4)$$

see [KKu]. Let  $(\lambda_n(\tau))_{n=1}^{pm}$  be eigenvalues of  $K_p(\tau)$ . We formulate our main result.

**Theorem 1.2.** *Let  $a_n > 0, b_n = b_n^*$  be real  $m \times m$  matrices for all  $n \in \mathbb{Z}$ .*

*i) Let  $c = 1$ . Then  $\sum_{n=1}^{mp} \lambda_n^2(\tau) = 2pm$  for some  $\tau \in \mathbb{S}^1$  iff  $J = J^0$ .*

*ii) Let  $c = 1$  and let  $\varkappa_1 \in \mathbb{R}$ . Then eigenvalues  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$  for all  $s \in \mathbb{N}_{mp}$  iff  $J = J^0$ .*

*iii) Let  $\varkappa_1, \varkappa_2 \in \mathbb{R}$  and  $\cos \varkappa_1 \neq \cos \varkappa_2$ . Let eigenvalues  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$  for all  $s \in \mathbb{N}_{mp}$  and  $\lambda_n(e^{i\varkappa_2}) = 2 \cos \frac{\varkappa_2 + 2\pi n_1}{p}, n \in \mathbb{N}_m$ , for some  $n_1 \in \mathbb{N}_p$ , and if  $e^{i2\varkappa_2} = 1$ , then additional eigenvalues  $\lambda_{n+m}(e^{i\varkappa_2}) = \lambda_1(e^{i\varkappa_2}) \neq \pm 2, n \in \mathbb{N}_m$ . It is possible iff  $J = J^0$ .*

**Remark.** 1) Note that the condition  $\cos \frac{\varkappa_2 + 2\pi n_1}{p} \neq \pm 1$  in iii) is associated with the unperturbed operator  $J^0$ , where the endpoints of the spectrum  $\sigma(J^0) = [-2, 2]$  have the multiplicity  $m$ , as the zeros of the determinant  $D_p(z, \pm 1)$ . Each point from  $(-2, 2)$  has the multiplicity  $2m$  as the zero of the determinant  $D_p(z, \tau), \tau \neq \pm 1$ .

2) Consider the case  $m = 2$  and  $\varkappa_1 = 0, \varkappa_2 = \pi$ . Let periodic eigenvalues  $\lambda_s(1) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$  for all  $s \in \mathbb{N}_{2p}$  and let anti-periodic eigenvalues  $\lambda_s(-1) = 2 \cos \frac{\pi(2n+1)}{p} \neq \pm 2, s \in \mathbb{N}_4$  for some  $n \in \mathbb{N}_p$ . Then we deduce that  $J = J^0$ .

3) Consider a self-adjoint matrix-valued Jacobi operator  $\tilde{J}$  acting on  $\ell^2(\mathbb{Z})^m$  and given by

$$(\tilde{J}y)_n = \tilde{a}_n y_{n+1} + \tilde{b}_n y_n + \tilde{a}_{n-1}^* y_{n-1}, \quad n \in \mathbb{Z}, \quad y_n \in \mathbb{C}^m, y = (y_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z})^m, m \geq 1,$$

where  $\tilde{a}_n$  ( $\det \tilde{a}_n \neq 0$ ),  $\tilde{b}_n = \tilde{b}_n^*, n \in \mathbb{Z}$  are  $p$ -periodic sequences of the complex  $m \times m$  matrices. Then the operator  $\tilde{J} = UJU^*$  where  $J$  is given by (1.2) and  $U = \text{diag}_{n \in \mathbb{Z}} u_n$  and the unitary matrices  $u_n$  have the forms  $u_0 = I_m; u_{n+1} = c_n^* u_n, n \geq 0; u_{n-1} = c_{n-1} u_{n+1}, n \leq 0$ . Here  $\tilde{a}_n = c_n a_n, a_n > 0$  and  $c_n$  are the unitary matrices.

## 2 Proof

We need the following results from [KKU].

**Lemma 2.1.** *The following identities and estimate are fulfilled:*

$$\sum_{n=1}^{mp} \lambda_{n,p}(\tau) = \sum_{n=1}^p \text{Tr } b_n, \quad \sum_{n=1}^{mp} \lambda_{n,p}^2(\tau) = \sum_{n=1}^p \text{Tr}(b_n^2 + 2a_n^2), \quad \text{all } \tau \in \mathbb{C} \setminus \{0\}, \quad (2.1)$$

$$\sum_{n=1}^{pm} \lambda_{n,p}^2(\tau) \geq 2pmc^{\frac{2}{pm}}, \quad \tau \in \mathbb{S}^1, \quad (2.2)$$

where the identity (2.2) holds true iff  $J = c^{\frac{1}{pm}} J^0$ , and here  $c = \det \prod_1^p a_n$ .

Recall the following identities for  $D_p(z, \tau) = \det(\mathcal{M}_p(z) - \tau I_{2m})$  from [KKU]

$$D_p(z, \tau) = \prod_{j=1}^{2m} (\tau_j(z) - \tau) = c(-\tau)^m \prod_{n=1}^{mp} (z - \lambda_n(\tau)) = c(-\tau)^m \det(z - K_p(\tau)), \quad (2.3)$$

where  $z, \tau \in \mathbb{C}$ ,  $\tau \neq 0$ .

**Proof of Theorem 1.1 Consider the first case**  $\sum_{n=1}^p \text{Tr } b_n = 0$ . Consider the  $p$ -periodic operator  $J$  as  $pk$ -periodic operator for some  $(k, \tau) \in \mathbb{N} \times \mathbb{S}^1$ . Let the eigenvalues of  $K_{pk}(\tau)$  be given by

$$\lambda_-(\tau) \equiv \lambda_{1,pk}(\tau) \leq \lambda_{2,pk}(\tau) \leq \dots \leq \lambda_+(\tau) \equiv \lambda_{pmk,pk}(\tau). \quad (2.4)$$

The eigenvalues  $\lambda_{\pm}(\tau)$  belong to  $\sigma(J)$ , then we have that  $[\lambda_-(\tau), \lambda_+(\tau)] \subset \sigma(J)$ . Using  $|\tau_j(z)| = 1$  for all  $(z, j) \in [\lambda_-(\tau), \lambda_+(\tau)] \times \mathbb{N}_{2m}$ , and (2.3), we obtain

$$|D_{pk}(z, \tau)| = \left| \prod_{j=1}^{2m} (\tau - \tau_j^k(z)) \right| \leq \prod_{j=1}^{2m} (|\tau| + |\tau_j^k(z)|) \leq 2^m, \quad \text{all } z \in [\lambda_-(\tau), \lambda_+(\tau)].$$

We have  $\det \prod_{n=1}^{pk} a_n = c^k = 1$ , since a sequence  $a_n$  is  $p$ -periodic. Then (2.3) gives

$$|D_{pk}(z, \tau)| = \left| \prod_{n=1}^{pmk} (z - \lambda_{n,pk}(\tau)) \right| \leq 2^m, \quad \text{all } z \in [\lambda_-(\tau), \lambda_+(\tau)]. \quad (2.5)$$

A sequence  $b_n$  is  $p$ -periodic, then

$$\sum_{n=1}^{pmk} \lambda_{n,pk}(\tau) = \sum_{n=1}^{pk} \text{Tr } b_n = k \sum_{n=1}^p \text{Tr } b_n = 0. \quad (2.6)$$

The relations (2.4), (2.5), (2.6) and Lemma 2.3 give that

$$(\lambda_{n,pk})_{n=1}^{mpk} \in \mathcal{P}_{mpk}(2^m), \quad \sum_{n=1}^{pmk} \lambda_{n,pk}^2(\tau) \leq 2pmkC_k, \quad C_k = 2^{\frac{2(m-1)}{pmk}}. \quad (2.7)$$

Then using (2.1) and  $a_n = a_{n+p}, b_n = b_{n+p}$  for all  $n \in \mathbb{Z}$ , we obtain

$$\sum_{n=1}^{pmk} \lambda_{n,pk}^2(\tau) = \sum_{n=1}^{pk} \text{Tr}(b_n^2 + a_n^2) = k \sum_{n=1}^p \text{Tr}(b_n^2 + a_n^2) = k \sum_{n=1}^{pm} \lambda_{n,p}^2 \leq k(2pm)C_k,$$

which yields  $\sum_{n=1}^p \lambda_{n,p}^2 \leq 2pm$ , since we take  $k \rightarrow +\infty$ . Thus Lemma 2.1 implies  $J = J^0$ .

**Consider the second case**  $\beta = \sum_{n=1}^p \text{Tr} b_n$ . Then a new operator  $J^1 = (J - \beta)$  satisfies  $\sum_{n=1}^p \text{Tr}(b_n - \beta I_m) = 0$ . Thus due to the first case we deduce that

$$\sigma(J) = \sigma(J^1) = [-2, 2] = [-x - \beta, x - \beta], \quad a_n = I_m, \quad b_n - \beta I_m = b_n^1 = 0$$

for  $x = \lambda_n^-$ , which yields  $x = 2, \beta = 0$  and the theorem has been proved. ■

**Lemma 2.2.** *i) Let  $\lambda$  be an eigenvalue of  $K_p(\tau)$  and have multiplicity  $m$  for some  $\tau \in \mathbb{S}^1 \setminus \{-1, 1\}$ . Then the multipliers have the form  $\tau_j(\lambda) = \tau, \tau_{j+m}(\lambda) = \tau^{-1}$  for all  $j \in \mathbb{N}_m$ .  
ii) Let  $\lambda$  be an eigenvalue of  $K_p(\tau)$  and have multiplicity  $2m$  for some  $\tau \in \{-1, 1\}$ . Then each  $\tau_j(\lambda) = \tau, j \in \mathbb{N}_{2m}$ .*

**Proof.** i) The matrix  $K_p(\tau)$  is self-adjoint. Let  $K_p(\tau)f^k = \lambda f^k$  for some orthogonal eigenvectors  $f^k = (f_n^k)_{n=1}^{mp}, k \in \mathbb{N}_m$ . If  $f_0^k = \tau^{-1} f_p^k$ , then the definition of the matrix  $\mathcal{M}_p$  gives  $\mathcal{M}_p(\lambda)(f_0^k, f_1^k)^\top = \tau(f_0^k, f_1^k)^\top$ . Note that  $\tilde{f}^k = (f_0^k, f_1^k)^\top$  define other components of vector  $f^k$ , since  $L(\tau)$  has special form, see (1.4). Then the vectors  $\tilde{f}^k, k \in \mathbb{N}_m$  are linearly independent vectors, since  $f^k$  are linearly independent vectors. Then  $\tau$  has multiplicity at least  $m$ . The matrix  $\mathcal{M}_p$  is symplectic, then  $\tau^{-1}$  is eigenvalue of  $\mathcal{M}_p$  and  $\tau^{-1}$  has multiplicity at least  $m$ . We obtain first statement, since  $\mathcal{M}_p$  is  $2m \times 2m$  matrix. The proof of ii) is similar. ■

**Proof of Theorem 1.2.** The statement i) follows from Lemma 2.1.

ii) **Sufficiency.** Recall that if  $m = 1$ , then  $\det(M_p^0 - \tau I_2) = \tau(\tau + \tau^{-1} - 2\mathcal{T}_p(z/2))$ , where  $\mathcal{T}_p(z) = \cos(p \arccos z)$  is the Chebyshev polynomial. Moreover, zeros of the polynomial  $\mathcal{T}_p(z) - \cos \varkappa_1$  are given by  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}, s \in \mathbb{N}_p$ .

Thus if  $J = J^0, m \geq 2$ , then corresponding monodromy operators  $M_p$  satisfies  $\det(M_p - \tau I_{2m}) = \tau^m \prod_{j=1}^m (\tau + \tau^{-1} - 2\mathcal{T}_p(z/2))$ , which yields at  $\tau = e^{i\varkappa_1}$

$$\det(M_p - e^{i\varkappa_1} I_{2m}) = \tau = e^{im\varkappa_1} 2^m (\cos \varkappa_1 - \mathcal{T}_p(z/2))^m = e^{im\varkappa_1} 2^m, \quad (2.8)$$

This implies  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}, s \in \mathbb{N}_{mp}$ .

**Necessity.** Let  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}, s \in \mathbb{N}_{mp}$ . Then the direct calculation implies  $\sum_{n=1}^{mp} \lambda_n^2(e^{i\varkappa_1}) = 2pm$ , and the statement i) gives  $J = J^0$ .

iii) **The sufficiency** is proved similar to the case ii).

**Necessity.** We consider only the case  $e^{2i\varkappa_2} \neq 1$ , the proof of the case  $e^{2i\varkappa_2} = 1$  is similar. Let  $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}, s \in \mathbb{N}_{mp}$ . Using (2.3), (2.8), we obtain

$$\prod_{j=1}^{2m} (e^{i\varkappa_1} - \tau_j(z)) = c(-1)^m e^{im\varkappa_1} \prod_{n=1}^{mp} (z - \lambda_n(e^{i\varkappa_1})) = c(-2)^m e^{im\varkappa_1} (\mathcal{T}_p(z/2) - \cos \varkappa_1)^m, \quad (2.9)$$

where  $c = \det \prod_{n=1}^p a_n$ . The eigenvalue  $\lambda_1(e^{i\kappa_2})$  has the multiplicity  $m$ , then using Lemma 2.2 and substituting  $z = \lambda_1(e^{i\kappa_2}) = 2 \cos \frac{\kappa_2 + 2\pi n_1}{p}$  and two multipliers  $e^{\pm i\kappa_2}$  (given by Lemma 2.2) into (2.9), we obtain

$$(e^{i\kappa_1} - e^{i\kappa_2})^m (e^{i\kappa_1} - e^{-i\kappa_2})^m = c 2^m e^{im\kappa_1} (\cos \kappa_2 - \cos \kappa_1)^m,$$

which yields  $c = 1$ . Then the statement ii) gives  $J = J^0$ . ■

**Lemma 2.3.** *For any  $r \geq 0$ ,  $s \geq 2$  the following identity holds true*

$$\sup_{x \in \mathcal{P}_s(r)} \sum_1^s x_n^2 = 2s \left(\frac{r}{2}\right)^{\frac{2}{s}},$$

$$\mathcal{P}_s(r) = \left\{ (x_n)_1^s : x_1 \leq \dots \leq x_s, \sum_{n=1}^s x_n = 0, \left| \prod_{n=1}^s (z - x_n) \right| \leq r, \text{ all } z \in [x_1, x_s] \right\} \subset \mathbb{R}^s.$$

**Proof.** In the proof we use arguments from [Ku]. Let  $\|x\|^2 = \sum x_n^2$ ,  $x = (x_n)_1^s \in \mathbb{R}^s$ . The set  $\mathcal{P}_s(r)$  is compact, then  $\sup_{x \in \mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2$  for some  $x^0 = (x_n^0)_1^s \in \mathcal{P}_s(r)$ . Introduce a polynomial  $p_0(z) = \prod_{n=1}^s (z - x_n^0)$ . Let  $x_0^0 = -\infty$ ,  $x_{s+1}^0 = +\infty$ . The polynomial  $p_0'$  has only real zeros  $\tilde{x}_n$ ,  $n \in \mathbb{N}_{s-1}$ . We will show that each  $|p_0(\tilde{x}_n)| = r$ ,  $n \in \mathbb{N}_{s-1}$ . Assume that there exist  $1 \leq n_1 < n_2 \leq s$  such that

$$x_{n_1-1}^0 < x_{n_1}^0 \leq x_{n_2}^0 < x_{n_2+1}^0, \quad \max_{z \in [x_{n_1}, x_{n_2}]} |p_0(z)| < r. \quad (2.10)$$

Introduce a polynomial  $p_\varepsilon(z) = \prod_{n=1}^s (z - x_n^\varepsilon)$ , where a vector  $x^\varepsilon = (x_n^\varepsilon)_{n=1}^s \in \mathbb{R}^s$  is given by

$$x_n^\varepsilon = x_n^0, \quad n \neq n_1, \quad n \neq n_2; \quad x_{n_1}^\varepsilon = x_{n_1}^0 - \varepsilon, \quad x_{n_2}^\varepsilon = x_{n_2}^0 + \varepsilon, \quad \varepsilon > 0. \quad (2.11)$$

Using  $x^0 \in \mathcal{P}_s(r)$  and (2.10), we obtain

$$x_1^\varepsilon \leq \dots \leq x_s^\varepsilon, \quad \sum_{n=1}^s x_n^\varepsilon = \sum_{n=1}^s x_n^0 = 0, \quad \max_{z \in [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]} |p_\varepsilon(z)| \leq r, \quad (2.12)$$

for sufficiently small  $\varepsilon > 0$ . We rewrite  $p_\varepsilon$  in the form

$$p_\varepsilon(z) = p_0(z) g_\varepsilon(z), \quad g_\varepsilon(z) = \frac{(z - x_{n_1}^\varepsilon)(z - x_{n_2}^\varepsilon)}{(z - x_{n_1}^0)(z - x_{n_2}^0)} = \frac{(z - x_{n_1}^0 + \varepsilon)(z - x_{n_2}^0 - \varepsilon)}{(z - x_{n_1}^0)(z - x_{n_2}^0)}. \quad (2.13)$$

Due to  $x_{n_1}^0 \leq x_{n_2}^0$  and  $\varepsilon > 0$  we deduce that  $|g_\varepsilon(z)| \leq 1$ ,  $z \in \mathbb{R} \setminus [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]$ . Then (2.13) yields  $|p_\varepsilon(z)| \leq |p_0(z)|$ ,  $z \in \mathbb{R} \setminus [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]$  and thus (2.12) gives  $|p_\varepsilon(z)| \leq r$ ,  $z \in [x_1^\varepsilon, x_s^\varepsilon]$ , and  $x^\varepsilon \in \mathcal{P}_s(r)$  for all sufficiently small  $\varepsilon > 0$ . There is an estimate

$$\|x^\varepsilon\|^2 = \sum_{n=1}^s (x_n^\varepsilon)^2 = \sum_{n=1}^s (x_n^0)^2 + 4\varepsilon(x_{n_2}^0 - x_{n_1}^0) + 2\varepsilon^2 > \|x^0\|^2, \quad (2.14)$$

since  $x_{n_1}^0 \leq x_{n_2}^0$  and  $\varepsilon > 0$ . But (2.14) and the condition  $x^\varepsilon \in \mathcal{P}_s(r)$  contradict the identity  $\sup_{x \in \mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2$ . Then the assumption (2.10) is not true and each  $|p_0(\tilde{x}_n)| = r, n \in \mathbb{N}_{s-1}$ . Note that only polynomials  $r\mathcal{T}_s(\alpha z + \beta)$ ,  $\alpha \in \mathbb{R} \setminus \{0\}$ ,  $\beta \in \mathbb{R}$ , (here  $\mathcal{T}_s$  are the Chebyshev polynomials, i.e.  $\mathcal{T}_s(\cos z) = \cos sz$ ) have this property, then  $p_0(z) = r\mathcal{T}_s(\alpha z + \beta)$  for some  $\alpha \in \mathbb{R} \setminus \{0\}$ ,  $\beta \in \mathbb{R}$ . We take  $\alpha, \beta$  such that  $p_0(z) = \prod_{n=1}^s (z - x_n^0)$ ,  $\sum_{n=1}^s x_n^0 = 0$  and then  $p_0(z) = r\mathcal{T}_s(zr^{-\frac{1}{s}}2^{\frac{1-s}{s}})$ , since (see [AS])

$$\mathcal{T}_s(z) = \frac{1}{2} \sum_{k=0}^{\lfloor \frac{s}{2} \rfloor} (-1)^k \frac{s}{s-k} C_k^{s-k} (2z)^{s-2k} = 2^{s-1} z^s - 2^{s-3} s z^{s-2} + o(z^{s-2}) \quad \text{as } z \rightarrow \infty,$$

where  $C_m^n = \frac{n!}{m!(n-m)!}$ . Then using Viète formulas we get

$$\mathcal{T}_s(z) = 2^{s-1} \prod_{n=1}^s (z - z_n) = 2^{s-1} \left( z^s - \xi z^{s-1} + \frac{1}{2} (\xi^2 - \eta) z^{s-2} + o(z^{s-2}) \right) \quad \text{as } z \rightarrow \infty,$$

where  $z_n, n \in \mathbb{N}_s$  are zeroes of  $\mathcal{T}_s$  and  $\xi = \sum_1^s z_n, \eta = \sum_1^s z_n^2$ . Then we get  $\xi = 0$  and  $\eta = \frac{s}{2}$ . This gives

$$\sup_{x \in \mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2 = r^{\frac{2}{s}} 2^{\frac{2s-2}{s}} \sum_1^s z_n^2 = r^{\frac{2}{s}} 2^{\frac{2s-2}{s}} \eta = 2s \left( \frac{r}{2} \right)^{\frac{2}{s}} \blacksquare$$

**Acknowledgments.** Evgeny Korotyaev was partly supported by DFG project BR691/23-1. The some part of this paper was written at the Math. Institute of Humboldt Univ., Berlin; Anton Kutsenko is grateful to the Institute for the hospitality.

## References

- [AS] Abramowitz, M.; Stegun, A., eds. Handbook of Mathematical Functions. N.Y.: Dover Publications Inc., 1992.
- [BGGK] Bättig, D.; Grebert, B.; Guillot, J.-C.; Kappeler, T. Fibration of the phase space of the periodic Toda lattice. J. Math. Pures Appl. 72 (1993), no. 6, 553–565.
- [BCK] Brüning, J.; Chelkak, D.; Korotyaev, E. Inverse spectral analysis for finite matrix-valued Jacobi operators, preprint 2006.
- [CG] Clark, S.; Gesztesy, F. On Weyl–Titchmarsh theory for singular finite difference Hamiltonian systems, J. Comput. Appl. Math., 171 (2004), 151–184.
- [CGR] Clark, S.; Gesztesy, F.; Renger, W. Trace formulas and Borg-type theorems for matrix-valued Jacobi and Dirac finite difference operators. J. Differential Equations 219 (2005), no. 1, 144–182.
- [GKM] Gesztesy, F.; Kiselev, A.; Makarov, K., A. Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators, Math. Nachr. 239–240 (2002), 103–145.

- [K] Korotyaev, E. Gap-length mapping for periodic Jacobi matrices, *Russ. J. Math. Phys.* 13(2006), no.1, 64-69.
- [KKr] Korotyaev, E.; Krasovsky, I. Spectral estimates for periodic Jacobi matrices, *Commun. Math. Phys.* 234 (2003), 517-532.
- [KKu] Korotyaev, E.; Kutsenko, A. Lyapunov functions for periodic matrix-valued Jacobi operators, preprint 2007, [www.arxiv.org](http://www.arxiv.org).
- [KKu1] Korotyaev, E.; Kutsenko, A. Inverse problem for the discrete 1D Schrödinger operator with small periodic potentials, *Commun. Math. Phys.* 261(2006), 673-692.
- [KKu2] Korotyaev, E.; Kutsenko, A. Marchenko-Ostrovski mappings for periodic Jacobi matrices, will be published in *Russ. J. Math. Phys.*
- [Ku] Kutsenko, A. Estimates of parameters for conformal mappings related to a periodic Jacobi matrix. *Journal of Mathematical Sciences*, 134 (2006), no. 4, 2295–2304.
- [vM] van Moerbeke, P. The spectrum of Jacobi matrices. *Invent. Math.* 37 (1976), no. 1, 45–81.
- [T] Teschl, G. *Jacobi Operators and Completely Integrable Nonlinear Lattices*, Mathematical Survey Monographs, vol. 72, American Mathematical Society, Providence, RI, 2000.