Density-density correlations of the ultra-cold Fermi gas in an optical lattice

T. Paananen,^{1,*} T. K. Koponen,² P. Törmä,^{2,3} and J.-P. Martikainen¹

 1 Department of Physical Sciences, University of Helsinki, PO Box 64, 00014 University of Helsinki, Finland ²Nanoscience Center, Department of Physics, PO Box 35, 40014 University of Jyväskylä, Finland

³Laboratory of Physics, Helsinki University of Technology,

P.O. Box 5100, 02015 HUT, Finland

(Dated: May 28, 2019)

Abstract

In this paper we study the density-density correlations of the two component Fermi gas in optical lattices. Three different type of phases, the BCS-state (Bardeen, Cooper, and Schieffer), the FFLO-state (Fulde, Ferrel, Larkin, and Ovchinnikov), and BP (breach pair) state, are considered. We show how these states differ in their density-density correlations. The density-density correlations are calculated not only at zero temperature, but also at non-zero temperatures paying particular attention to how much the finite temperature effects might complicate the detection of different phases. Since one-dimensional systems have been shown to be very promising candidates to observe FFLO states, we apply our results also to the computation of correlation signals in a one-dimensional lattice. We find that the density-density correlations reveal important information about the structure of the underlying order parameter as well as about the quasiparticle dispersions.

[∗]Electronic address: Tomi.Paananen@helsinki.fi

I. INTRODUCTION

Recent studies in the experiments on ultracold Fermi gases have shown great potential for elucidating long-standing problems in many different fields of physics related to strongly correlated fermions. For instance, in recent experiments $\begin{bmatrix} 1, 2, 3, 4, 5 \end{bmatrix}$ spin-density imbalanced, or polarized, Fermi gases were considered. Among other things, such systems make it possible to study pairing with mismatched Fermi surfaces, potentially leading to non-BCS type pairing such as that appearing in FFLO-states [\[6,](#page-22-5) [7](#page-22-6)] or BP-states [\[8,](#page-22-7) [9](#page-22-8)] (Sarma-states). These possibilities have been considered extensively in condensed-matter, nuclear, and highenergy physics $|10|$.

In optical lattices it is possible to study many different physical problems with close analogs in the field of solid state physics. However, in contrast to many solid state systems, ultra-cold gases in optical lattices provide a very clean environment. In other words, these systems have very few imperfections and if imperfections are of interest, they can be introduced in a more controlled manner. In addition, since optical lattices are made with lasers, the lattice geometry is easy to vary [\[11,](#page-22-10) [12,](#page-22-11) [13](#page-22-12), [14\]](#page-22-13) by changing the properties of the intersecting laser beams. For these reasons optical lattices enable one to investigate various quantum many-body physics problems, such as those related to Mott insulators, phase coherence, and superfluidity. Indeed, the possibility of a superfluid alkali atom Fermi gas in an optical lattice has been recently studied both theoretically [\[15,](#page-22-14) [16,](#page-22-15) [17,](#page-22-16) [18](#page-22-17), [19](#page-22-18), [20,](#page-22-19) [21\]](#page-22-20), as well as experimentally [\[22\]](#page-22-21). Furthermore, Feshbach resonances, molecules of fermionic atoms, and p-wave interactions in a lower dimensional fermionic systems have been studied using optical lattices [\[23](#page-22-22), [24](#page-22-23), [25](#page-22-24)].

Density-density correlations tell us how strongly the atomic densities at different positions are correlated. Often it is useful to focus on the deviation of the atomic density from its mean value and subtract the background away. In such cases one studies noise correlations i.e how the density fluctuations at different positions are correlated. Measuring noise correlations is a promising way to observe different phases in an optical lattice since while densities can be very similar for different phases, the noise correlations can still be very different. As an example, one can mention the famous phase transition between the superfluid Bose gas and the Mott insulator [\[26](#page-22-25), [27\]](#page-22-26). In the Mott insulator phase one can see the Bragg peaks in the noise correlations, but the noise correlations vanish for a Bose-Einstein condensate. Experimentally noise correlations have been used as an indicator for different phases in optical lattices for bosons [\[27\]](#page-22-26) and to observe fermionic anti-bunching for ideal fermions [\[28\]](#page-22-27). Also, noise correlations have been used to detect pairing correlations in an interacting Fermi gas in a harmonic trap [\[29](#page-23-0)].

In this paper we study noise correlations in a cloud of ultra-cold two component Fermi gas at finite temperatures. We are motivated by the fact that via noise correlations one can see subtle correlation effects which are not visible in the lower order correlations functions. Not only are the noise correlation signals very different for a superfluid and a normal Fermi gas, but noise correlations also differ between different types of paired states. While we compute density-density correlations also at zero temperature, special attention is payed to finite temperature calculations, since finite temperature effects have not been extensively discussed before. We find that although finite temperature effects do smooth out some sharp features present in zero temperature calculations, qualitative and clear quantitative differences between phases can still exist at higher temperatures. Also, we show how regions of gapless excitations are reflected in the density-density correlations. For instance, gapless excitations show up as absence of correlations for certain momenta. Moreover, they are also reflected in the value of the maximum correlation peak height, this quantitative signature persists at finites temperature as well.

Reducing the dimensionality of the system from three dimensions into one has been shown to favor the FFLO-type modulated order parameters in free space [\[30](#page-23-1), [31\]](#page-23-2) as well as in a lattice [\[32\]](#page-23-3). This effect of dimensionality, together with the general tendency of a lattice geometry to favour FFLO state due to nesting of the Fermi surfaces [\[21\]](#page-22-20), makes one dimensional optical lattices a promising system to study the FFLO state. For this reason, we apply our general results on density-density correlations also to the one-dimensional (1D) system. Very recently a similar one-dimensional problem was also discussed at zero temperature using the DMRG algorithm [\[33\]](#page-23-4). Here we compute the density-density correlations of the one-dimensional system at representative points of the finite temperature phase diagram of a polarized system. The noise correlations in a one-dimensional system turn out to show differences between different states which are straightforward to interpret and contain useful information on the underlying pair-wavefunction as well as on the structure of the quasiparticle dispersions.

This paper is organized as follows. In Sec. [II](#page-2-0) we discuss the physical system and present the Hamiltonian of the system. In this section different paired phases are also discussed In Sec. [III](#page-4-0) we proceed to compute the noise correlation functions for different paired phases at zero temperature and in Sec. [IV](#page-12-0) computations are generalized to non-zero temperatures. In Sec. [V](#page-18-0) noise correlations in a one dimensional lattice are discussed while making a clear connection with the computed 1D phase-diagram. We end with some concluding remarks in Sec. [VI.](#page-19-0)

II. THE HAMILTONIAN OF THE SYSTEM

We assume in our calculation a three dimensional cubic optical lattice. The system is composed of a two component Fermi gas where components are different hyperfine states (for concreteness we assume ⁶Li atoms). In terms of the field operators $\hat{\Psi}_{\sigma}(\mathbf{r})$ the Hamiltonian of the system is given by

$$
\hat{H} = \sum_{\sigma=\uparrow,\downarrow} \int d\mathbf{r} \,\hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r}) \left(-\frac{\hbar^{2}\nabla^{2}}{2m_{\sigma}} + V_{\sigma}(\mathbf{r}) \right) \hat{\Psi}_{\sigma}(\mathbf{r}) + \n\int \int d\mathbf{r} \,d\mathbf{r}' \hat{\Psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\Psi}_{\downarrow}^{\dagger}(\mathbf{r}') g(\mathbf{r}, \mathbf{r}') \hat{\Psi}_{\downarrow}(\mathbf{r}') \hat{\Psi}_{\uparrow}(\mathbf{r}) - \sum_{\sigma=\uparrow,\downarrow} \mu_{\sigma} \hat{N}_{\sigma} \n= \sum_{\sigma=\uparrow,\downarrow} \int d\mathbf{r} \,\hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r}) \left(-\frac{\hbar^{2}\nabla^{2}}{2m_{\sigma}} + V_{\sigma}(\mathbf{r}) - \mu_{\sigma} \right) \hat{\Psi}_{\sigma}(\mathbf{r}) + \n\int \int d\mathbf{r} \,d\mathbf{r}' \hat{\Psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\Psi}_{\downarrow}^{\dagger}(\mathbf{r}') g(\mathbf{r}, \mathbf{r}') \hat{\Psi}_{\downarrow}(\mathbf{r}') \hat{\Psi}_{\uparrow}(\mathbf{r}),
$$
\n(1)

where $\hbar = h/(2\pi)$ and h is the Planck constant. μ_{σ} is the chemical potential of the component σ and the lattice potential is given by $V_{\sigma}(\mathbf{r}) = sE_r \sum_{i=1}^3 \sin^2(kx_i)$, where s is the lattice depth and $E_r = \hbar^2 k^2/(2m)$ is the recoil energy $(k = \pi/d)$ and d is the lattice constant). In the usual way the interaction between atoms is modelled by a contact potential

$$
g(\mathbf{r}, \mathbf{r}') = \frac{4\pi\hbar^2 a}{m} \delta(\mathbf{r} - \mathbf{r}'),
$$

where a is the s-wave scattering length. The number operator of the component σ

$$
\hat{N}_{\sigma} = \int d\mathbf{r} \,\hat{n}_{\sigma}(\mathbf{r}) = \int d\mathbf{r} \,\hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r})\hat{\Psi}_{\sigma}(\mathbf{r}),
$$

is expressed in terms of the density operator $\hat{n}_{\sigma}(\mathbf{r}) = \hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r}) \hat{\Psi}_{\sigma}(\mathbf{r})$. In our case the fieldoperators are fermionic which implies fermionic anticommutation relations

$$
\{\hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r}),\hat{\Psi}_{\beta}(\mathbf{r}')\} = \hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r})\hat{\Psi}_{\beta}(\mathbf{r}') + \hat{\Psi}_{\beta}(\mathbf{r}')\hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r}) = \delta_{\alpha\beta}\delta(\mathbf{r} - \mathbf{r}')
$$
\n
$$
\{\hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r}),\hat{\Psi}_{\beta}^{\dagger}(\mathbf{r}')\} = \{\hat{\Psi}_{\alpha}(\mathbf{r}),\hat{\Psi}_{\beta}(\mathbf{r}')\} = 0,
$$
\n(2)

where $\delta_{\alpha\beta}$ is the Kronecker delta, and $\delta(\mathbf{r} - \mathbf{r}')$ is the Dirac delta-function.

A. The Hubbard model

In a sufficiently deep lattice we can expand the field operators in terms of the well localized (lowest band) Wannier functions as

$$
\hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r}) = \sum_{i} w_{i}^{*}(\mathbf{r}) \hat{c}_{\sigma,i}^{\dagger}
$$
\n
$$
\hat{\Psi}_{\sigma}(\mathbf{r}) = \sum_{i} w_{i}(\mathbf{r}) \hat{c}_{\sigma,i},
$$
\n(3)

where $w_i(\mathbf{r})$ is Wannier function centered around a lattice point *i*, and $\hat{c}_{\sigma,i}$ and $\hat{c}_{\sigma,i}^{\dagger}$ are fermionic annihilation- and creation operators which annihilate and create fermions of component σ at site *i*.

It is assumed that only tunneling between the nearest neighbours is of importance. This assumption of tight binding is reasonable when the Wannier functions are well localized. In other words, overlap integrals between the next nearest neighbour Wannier functions are small compared to overlap integrals between the nearest neighbour Wannier functions. This always happens when the lattice is deep enough [\[34](#page-23-5)]. Using this approximation one finds the Hubbard Hamiltonian [\[35\]](#page-23-6)

$$
\hat{H} = \sum_{n} \left(-\mu_{\uparrow} \hat{c}_{\uparrow,n}^{\dagger} \hat{c}_{\uparrow,n} - \mu_{\downarrow} \hat{c}_{\downarrow,n}^{\dagger} \hat{c}_{\downarrow,n} \right) + U \sum_{n} \hat{c}_{\uparrow,n}^{\dagger} \hat{c}_{\downarrow,n} \hat{c}_{\downarrow,n} \hat{c}_{\uparrow,n} \n- \left(J_x \sum_{\langle n,m \rangle_x} + J_y \sum_{\langle n,m \rangle_y} + J_z \sum_{\langle n,m \rangle_z} \right) \left(\hat{c}_{\uparrow,m}^{\dagger} \hat{c}_{\uparrow,n} + \hat{c}_{\downarrow,n}^{\dagger} \hat{c}_{\downarrow,n} \right),
$$
\n(4)

where $\langle m, n \rangle_\alpha$ means a sum over the nearest neighbours in the α-direction and hopping strength is defined by

$$
J_{\alpha} = -\int d\mathbf{r} \, w_n^*(\mathbf{r}) \left(-\frac{\hbar^2 \nabla^2}{2m} + sE_r \sum_i \sin^2(kx_i) \right) w_{n \pm d\hat{x}_{\alpha}}(\mathbf{r}).
$$

Furthermore, the dominant on site interaction coupling strength is given by

$$
U = \frac{4\pi\hbar^2 a}{m} \int d\mathbf{r} |w_i(\mathbf{r})|^4.
$$

In our calculations we have approximated the Wannier functions with a harmonic oscillator ground state in each potential well.

B. The ground state ansatz at zero temperature

On a mean-field level in the homogeneous system the ground state ansatz at zero temperature which includes the possibility of the BCS-state [\[36\]](#page-23-7), the breach pair(BP)-state [\[8,](#page-22-7) [9\]](#page-22-8), and the plane wave FFLO-state [\[6](#page-22-5), [7](#page-22-6)] can be expressed as [\[37\]](#page-23-8)

$$
|\Psi_{GS}\rangle = \prod_{\mathbf{k}\in G_3} \hat{c}_{\downarrow,-\mathbf{k}+\mathbf{q}}^{\dagger} \prod_{\mathbf{k}\in G_2} \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \prod_{\mathbf{k}\in G_1} (u_{\mathbf{k},\mathbf{q}} + v_{\mathbf{k},\mathbf{q}} \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k}+\mathbf{q}}^{\dagger}) |0\rangle, \tag{5}
$$

where $|0\rangle$ the vacuum state. In area G_1 both quasiparticle dispersions $E_{\uparrow,\mathbf{k},\mathbf{q}},E_{\downarrow,\mathbf{k},\mathbf{q}}\geq 0$ are positive, in area G_2 $E_{\downarrow,\mathbf{k},\mathbf{q}} \ge 0$ and $E_{\uparrow,\mathbf{k},\mathbf{q}} < 0$, and in area G_3 $E_{\uparrow,\mathbf{k},\mathbf{q}} \ge 0$ and $E_{\downarrow,\mathbf{k},\mathbf{q}} < 0$. Area G_1 is populated by pairs with a quasi-momentum q. Area G_2 is populated by atoms of the component \uparrow in the quasi-momentum states $\mathbf{k} \in G_2$ while the area G_3 is populated by \downarrow -atoms in the quasi-momentum states $-\mathbf{k} + \mathbf{q}$. By diagonalizing the mean-field theory one finds the quasiparticle dispersions

$$
E_{\downarrow,\mathbf{k},\mathbf{q}} = \sqrt{\frac{\left(\frac{\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2}\right)^2 + \Delta^2} - \frac{\epsilon_{\uparrow,\mathbf{k}} - \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2}}
$$
\n
$$
E_{\uparrow,\mathbf{k},\mathbf{q}} = \sqrt{\frac{\left(\frac{\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2}\right)^2 + \Delta^2} + \frac{\epsilon_{\uparrow,\mathbf{k}} - \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2}},
$$
\n(6)

where Δ is the pairing gap and the single particle dispersions are

$$
\epsilon_{\sigma,\mathbf{k}} = \sum_{\alpha=1}^{3} 2J_{\alpha}(1 - \cos(k_{\alpha}d)) - \mu_{\sigma}.
$$

The quasiparticle amplitudes $u_{\mathbf{k},\mathbf{q}}$ and $v_{\mathbf{k},\mathbf{q}}$ are given by

$$
u_{\mathbf{k},\mathbf{q}}^2 = \frac{1}{2} \left(1 + \frac{\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2\sqrt{[(\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q})}/2]^2 + \Delta^2}} \right)
$$

$$
v_{\mathbf{k},\mathbf{q}}^2 = \frac{1}{2} \left(1 - \frac{\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q}}}{2\sqrt{[(\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,-\mathbf{k}+\mathbf{q})}/2]^2 + \Delta^2}} \right).
$$
 (7)

Based on the properties of this ansatz, it can be classified further. When the wavevector $q \neq 0$, the ground state is called the FFLO-state which breaks the translational symmetry. If $\mathbf{q} = 0$ and both quasiparticle dispersions $E_{\uparrow,\mathbf{k}}$, $E_{\downarrow,\mathbf{k}}$ are positive for all **k**-vectors, the ground state is called BCS-state, and finally if $q = 0$ and $E_{\sigma, k} < 0$ for some k-vectors, the ground state is called the BP-state. This last case involves phase separation in the momentum space.

III. NOISE CORRELATIONS

Our aim is to compute the noise correlation functions which are defined by [\[26](#page-22-25)]

$$
G_{\alpha\beta}(\mathbf{r},\mathbf{r}') = \langle \hat{n}_{\alpha}(\mathbf{r},t)\hat{n}_{\beta}(\mathbf{r}',t) \rangle - \langle \hat{n}_{\alpha}(\mathbf{r},t)\rangle \langle \hat{n}_{\beta}(\mathbf{r}',t) \rangle = \langle \hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r},t)\hat{\Psi}_{\alpha}(\mathbf{r},t)\hat{\Psi}_{\beta}(\mathbf{r}',t)\hat{\Psi}_{\beta}(\mathbf{r}',t) \rangle - \langle \hat{n}_{\alpha}(\mathbf{r},t)\rangle \langle \hat{n}_{\beta}(\mathbf{r}',t) \rangle = \langle \hat{\Psi}_{\alpha}^{\dagger}(\mathbf{r},t)\hat{\Psi}_{\beta}^{\dagger}(\mathbf{r}',t)\hat{\Psi}_{\beta}(\mathbf{r}',t)\rangle + \delta_{\alpha\beta}\delta(\mathbf{r}-\mathbf{r}')\langle \hat{n}_{\alpha}(\mathbf{r},t)\rangle - \langle \hat{n}_{\alpha}(\mathbf{r},t)\rangle \langle \hat{n}_{\beta}(\mathbf{r}',t) \rangle, \tag{8}
$$

where α and β are component indeces, **r** and **r'** are the positions, and t is time. The term with Dirac's delta function is due to the normal ordering of the density-density expectation value.

Often in experiments a gas which has been trapped is released and the gas expands freely. The free expansion means that interactions between the atoms are so weak that during the expansion collisions between atoms can be ignored. Because the Wannier functions $w_i(\mathbf{r}, t)$ are strongly localized at a lattice point i at time $t = 0$, at long times (the saddle point approximation of the free evolution amounts to the limit $t \gg md^2/\hbar$ the Wannier functions are essentially plane waves. In other words at long times

$$
w_i(\mathbf{r},t) \sim e^{-iQ(\mathbf{r}) \cdot \mathbf{R}_i},
$$

where $\mathbf{R}_i = d(i_x, i_y, i_z)$ (i_n are integers) is a lattice vector and

$$
Q(\mathbf{r}) = \frac{m\mathbf{r}}{\hbar t}.
$$

Thus the densities become

$$
n_{\alpha}(\mathbf{r},t) = \langle \hat{n}_{\alpha}(\mathbf{r},t) \rangle = A(t)^{2} \sum_{i,j} e^{iQ(\mathbf{r}) \cdot \mathbf{R}_{ij}} \langle \hat{c}^{\dagger}_{\alpha,i} \hat{c}_{\alpha,j} \rangle, \tag{9}
$$

where $\mathbf{R}_{ij} = \mathbf{R}_i - \mathbf{R}_j$ and $A(t)$ is a time dependent scale factor which depends on the Wannier functions. The correlation functions in Eq. [\(8\)](#page-4-1) now become

$$
G_{\alpha\beta}(\mathbf{r}, \mathbf{r}') = A(t)^4 \sum_{i,j,m,n} e^{iQ(\mathbf{r}) \cdot \mathbf{R}_{ij} + iQ(\mathbf{r}') \cdot \mathbf{R}_{mn}} \langle \hat{c}^{\dagger}_{\alpha,i} \hat{c}^{\dagger}_{\beta,m} \hat{c}_{\beta,n} \hat{c}_{\alpha,j} \rangle + \delta_{\alpha\beta} \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\alpha}(\mathbf{r}, t) \rangle - \langle \hat{n}_{\alpha}(\mathbf{r}, t) \rangle \langle \hat{n}_{\beta}(\mathbf{r}', t) \rangle.
$$
\n(10)

A. The noise correlations of the BCS-state at zero temperature

We review here the noise correlations of the BCS-state. The usual unpolarized BCS ground state is extracted from the ansatz in Eq. [\(5\)](#page-4-2) by setting both chemical potentials to be equal and $q = 0$. Because both quasiparticle dispersions are now positive, the BCS ground state at zero temperature is given by

$$
|\Psi_{BCS}\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}^{\dagger}_{\uparrow,\mathbf{k}} \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}})|0\rangle.
$$
 (11)

Because the BCS-ground state is presented in momentum space, for simplicity we express the densities in Eq. [\(9\)](#page-5-0) in terms of the operators in the momentum space

$$
\langle \hat{n}_{\alpha}(\mathbf{r},t) \rangle = \frac{A(t)^2}{M^2} \sum_{i,j,\mathbf{k}_1,\mathbf{k}_2} e^{iQ(\mathbf{r}) \cdot \mathbf{R}_{ij}} e^{-i\mathbf{k}_1 \cdot \mathbf{R}_i + i\mathbf{k}_2 \cdot \mathbf{R}_j} \langle \hat{c}^{\dagger}_{\alpha,\mathbf{k}_1} \hat{c}_{\alpha,\mathbf{k}_2} \rangle \tag{12}
$$

and similarly for the noise correlations in Eq. [\(10\)](#page-5-1)

$$
G_{\alpha\beta}(\mathbf{r}, \mathbf{r}') = \frac{A(t)^4}{M^4} \sum_{\substack{i,j,m,n,\\ \mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4}} \left(e^{iQ(\mathbf{r}) \cdot \mathbf{R}_{ij} + iQ(\mathbf{r}') \cdot \mathbf{R}_{mn}} e^{-i\mathbf{k}_1 \cdot \mathbf{R}_i + i\mathbf{k}_2 \cdot \mathbf{R}_j - i\mathbf{k}_3 \cdot \mathbf{R}_m + i\mathbf{k}_4 \cdot \mathbf{R}_n} \times \left(\hat{c}^{\dagger}_{\alpha, \mathbf{k}_1} \hat{c}^{\dagger}_{\beta, \mathbf{k}_3} \hat{c}_{\beta, \mathbf{k}_4} \hat{c}_{\alpha, \mathbf{k}_2} \right) \right)
$$
\n
$$
+ \delta_{\alpha\beta} \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\alpha}(\mathbf{r}, t) \rangle - \langle \hat{n}_{\alpha}(\mathbf{r}, t) \rangle \langle \hat{n}_{\beta}(\mathbf{r}', t) \rangle, \tag{13}
$$

where M is the number of lattice sites. Now we need the expectation values $\langle \hat{c}_c^\dagger$ $\overset{\dagger}{\alpha},\mathbf{k}_1 \overset{\dagger}{c}^{\dagger}_\beta$ $\rangle^{\dagger}_{\beta, \mathbf{k}_3} \hat{c}_{\beta, \mathbf{k}_4} \hat{c}_{\alpha, \mathbf{k}_2} \rangle$ and $\langle \hat{c}^{\dagger}_{\alpha} \rangle$ $\langle \delta_{\alpha,\mathbf{k}_1} \hat{c}_{\alpha,\mathbf{k}_2} \rangle$. Because

$$
\langle 0|\hat{c}_{\downarrow,-\mathbf{k}_1}\hat{c}_{\uparrow,\mathbf{k}_1}(\hat{c}_{\uparrow,\mathbf{k}_1}^{\dagger}\hat{c}_{\uparrow,\mathbf{k}_2})\hat{c}_{\uparrow,\mathbf{k}_1}^{\dagger}\hat{c}_{\downarrow,-\mathbf{k}_1}^{\dagger}|0\rangle = \langle 0|\hat{c}_{\downarrow,\mathbf{k}_2}\hat{c}_{\uparrow,-\mathbf{k}_2}(\hat{c}_{\downarrow,\mathbf{k}_1}^{\dagger}\hat{c}_{\downarrow,\mathbf{k}_2})\hat{c}_{\uparrow,-\mathbf{k}_2}^{\dagger}\hat{c}_{\downarrow,\mathbf{k}_2}^{\dagger}|0\rangle = \delta_{\mathbf{k}_1\mathbf{k}_2},
$$

and $v_{\mathbf{k}} = v_{-\mathbf{k}}$, we find

$$
\langle \Psi_{BCS} | \hat{c}_{\alpha, \mathbf{k}_1}^{\dagger} \hat{c}_{\alpha, \mathbf{k}_2} | \Psi_{BCS} \rangle = \delta_{\mathbf{k}_1 \mathbf{k}_2} |v_{\mathbf{k}_1}|^2. \tag{14}
$$

Using this we can calculate the densities in Eq. [\(12\)](#page-5-2) and find

$$
\langle \hat{n}_{\alpha}(\mathbf{r},t) \rangle = \frac{A(t)^2}{M^2} \sum_{i,j,\mathbf{k}} e^{i(Q(\mathbf{r}) - \mathbf{k}) \cdot \mathbf{R}_{ij}} |v_{\mathbf{k}}|^2.
$$
 (15)

Now the number of particles is the same for both the components as it should be. When the lattice is large, i.e. when the lattice size $M \gg 1$, the exponential terms only add constructively when $Q(\mathbf{r}) - \mathbf{k} = 0$ and we can approximate the densities as

$$
\langle \hat{n}_{\alpha}(\mathbf{r},t) \rangle = A(t)^2 \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t}{m} \mathbf{k}\right) |v_{\mathbf{k}}|^2.
$$
 (16)

Non-vanishing four operator expectation values are

$$
\langle \Psi_{BCS} | \hat{c}^{\dagger}_{\uparrow,\mathbf{k}} \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}'} \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}'} \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}'} \hat{c}^{\dagger}_{\uparrow,\mathbf{k}} | \Psi_{BCS} \rangle = \delta_{\mathbf{k}\mathbf{k}'} |v_{\mathbf{k}}|^2 + (1 - \delta_{\mathbf{k}\mathbf{k}'}) |v_{\mathbf{k}'}|^2 |v_{\mathbf{k}}|^2
$$
\n
$$
\langle \Psi_{BCS} | \hat{c}^{\dagger}_{\uparrow(\downarrow),\mathbf{k}} \hat{c}^{\dagger}_{\uparrow(\downarrow),\mathbf{k}'} \hat{c}^{\dagger}_{\uparrow(\downarrow),\mathbf{k}} | \Psi_{BCS} \rangle = (1 - \delta_{\mathbf{k}\mathbf{k}'}) |v_{\mathbf{k}'}|^2 |v_{\mathbf{k}}|^2. \tag{17}
$$

Combining these results we can compute the noise correlations of the BCS-state at zero temperature

$$
G_{\uparrow\downarrow}(\mathbf{r}, \mathbf{r}') = G_{\downarrow\uparrow}(\mathbf{r}, \mathbf{r}') = \frac{A(t)}{M^4} \sum_{i,j,m,n,\mathbf{k}} |u_{\mathbf{k}}|^2 |v_{\mathbf{k}}|^2 e^{i(Q(\mathbf{r}) - \mathbf{k}) \cdot \mathbf{R}_{ij} + i(Q(\mathbf{r}') + \mathbf{k}) \cdot \mathbf{R}_{mn}}
$$

\n
$$
G_{\sigma\sigma}(\mathbf{r}, \mathbf{r}') = \frac{-A(t)}{M^4} \sum_{i,j,m,n,\mathbf{k}} |v_{\mathbf{k}}|^4 e^{i(Q(\mathbf{r}) - \mathbf{k}) \cdot \mathbf{R}_{ij} + i(Q(\mathbf{r}') - \mathbf{k}) \cdot \mathbf{R}_{mn}}
$$
(18)
\n
$$
+ \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\sigma}(\mathbf{r}, t) \rangle.
$$

Again for large lattices these equations can be approximated in the following way

$$
G_{\uparrow\downarrow}(\mathbf{r}, \mathbf{r}') = G_{\downarrow\uparrow}(\mathbf{r}, \mathbf{r}') = A(t) \sum_{\mathbf{k}} |u_{\mathbf{k}}|^2 |v_{\mathbf{k}}|^2 \delta \left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m} \right) \delta \left(\mathbf{r}' + \frac{\hbar t \tilde{\mathbf{k}'}}{m} \right)
$$

$$
G_{\sigma\sigma}(\mathbf{r}, \mathbf{r}') = -A(t)^4 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^4 \delta \left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m} \right) \delta \left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}'}}{m} \right)
$$

$$
+ \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\sigma}(\mathbf{r}, t) \rangle,
$$
 (19)

where $\tilde{\mathbf{k}} = \mathbf{k} + \sum_{\alpha=1}^{3} 2n_{\alpha} \pi \hat{x}_{\alpha}/d$ and $\tilde{\mathbf{k}}' = \mathbf{k} + \sum_{\alpha=1}^{3} 2m_{\alpha} \pi \hat{x}_{\alpha}/d$, where **k** is a lattice momentum, \hat{x}_{α} are orthogonal unit vectors and n_i and m_i are integers. Because

$$
|u_{\mathbf{k}}|^2|v_{\mathbf{k}}|^2 = \frac{\Delta^2}{4([(\epsilon_{\uparrow,\mathbf{k}} + \epsilon_{\downarrow,\mathbf{k}})/2]^2 + \Delta^2)},
$$

FIG. 1: [Color online] In (a) we show a cut of the BCS noise correlations between components in the $z = 0$ plane. In (b) we show the integrated BCS correlations between the components. In (c) we demonstrate the noise correlation of a single component in the plane $z = 0$. As parameters we used the average filling fraction $(n_1 + n_1)/2 = 0.30$, polarization $P = (n_1 - n_1)/(n_1 + n_1) = 0.0$, $\Delta/(2J) = 1.03$ and $U/(2J) = -3.0$. All these examples are calculated at $T = 0$. In (a) and (b) we choose $\mathbf{r}' = -\mathbf{r}$ and in (c) we choose $y' = y$, $z = z' = 0$, and $x' = x + \hbar t 2\pi/(md)$. Color-coding is such that warm colors imply high peaks and cold colors low.

the noise correlation between different components vanishes with vanishing gap Δ and is thus identically zero in the normal state. In other words in the normal state in the mean field level correlations between different components vanish:

$$
G_{\uparrow\downarrow}(\mathbf{r},\mathbf{r}') = G_{\downarrow\uparrow}(\mathbf{r},\mathbf{r}') = 0.
$$

One can see that the noise correlations in the BCS-state differ crucially from this. Because in the BCS-state momenta **k** and $-k$ are correlated, the noise correlation signal is pronounced only if

$$
\mathbf{r} + \mathbf{r}' = \sum_{i=1}^{3} \frac{\hbar 2n_i \pi t \hat{x}_i}{md}
$$

and

$$
\mathbf{r} = \frac{\hbar t \mathbf{k}}{m} + \sum_{i=1}^{3} \frac{\hbar 2 m_i \pi t \hat{x}_i}{m d},
$$

where **k** is a lattice momentum. The continuum result $[26]$ is similar, but peaks appear when $\mathbf{r} + \mathbf{r}' = 0$. In the continuum limit our result reduces to the known result.

On the other hand correlations of the single component show a hole when

$$
\mathbf{r} - \mathbf{r}' = \sum_{i=1}^{3} \frac{2\hbar n_i \pi t \hat{x}_i}{md}
$$

and $n_i \neq 0$ for at least one i [\[38](#page-23-9)]. The reason for this is the fermionic antibunching related to Pauli blocking i.e. two or more identically fermions cannot be in the same momentum-state. This fermionic antibunching was also measured in the noise correlations of an ideal Fermi gas [\[28](#page-22-27)].

In Fig. [1](#page-7-0) we demonstrate the BCS noise correlations at $T = 0$. In Fig. 1 (a) we show a cut in the $z = 0$ plane of the BCS noise correlation between the components while in Fig. [1](#page-7-0) (b) we plot the column integrated BCS-state noise correlations

$$
G_{\uparrow,\downarrow}(x,y) = \int dz \, G_{\uparrow,\downarrow}(x,y,z,-x+\hbar t q_x/m, -y+\hbar t q_y/m, -z+\hbar t q_z/m).
$$

As one can see from the Fig. [1,](#page-7-0) the noise correlations reflect underlying lattice structure and strongly non-spherical structure of the Fermi surface. We can see also from Fig. [1](#page-7-0) (a) and (b) that the peaks are highest near the Fermi surfaces where $\epsilon_{\uparrow,k} + \epsilon_{\downarrow,-k} = 0$. In the weakly interacting BCS limit correlations are strongly peaked at the Fermi surface, but the distribution of the peak heights becomes broad in the BEC limit, and the signal approaches the interference signal of the Bose-Einstein condensate.

In Fig. [1](#page-7-0) (c) we show the BCS state correlations of a single component in the $z = 0$ plane. As is clear, the BCS-state noise correlation of a single component shows similar antibunching behavior as the ideal Fermi gas noise correlation. If $\mathbf{r} = \mathbf{r}' + \sum_{i=1}^{3} 2\hbar n_i \pi t \hat{x}_i / (md)$ and and $n_i \neq 0$ for at least one i, then the noise correlation of a single component shows the holes. However, the BCS-state noise correlation of a single component would also show the bunching peak when $\mathbf{r} = \mathbf{r}'$. This result differs from the ideal gas result where such a peak is absent.

B. The noise correlations of the BP-state at zero temperature

The ground state in Eq. [\(5\)](#page-4-2) is the BP-state if $q = 0$ and one of the quasiparticle dispersions $E_{\sigma, \mathbf{k}}$ is negative for some values of the momentum **k**. This implies that in the BP-state $|\delta \mu| = |(\mu_{\uparrow} - \mu_{\downarrow})| > 2\Delta > 0$. The BP-state is a gapless state, because one of the quasiparticle dispersions changes its sign when the momentum varies. The BP ground state at zero-temperature is

$$
|\Psi_{BP}\rangle = \prod_{\mathbf{k}\in G_3} \hat{c}_{\downarrow,-\mathbf{k}}^{\dagger} \prod_{\mathbf{k}\in G_2} \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \prod_{\mathbf{k}\in G_1} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k}}^{\dagger}) |0\rangle.
$$
 (20)

In the area G_1 both quasiparticle dispersions $E_{\uparrow,\mathbf{k},\mathbf{q}},E_{\downarrow,\mathbf{k},\mathbf{q}}\geq 0$, in the area G_2 $E_{\downarrow,\mathbf{k}}\geq 0$ and $E_{\uparrow,\mathbf{k}} < 0$, and in the area G_3 $E_{\uparrow,\mathbf{k}} \ge 0$ and $E_{\downarrow,\mathbf{k}} < 0$.

Now we can calculate the required expectation values similarly as before. We find

$$
\langle \Psi_{BP} | \hat{c}^{\dagger}_{\uparrow(\downarrow),\mathbf{k}} \hat{c}_{\uparrow(\downarrow),\mathbf{k'}} | \Psi_{BP} \rangle = \delta_{\mathbf{k}\mathbf{k'}} [(1 - \theta(E_{\uparrow(\downarrow),\mathbf{k}}) + \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) | v_{\mathbf{k}} |^2]
$$
(21)

so that the densities in the BP-state (in a large lattice) become

$$
\langle \hat{n}_{\uparrow(\downarrow)}(\mathbf{r},t) \rangle = A(t)^2 \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t \mathbf{k}}{m}\right) \left[(1 - \theta(E_{\uparrow(\downarrow),\mathbf{k}}) + \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) |v_{\mathbf{k}}|^2 \right]. \tag{22}
$$

Non-vanishing expectation values in the noise correlations in Eq. [\(13\)](#page-5-3) are in turn given by

$$
\langle \Psi_{BP} | \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k'}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k'}} \hat{c}_{\uparrow,\mathbf{k}} | \Psi_{BP} \rangle =
$$
\n
$$
(1 - \delta_{\mathbf{k}\mathbf{k'}}) [(1 - \theta(E_{\uparrow,\mathbf{k}})) (1 - \theta(E_{\downarrow,\mathbf{k'}})) + \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k'}}) \theta(E_{\downarrow,\mathbf{k'}}) |v_{\mathbf{k}}|^2 |v_{\mathbf{k'}}|^2
$$
\n
$$
+ \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) (1 - \theta(E_{\downarrow,\mathbf{k'}})) |v_{\mathbf{k}}|^2 + \theta(E_{\uparrow,\mathbf{k'}}) \theta(E_{\downarrow,\mathbf{k'}}) (1 - \theta(E_{\uparrow,\mathbf{k'}})) |v_{\mathbf{k'}}|^2
$$
\n
$$
+ \delta_{\mathbf{k}\mathbf{k'}} \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) |v_{\mathbf{k}}|^2
$$
\n
$$
\langle \Psi_{BP} | \hat{c}_{\uparrow(\downarrow),\mathbf{k}}^{\dagger} \hat{c}_{\uparrow(\downarrow),\mathbf{k'}}^{\dagger} \hat{c}_{\uparrow(\downarrow),\mathbf{k'}} \hat{c}_{\uparrow(\downarrow),\mathbf{k'}} | \Psi_{BP} \rangle =
$$
\n
$$
(1 - \delta_{\mathbf{k}\mathbf{k'}}) [(1 - \theta(E_{\uparrow(\downarrow),\mathbf{k}})) (1 - \theta(E_{\uparrow(\downarrow),\mathbf{k'}})) + \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) |\psi_{\mathbf{k}}|^2 |v_{\mathbf{k'}}|^2
$$
\n
$$
+ \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) (1 - \theta(E_{\uparrow(\downarrow),\mathbf{k'}})) |v_{\mathbf{k}}|^2 + \theta(E_{\uparrow,\mathbf{k'}}) \theta(E_{\downarrow,\mathbf{k'}}) (1 - \theta(E_{\uparrow,\mathbf{k}})) |v_{\mathbf{k'}}|^2]].
$$
\n(23)

Subtracting the densities cancels out most of these terms and the noise correlations of the BP-state at zero temperature are given by

$$
G_{\uparrow\downarrow(\downarrow\uparrow)}(\mathbf{r},\mathbf{r}') = \frac{A(t)^4}{M^2} \sum_{i,j,m,n,\mathbf{k}} e^{i(Q(\mathbf{r})-\mathbf{k})\cdot\mathbf{R}_{ij}+i(Q(\mathbf{r}')+\mathbf{k})\cdot\mathbf{R}_{mn}}\theta(E_{\uparrow,\mathbf{k}})\theta(E_{\downarrow,\mathbf{k}})|u_{\mathbf{k}}|^2|v_{\mathbf{k}}|^2
$$

\n
$$
G_{\sigma\sigma}(\mathbf{r},\mathbf{r}') = \frac{-A(t)^4}{M^2} \sum_{i,j,m,n,\mathbf{k}} \left[e^{i(Q(\mathbf{r})-\mathbf{k})\cdot\mathbf{R}_{ij}+i(Q(\mathbf{r}')-\mathbf{k})\cdot\mathbf{R}_{mn}} \left(\theta(E_{\uparrow,\mathbf{k}})\theta(E_{\downarrow,\mathbf{k}})|v_{\mathbf{k}}|^4 \right) \right]
$$

\n+ $(1-\theta(E_{\sigma,\mathbf{k}})) \bigg) \bigg] + \delta(\mathbf{r}-\mathbf{r}')\langle \hat{n}_{\sigma}(\mathbf{r},t) \rangle.$ (24)

Again for large lattices we get strongly peaked behaviour and

$$
G_{\uparrow\downarrow(\downarrow\uparrow)}(\mathbf{r},\mathbf{r}') = A(t)^4 \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m}\right) \delta\left(\mathbf{r}' + \frac{\hbar t \tilde{\mathbf{k}'}}{m}\right) \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) |u_{\mathbf{k}}|^2 |v_{\mathbf{k}}|^2
$$

$$
G_{\sigma\sigma}(\mathbf{r},\mathbf{r}') = -A(t)^4 \sum_{\mathbf{k}} \left[\delta\left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m}\right) \delta\left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}'}}{m}\right) \theta(E_{\uparrow,\mathbf{k}}) \theta(E_{\downarrow,\mathbf{k}}) |v_{\mathbf{k}}|^4 \right]
$$
(25)
+
$$
(1 - \theta(E_{\sigma,\mathbf{k}})) \left] + \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\sigma}(\mathbf{r},t) \rangle.
$$

In the correlation between different components there are now areas where peaks are absent. The reason for this is that the BP-state has phase separation in the momentum space. Because in the BP-state there is no pairing between momenta **k** and $-\mathbf{k}$, if $E_{\uparrow,\mathbf{k}} < 0$ or $E_{\downarrow,\mathbf{k}} < 0$, in contrast to the BCS-state, there are no correlations between the points

$$
\mathbf{r} + \mathbf{r}' = \sum_{i=1}^{3} \frac{\hbar 2 n_i \pi t \hat{x}_i}{m d},
$$

in this region.

In Fig. [2](#page-10-0) we demonstrate the BP-state noise correlations at $T = 0$. In Fig. 2 (a) we have plotted the BP-state noise correlation at the $z = 0$ plane while in Fig. (b) we show the column integrated BP noise correlation. From Fig. [2](#page-10-0) (a) one can see clear areas without correlation peaks in the BP noise correlation due to the fact that these areas are populated only by atoms of the majority component which are unpaired. From the integrated correlations in Fig. [2](#page-10-0) (b) we can see that the peaks are now relatively low in the area between the Fermi surfaces. This differs substantially from the integrated BCS noise correlation (see Fig. [1](#page-7-0) (b)).

Results in this section are intended mainly as a reference since the BP-state is not expected to be the ground state at zero temperature. Depending on polarization the ground state at zero temperature can be the BCS-state, phase separation [\[39\]](#page-23-10), or the FFLO type state [\[21](#page-22-20)]. However, at non-zero temperatures the BP-state can be the minimum of the free energy.

C. The noise correlations of the plane wave FFLO-state at zero temperature

The FFLO ground state is presented in Eq. [\(5\)](#page-4-2). Contrary to the BCS-state and the BPstate, in the FFLO-state the parameter $q \neq 0$ and the translational invariance is broken. In

FIG. 2: [Color online] In this figure we demonstrate examples of the BP noise correlations between the components at $T = 0$. In (a) we demonstrate a cut in the $z = 0$ plane while in (b) we show the column integrated BP noise correlation. Parameters we used were $\Delta/(2J) = 0.318$, $(\mu_{\uparrow} + \mu_{\downarrow})/2.0 = 2.18$, and $\delta\mu = \mu_{\uparrow} - \mu_{\downarrow} = 1.14$. We have chosen $\mathbf{r}' = -\mathbf{r}$ and color-coding is such that warm colors imply high peaks and cold colors low.

this case the order parameter is given by $\Delta(\mathbf{R}_i) = \Delta_0 \exp(i\mathbf{q} \cdot \mathbf{R}_i)$, where \mathbf{R}_i is a lattice vector. In the FFLO-state the other Fermi surface has been effectively shifted by the wavevector q.

The densities in the FFLO-state are (in the limit of large lattices) given by

$$
\langle \hat{n}_{\uparrow}(\mathbf{r},t) \rangle = A(t)^{2} \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t \mathbf{k}}{m}\right) \left[(1 - \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) + \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) |v_{\mathbf{k},\mathbf{q}}|^{2} \right]
$$

$$
\langle \hat{n}_{\downarrow}(\mathbf{r},t) \rangle = A(t)^{2} \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t(\mathbf{q} - \mathbf{k})}{m}\right) \left[(1 - \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) + \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) |v_{\mathbf{k},\mathbf{q}}|^{2} \right].
$$
 (26)

In the same way that we computed the noise correlations in the BCS- and BP-states, we

find the noise correlation in the FFLO-state as

$$
G_{\uparrow\downarrow}(\mathbf{r},\mathbf{r}') = A(t)^{4} \sum_{\mathbf{k}} \delta\left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m}\right) \delta\left(\mathbf{r}' + \frac{\hbar t(\tilde{\mathbf{k}}' - \mathbf{q})}{m}\right) \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) |u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2}
$$

\n
$$
G_{\downarrow\uparrow}(\mathbf{r},\mathbf{r}') = A(t)^{4} \sum_{\mathbf{k}} \delta\left(\mathbf{r} + \frac{\hbar t(\tilde{\mathbf{k}} - \mathbf{q})}{m}\right) \delta\left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}}'}{m}\right) \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) |u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2}
$$

\n
$$
G_{\uparrow\uparrow}(\mathbf{r},\mathbf{r}') = -A(t)^{4} \sum_{\mathbf{k}} \left[\delta\left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m}\right) \delta\left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}}'}{m}\right) \theta(E_{\uparrow,\mathbf{k},\mathbf{q}}) \theta(E_{\downarrow,\mathbf{k},\mathbf{q}}) |v_{\mathbf{k},\mathbf{q}}|^{4} \right]
$$
(27)
\n
$$
+ (1 - \theta(E_{\uparrow,\mathbf{k},\mathbf{q}})) \right] + \delta(\mathbf{r} - \mathbf{r}') \langle \hat{n}_{\uparrow}(\mathbf{r},t) \rangle
$$

\n
$$
G_{\downarrow\downarrow}(\mathbf{r},\mathbf{r}') = -A(t)^{4} \sum_{\mathbf{k}} \left[\delta\left(\mathbf{r} - \frac{\hbar t(\tilde{\mathbf{k}} - \mathbf{q})}{m}\right) \delta\left(\mathbf{r}' - \frac{\hbar t(\tilde{\mathbf{k}}' - \mathbf{q})}{m}\right) \cdot \theta(E_{\uparrow,\mathbf{k},\mathbf{
$$

Because in the FFLO-state momenta **k** and $-\mathbf{k} + \mathbf{q}$ are correlated, the points

$$
\mathbf{r} + \mathbf{r}' + \frac{\hbar t}{m} \mathbf{q} = \sum_{i=1}^{3} \frac{\hbar 2 n_i \pi t \hat{x}_i}{m d}
$$

are now correlated in the expanded cloud. Since the FFLO-state arises only when there is some polarization in the system, at $T = 0$ the FFLO-state is always a gapless state i.e. one of the quasiparticle dispersions $E_{\sigma,\mathbf{k},\mathbf{q}}$ changes its sign when momentum **k** varies.

Multimode FFLO-states also leave clear signatures on the noise correlations. For example, for the two mode FFLO-state i.e. the state where the gap is given by $\Delta(\mathbf{R_i}) = \Delta_0 \cos(\mathbf{q} \cdot \mathbf{R_i})$ (\mathbf{R}_i) is a lattice vector), the order parameter can be written as

$$
\Delta(\mathbf{R}_{i}) = \Delta_{0} \cos(\mathbf{q} \cdot \mathbf{R}_{i}) = \Delta_{0} \left(\frac{e^{i\mathbf{q} \cdot \mathbf{R}_{i}} + e^{-i\mathbf{q} \cdot \mathbf{R}_{i}}}{2} \right)
$$

= $-U \langle \hat{c}_{\uparrow,i} \hat{c}_{\downarrow,i} \rangle = \frac{-U}{M} \sum_{\mathbf{k},\mathbf{k}'} e^{-i(\mathbf{k} + \mathbf{k}') \cdot \mathbf{R}_{i}} \langle \hat{c}_{\uparrow,\mathbf{k}} \hat{c}_{\downarrow,\mathbf{k}'} \rangle$
= $e^{i\mathbf{q} \cdot \mathbf{R}_{i}} \frac{-U}{M} \sum_{\mathbf{k}} \langle \hat{c}_{\uparrow,\mathbf{k}} \hat{c}_{\downarrow,-\mathbf{k} - \mathbf{q}} \rangle + e^{-i\mathbf{q} \cdot \mathbf{R}_{i}} \frac{-U}{M} \sum_{\mathbf{k}} \langle \hat{c}_{\uparrow,\mathbf{k}} \hat{c}_{\downarrow,-\mathbf{k} + \mathbf{q}} \rangle.$

Therefore the only non-vanishing expectation values in the momentum space are $\langle \hat{c}_{\uparrow,\mathbf{k}} \hat{c}_{\downarrow,-\mathbf{k+q}} \rangle$ and $\langle \hat{c}_{\uparrow,\mathbf{k}} \hat{c}_{\downarrow,-\mathbf{k}-\mathbf{q}} \rangle$. This implies that an \uparrow -atom in the momentum state **k** is paired with \downarrow atoms in the momentum states $-k+q$ and $-k-q$. Because in the single mode FFLO-state points **r** and $-\mathbf{r} + \hbar t\mathbf{q}/m$ are correlated after free expansion, in the two mode FFLOstate **r** is correlated with $-\mathbf{r} + \hbar t \mathbf{q}/m$ and $-\mathbf{r} - \hbar t \mathbf{q}/m$ after free expansion. For this reason with the two mode FFLO-state one can see pronounced correlation peaks, when $\mathbf{r} + \mathbf{r}' \pm \hbar t \mathbf{q}/m = 0$. Actually for the multimode FFLO-state, one could use noise correlations to perform a "Fourier analysis" of the periodic order parameter and probe more complicated spatial dependencies than those discussed here.

FIG. 3: [Color online] In this figure we show the FFLO-state noise correlation $(G_{\uparrow\downarrow})$ at the temperature $T = 0$. In (a) we show a cut in the $z = 0$ plane while (b) shows the column integrated signal. Parameters we used were $q_x = 0.25(\pi/d), q_y = q_z = 0$, polarization $P = (n_\uparrow - n_\downarrow)/(n_\uparrow + n_\downarrow) = 0.168$, $\Delta/(2J) = 0.16$, and $U/(2J) = -1.86$. We choose $\mathbf{r} + \mathbf{r}' - (\hbar t/m)\mathbf{q} = 0$ and color-coding is again such that warm colors imply high peaks and cold colors low.

In Fig. [3](#page-12-1) we show the FFLO-state noise correlations $(G_†)$ at zero temperature. Fig. 3 (a) shows the FFLO-state noise correlation in the $z = 0$ plane, and Fig. [3](#page-12-1) (b) demonstrates the integrated FFLO noise correlation. As one can see, the Fermi surface shift creates asymmetry which remains also in the integrated signal. Gapless regions of the FFLO-state are again reflected as areas without correlation peaks and this is clearly presented in Fig. [3](#page-12-1) (a).

IV. NOISE CORRELATIONS AT NON-ZERO TEMPERATURE

Until now we have assumed zero temperature, but now we generalize our computations to non-zero temperatures. At non-zero temperatures it is useful to write the lowest energy state by using fermionic quasiparticle operators $\hat{\gamma}_{\sigma,\mathbf{k},\mathbf{q}}$ and $\hat{\gamma}_{\sigma}^{\dagger}$ $\sigma_{\alpha,\mathbf{k},\mathbf{q}}$. Creation and annihilation operators can be written in terms of these operators as

$$
\hat{c}_{\uparrow,\mathbf{k}} = u_{\mathbf{k},\mathbf{q}} \hat{\gamma}_{\uparrow,\mathbf{k},\mathbf{q}} - v_{\mathbf{k},\mathbf{q}} \hat{\gamma}_{\downarrow,\mathbf{k},\mathbf{q}}^{\dagger}
$$
\n
$$
\hat{c}_{\downarrow,-\mathbf{k}+\mathbf{q}} = u_{\mathbf{k},\mathbf{q}} \hat{\gamma}_{\downarrow,\mathbf{k},\mathbf{q}} + v_{\mathbf{k},\mathbf{q}} \hat{\gamma}_{\uparrow,\mathbf{k},\mathbf{q}}^{\dagger}.
$$
\n(28)

Now the lowest energy state ansatz is given by

$$
|\Psi_{GS}\rangle = \prod_{\mathbf{k}} \hat{\gamma}^{\dagger}_{\uparrow,\mathbf{k},\mathbf{q}} \hat{\gamma}^{\dagger}_{\downarrow,\mathbf{k},\mathbf{q}} |0\rangle \tag{29}
$$

and the lowest energy state is an ideal gas of quasiparticles. In other words the state in Eq. [\(5\)](#page-4-2) is the vacuum state for these quasiparticle operators. Therefore one has

$$
\langle \hat{\gamma}_{\alpha,\mathbf{k}}^{\dagger} \hat{\gamma}_{\beta,\mathbf{q}}^{\dagger} \rangle = \langle \hat{\gamma}_{\alpha,\mathbf{k}} \hat{\gamma}_{\beta,\mathbf{q}} \rangle = 0
$$

$$
\langle \hat{\gamma}_{\alpha,\mathbf{k}}^{\dagger} \hat{\gamma}_{\beta,\mathbf{q}} \rangle = \delta_{\alpha\beta} \delta_{\mathbf{kq}} f(E_{\alpha,\mathbf{k}}),
$$
 (30)

where the Fermi-Dirac distribution is given by

$$
f(E) = \frac{1}{e^{E/(k_B T)} + 1},
$$

and k_B is the Boltzmann constant.

The two-operator expectation values in the densities in Eq. [\(12\)](#page-5-2) at non-zero temperatures are

$$
\langle \hat{c}^{\dagger}_{\uparrow,\mathbf{k}} \hat{c}^{\dagger}_{\uparrow,\mathbf{k}'} \rangle = \delta_{\mathbf{k}\mathbf{k}'}(|u_{\mathbf{k},\mathbf{q}}|^2 f(E_{\uparrow,\mathbf{k},\mathbf{q}}) + |v_{\mathbf{k},\mathbf{q}}|^2 (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}}))) \langle \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}+\mathbf{q}} \hat{c}^{\dagger}_{\downarrow,-\mathbf{k}'+\mathbf{q}} \rangle = \delta_{\mathbf{k}\mathbf{k}'}(|u_{\mathbf{k},\mathbf{q}}|^2 f(E_{\downarrow,\mathbf{k},\mathbf{q}}) + |v_{\mathbf{k},\mathbf{q}}|^2 (1 - f(E_{\uparrow,\mathbf{k},\mathbf{q}}))),
$$
\n(31)

while non-vanishing four-operator expectation values in the noise correlations in Eq. [\(13\)](#page-5-3) at non-zero temperatures are

$$
\langle \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k'+q}}^{\dagger} \hat{c}_{\downarrow,-\mathbf{k'+q}} \hat{c}_{\uparrow,\mathbf{k}} \rangle = |u_{\mathbf{k},\mathbf{q}}|^{2} |u_{\mathbf{k'},\mathbf{q}}|^{2} f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\downarrow,\mathbf{k'},\mathbf{q}})
$$

+ |u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k'},\mathbf{q}}|^{2} f(E_{\uparrow,\mathbf{k},\mathbf{q}}) (1 - f(E_{\uparrow,\mathbf{k'},\mathbf{q}})) + |u_{\mathbf{k'},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} f(E_{\downarrow,\mathbf{k'},\mathbf{q}}) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}}))
+ |v_{\mathbf{k'},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} (1 - f(E_{\uparrow,\mathbf{k'},\mathbf{q}})) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}}))
+ \delta_{\mathbf{k}} \langle \{ |u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} [f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\downarrow,\mathbf{k},\mathbf{q}}) + (1 - f(E_{\uparrow,\mathbf{k},\mathbf{q}})) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}}))] \}
\langle \hat{c}_{\uparrow,\mathbf{k}}^{\dagger} \hat{c}_{\uparrow,\mathbf{k'}}^{\dagger} \hat{c}_{\uparrow,\mathbf{k'}} \hat{c}_{\uparrow,\mathbf{k'}} \hat{c}_{\uparrow,\mathbf{k}} \rangle = |u_{\mathbf{k},\mathbf{q}}|^{2} |u_{\mathbf{k'},\mathbf{q}}|^{2} f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\uparrow,\mathbf{k'},\mathbf{q}}) (32)
+ |u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k'},\mathbf{q}}|^{2} f(E_{\uparrow,\mathbf{k},\mathbf{q}}) (1 - f(E_{\downarrow,\mathbf{k'},\mathbf{q}})) + |u_{\mathbf

Because the densities again cancel out most of the terms, the noise correlations turn out to

be quite simple

$$
G_{\uparrow\downarrow}(\mathbf{r},\mathbf{r}') = A(t)^{4} \sum_{\mathbf{k}} \left[\delta \left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m} \right) \delta \left(\mathbf{r}' + \frac{\hbar t (\tilde{\mathbf{k}}' - \mathbf{q})}{m} \right) \right]
$$

\n
$$
[f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\downarrow,\mathbf{k},\mathbf{q}}) + (1 - f(E_{\uparrow,\mathbf{k},\mathbf{q}})) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}})) ||u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} \right]
$$

\n
$$
G_{\downarrow\uparrow}(\mathbf{r},\mathbf{r}') = A(t)^{4} \sum_{\mathbf{k}} \left[\delta \left(\mathbf{r} + \frac{\hbar t (\tilde{\mathbf{k}} - \mathbf{q})}{m} \right) \delta \left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}}'}{m} \right) \right]
$$

\n
$$
[f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\downarrow,\mathbf{k},\mathbf{q}}) + (1 - f(E_{\uparrow,\mathbf{k},\mathbf{q}})) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}})) ||u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} \right]
$$

\n
$$
G_{\uparrow\uparrow}(\mathbf{r},\mathbf{r}') = A(t)^{4} \sum_{\mathbf{k}} \left[\delta \left(\mathbf{r} - \frac{\hbar t \tilde{\mathbf{k}}}{m} \right) \delta \left(\mathbf{r}' - \frac{\hbar t \tilde{\mathbf{k}}'}{m} \right) \right]
$$
(33)
\n
$$
[f(E_{\uparrow,\mathbf{k},\mathbf{q}}) f(E_{\downarrow,\mathbf{k},\mathbf{q}}) + (1 - f(E_{\uparrow,\mathbf{k},\mathbf{q}})) (1 - f(E_{\downarrow,\mathbf{k},\mathbf{q}})) ||u_{\mathbf{k},\mathbf{q}}|^{2} |v_{\mathbf{k},\mathbf{q}}|^{2} \right]
$$

\n<math display="</math>

At non-zero temperatures the state is the BCS-state if $q = 0$ and the densities are equal, the BP-state if $q = 0$ and the densities are different. Since at finite temperatures the system can support polarization due to thermal effects even if both quasiparticle dispersions are always positive, we call the state gapless BP-state if $q = 0$ and one of the quasiparticle dispersions changes its sign.

Fig. [4](#page-15-0) demonstrates why the noise correlations are a better indicator of the superfluidity than densities and how the BCS-state noise correlation between the components changes quite substantially when the temperature rises and the gap is reduced. Fig. [4](#page-15-0) (a) shows a cut of the BCS-state density in the $z = 0$ plane when the temperature is $k_BT/(2J) = 0.56$ (i.e. just below the critical temperature). Fig. [4](#page-15-0) (b) shows a similar result just above the critical temperature. In Fig. [4](#page-15-0) (c) and (d) we compare the column integrated BCS-state noise correlations between the components at the temperatures $k_BT/(2J) = 0.0$, $k_BT/(2J) = 0.56$, respectively. By comparing Fig. [4](#page-15-0) (a) and (b), one can see that the densities are almost the same for the BCS-state and the normal-state. On the other hand in the normal-state the noise correlation is identically zero. Thermal effects are strongly present in the results in Fig. [4](#page-15-0) (d). This can be seen from the fact that the correlations peak in two areas, close to the center of the figure as well as near the Fermi surface. In the zero temperature result in Fig. [4](#page-15-0) (c) there are high peaks only near the Fermi surface. The reason for this difference between low and higher temperatures is that when the temperature rises the peak heights around the Fermi surface are reduced with reduced gap while closer to the center the thermal effects are relatively weaker.

Fig. [5](#page-16-0) demonstrates how the temperature contributes to the gapless BP-state superfluid

FIG. 4: [Color online] Figure (a) shows a cut of the BCS-state density in the $z = 0$ plane when the temperature is $k_BT/(2J) = 0.56$ (just below T_c) while figure (b) shows a cut of the normal-state density in the $z = 0$ plane when the temperature is $k_BT/(2J) = 0.58$ (just above T_c). Figures (c) and (d) show the integrated BCS-state noise correlations between components at temperatures $k_BT/(2J) = 0.0$ and $k_BT/(2J) = 0.56$, respectively. The other parameters used were $n_† = n_{\perp} =$ 0.30, $U/(2J) = 3.0$, in figures (a) and (d) $\Delta/(2J) = 0.10$ and in figure (c) $\Delta/(2J) = 1.03$. Colorcoding is again such that warm colors imply high peaks and cold colors low.

noise correlations. As one can see, when temperature rises the peakless areas become smaller and eventually vanish. The reason for this is in the temperature fluctuations. These figures show also that the peakless areas vanish only gradually, when the temperature becomes non-zero.

Finally, Fig. [6](#page-17-0) demonstrates how temperature influences the noise correlations of the FFLO state. The states correspond to minima of the free energy at their respective temperatures. Figs. [6](#page-17-0) (a) and (b) show an example of what happens to the correlation between components on the $z = 0$ plane when temperature rises from $k_BT/(2J) = 0.0475$ to $k_BT/(2J) = 0.095$. Figs. [6](#page-17-0) (c) and (d) demonstrate the same for the column integrated noise correlations. As one can see from the figures, sharp areas without correlation peaks

FIG. 5: [Color online] This figure demonstrates how temperature contributes to the noise correlations of the gapless BP-state. In these figures the temperature gradually increases, while other things remain equal. The last figure corresponds to the minimum of the grand potential with polarization $P = (n_1 - n_1)/(n_1 + n_1) = 0.40$, average filling fraction $(n_1 + n_1)/2 = 0.30$, $\Delta/(2J) = 0.318$ and coupling strength $U/(2J) = -3.0$. We have chosen $\mathbf{r} + \mathbf{r}' = 0$. All figures show the cut in the $z = 0$ plane. Color-coding is such that warm colors imply high peaks and cold colors low.

again disappear with increasing temperatures. However, the shift in the positions of the peaks persists even at non-zero temperatures.

It might not be easy to detect the gapless states via noise correlations because at the temperature regime where the gapless BP-state becomes the lowest energy state, the noise correlations of the gapless BP-state and non-gapless BP-state appear quite similar. On the other hand, by monitoring the peak heights at $z = 0$ (and $\mathbf{r}' = -\mathbf{r}$), one can identify the gapless states. The reason for this is that in the gapless BP-state the maximum peak height is always smaller than $A(t)^4/8$ while in the-non gapless state the maximum is always bigger than this. This can be seen from the fact that the term $|u_{\mathbf{k}}|^2 |v_{\mathbf{k}}|^2$ peaks on the Fermi surface corresponding to the average chemical potential. On the other hand, near this surface the

FIG. 6: [Color online] Finite temperature noise correlations $(G_{\uparrow\downarrow})$ of the FFLO state. Figures (a) and (b) show cuts in the $z = 0$ plane, while figures (c) and (d) are the column integrated correlations. Figures (a) and (c) were calculated at $k_BT/(2J) = 0.0475$ and figures (b) and (d) were calculated $k_BT/(2J) = 0.095$. In all figures we used the coupling strength $U/(2J) = -1.86$. In figures (a) and (c) $P = 0.208$, $(n_1 + n_1)/2 = 0.46$, $q_x = 0.26(\pi/d)$, $q_y = q_z = 0$, and $\Delta/(2J) =$ 0.14. In figures (b) and (d) $P = 0.24$, $(n_1 + n_1)/2 = 0.37$, $q_x = 0.21(\pi/d)$, $q_y = q_z = 0$, and $\Delta/(2J) = 0.046$. We have chosen $\mathbf{r} + \mathbf{r}' - (\hbar t/m)\mathbf{q} = 0$. Color-coding is again such that warm colors imply high peaks and cold colors low.

peak heights are

$$
|u_{\mathbf{k}}|^2 |v_{\mathbf{k}}|^2 [f(E_{\uparrow,\mathbf{k}}) f(E_{\downarrow,\mathbf{k}}) + (1 - f(E_{\uparrow,\mathbf{k}})) (1 - f(E_{\downarrow,\mathbf{k}}))]
$$

\n
$$
\approx \frac{1}{4} [f(\Delta + \delta \mu/2) f(\Delta - \delta \mu/2) + (1 - f(\Delta - \delta \mu/2)) (1 - f(\Delta + \delta \mu/2))],
$$

where $\delta \mu = \mu_{\uparrow} - \mu_{\downarrow}$ and then the term in the square brackets is bigger than 1/2, when $\Delta \pm \delta \mu/2 > 0$ (and the state is gapped) and otherwise always lower than 1/2.

In Fig. [7](#page-18-1) we show the height of the maximum of peak height as a function of the polar-ization (Fig. [7](#page-18-1) (a)) and as a function of chemical potential difference (Fig. 7 (b)). Fig. 7 (a)

FIG. 7: [Color online] This figure demonstrates how the polarization contributes to the maximum peak height. (a) shows the maximum peak height as a function of the polarization while (b) shows the maximum peak height as a function of the chemical potential difference $(\mu_{\uparrow} - \mu_{\downarrow})/(2\Delta)$ $\delta\mu/(2\Delta)$. We used the parameters $k_BT/(2J) = 0.45$ and an average filling fraction $(n_T + n_L)/2 =$ 0.30. We have canceled the time dependence of the maximum peak height by dividing the maximum peak height by the scaling factor $A(t)^4$. The horizontal lines show value $1/8$.

shows that when the polarization P rises then the maximum peak height becomes smaller and drops suddenly to zero when the gas becomes normal. With our parameters this happens when the polarization is about 0.35. By inspecting Fig. [7](#page-18-1) (b) it is clear that when the BP-state is gapless, i. e. when $\delta \mu > 2\Delta$, the maximum peak height is indeed smaller than 1/8, a result which was made plausible above.

V. NOISE CORRELATIONS IN A ONE-DIMENSIONAL LATTICE

We call the lattice one-dimensional (1D) when $J_x \gg J_y = J_z$. In other words a 1D lattice can be realized with three dimensional (3D) lattice with a very strong confinement in two directions and weaker confinement in the third one. Since the FFLO type states are more favorable in 1D than they are in 3D [\[30,](#page-23-1) [31](#page-23-2), [32\]](#page-23-3) and the differences between different phases can be more pronounced in 1D than in 3D, we now consider the noise correlations in 1D lattices. We assume that the gas in a 1D lattice is released only one direction (in our case x-direction). The released gas creates a set of almost identical one dimensional tubes.

Fig. [8](#page-19-1) shows an example of the phase diagram in a 1D lattice [\[32](#page-23-3)]. It is clear that the FFLO region is remarkably large and occupies the entire superfluid region at $T = 0$. On the other hand the expected region for phase separation between the normal gas and the BCSstate (red) is dramatically smaller than in 3D [\[21](#page-22-20)]. When the temperature rises there are phase transitions between the FFLO-state and, depending on polarization, phase separation, the BCS/BP states, or the normal gas phases. On Fig [8](#page-19-1) we also denote by dashed lines those values of temperature and polarization we have used in our following noise correlation computations.

Fig. [9](#page-20-0) demonstrates how temperature contributes to the noise correlations in a 1D lattice. As is clear, when the temperature becomes non-zero the gapless region does not vanish suddenly. Also, there is a phase transition between the FFLO-state and the gap-

FIG. 8: [Color online] The phase diagram of Fermi gas in an effectively 1D-lattice [\[32\]](#page-23-3). Colors: BCS/PB states=blue, FFLO=yellow, phase separation=red, and normal gas=white. The parameters where such that the average filling fraction $(n_{\uparrow}+n_{\downarrow})/2=0.2$, $J_x=0.07E_r$, and $U=-0.2E_r$. The dashed lines show the values $P = 0.40$ and $k_BT/(2J_x) = 0.104$ used in Fig. [9](#page-20-0) and in Fig. [10](#page-21-0)

less BP-state when the temperature is (with our parameters) between $k_BT/(2J_x) = 0.095$ and $k_BT/(2J_x) = 0.104$. One can identify this phase transition from the fact that in the FFLO-state the noise correlation is not symmetric with respect to $x = 0$, while the gapless BP-state noise correlation is symmetric. For the two-mode FFLO-state we expect a mirror image of the FFLO-state noise correlation shown here to appear. This mirror image would correspond to correlations with the pair momenta $-q_x$. The figure which has been calculated at highest temperature $(k_BT/(2J_x) = 0.104)$ shows that the gapless BP-state noise correlation is again always below 1/8.

In Fig. [10](#page-21-0) we show how the polarization contributes to the noise correlations in a 1D lattice. When the polarization is between $P = 0.32$ and $P = 0.35$ the non-gapless state becomes the gapless and as one can see that the maximum of the gapped BP-state noise correlation is always bigger than $1/8$. When the polarization is between $P = 0.35$ to $P = 0.48$, there is a phase transition from the BP-superfluid state to the FFLO-state which is visible in the pronounced asymmetry.

VI. CONCLUSIONS

In this paper we have presented, at the mean field level, the noise correlations of the two component Fermi gas in an optical lattice. We have shown that the noise correlations are a promising way to detect different phases in optical lattices. The different superfluid phases (BCS, FFLO, and BP) can be distinguished via the noise correlations and by mapping the correlations more extensively, a "Fourier analysis" of the multimode FFLO-state is possible and could be used to reveal the structure of the more complicated periodic order parameters. We computed the noise correlations also at non-zero temperatures and demonstrated that the differences between the correlations of different states can persist at finite temperatures and some qualitative features are insensitive to temperature. Also, regions of gapless quasiparticle dispersions can be visible in the noise correlation signals.

Density-density correlations of a one-dimensional system were also discussed quite ex-

FIG. 9: [Color online] This figure demonstrates how temperature contributes to the noise correlations in a 1D lattice. In these figures we gradually increase the temperature while keeping a polarization $P = 0.40$ and an average filling fraction $(n_1 + n_1)/2 = 0.20$ fixed. Figures in the upper row and bottom left show the FFLO-state noise correlations. The figure in bottom right shows the gapless BP-state noise correlation. In the first figure the dashed lines show the gapless region and in the bottom right figure the dashed line shows the value 1/8. We choose $x + x' + \hbar t q_x/m = 0$ in all these figures. In the BP-state $q_x = 0$.

tensively and were shown to contain very clear information on the structure of the pairwavefunction as well as on the quasi-particle dispersions. In one-dimensional systems, order parameter modulations are favored and possible problems with signals becoming smoothed out by column integration can be avoided. For this reason they are very attractive candidates to observe FFLO-type states experimentally. However, how to balance the requirements of sufficiently reduced dimensionality with sufficient phase coherence [\[31](#page-23-2)] is to a large extent still an open question.

It is physically possible that in a three-dimensional system a phase-separation occurs between the normal state and the BCS-state if the system is polarized [\[39](#page-23-10), [40](#page-23-11), [41](#page-23-12)]. This means that the normal state exists in one part of the lattice and the BCS-state exists in

FIG. 10: [Color online] This figure shows how polarization contributes to the noise correlations in a 1D lattice. Here we gradually increase the polarization while keeping the temperature fixed at $k_BT/(2J_x) = 0.104$ and an average filling fraction at $(n_† + n_{\downarrow})/2 = 0.20$. The upper left figure shows the BCS-state noise correlation and upper right figure shows the gapped BP-state noise correlations. The bottom left figure shows the gapless BP-state noise correlation and the figure on the bottom right is the FFLO-state noise correlation. The dashed line shows the value 1/8. We choose $x + x' + \hbar t q_x/m = 0$ in all these figures. In the BCS- and the BP-states $q_x = 0$.

another part of the lattice, typically in the core if a harmonic trap potential is included. However, other more exotic possibilities also exist [\[32\]](#page-23-3). This phase separation can be seen via the noise correlations. Because the noise correlation of the normal state is identically zero, correlation peaks can be seen only in parts of the freely expanded cloud. If the BCS-state is in the core, the peaks can be seen at the center of the cloud while closer to the edges of the cloud the peaks disappear even if the background density is still non-zero.

Acknowledgements This work was supported by the National Graduate School in Materials Physics, Academy of Finland (Projects Nos. 115020, 213362, 121157, 207083) and conducted as a part of a EURYI scheme award. See www.esf.org/euryi.

- [1] M. W. Zwierlein, A. Schirotzek, C. H. Schunck, and W. Ketterle, Science 311, 492 (2006).
- [2] G. B. Partridge, W. Li, R. I. Kamar, Y. Liao, and R. G. Hulet, Science 311, 506 (2006).
- [3] M. W. Zwierlein, C. H. Schunck, A. Schirotzek, and W. Ketterle, Nature 442, 54 (2006).
- [4] Y. Shin, M. W. Zwierlein, C. H. Schunck, A. Schirotzek, and W. Ketterle, Phys. Rev. Lett. 97, 030401 (2006).
- [5] G. B. Partridge, W. Li, Y. Liao, R. G. Hulet, M. Haque, and H. T. C. Stoof, Phys. Rev. Lett. 97, 190407 (2006).
- [6] P. Fulde and R. A. Ferrell, Phys. Rev. 135, A550 (1964).
- [7] A. I. Larkin and Y. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. 47, 1136 (1964).
- [8] G. Sarma, J. Phys. Chem. Solids 24, 1029 (1963).
- [9] W. V. Liu and F. Wilczek, Phys. Rev. Lett. 90, 047002 (2003).
- [10] R. Casalbuoni and G. Nardulli, Rev. Mod. Phys. 76, 263 (2004).
- [11] C. Orzel, A. K. Tuchman, M. L. Fenselau, M. Yasuda, and M. A. Kasevich, Science 291, 2386 (2001).
- [12] M. Greiner, I. Bloch, O. Mandel, T. W. Hänsch, and T. Esslinger, Phys. Rev. Lett. 87, 160405 (2001).
- [13] S. Burger, F. S. Cataliotti, C. Fort, F. Minardi, M. Inguscio, M. L. Chiofalo, and M. P. Tosi, Phys. Rev. Lett. 86, 4447 (2001).
- [14] Z. Hadzibabic, S. Stock, B. Battelier, V. Bretin, and J. Dalibard, Phys. Rev. Lett. 93, 180403 (2004).
- [15] W. Hofstetter, J. I. Cirac, P. Zoller, E. Demler, and M. D. Lukin, Phys. Rev. Lett. 89, 220407 (2002).
- [16] G. Orso and G. Shlyapnikov, Phys. Rev. Lett. 95, 260402 (2005).
- [17] L. P. Pitaevskii, S. Stringari, and G. Orso, Phys. Rev. A 71, 053602 (2005).
- [18] M. Iskin and C. A. R. S. de Melo, Phys. Rev. Lett. 99, 080403 (2007).
- [19] T. Koponen, J.-P. Martikainen, J. Kinnunen, and P. Törmä, Phys. Rev. A **73**, 033620 (2006).
- [20] T. Koponen, J. Kinnunen, J.-P. Martikainen, L. M. Jensen, and P. Törmä, New J. Phys. 8, 179 (2006).
- [21] T. K. Koponen, T. Paananen, J.-P. Martikainen, and P. Törmä, Phys. Rev. Lett. 99, 120403 (2007).
- [22] J. K. Chin, D. E. Miller, Y. Liu, C. Stan, W. Setiawan, C. Sanner, K. Xu, and W. Ketterle, Nature 443, 961 (2006).
- [23] M. Köhl, H. Moritz, T. Stöferle, K. Günter, and T. Esslinger, Phys. Rev. Lett. 94, 080403 $(2005).$
- [24] T. Stöferle, H. Moritz, K. Günter, M. Köhl, and T. Esslinger, Phys. Rev. Lett. $96, 030401$ (2005).
- [25] K. Günter, T. Stöferle, H. Moritz, M. Köhl, and T. Esslinger, Phys. Rev. Lett. 95, 230401 (2005).
- [26] E. Altman, E. Demler, and M. D. Lukin, Phys. Rev. A 70, 013603 (2004).
- [27] S. Foelling, F. Gerbier, A. Widera, O. Mandel, T. Gericke, and I. Bloch, Nature 434, 481 (2005).
- [28] T. Rom, T. Best, D. van Oosten, U. Schneider, S. Foelling, B. Paredes, and I. Bloch, Nature

444, 733 (2006).

- [29] M. Greiner, C. A. Regal, J. T. Stewart, and D. S. Jin, Phys. Rev. Lett. **94**, 110401 (2005).
- [30] X.-J. Liu, H. Hu, and P. D. Drummond, Phys. Rev. A 76, 043605 (2007).
- [31] M. M. Parish, S. K. Baur, E. J. Mueller, and D. A. Huse (2007), arXiv/0709.1120.
- [32] T. K. Koponen, T. Paananen, J.-P. Martikainen, M. R. Bakhtiari, and P. Törmä (2007), arXiv/0711.4065v1.
- [33] A. Luscher, R. M. Noack, and A. Laeuchli (2007), arXiv/0712.1808v2.
- [34] I. Bloch, J. Dalibard, and W. Zwerger (2007), arxiv/0704.3011.
- [35] D. Jaksch, C. Bruder, J. Cirac, C. W. Gardiner, and P. Zoller, Phys. Rev. Lett. 81, 3108 (1998).
- [36] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. 108, 1175 (1957).
- [37] D. E. Sheehy and L. Radzihovsky, Annals of Physics 322, 1790 (2007).
- [38] B. M. Andersen and G. M. Bruun, Phys. Rev. A 76, 041602(R) (2007).
- [39] P. F. Bedaque, H. Caldas, and G. Rupak, Phys. Rev. Lett. 91, 247002 (2003).
- [40] J.-P. Martikainen, Phys. Rev. A 74, 013602 (2006).
- [41] M. Haque and H. Stoof, Phys. Rev. A **74**, 011602(R) (2006).