

Operator Coupling of Gauge Fields and Unparticles

A. Lewis Licht
Dept. of Physics
U. of Illinois at Chicago
Chicago, Illinois 60607
licht@uic.edu

We show that it is possible to couple gauge fields to unparticles without the use of path integrals in the unparticle effective action. This is done by treating the unparticle field as a vector in an abstract Hilbert space, and the gauge field as a linear operator on that space.

INTRODUCTION

Terning et al [1] have used a path integral in Georgi's [2] [3] unparticle effective action to couple gauge fields to the unparticles. We have shown in a related work [4] that if done rigorously, this leads to a formula for the gauge-unparticle vertex that is much more complicated than that found in Ref. [1]. Here we show that there is an operator formulism, not involving path integrals, that does lead to Terning et al's result.

OPERATOR POWERS AS BRANCH CUT INTEGRALS

The unparticle formulism relies very heavily on the representation of operator powers as branch cut integrals. Given the complex function $f(z) = z^n$, we can express it as a contour integral,

$$z^n = \frac{1}{2\pi i} \oint dz' \frac{z'^n}{z' - z} \quad (1)$$

If $-1 < n < 0$ we can expand the contour to wrap around the positive x-axis and we can discard the loop integral at infinity. Then

$$z^n = -\frac{e^{\pi i n}}{\pi} \sin(\pi n) \int_0^\infty dx \frac{x^n}{x - z} \quad (2)$$

If n is not in this range, say $n = k - \delta$, where k is an integer and δ is in the range, one could still use the branch integral to express $z^{-\delta}$, and just multiply by z^k . In many cases it should be possible to start with n in the negative unit range, do the branch cut integral, and analytically continue to the desired value of n .

Georgi [2] [3] has for the propagator of his unparticle field;

$$\begin{aligned} \langle 0 | T (\Phi_u^\dagger(x) \Phi_u(0)) | 0 \rangle &= S(x, 0) \\ &= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \frac{A_{du}}{2\pi i} \int_0^\infty dM^2 \frac{(M^2)^{d_u-2}}{M^2 - p^2 - i\varepsilon} \\ &= \int \frac{d^4 p}{(2\pi)^4} e^{-ipx} \frac{A_{du} e^{-\pi i d_u}}{-2i \sin(\pi d_u)} (p^2 + i\varepsilon)^{d_u-2} \end{aligned} \quad (3)$$

Where

$$A_{du} = \frac{16\pi^{5/2}}{(2\pi)^{2d_u}} \frac{\Gamma(d_u + 1/2)}{\Gamma(d_u - 1) \Gamma(2d_u)} \quad (4)$$

The quadratic part of the unparticle action integral is then nonlocal,

$$I = \int d^4 x d^4 y \Phi_u^\dagger(x) K(x, y) \Phi_u(y) \quad (5)$$

Where K and S are inverse operators,

$$\int d^4 z K(x, z) S(z, y) = i\delta^4(x - y) \quad (6)$$

The Kernel K can be expressed as:

$$K(x, y) = \frac{2 \sin(\pi d_u) e^{\pi i d_u}}{A_{d_u}} \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} (p^2 + i\varepsilon)^{2-d_u} \quad (7)$$

We can express K in terms of a branch cut integral

$$K(x, y) = \frac{2 \sin^2(\pi d_u)}{\pi A_{d_u}} \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \int_0^\infty dM^2 \frac{(M^2)^{2-d_u}}{M^2 - p^2 - i\varepsilon} \quad (8)$$

Where we have assumed an analytical continuation to the appropriate value of d_u .

THE OPERATOR FORMULISM

We wish to have the action invariant under the gauge transformation:

$$\begin{aligned} \Phi_u(x) &\rightarrow e^{ig\Lambda(x)} \Phi_u(x) \\ A_\mu(x) &\rightarrow A_\mu(x) + \partial_\mu \Lambda(x) \end{aligned} \quad (9)$$

We do this by writing the action integral as an inner product in a Hilbert space, and formulating derivatives and functional multiplication as operators on that Hilbert space.

We consider the functions $\Phi_u(x)$ as corresponding to Hilbert space vectors,

$$\Phi_u(x) = \langle x | \Phi_u \rangle \quad (10)$$

where the $|x\rangle$ are eigenkets of a position operator:

$$\begin{aligned} X^\mu |x\rangle &= x^\mu |x\rangle \\ \langle x | x' \rangle &= \delta^4(x - x') \end{aligned} \quad (11)$$

We define a momentum operator P_ν so that

$$[X^\mu, P_\nu] = -i\delta_\nu^\mu \quad (12)$$

The eigenkets of the momentum operator satisfy

$$\begin{aligned} P_\nu |p_\nu\rangle &= p_\nu |p\rangle \\ \langle p | p' \rangle &= \delta^4(p - p') \\ \langle x | p \rangle &= \frac{e^{-ipx}}{(2\pi)^2} \end{aligned} \quad (13)$$

It should be noted that X^μ represents an abstract point in 4-dimensional Minkowski space, and P_ν is the translation operator for such points, They do not refer to the position of an actual particle.

Let

$$K_0 = \frac{2 \sin^2(\pi d_u)}{\pi A_{d_u}} \quad (14)$$

The action can now be expressed as

$$I = \langle \Phi_u | K | \Phi_u \rangle \quad (15)$$

where the operator K is

$$K = K_0 \int_0^\infty dM^2 \frac{(M^2)^{2-d_u}}{M^2 - P^2 - i\varepsilon} \quad (16)$$

The gauge transformation can now be expressed as a transformation of vectors and operators:

$$\begin{aligned} |\Phi_u\rangle &\rightarrow e^{ig\Lambda} |\Phi_u\rangle \\ A_\mu &\rightarrow A_\mu - i [P_\mu, \Lambda] \end{aligned} \quad (17)$$

Here we have

$$\begin{aligned} \Lambda &= \int d^4x |x\rangle \Lambda(x) \langle x| \\ A_\mu &= \int d^4x |x\rangle A_\mu(x) \langle x| \\ -i [P_\mu, \Lambda] &= \int d^4x |x\rangle \partial_\mu \Lambda(x) \langle x| \end{aligned} \quad (18)$$

The combination

$$D_\mu = P_\mu - gA_\mu \quad (19)$$

is then invariant under the gauge transformation:

$$D_\mu = e^{-ig\Lambda} D_\mu e^{+ig\Lambda} \quad (20)$$

Replacing P_μ in K by D_μ then gives us a gauge invariant action.

Let

$$G_M(p^2) = \frac{1}{M^2 - P^2 - i\varepsilon} \quad (21)$$

To first order in g, the action is

$$I = I_0 - K_0 g \int_0^\infty dM^2 (M^2)^{2-d_u} \langle \Phi_u | \frac{1}{M^2 - P^2 - i\varepsilon} \{A_\mu, P^\mu\} \frac{1}{M^2 - P^2 - i\varepsilon} | \Phi_u \rangle \quad (22)$$

It is most convenient to evaluate this in momentum space. With

$$\begin{aligned} A_\mu &= \int d^4k d^4l d^4x |k\rangle \langle k|x\rangle A_\mu(x) \langle x|l\rangle \langle l| \\ &= \int d^4l d^4q |l+q\rangle A_\mu(q) \langle l| \end{aligned} \quad (23)$$

where

$$A_\mu(q) = \int \frac{d^4x}{(2\pi)^4} e^{iqx} A_\mu(x) \quad (24)$$

and with also

$$\Phi_u(p) = \int \frac{d^4x}{(2\pi)^4} \Phi_u(x) \quad (25)$$

so that

$$|\Phi_u\rangle = (2\pi)^2 \int d^4p |p\rangle \Phi_u(p) \quad (26)$$

then

$$I_1 = -K_0 g \int_0^\infty dM^2 (M^2)^{2-d_u} \int d^4p' d^4q d^4p (2\pi)^4 \delta(p' - p - q) \Phi_u^*(p') D_M(p') A_\mu(q) (p'^\mu + p^\mu) D_M(p) \Phi_u(p) \quad (27)$$

Defining the vertex function as

$$ig\Gamma^\mu(p', q, p) (2\pi)^4 \delta^4(p' - q - p) = \frac{i\delta^3 I}{\delta\Phi_u^\dagger(p') \delta A_\mu(q) \delta\Phi_u(p)} \quad (28)$$

We get

$$\begin{aligned}
 i\Gamma^\mu &= -i \frac{2 \sin(\pi d_u)}{A_{du}} e^{i\pi d_u} \frac{p'^\mu + p^\mu}{p'^2 - p^2} \left((p'^2)^{2-d_u} - (p^2)^{2-d_u} \right) \\
 &= \frac{2p^\mu + q^\mu}{2p \cdot q + q^2} [S^{-1}(p') - S^{-1}(p)]
 \end{aligned} \tag{29}$$

The same as Terning et al's result. [1]

CONCLUSIONS

We have found an operator method that can be used to introduce gauge fields into the unparticle effective action. This method yields results for the vertex that are considerably simpler than those obtained by rigorous application of the path integral method and are the same as those obtained from the looser path integral method of Terning et al. [1] We would like to point out that there are however ordering problems with the operator method. An action of the form

$$I = \int d^4x d^4y \partial_{x\mu} \Phi_u^\dagger(x) K(x, y) \partial_y^\mu \Phi_u(y) \tag{30}$$

where K here involves one less power of P^2 , can be shown to yield a quite different vertex.

ACKNOWLEDGEMENTS

I would like to express my thanks to Wai-Yee Keung for interesting me in this problem.

-
- [1] Giacomo Cacciapaglia, Guido Marandella and John Terning, *Colored Unparticles*, arXiv:0708.005 [hep-ph].
 - [2] Howard Georgi, *Unparticle Physics*, arXiv:hep-ph/0703260.
 - [3] Howard Georgi, *Another Odd Thing About Unparticle Physics* arXiv:0704.2457 [hep-ph].
 - [4] A. Lewis Licht, *Gauge Fields and Unparticles* arXiv:0801.0892 [hep-th]