Quasi-orthogonal subalgebras of matrix algebras@

Hiromichi Ohno

1 Introduction

In the theory of quantum mechanics, an *n*-level system is described by the algebra $M_n = M_n(\mathbb{C})$ of $n \times n$ complex matrices. The matrix algebra of a composite system consisting of an *n*-level and an *m*-level system is $M_n \otimes M_m \simeq M_{nm}$. A subalgebra of M_k corresponds to a subsystem of a *k*-level quantum system. In this paper, subalgebras contain the identity and are closed under the adjoint operation of matrices, that is, they are unital *-subalgebras. The algebra M_n can be endowed by the inner product $\langle A, B \rangle = \text{Tr}(A^*B)$ and it becomes a Hilbert space. Two subalgebras \mathcal{A}_1 and \mathcal{A}_2 are called quasi-orthogonal if $\mathcal{A}_1 \oplus \mathbb{C}I \perp \mathcal{A}_2 \oplus \mathbb{C}I$.

We consider pairwise quasi-orthogonal subalgebras $\mathcal{A}_1, \mathcal{A}_2, \ldots, \mathcal{A}_l$ in $M_{p^{kn}}$ which are isomorphic to M_{p^k} for $k \ge 1$, $n \ge 2$ and a prime number p with $p \ge 3$. The aim of this paper is to obtain the maximum l. The case p = 2, n = 2 and k = 1 is shown in [3, 6, 8] and the maximum is 4.

The motivations of this problem are followings. If a total system $M_n \otimes M_m$ has a statistical operator ρ , we can reconstruct the reduced density $\rho_n^{(1)} = \text{Tr}_m \rho$ in the subsystem M_n , where Tr_m is a partial trace onto M_n . In order to get more information, we change the density ρ by an interaction. For a Hamiltonian H, the new state is

$$e^{iH}\rho e^{-iH} = W_1\rho W_1^*$$

after the interaction. The new reduced density is $\rho_n^{(2)} = \text{Tr}_m W_1 \rho W_1^*$. By using other interactions, we have a sequence of reduced states $\rho_n^{(1)}, \rho_n^{(2)}, \ldots, \rho_n^{(k)}$. We want to determine the minimum k such that this sequence of reduced densities determines ρ .

Another reason comes from the relation to the mutually unbiased bases (MUB) problem. Given a orthonormal basis of an *n*-dimensional Hilbert space \mathcal{H} , the linear operators diagonal in this basis form a maximal Abelian subalgebra of $M_n \simeq \mathcal{B}(\mathcal{H})$. Conversely if $|e_i\rangle\langle e_i|$ are minimal projections in a maximal Abelian subalgebra, then $(|e_i\rangle)_i$ is a orthonormal basis. Two maximal Abelian subalgebras of M_n are quasiorthogonal if and only if the two corresponding bases (ξ_i) and (ζ_j) are mutually unbiased (see Proposition 2.2), that is,

$$|\langle \xi_i, \zeta_j \rangle|^2 = \frac{1}{n}$$
 $(1 \le i, j \le n).$

Mutually unbiased bases are very interesting from many point of view and the maximal number of such bases is not known for arbitrary n. We want to study mutually unbiased bases in terms of quasi-orthogonal subalgebras (see [1, 2, 4]).

In Section 2, we consider quasi-orthogonal subalgebras in M_{p^n} which are isomorphic to M_p . In Section 3, quasi-orthogonal subalgebras in $M_{p^{k_n}}$ which are isomorphic to M_{p^k} are investigated.

2 Quasi-orthogonal subalgebras in M_{p^n}

 \mathcal{A} is a finite dimensional C^* -algebra with the usual trace Tr and is considered as a Hilbert space under the inner product

$$\langle A, B \rangle = \operatorname{Tr}(A^*B)$$

for any $A, B \in \mathcal{A}$.

Definition 2.1 Two subalgebras \mathcal{A}_1 and \mathcal{A}_2 of \mathcal{A} are called *quasi-orthogonal* if

$$\mathcal{A}_1 \ominus \mathbb{C}I \perp \mathcal{A}_2 \ominus \mathbb{C}I.$$

The equivalent conditions of this definition are followings [5]: (a) For any $A_1 \in \mathcal{A}_1$ and $A_2 \in \mathcal{A}_2$,

$$\operatorname{Tr}(A_1 A_2) = \frac{\operatorname{Tr}(A_1) \operatorname{Tr}(A_2)}{\operatorname{Tr}(I)}.$$

(b) For any $A_1 \in \mathcal{A}_1$ and $A_2 \in \mathcal{A}_2$ with $\operatorname{Tr}(A_1) = \operatorname{Tr}(A_2) = 0$,

 $\operatorname{Tr}(A_1 A_2) = 0.$

The theory of quasi-orthogonal subalgebras is related to the theory of mutually unbiased bases. This is stated as follows:

Proposition 2.2 The maximal number of mutually unbiased bases in \mathbb{C}^d is equal to the maximal number of pairwise quasi-orthogonal subalgebras in M_d which are isomorphic to \mathbb{C}^d .

Proof. For orthonormal bases $\{\xi_i\}_{1\leq i\leq d}$ and $\{\zeta_j\}_{1\leq j\leq d}$ of \mathbb{C}^d , define minimal projections in M_d by $P_i = |\xi_i\rangle\langle\xi_i|$ and $Q_j = |\zeta_j\rangle\langle\zeta_j|$. Since $\operatorname{Tr}(P_iQ_j) = |\langle\xi_i,\zeta_j\rangle|^2$, the condition

$$|\langle \xi_i, \zeta_j \rangle|^2 = \frac{1}{d}$$
 $(1 \le i, j \le d)$

is equivalent to the condition

$$\operatorname{Tr}(P_i Q_j) = \frac{1}{d} = \frac{\operatorname{Tr}(P_i) \operatorname{Tr}(Q_j)}{\operatorname{Tr}(I)} \qquad (1 \le i, j \le d).$$

From (a), we conclude that $\{\xi_i\}$ and $\{\zeta_j\}$ are mutually unbiased bases if and only if the algebras generated by $\{P_i\}$ and $\{Q_j\}$ are quasi-orthogonal. This prove the assertion.

It is known that there exist d + 1 mutually unbiased bases in \mathbb{C}^d if d is a power of a prime number. Therefore there are d + 1 quasi-orthogonal subalgebras in M_d which are isomorphic to \mathbb{C}^d if d is a power of a prime number.

Now we consider the quasi-orthogonal subalgebras in $M_{p^2} = M_p \otimes M_p$ which are isomorphic to M_p , where p is a prime number with $p \ge 3$. We will show that we can construct $p^2 + 1$ pairwise quasi-orthogonal subalgebras.

Define the unitary operators W and S in M_p by

$$W = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda & 0 & \cdots & 0 \\ 0 & 0 & \lambda^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda^{p-1} \end{bmatrix}, \qquad S = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix},$$

where $\lambda = e^{2\pi i/p}$. Then these two unitary operators have following properties:

- i. $S^p = W^p = I$.
- ii. The set $\{S^i W^j\}_{0 \le i,j \le p-1}$ is a natural orthogonal basis of M_p .
- iii. Since $SW = \lambda^{-1}WS$, we have

$$S^{k_1}W^{l_1}S^{k_2}W^{l_2} = \lambda^{k_2l_1}S^{k_1+k_2}W^{l_1+l_2}.$$
(1)

iv. From (1), $S^{k_1}W^{l_1}$ and $S^{k_2}W^{l_2}$ commute if and only if $k_1l_2 = k_2l_1 \mod p$.

We consider the commutativity condition iv in the context of a vector space over Z_p , where Z_p is a finite field with p elements. Let $Z_p^4 = \{(k_1, l_1, k_2, l_2) | k_1, l_1, k_2, l_2 \in Z_p\}$ be a vector space over Z_p and define a natural homomorphism π (up to scalar multiple) from Z_p^4 to $M_p \otimes M_p$ by

$$\pi(k_1, l_1, k_2, l_2) = S^{k_1} W^{l_1} \otimes S^{k_2} W^{l_2}.$$

We denote a *symplectic product* by

$$u \circ u' = k_1 l'_1 - k'_1 l_1 + k_2 l'_2 - k'_2 l_2 \mod p,$$

where $u = (k_1, l_1, k_2, l_2)$ and $u' = (k'_1, l'_1, k'_2, l'_2)$. From (1),

$$\pi(u)\pi(u') = \lambda^{-u \circ u'}\pi(u')\pi(u).$$
(2)

Hence $\pi(u)$ and $\pi(u')$ commute if and only if their symplectic product equals zero.

Lemma 2.3 If $\pi(u)$ and $\pi(u')$ are not commutative for $u = (k_1, l_1, k_2, l_2), u' = (k'_1, l'_1, k'_2, l'_2) \in Z_p^4$, then the algebra \mathcal{A} generated by $\pi(u)$ and $\pi(u')$ is isomorphic to M_p .

Proof. From the assumption, $u \circ u' \neq 0$. Define a map ρ from $\{S, W^{u \circ u'}\}$ to \mathcal{A} by

$$\rho(S) = \pi(u), \qquad \rho(W^{u \circ u'}) = \pi(u').$$

From (2) and $SW^{u\circ u'} = \lambda^{-u\circ u'}W^{u\circ u'}S$, the commutativity condition of $\pi(u)$, $\pi(u')$ and that of S, $W^{u\circ u'}$ are same. Therefore ρ can be extended to an isomorphism from M_p generated by S and $W^{u\circ u'}$ to \mathcal{A} .

From this lemma, we need to find such u and u'. Let D be a non-zero integer in Z^p with the requirement that $D \neq k^2 \mod p$ for all k in Z_p , i.e., D is not a quadratic residue of p. For any $a_0, a_1 \in Z_p$, we define a subspace of Z_p^4 by

$$C_{a_0,a_1} = \{b_0(1, a_1, 0, a_0) + b_1(0, a_0, -1, a_1D) \mid b_0, b_1 \in \mathbb{Z}_p\},\$$

where scalar multiplication and addition are defined by a natural way. Moreover put

$$C_{\infty} = \{b_0(0, 1, 0, 0) + b_1(0, 0, 0, 1) \mid b_0, b_1 \in \mathbb{Z}_p\}.$$

Lemma 2.4 The only vector common to any pair of above subspaces is (0,0,0,0). In particular, the subspaces expect (0,0,0,0) partition $Z_p^4 \setminus \{(0,0,0,0)\}$.

Proof. Since there are p^2+1 subspaces and each subspace has p^2 elements, it is enough to prove that the intersection of any two subspaces is $\{(0,0,0,0)\}$. It is easy to see $C_{a_0,a_1} \cap C_{\infty} = \{(0,0,0,0)\}$. Therefore we prove that $C_{a_0,a_1} \cap C_{a'_0,a'_1} = \{(0,0,0,0)\}$ if $a_0 \neq a'_0$ or $a_1 \neq a'_1$.

Assume $b_0(1, a_1, 0, a_0) + b_1(0, a_0, -1, a_1D) = b'_0(1, a'_1, 0, a'_0) + b'_1(0, a'_0, -1, a'_1D)$, then from the first and third components we have $b_0 = b'_0$ and $b_1 = b'_1$. Similarly, from the second and fourth components we are led to the equations

$$a_1b_0 + a_0b_1 = a'_1b_0 + a'_0b_1$$

$$a_0b_0 + a_1b_1D = a'_0b_0 + a'_1b_1D.$$

These equations can be rewritten as a matrix equation

$$\begin{bmatrix} b_1 & b_0 \\ b_0 & b_1D \end{bmatrix} \begin{bmatrix} a_0 - a'_0 \\ a_1 - a'_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

If $b_0 = b_1 = 0$, then the common element is (0, 0, 0, 0). Therefore we assume $b_0 \neq 0$ or $b_1 \neq 0$. Then the above matrix is invertible, indeed

$$\begin{bmatrix} b_1 & b_0 \\ b_0 & b_1 D \end{bmatrix}^{-1} = (b_1^2 D - b_0^2)^{-1} \begin{bmatrix} b_1 D & -b_0 \\ -b_0 & b_1 \end{bmatrix}$$

Here we use that $b_1^2 D \neq b_0^2 \mod p$ from the assumption of D. This implies $a_0 = a'_0$ and $a_1 = a'_1$ which is a contradiction.

Since $(1, a_1, 0, a_0) \circ (0, a_0, -1, a_1 D) = 2a_0$, if $a_0 \neq 0$ then the algebra span $\{\pi(C_{a_0, a_1})\}$ generated by $\pi(1, a_1, 0, a_0)$ and $\pi(0, a_0, -1, a_1 D)$ is isomorphic to M_p by Lemma 2.3.

But if $a_0 = 0$, then the algebra is commutative and hence span{ $\pi(C_{a_0,a_1})$ } $\simeq \mathbb{C}^{p^2}$. Therefore we need to mix the subspaces with $a_0 = 0$.

For any $a \in Z_p$, define subspaces by

$$D_a = \{b_0(1, 1, -a, aD) + b_1(1, 2, -a, 2aD) | b_0, b_1 \in Z_p\}, D_{\infty} = \{b_0(0, 0, 1, 0) + b_1(0, 0, 0, 1) | b_0, b_1 \in Z_p\}.$$

Lemma 2.5 The only vector common to any pair of above subspaces is (0, 0, 0, 0). Moreover we have

$$\bigcup_{a\in Z_p} D_a \cup D_{\infty} = \bigcup_{a_1\in Z_p} C_{0,a_1} \cup C_{\infty}.$$

Proof. The proof of the first assertion is same as Lemma 2.4. To show the second assertion, it is enough to prove $D_a, D_{\infty} \subset \bigcup_{a_1 \in Z_p} C_{0,a_1} \cup C_{\infty}$. Indeed, since each subspaces has p^2 elements and the intersection is trivial, the numbers of elements in both sets are equal.

First consider a element $(0, 0, b_0, b_1)$ in D_{∞} . If $b_0 = 0$, then $(0, 0, 0, b_1) \in C_{\infty}$. If $b_0 \neq 0$, then

$$(0,0,b_0,b_1) = -b_0(0,0,-1,-b_0^{-1}b_1D^{-1}D) \in C_{0,-b_0^{-1}b_1D^{-1}}.$$

Hence $D_{\infty} \subset \bigcup_{a_1 \in \mathbb{Z}_p} C_{0,a_1} \cup C_{\infty}$. Next consider the element $b_0(1, 1, -a, aD) + b_1(1, 2, -a, 2aD)$ in D_a . If $b_0 + b_1 = 0$, then

$$b_0(1, 1, -a, aD) + b_1(1, 2, -a, 2aD) = (0, b_0 + 2b_1, 0, ab_0D + 2ab_1D) \in C_{\infty}.$$

If $b_0 + b_1 \neq 0$, then

$$b_0(1, 1, -a, aD) + b_1(1, 2, -a, 2aD)$$

= $(b_0 + b_1) (1, (b_0 + b_1)^{-1}(b_0 + 2b_1), 0, 0)$
+ $a(b_0 + b_1) (0, 0, -1, (b_0 + b_1)^{-1}(b_0 + 2b_1)D)$
 $\in C_{0,(b_0+b_1)^{-1}(b_0+2b_1)}.$

Therefore we obtain $D_a \subset \bigcup_{a_1 \in Z_n} C_{0,a_1} \cup C_{\infty}$.

Since $(1, 1, -a, aD) \circ (1, 2, -a, 2aD) = 1 - a^2D \neq 0$ by the assumption of D and $(0, 0, 1, 0) \circ (0, 0, 0, 1) = 1$, we obtain

$$\operatorname{span}\{\pi(D_a)\} \simeq M_p,$$
$$\operatorname{span}\{\pi(D_\infty)\} \simeq M_p$$

by Lemma 2.3. Consequently we have the next theorem.

Theorem 2.6 There are $p^2 + 1$ pairwise quasi-orthogonal subalgebras in M_{p^2} which are isomorphic to M_p .

Example 2.7 Consider a 9-level quantum system $M_3 \otimes M_3$. We list 10 pairwise quasi-orthogonal subalgebras in M_9 which are isomorphic to M_3 :

$$\begin{split} & \text{span}\{\pi(C_{1,0})\} = \text{span}\{I \otimes I, S \otimes W, S^2 \otimes W^2, W \otimes S^2, SW \otimes S^2W, \\ S^2W \otimes S^2W^2, W^2 \otimes S, SW^2 \otimes SW, S^2W^2 \otimes SW^2\}, \\ & \text{span}\{\pi(C_{1,1})\} = \text{span}\{I \otimes I, SW \otimes W, S^2W^2 \otimes W^2, W \otimes S^2W^2, SW^2 \otimes S^2, \\ S^2 \otimes S^2W, W^2 \otimes SW, S \otimes SW^2, S^2W \otimes S\}, \\ & \text{span}\{\pi(C_{1,2})\} = \text{span}\{I \otimes I, SW^2 \otimes W, S^2W \otimes W^2, W \otimes S^2W, S \otimes S^2W^2, \\ S^2W^2 \otimes S^2, W^2 \otimes SW^2, SW \otimes S, S^2 \otimes SW\}, \\ & \text{span}\{\pi(C_{2,0})\} = \text{span}\{I \otimes I, S \otimes W^2, S^2 \otimes W, W^2 \otimes S^2, SW^2 \otimes S^2W^2, \\ S^2W^2 \otimes S^2W, W \otimes S, SW \otimes SW^2, S^2W \otimes SW\}, \\ & \text{span}\{\pi(C_{2,1})\} = \text{span}\{I \otimes I, SW \otimes W^2, S^2W^2 \otimes W, W^2 \otimes S^2W^2, S \otimes S^2W, \\ S^2W \otimes S^2, W \otimes SW, SW^2 \otimes S, S^2 \otimes SW^2\}, \\ & \text{span}\{\pi(C_{2,2})\} = \text{span}\{I \otimes I, SW^2 \otimes W^2, S^2W \otimes W, W^2 \otimes S^2W, SS^2W \otimes S^2, \\ S^2 \otimes S^2W^2, W \otimes SW^2, S \otimes SW, S^2W^2 \otimes S^2W, SW \otimes S^2, \\ S^2 \otimes S^2W^2, W \otimes SW^2, S \otimes SW, S^2W^2 \otimes S^2W, SW \otimes S^2, \\ & S^2 \otimes S^2W^2, W \otimes SW^2, S \otimes SW, S^2W^2 \otimes S^2W, SW \otimes S^2, \\ & Span\{\pi(D_0)\} = \text{span}\{I \otimes I, SW \otimes I, S^2W^2 \otimes SW, SW^2 \otimes S^2W, S^2 \otimes S^2, \\ & W \otimes W^2, S^2W \otimes SW^2, W^2 \otimes W, SW^2 \otimes S^2W, S^2 \otimes S^2, \\ & W \otimes W^2, S^2W \otimes SW^2, W^2 \otimes W, SW^2 \otimes SW^2, S^2 \otimes S^2W, S^2 \otimes S^2, \\ & W \otimes W^2, S^2W \otimes SW^2, S^2W^2 \otimes SW, SW^2 \otimes S^2W, S^2 \otimes S^2, \\ & W \otimes W^2, S^2W \otimes SW^2, W^2 \otimes W, SW^2 \otimes S^2W^2, S^2 \otimes S^2W, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W^2, S^2W^2 \otimes SW^2, SW^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W^2, S^2W^2 \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W^2, S^2W^2 \otimes SW^2, SW^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, W^2 \otimes W^2, S \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S^2 \otimes SW^2, S^2 \otimes S^2, \\ & W \otimes W, S^2W \otimes S^2W, S^2W^2 \otimes SW^2, S^2 \otimes SW^$$

where we denote D = 2.

Next, we consider pairwise quasi-orthogonal subalgebras in M_{p^n} which are isomorphic to M_p . The dimensions of $M_{p^n} \oplus \mathbb{C}I$ and $M_p \oplus \mathbb{C}I$ are $p^{2n} - 1$ and $p^2 - 1$, respectively. Hence such subalgebras are at most $N_n := p^{2n} - 1/p^2 - 1$. We construct N_n pairwise quasi-orthogonal subalgebras.

Theorem 2.8 M_{p^n} contains N_n pairwise quasi-orthogonal subalgebras which are isomorphic to M_p .

Proof. The case n = 1 is trivial and the case n = 2 is already proven in Theorem 2.6. Assume $M_{p^{n-2}}$ contains N_{n-2} pairwise quasi-orthogonal subalgebras which are isomorphic to M_p . Let $\mathcal{A} \simeq M_p \subset M_{p^{n-2}}$ be one of them. Then we can assume that \mathcal{A} is generated by W and S. On the other hand, M_{p^2} contains $p^2 + 1$ quasi-orthogonal subalgebras which are isomorphic to $\mathbb{C}^{2p} \simeq \mathbb{C}^p \otimes \mathbb{C}^p$ from Proposition 2.2 and the theory of mutually unbiased bases. Let $\mathcal{C} \simeq \mathbb{C}^p \otimes \mathbb{C}^p \subset M_{p^2}$ be one of them. Suppose \mathcal{C} is generated by $W \otimes I$ and $I \otimes W$. We define a homomorphism $\hat{\pi}$ (up to scalar multiple) from Z_p^4 to $\mathcal{A} \otimes \mathcal{C}$ by

$$\hat{\pi}(k,l,m_1,m_2) = S^k W^l \otimes W^{m_1} \otimes W^{m_2}.$$

For any $a_0, a_1 \in \mathbb{Z}_p$, let

$$\hat{C}_{a_0,a_1} = \{ b_0(1,0,a_0,a_1) + b_1(0,1,a_1D,a_0) \mid b_0, b_1 \in \mathbb{Z}_p \}$$

and

$$\hat{C}_{\infty} = \{ b_0(0,0,1,0) + b_1(0,0,0,1) \mid b_0, b_1 \in \mathbb{Z}_p \}$$

From a similar method of the proof of Lemma 2.4, the only vector common to any pair of above subspaces is (0, 0, 0, 0). Moreover since W and S satisfy (1) and C is commutative, we have

$$\hat{\pi}(1,0,a_0,a_1)\hat{\pi}(0,1,a_1D,a_0) = \lambda^{-1}\hat{\pi}(0,1,a_1D,a_0)\hat{\pi}(1,0,a_0,a_1).$$

Therefore the algebra span{ $\hat{\pi}(\hat{C}_{a_0,a_1})$ } generated by $\hat{\pi}(1,0,a_0,a_1)$ and $\hat{\pi}(0,1,a_1D,a_0)$ is isomorphic to the algebra generated by S and W, that is, M_p . Furthermore the algebra span{ $\hat{\pi}(\hat{C}_{\infty})$ } generated by $\hat{\pi}(0,0,1,0)$ and $\hat{\pi}(0,0,0,1)$ is $I \otimes \mathcal{C}$. Consequently $\mathcal{A} \otimes \mathcal{C}$ can be decomposed by $\mathcal{A} \otimes I = \text{span}\{\hat{\pi}(\hat{C}_{0,0})\}, I \otimes \mathcal{C}$ and $p^2 - 1$ pairwise quasiorthogonal subalgebras which are isomorphic to M_p . We denote the $p^2 - 1$ pairwise quasi-orthogonal subalgebras by $\{\mathcal{B}^k_{\mathcal{A},\mathcal{C}}\}_{k=1}^{p^2-1}$.

Let $\mathcal{A}_1, \ldots, \mathcal{A}_{N_{n-2}}$ be pairwise quasi-orthogonal subalgebras in $M_{p^{n-2}}$ which are isomorphic to M_p and let $\mathcal{C}_1, \ldots, \mathcal{C}_{p^2+1}$ be pairwise quasi-orthogonal subalgebras in M_{p^2} which are isomorphic to \mathbb{C}^{p^2} . For each $\mathcal{A}_{i_1}, \mathcal{A}_{i_2} \subset M_{p^{n-2}}$ and $\mathcal{C}_{j_1}, \mathcal{C}_{j_2} \subset M_{p^2}$ with $i_1 \neq i_2$ or $j_1 \neq j_2$, we have

$$(\mathcal{A}_{i_1} \ominus \mathbb{C}I) \otimes (\mathcal{C}_{j_1} \ominus \mathbb{C}I) \perp \mathcal{A}_{i_2} \otimes \mathcal{C}_{j_2}.$$

This implies $\mathcal{B}_{\mathcal{A}_{i_1},\mathcal{C}_{j_1}}^{k_1} \ominus \mathbb{C}I \perp \mathcal{B}_{\mathcal{A}_{i_2},\mathcal{C}_{j_2}}^{k_2} \ominus \mathbb{C}I$ if $i_1 \neq i_2$ or $j_1 \neq j_2$ or $k_1 \neq k_2$. Furthermore $\mathcal{B}_{\mathcal{A}_{i_1},\mathcal{C}_{j_1}}^k$ is quasi-orthogonal with $\mathcal{A}_{i_2} \otimes I$ and $I \otimes \mathcal{C}_{j_2}$ for all i_2, j_2 . In consequence, M_{p^n} is decomposed by $M_{p^{n-2}} \otimes I_{M_{p^2}} = \operatorname{span}\{\mathcal{A}_i \otimes I \mid i\}$ and $I_{M_{p^{n-2}}} \otimes M_{p^2} = \operatorname{span}\{I \otimes \mathcal{C}_j \mid j\}$ and $\{\mathcal{B}_{\mathcal{A}_i,\mathcal{C}_j}^k\}_{i,j,k}$. From Theorem 2.6, M_{p^2} contains $p^2 + 1$ pairwise quasi-orthogonal subalgebras which are isomorphic to M_p . Therefore we get

$$N_{n-2} + p^2 + 1 + (p^2 - 1)(p^2 + 1)N_{n-2} = N_n$$

pairwise quasi-orthogonal subalgebras in M_{p^n} which are isomorphic to M_p .

3 Quasi-orthogonal subalgebras in $M_{p^{kn}}$

In this section, we consider quasi-orthogonal subalgebras in $M_{p^{kn}}$ which are isomorphic to M_{p^k} . First we consider the case n = 2.

 $GF(p^k)$ denotes a finite field with p^k elements. Up to isomorphisms, $GF(p^k)$ is unique and is defined using a polynomial

$$f(x) = c_0 + \dots + c_{k-1}x^{k-1} + x^k$$

that is irreducible over the field Z_p . Then we can write

$$GF(p^k) = \{a(t) = a_0 + a_1t + \dots + a_{k-1}t^{k-1} \mid a_i \in Z_p, \ 0 \le i \le k-1\},\$$

where t satisfies f(t) = 0. Let

$$GF(p^k)^4 = \{a = (a^{(1)}(t), a^{(2)}(t), a^{(3)}(t), a^{(4)}(t)) \mid a^{(i)}(t) \in GF(p^k), 1 \le i \le 4\}$$

and define a symplectic product:

$$a \circ b = a^{(1)}(t)b^{(2)}(t) - a^{(2)}(t)b^{(1)}(t) + a^{(3)}(t)b^{(4)}(t) - a^{(4)}(t)b^{(3)}(t)$$

for $a = (a^{(1)}(t), a^{(2)}(t), a^{(3)}(t), a^{(4)}(t))$ and $b = (b^{(1)}(t), b^{(2)}(t), b^{(3)}(t), b^{(4)}(t))$ in $GF(p^k)^4$. Similarly, symplectic product on Z_p^{4k} is denoted by

$$u \circ v = \sum_{i=1}^{k} u_i^{(1)} v_i^{(2)} - u_i^{(2)} v_i^{(1)} + u_i^{(3)} v_i^{(4)} - u_i^{(4)} v_i^{(3)}$$

for $u = (u_i^{(j)})_{1 \le i \le k, 1 \le j \le 4}$ and $v = (v_i^{(j)})_{1 \le i \le k, 1 \le j \le 4}$ in $Z_p^{4k} = (Z_p^k)^4$. From [4], there exist a linear functional φ on $GF(p^k)$ and a linear isomorphism π_1 from $GF(p^k)^4$ to Z_p^{4k} such that

$$\varphi(a \circ b) = \pi_1(a) \circ \pi_1(b), \tag{3}$$

where we consider $GF(p^k)^4$ and Z_p^{4k} as vector spaces over Z_p . Define a natural homomorphism (up to scalar multiple) π_2 from Z_p^{4k} to $M_{p^{2k}} = \bigotimes_{i=1}^k M_p \otimes \bigotimes_{i=1}^k M_p$ by

$$\pi_2(u) = \bigotimes_{i=1}^k S^{u_i^{(1)}} W^{u_i^{(2)}} \otimes \bigotimes_{i=1}^k S^{u_i^{(3)}} W^{u_i^{(4)}}.$$

By (1), we have

$$\pi_2(u)\pi_2(v) = \lambda^{-u \circ v}\pi_2(v)\pi_2(u).$$

Lemma 3.1 If $a \circ b \neq 0$ for $a, b \in GF(p^k)^4$, then the algebra \mathcal{A} generated by

$$\{\pi_2\pi_1(p(t)a), \pi_2\pi_1(q(t)b) \mid p(t), q(t) \in GF(p^k)\}$$

is isomorphic to M_{p^k} , where p(t)a is defined by

$$p(t)(a_1(t), a_2(t), a_3(t), a_4(t)) = (p(t)a_1(t), p(t)a_2(t), p(t)a_3(t), p(t)a_4(t)).$$

Proof. If $\{p(t)a \mid p(t) \in GF(p^k)\} \cap \{q(t)b \mid q(t) \in GF(p^k)\} \neq \{(0,0,0,0)\}$, there exists $r(t) \in GF(p^k)$ such that a = r(t)b. This shows $a \circ b = r(t)(b \circ b) = 0$. Hence we can assume that $\{p(t)a\} \cap \{q(t)b\} = \{(0,0,0,0)\}$.

For $q(t) \in GF(p^k)$, let $\psi_{q(t)}$ be a functional on $GF(p^k)$ defined by

$$\psi_{q(t)}(p(t)) = \pi_1(p(t)a) \circ \pi_1(q(t)b),$$

where we consider $GF(p^k)$ as a vector space over Z_p . By (3), we have

$$\psi_{q(t)}(p(t)) = \varphi(p(t)a \circ q(t)b) = \varphi(p(t)q(t)(a \circ b)).$$

Since $a \circ b \neq 0$, $\psi_{q^{(1)}(t)} = \psi_{q^{(2)}(t)}$ implies $q^{(1)}(t) = q^{(2)}(t)$. Therefore we obtain $\{\psi_{q(t)} | q(t) \in GF(p^k)\} = GF(p^k)^*$, where $GF(p^k)^*$ is a dual space of $GF(p^k)$.

Let $\{p^{(i)}(t)\}_{1 \le i \le k}$ be a basis of the vector space $GF(p^k)$. Then there exists a basis $\{q^{(j)}\}_{1 \le j \le k} \subset GF(p^k)$ such that

$$\varphi(p^{(i)}(t)a \circ q^{(j)}(t)b) = \psi_{q^{(j)}(t)}(p^{(i)}(t)) = \delta_{ij}.$$

For $1 \leq i, j \leq k$, let $S_i = I^{\otimes i-1} \otimes S \otimes I^{\otimes k-i}$ and $W_j = I^{\otimes j-1} \otimes W \otimes I^{\otimes k-j}$ in $M_{p^k} = \bigotimes^k M_p$. We define a map ρ from $\{S_i, W_j \mid 1 \leq i, j \leq k\}$ to \mathcal{A} by

$$\rho(S_i) = \pi_2 \pi_1(p^{(i)}(t)a),
\rho(W_j) = \pi_2 \pi_1(q^{(j)}(t)b).$$

Then the commutativity condition of $\{S_i, W_j\}$, that is,

$$\begin{array}{rcl} S_i W_j &=& \lambda^{-\delta_{ij}} W_j S_i, \\ S_i S_j &=& S_j S_i, \\ W_i W_j &=& W_j W_i \end{array}$$

and the commutativity condition of $\{\pi_2\pi_1(p^{(i)}(t)a), \pi_2\pi_1(q^{(j)}(t)b) \mid 1 \leq i, j \leq k\}$, that is,

$$\begin{aligned} \pi_2 \pi_1(p^{(i)}(t)a) \cdot \pi_2 \pi_1(q^{(j)}(t)b) &= \lambda^{-\pi_1(p^{(i)}(t)a)\circ\pi_1(q^{(j)}(t)b)} \cdot \pi_2 \pi_1(q^{(j)}(t)b) \cdot \pi_2 \pi_1(p^{(i)}(t)a) \\ &= \lambda^{-\varphi(p^{(i)}(t)a\circ q^{(j)}(t)b)} \cdot \pi_2 \pi_1(q^{(j)}(t)b) \cdot \pi_2 \pi_1(p^{(i)}(t)a) \\ &= \lambda^{-\delta_{ij}} \cdot \pi_2 \pi_1(q^{(j)}(t)b) \cdot \pi_2 \pi_1(p^{(i)}(t)a), \\ \pi_2 \pi_1(p^{(i)}(t)a) \cdot \pi_2 \pi_1(p^{(j)}(t)a) &= \pi_2 \pi_1(p^{(j)}(t)a) \cdot \pi_2 \pi_1(p^{(i)}(t)a), \\ \pi_2 \pi_1(q^{(i)}(t)b) \cdot \pi_2 \pi_1(q^{(j)}(t)b) &= \pi_2 \pi_1(q^{(j)}(t)b) \cdot \pi_2 \pi_1(q^{(i)}(t)b) \end{aligned}$$

are same. Hence ρ can be extended to an isomorphism from M_{p^k} generated by $\{S_i, W_j\}_{1 \leq i, j \leq k}$ to \mathcal{A} .

From this lemma, we need to find such a and b. Let D be a non-zero element in $GF(p^k)$ with the requirement that $D \neq p(t)^2$ for all p(t) in $GF(p^k)$. For any $a(t), b(t) \in GF(p^k)$, define a subspace of $GF(p^k)^4$ by

$$C_{a(t),b(t)} = \{ p(t)(1,b(t),0,a(t)) + q(t)(0,a(t),-1,b(t)D) \mid p(t),q(t) \in GF(p^k) \}.$$

Moreover, put

$$C_{\infty} = \{ p(t)(0, 1, 0, 0) + q(t)(0, 0, 0, 1) \mid p(t), q(t) \in GF(p^k) \}.$$

Lemma 3.2 The only vector common to any pair of above subspaces is (0,0,0,0). In particular, the subspaces expect (0,0,0,0) partition $GF(p^k)^4 \setminus \{(0,0,0,0)\}$.

Proof. This proof is same as Lemma 2.4.

Since $(1, b(t), 0, a(t)) \circ (0, a(t), -1, b(t)D) = 2a(t)$, if $a(t) \neq 0$ then

$$\operatorname{span}\{\pi_2\pi_1(C_{a(t),b(t)})\} \simeq M_{p^k}$$

by Lemma 3.1. But if a(t) = 0, the algebra is commutative and hence span $\{\pi_2 \pi_1(C_{a(t),b(t)})\} \simeq \mathbb{C}^{p^{2k}}$. Therefore we need to mix the subspaces with a(t) = 0.

For any $a(t) \in GF(p^k)$, define subspaces by

$$D_{a(t)} = \{ p(t)(1, 1, -a(t), a(t)D) + q(t)(1, 2, -a(t), 2a(t)D) | p(t), q(t) \in GF(p^k) \}$$

$$D_{\infty} = \{ p(t)(0, 0, 1, 0) + q(t)(0, 0, 0, 1) | p(t), q(t) \in GF(p^k) \}.$$

Lemma 3.3 The only vector common to any pair of above subspaces is (0,0,0,0). Moreover we have

$$\bigcup_{a(t)\in GF(p^k)} D_{a(t)} \cup D_{\infty} = \bigcup_{b(t)\in GF(p^k)} C_{0,b(t)} \cup C_{\infty}.$$

Proof. This proof is same as Lemma 2.5.

Since $(1, 1, -a(t), a(t)D) \circ (1, 2, -a(t), 2a(t)D) = 1 - a(t)^2 D \neq 0$ by the assumption of D and $(0, 0, 1, 0) \circ (0, 0, 0, 1) = 1$, we have

by Lemma 3.1. Consequently we have the next theorem.

Theorem 3.4 There are $p^{2k} + 1$ pairwise quasi-orthogonal subalgebras in $M_{p^{2k}}$ which are isomorphic to M_{p^k} .

Next we consider pairwise quasi-orthogonal subalgebras in $M_{p^{kn}}$ which are isomorphic to M_{p^k} . The dimensions of $M_{p^{kn}} \oplus \mathbb{C}I$ and $M_{p^k} \oplus \mathbb{C}I$ are $p^{2kn} - 1$ and $p^{2k} - 1$, respectively. Hence such subalgebras are at most $N_{(k,n)} = p^{2kn} - 1/p^{2k} - 1$. We construct $N_{(k,n)}$ pairwise quasi-orthogonal subalgebras.

Theorem 3.5 $M_{p^{kn}}$ contains $N_{(k,n)}$ pairwise quasi-orthogonal subalgebras which are isomorphic to M_{n^k} .

Proof. The case n = 1 is trivial and the case n = 2 is already proven in Theorem 3.4. Assume $M_{p^{k(n-2)}}$ contains $N_{(k,n-2)}$ pairwise quasi-orthogonal subalgebras which are isomorphic to M_{p^k} . Let $\mathcal{A} \simeq M_{p^k} \subset M_{p^{k(n-2)}}$ be one of them. Then we can assume that \mathcal{A} is generated by $\{S_i, W_j\}_{1 \leq i,j \leq k}$. On the other hand, $M_{p^{2k}}$ contains $N_{(k,2)} =$ $p^{2k} + 1$ quasi-orthogonal subalgebras which are isomorphic to $\mathbb{C}^{p^{2k}} \simeq \mathbb{C}^{p^k} \otimes \mathbb{C}^{p^k}$ from Proposition 2.2 and the theory of mutually unbiased bases. Let $\mathcal{C} \simeq \mathbb{C}^{p^k} \otimes \mathbb{C}^{p^k} \subset M_{p^{2k}}$ be one of them. Suppose \mathcal{C} is generated by $\{W_i \otimes I, I \otimes W_j\}_{1 \leq i,j \leq k}$.

Define a homomorphism $\hat{\pi}_2$ (up to scalar multiple) from Z_p^{4k} to $\mathcal{A} \otimes \mathcal{C}$ by

$$\hat{\pi}_2(u_i^{(j)}) = \bigotimes_{i=1}^k S^{u_i^{(1)}} W^{u_i^{(2)}} \otimes \bigotimes_{i=1}^k W^{u_i^{(3)}} \otimes \bigotimes_{i=1}^k W^{u_i^{(4)}}$$

for $(u_i^{(j)})_{1 \le i \le k, 1 \le j \le 4} \in \mathbb{Z}_p^{4k}$. We denote other symplectic products by

$$a\hat{\circ}b = a^{(1)}(t)b^{(2)}(t) - a^{(2)}(t)b^{(1)}(t)$$

for $a = (a^{(1)}(t), a^{(2)}(t), a^{(3)}(t), a^{(4)}(t))$ and $b = (b^{(1)}(t), b^{(2)}(t), b^{(3)}(t), b^{(4)}(t))$ in $GF(p^k)^4$, and

$$\hat{u \circ v} = \sum_{i=1}^{k} u_i^{(1)} v_i^{(2)} - u_i^{(2)} v_i^{(1)}$$

for $u = (u_i^{(j)})_{1 \le i \le k, 1 \le j \le 4}$ and $v = (v_i^{(j)})_{1 \le i \le k, 1 \le j \le 4}$ in Z_p^{4k} . Since \mathcal{C} is commutative, we have

$$\hat{\pi}_2(u)\hat{\pi}_2(v) = \lambda^{-u\hat{\circ}v}\hat{\pi}_2(v)\hat{\pi}_2(u).$$

Moreover from [4], there exists a homomorphism $\hat{\pi}_1$ from $GF(p^k)^4$ to Z_p^{4k} such that

$$\varphi(a \circ b) = \hat{\pi}_1(a) \circ \hat{\pi}_1(b).$$

Here for elements $a = (a^{(1)}(t), a^{(2)}(t), 0, 0)$ and $b = (0, 0, b^{(3)}(t), b^{(4)}(t))$ in $GF(p^k)^4$, we can assume

$$\hat{\pi}_1(a) = ((u_i^{(1)}), (u_j^{(2)}), 0, 0)$$

$$\hat{\pi}_1(b) = (0, 0, (v_i^{(3)}), (v_j^{(4)}))$$

for some $(u_i^{(1)}), (u_j^{(2)}), (v_i^{(3)}), (v_j^{(4)}) \in Z_p^k$. Then we can prove that the algebra generated by $\hat{\pi}_2 \hat{\pi}_1(a)$ and $\hat{\pi}_2 \hat{\pi}_1(b)$ is isomorphic to M_{p^k} , if $a \hat{\circ} b \neq 0$, by using a similar proof of Lemma 3.1.

For any $a(t), b(t) \in GF(p^k)$, let

$$\hat{C}_{a(t),b(t)} = \{ p(t)(1,0,a(t),b(t)) + q(t)(0,1,b(t)D,a(t)) \mid p(t),q(t) \in GF(p^k) \}$$

and

$$\hat{C}_{\infty} = \{ p(t)(0,0,1,0) + q(t)(0,0,0,1) \mid p(t), q(t) \in GF(p^k) \}.$$

From the similar method of the proof of Lemma 2.4, the only vector common to any pair of above subspaces is (0, 0, 0, 0). Since $(1, 0, a(t), b(t)) \circ (0, 1, b(t)D, a(t)) = 1$, we obtain

$$\operatorname{span}\{\hat{\pi}_2\hat{\pi}_1(\hat{C}_{a(t),b(t)})\} \simeq M_{p^k}.$$

Moreover we have

$$\operatorname{span}\{\hat{\pi}_2\hat{\pi}_1(\hat{C}_\infty)\} = I \otimes \mathcal{C} \simeq \mathbb{C}^{p^{2k}}.$$

Consequently $\mathcal{A} \otimes \mathcal{C}$ can be decomposed by $\mathcal{A} \otimes I = \operatorname{span}\{\hat{\pi}_2 \hat{\pi}_1(\hat{C}_{0,0})\}, I \otimes \mathcal{C} = \operatorname{span}\{\hat{\pi}_2 \hat{\pi}_1(\hat{C}_{\infty})\}\$ and $p^{2k} - 1$ pairwise quasi-orthogonal subalgebras which are isomorphic to M_{p^k} . We denote the $p^{2k} - 1$ pairwise quasi-orthogonal subalgebras by $\{\mathcal{B}_{\mathcal{A},\mathcal{C}}^k\}_{k=1}^{p^{2k}-1}$.

Let $\mathcal{A}_1, \ldots, \mathcal{A}_{N_{(k,n-2)}}$ be pairwise quasi-orthogonal subalgebras in $M_{p^{k(n-2)}}$ which are isomorphic to M_{p^k} and let $\mathcal{C}_1, \ldots, \mathcal{C}_{N_{(k,2)}}$ be pairwise quasi-orthogonal subalgebras in $M_{p^{2k}}$ which are isomorphic to $\mathbb{C}^{p^{2k}}$. Since

$$(\mathcal{A}_{i_1} \ominus \mathbb{C}I) \otimes (\mathcal{C}_{j_1} \ominus \mathbb{C}I) \perp \mathcal{A}_{i_2} \otimes \mathcal{C}_{j_2}$$

for $i_1 \neq i_2$ or $j_1 \neq j_2$, $\mathcal{B}_{\mathcal{A}_{i_1},\mathcal{C}_{j_1}}^{k_1}$ and $\mathcal{B}_{\mathcal{A}_{i_2},\mathcal{C}_{j_2}}^{k_2}$ are quasi-orthogonal if $i_1 \neq i_2$ or $j_1 \neq j_2$ or $k_1 \neq k_2$. Furthermore, $\mathcal{B}_{\mathcal{A}_{i_1},\mathcal{C}_{j_1}}^{k_1}$ is quasi-orthogonal with $\mathcal{A}_{i_2} \otimes I_{M_{p^{2k}}}$ and $I_{M_{p^{k(n-2)}}} \otimes \mathcal{C}_{j_2}$ for any i_2 and j_2 . Therefore we can decompose $M_{p^{kn}}$ by $M_{p^{k(n-2)}} \otimes I_{M_{p^{2k}}} = \operatorname{span}\{\mathcal{A}_i \otimes I\}$, $I_{M_{p^{k(n-2)}}} \otimes M_{p^{2k}} = \operatorname{span}\{I \otimes \mathcal{C}_j\}$ and $\{\mathcal{B}_{\mathcal{A}_i,\mathcal{C}_j}^k\}_{i,j,k}$. In consequence, we get

$$N_{(k,n-2)} + N_{(k,2)} + (p^{2k} - 1)N_{(k,2)}N_{(k,n-2)} = N_{(k,n)}$$

pairwise quasi-orthogonal subalgebras in $M_{p^{kn}}$ which are isomorphic to M_{p^k} .

References

- [1] S. Bandyopadhyay, P.O. Boykin, V. Roychowdhury and F. Vatan, A new proof for the existence of mutually unbiased bases, Algoritmica **34** (2002), 512–528.
- [2] P.O Boykin, M. Sitharam, P.H. Tiep and P. Wocjan, Mutually unbiased bases and orthogonal decompositions of Lie algebras, Quantum Inf. comput. 7(4) (2007), 371-382.
- [3] H. Ohno, D. Petz and A. Szántó, Quasi-orthogonal subalgebras of 4 × 4 matrices, Linear Alg. Appl. 425 (2007), 109-118.
- [4] O. Pittenger and M. H. Rubin, Mutually unbiased bases, generalized spin matrices and separability, Linear Alg. Appl. 390 (2004), 255-278.
- [5] S. Popa, Orthogonal pairs of *-subalgebras in finite von Neumann algebras, J. Oper. Theory 9 (1983), 253-268.

- [6] D. Petz, Complementarity in quantum systems, Rep. Math. Phys. 59 (2007), 209-224.
- [7] D. Petz, Quantum information and quantum statistics, Springer, 2007.
- [8] D. Petz and J. Kahn, Complementary reductions for two qubits, J. Math. Phys. 48 (2007), 012107.