

Fluctuations of statistics among subregions of a turbulence velocity field

Hideaki Mouri^a and Akihiro Hori^b

Meteorological Research Institute, Nagamine, Tsukuba 305-0052, Japan

Masanori Takaoka^c

Department of Mechanical Engineering, Doshisha University, Kyotanabe, Kyoto 610-0321, Japan

To study subregions of a turbulence velocity field, a long record of velocity data of grid turbulence is divided into smaller segments. For each segment, we calculate statistics such as the mean rate of energy dissipation and the mean energy at each scale. Their values significantly fluctuate, in lognormal distributions at least as a good approximation. Each segment is not under equilibrium between the mean rate of energy dissipation and the mean rate of energy transfer that determines the mean energy. These two rates still correlate among segments when their length exceeds the correlation length. Also between the mean rate of energy dissipation and the mean total energy, there is a correlation characterized by the Reynolds number for the whole record, implying that the large-scale flow affects each of the segments.

I. INTRODUCTION

For locally isotropic turbulence, Kolmogorov¹ considered that small-scale statistics are uniquely determined by the kinematic viscosity ν and the mean rate of energy dissipation $\langle \varepsilon \rangle$. The Kolmogorov velocity $u_K = (\nu \langle \varepsilon \rangle)^{1/4}$ and the Kolmogorov length $\eta = (\nu^3 / \langle \varepsilon \rangle)^{1/4}$ determine the statistics of velocity increment $\delta u_r = u(x+r) - u(x)$ at scale r as

$$\frac{\langle \delta u_r^n \rangle}{u_K^n} = F_n \left(\frac{r}{\eta} \right) \quad \text{for } n = 2, 3, 4, \dots \quad (1)$$

Here $\langle \cdot \rangle$ denotes an average over position x , and F_n is a universal function. The universality is known to hold well. While $\langle \delta u_r^n \rangle$ at each r is different in different velocity fields, $\langle \varepsilon \rangle$ and hence u_K^n and η are accordingly different. That is, $\langle \varepsilon \rangle$ is in equilibrium with the mean rate of energy transfer that determines $\langle \delta u_r^n \rangle$.

However, the universality of small-scale statistics might not be exact. To argue against the exact universality, Landau² pointed out that the local rate of energy dissipation ε fluctuates over large scales. This fluctuation is not universal and is always significant.^{3,4,5} In fact, the large-scale flow or the configuration for turbulence production appears to affect some small-scale statistics.^{5,6,7,8,9,10}

Obukhov¹¹ discussed that Kolmogorov's theory¹ still holds in an ensemble of "pure" subregions where ε is constant at a certain value. Then, the ε value represents the rate of energy transfer averaged over those subregions. For the whole region, small-scale statistics reflect the large-scale flow through the large-scale fluctuation of the ε value. The idea that turbulence consists of some elementary subregions is of interest even now.⁸ We study

statistics among subregions in terms of the effect of large scales on small scales, by using a long record of velocity data obtained in grid turbulence.

II. EXPERIMENT

The experiment was done in a wind tunnel of the Meteorological Research Institute. Its test section had the size of 18, 3, and 2 m in the streamwise, spanwise, and floor-normal directions. We placed a grid across the entrance to the test section. The grid consisted of two layers of uniformly spaced rods, with axes in the two layers at right angles. The cross section of the rods was $0.04 \times 0.04 \text{ m}^2$. The separation of the axes of adjacent rods was 0.20 m.

On the tunnel axis at 4 m downstream of the grid, we simultaneously measured the streamwise ($U + u$) and spanwise (v) velocities. Here U is the average while $u(t)$ and $v(t)$ are fluctuations as a function of time t . We used a hot-wire anemometer with a crossed-wire probe. The wires were made of platinum-plated tungsten, 5 μm in diameter, 1.25 mm in sensing length, 1 mm in separation, oriented at $\pm 45^\circ$ to the streamwise direction, and 280 $^\circ\text{C}$ in temperature. The signal was linearized, low-pass filtered at 35 kHz, and then digitally sampled at $f_s = 70 \text{ kHz}$. We obtained as long as 4×10^8 data.

The calibration coefficient, with which the flow velocity is proportional to the anemometer signal, depends on the condition of the hot wires and thereby varied slowly in time. We determine the coefficient so as to have $U = 21.16 \text{ m s}^{-1}$ for each segment with 4×10^6 data. Within each segment, the coefficient varied by $\pm 0.4\%$ at most. Also varied slowly in time the flow temperature and hence the kinematic viscosity ν . We adopt $\nu = 1.42 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ based on the mean flow temperature, 11.8 $^\circ\text{C}$. The temperature variation, $\pm 1.2 \text{ }^\circ\text{C}$, corresponds to the ν variation of $\pm 0.7\%$. These variations are small and ignored here.

Taylor's frozen-eddy hypothesis, i.e., $x = -Ut$, is used to obtain $u(x)$ and $v(x)$ from $u(t)$ and $v(t)$. This hypoth-

^aElectronic address: hmouri@mri-jma.go.jp

^bAlso at Meteorological and Environmental Sensing Technology, Inc., Nanpeidai, Ami 300-0312, Japan

^cElectronic address: mtakaoka@mail.doshisha.ac.jp

TABLE I: Turbulence parameters: mean energy dissipation rate $\langle \varepsilon \rangle$, rms velocity fluctuations $\langle u^2 \rangle^{1/2}$ and $\langle v^2 \rangle^{1/2}$, Kolmogorov velocity u_K , flatness factors $\langle u^4 \rangle / \langle u^2 \rangle^2$ and $\langle v^4 \rangle / \langle v^2 \rangle^2$, correlation lengths L_u , L_v , and L_ε , Taylor microscale λ , Kolmogorov length η , and microscale Reynolds number Re_λ .

Quantity	Value
$\langle \varepsilon \rangle = 15\nu \langle (\partial_x v)^2 \rangle / 2$	$7.98 \text{ m}^2 \text{ s}^{-3}$
$\langle u^2 \rangle^{1/2}$	1.10 m s^{-1}
$\langle v^2 \rangle^{1/2}$	1.06 m s^{-1}
$u_K = (\nu \langle \varepsilon \rangle)^{1/4}$	0.103 m s^{-1}
$\langle u^4 \rangle / \langle u^2 \rangle^2$	3.02
$\langle v^4 \rangle / \langle v^2 \rangle^2$	3.00
$L_u = \int_0^\infty \langle u(x+r)u(x) \rangle dr / \langle u^2 \rangle$	17.9 cm
$L_v = \int_0^\infty \langle v(x+r)v(x) \rangle dr / \langle v^2 \rangle$	4.69 cm
$L_\varepsilon = \int_0^\infty \langle \varepsilon(x+r)\varepsilon(x) - \langle \varepsilon \rangle^2 \rangle dr / \langle \varepsilon^2 - \langle \varepsilon \rangle^2 \rangle$	0.469 cm
$\lambda = [2 \langle v^2 \rangle / \langle (\partial_x v)^2 \rangle]^{1/2}$	0.548 cm
$\eta = (\nu^3 / \langle \varepsilon \rangle)^{1/4}$	0.0138 cm
$\text{Re}_\lambda = \langle v^2 \rangle^{1/2} \lambda / \nu$	409

esis requires a small value of $\langle u^2 \rangle^{1/2} / U$. The value in our experiment, 0.05, is small enough. Since $u(t)$ and $v(t)$ are stationary, $u(x)$ and $v(x)$ are homogeneous, although grid turbulence decays along the streamwise direction in the wind tunnel. We are mostly interested in scales up to about the typical scale for energy-containing eddies, which is much less than the tunnel size. Over such scales, fluctuations of $u(x)$ and $v(x)$ correspond to spatial fluctuations that were actually present in the wind tunnel.¹² Those over the larger scales do not. They have to be interpreted as fluctuations over long timescales described in terms of large length scales.¹³

Turbulence parameters are listed in Table I. Here and hereafter, $\langle \cdot \rangle$ is used to denote an average over the whole record. The derivative was obtained as $\partial_x v = [8v(x + \delta x) - 8v(x - \delta x) - v(x + 2\delta x) + v(x - 2\delta x)] / 12\delta x$ with $\delta x = U / f_s$. The local rate of energy dissipation was obtained as $\varepsilon = 15\nu(\partial_x v)^2 / 2$ instead of usual $15\nu(\partial_x u)^2$, in order to avoid possible spurious correlations with δu_r over small r for analyses in the next section.

Figure 1 shows $\langle \delta u_r^2 \rangle / u_K^2$, the u , v , and ε correlations, and also the correlation lengths L_u , L_v , and L_ε . We see the inertial range, albeit narrow, where $\langle \delta u_r^2 \rangle$ roughly scales with $r^{2/3}$. The u and v correlations are significant up to $r \simeq 10^4 \eta$, which corresponds to the scale of largest eddies. The correlation length L_u corresponds to the typical scale for energy-containing eddies. Since ε belongs to small scales, its correlation decays quickly.

III. RESULTS AND DISCUSSION

The data record is now divided into segments with length R . They correspond to subregions considered by

Obukhov.¹¹ For each segment, we have statistics such as

$$\begin{aligned} (\partial_x v)_R^2(x) &= \frac{1}{R} \int_{x-R/2}^{x+R/2} \left[\frac{\partial v(x')}{\partial x'} \right]^2 dx', \\ \delta u_{r,R}^n(x) &= \frac{1}{R-r} \int_{x-R/2}^{x+R/2-r} \delta u_r^n(x') dx', \\ v(x+r)v(x)_R &= \frac{1}{R-r} \int_{x-R/2}^{x+R/2-r} v(x'+r)v(x') dx'. \end{aligned} \quad (2)$$

Here x is the center of the segment, and $r < R$. The mean rate of energy dissipation is $\varepsilon_R = 15\nu(\partial_x v)_R^2 / 2$, which yields the Kolmogorov velocity $u_{K,R} = (\nu \varepsilon_R)^{1/4}$ and the Kolmogorov length $\eta_R = (\nu^3 / \varepsilon_R)^{1/4}$. We also have the mean total energy, $v_R^2 = v(x+r)v(x)_R$ for $r = 0$, and the microscale Reynolds number, $\text{Re}_{\lambda,R} = 2^{1/2} v_R^2 / \nu [(\partial_x v)_R^2]^{1/2}$. The mean rate of energy transfer, however, is not available from our experimental data. Fig. 1(a) shows an example of $\delta u_{r,R}^2 / u_{K,R}^2$, which differs from $\langle \delta u_r^2 \rangle / u_K^2$.

A. Distribution of fluctuation

Over a range of R , we study statistics of $\delta u_{r,R}^2 / u_{K,R}^2$ among segments. The scale r is fixed at $10\eta_R$ in the dissipation range and $100\eta_R$ in the inertial range. Since




FIG. 1: (a) $\langle \delta u_r^2 \rangle / u_K^2$ as a function of r/η and $\delta u_{r,R}^2 / u_{K,R}^2$ for a segment with $R = 10^3 \eta$ as a function of r/η_R . We indicate L_u , L_v , L_ε , and the inertial range. (b) $\langle u(x+r)u(x) \rangle / \langle u^2 \rangle$, $\langle v(x+r)v(x) \rangle / \langle v^2 \rangle$, and $\langle \varepsilon(x+r)\varepsilon(x) - \langle \varepsilon \rangle^2 \rangle / \langle \varepsilon^2 - \langle \varepsilon \rangle^2 \rangle$ as a function of r/η .

δu_r is available only at discrete scales r that are multiples of the sampling interval U/f_s , $\delta u_{10\eta R,R}^2$ and $\delta u_{100\eta R,R}^2$ are obtained through interpolation by incorporating the fluctuation of η_R among segments.

Figure 2(a) shows the standard deviation. The fluctuation of $\delta u_{r,R}^2/u_{K,R}^2$ at a fixed r/η_R is significant even when R is large. In individual segments, Kolmogorov's theory¹ does not hold. The mean rate of energy transfer that determines $\delta u_{r,R}^2$ is not in equilibrium with the mean rate of energy dissipation ε_R that determines $u_{K,R}^2$ and η_R . The degree of this nonequilibrium fluctuates among segments and thereby induces the observed fluctuation of $\delta u_{r,R}^2/u_{K,R}^2$.

The mean rate of energy transfer also fluctuates among




FIG. 2: Statistics of $\delta u_{10\eta R,R}^2/u_{K,R}^2$ (filled triangles down), $\delta u_{100\eta R,R}^2/u_{K,R}^2$ (filled triangles up), ε_R (open circles), v_R^2 (open squares), and $\text{Re}_{\lambda,R}$ (open diamonds) as a function of R/η . (a) Standard deviation normalized by the average. The dotted line indicates the $R^{-1/2}$ scaling. (b) Skewness factor. (c) Flatness factor. Also shown are the skewness and flatness factors of $\ln(\delta u_{10\eta R,R}^2/u_{K,R}^2)$ (open triangle down), $\ln(\delta u_{100\eta R,R}^2/u_{K,R}^2)$ (open triangles up), $\ln \varepsilon_R$ (filled circles), $\ln v_R^2$ (filled squares), and $\ln \text{Re}_{\lambda,R}$ (filled diamonds).




FIG. 3: (a) Probability densities of $X_R = \delta u_{100\eta R,R}^2/u_{K,R}^2$ (triangles up) and $\ln(\delta u_{100\eta R,R}^2/u_{K,R}^2)$ (triangles down) at $R = 10^3\eta$ (open triangles) and $10^5\eta$ (filled triangles) as a function of $(X_R - \langle X_R \rangle) / (\langle X_R - \langle X_R \rangle \rangle^2)^{1/2}$. Those of $X_R = \delta u_{100\eta R,R}^2/u_{K,R}^2$ are shifted by a factor of 10^2 . The dotted lines denote the Gaussian distribution. (b) Same as (a) but for $X_R = \varepsilon_R$ (triangles up) and $\ln \varepsilon_R$ (triangles down).

scales in each segment, which does not necessarily have the inertial-range scaling $\delta u_{r,R}^2 \propto r^{2/3}$ [Fig. 1(a)].¹⁴ If each segment had this scaling, the fluctuation of $\delta u_{r,R}^2/u_{K,R}^2$ among segments at a fixed r/η_R in the inertial range would correspond to the fluctuation of the Kolmogorov constant $\delta u_{r,R}^2/(r\varepsilon_R)^{2/3}$.

Figures 2(b) and 2(c) show the skewness and flatness factors of $\ln(\delta u_{r,R}^2/u_{K,R}^2)$ (open triangles). They are close to the Gaussian values of 0 and 3. Thus, at least as a good approximation, the distribution of $\delta u_{r,R}^2/u_{K,R}^2$ is lognormal. This is also the case in ε_R (filled circles) and at $R \gtrsim 10^3\eta \simeq L_u$ in v_R^2 and $\text{Re}_{\lambda,R}$ (filled squares and diamonds), while the mean rate of energy transfer should not have a lognormal distribution because it changes its sign. Examples of the probability density functions are shown in Fig. 3. The lognormal distribution of ε_R was discussed as a tentative model by Obukhov.¹¹

A lognormal distribution stems from some multiplicative stochastic process, e.g., a product of many independent stochastic variables with similar variances. To its logarithm, if not too far from the average, the central limit theorem applies. For the lognormal distributions observed here, the process is related with the energy transfer. While the mean energy transfer is to a smaller scale and is significant between scales in the inertial range alone, the local energy transfer is either to




FIG. 4: (a) $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ at $R = 10^2 \eta$ as a function of r/η_R (solid lines) and $\langle \delta u_r^2 \rangle / u_K^2$ as a function of r/η (dotted line). The inset shows the dependence of $\langle \delta u_{10\eta,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ on $\langle \varepsilon_R \rangle_\varepsilon / \langle \varepsilon \rangle$. The horizontal dotted line indicates the value of $\langle \delta u_{10\eta}^2 \rangle / u_K^2$. (b) Same as (a) but at $R = 10^3 \eta$.

a smaller or larger scale and is significant between all scales.^{5,10,15} Any scale is thereby affected by itself and by many other scales. They involve large scales because the lognormal distributions are observed up to large R . There is no dominant effect from a few specific scales, in order for the central limit theorem to be applicable.

At $R \gtrsim 10^5 \eta \simeq 10^2 L_u$, there are alternative features.^{5,16} The standard deviations scale with $R^{-1/2}$ [Fig. 2(a)]. The skewness and flatness factors of $\delta u_{r,R}^2 / u_{K,R}^2$, ε_R , v_R^2 , and $\text{Re}_{\lambda,R}$ are close to the Gaussian values [Figs. 2(b) and 2(c): filled triangles, open circles, open squares, and open diamonds; see also Fig. 3]. Their distributions are regarded as Gaussian rather than lognormal, although this has to be confirmed in future using high-order moments or probability densities at the tails for the larger number of segments. The $R^{-1/2}$ scaling and Gaussian distribution are typical of fluctuations in thermodynamics and statistical mechanics,¹⁷ for which no correlation is significant at scales of interest as in our case at $r \gtrsim 10^5 \eta$ [Fig. 1(b)].

B. Correlation between fluctuations

The fluctuations among segments at $R \gtrsim L_u$ have interesting correlations. Since these correlations are weak, they are extracted by following Obukhov,¹¹ i.e., by av-




FIG. 5: (a) $\langle v_R^2 \rangle_\varepsilon / \langle v^2 \rangle$ as a function of $\langle \varepsilon_R \rangle_\varepsilon / \langle \varepsilon \rangle$ at $R = 10^2 \eta$ (triangles), $10^3 \eta$ (circles), and $10^4 \eta$ (squares). Along the dotted lines, the microscale Reynolds number $15^{1/2} \langle v_R^2 \rangle_\varepsilon / (\nu \langle \varepsilon_R \rangle_\varepsilon)^{1/2}$ is constant. (b) $\langle \text{Re}_{\lambda,R} \rangle_\varepsilon$ as a function of $\langle \varepsilon_R \rangle_\varepsilon / \langle \varepsilon \rangle$. The symbols are the same as in (a).

eraging over segments with similar ε_R . Specifically, we use conditional averages, denoted by $\langle \cdot \rangle_\varepsilon$, for ranges of ε_R separated at $\langle \varepsilon \rangle / 4$, $\langle \varepsilon \rangle / 2$, $\langle \varepsilon \rangle$, $2\langle \varepsilon \rangle$, and $4\langle \varepsilon \rangle$.

Figure 4 shows $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ as a function of r/η_R . When $R = 10^3 \eta \simeq L_u$ [Fig. 4(b)], $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ is independent of $\langle \varepsilon_R \rangle_\varepsilon$. The former varies only by a factor of 1.5 while the latter varies by a factor of 20. When $R = 10^2 \eta \simeq 10^{-1} L_u$ [Fig. 4(a)], $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ is not independent of $\langle \varepsilon_R \rangle_\varepsilon$.

The implication of the above result is that the mean rate of energy transfer that determines $\delta u_{r,R}^2$ correlates with the mean rate of energy dissipation ε_R that determines $u_{K,R}^2$ and η_R among segments with $R \gtrsim L_u$. Here L_u is the typical scale for energy-containing eddies. Most of the energy of such an eddy is transferred through scales and dissipated within its own volume. Thus, each energy-containing eddy tends toward equilibrium between the mean rates of energy transfer and dissipation. This tendency does not exist at $R \lesssim L_u$.¹⁸ Within an energy-containing eddy, the spatial distribution of ε is not homogeneous. In fact, the ε correlation is significant at $r \lesssim L_u$ [Fig. 1(b)].

Therefore, in order for statistics such as $\delta u_{r,R}^2 / u_{K,R}^2$ and $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$ to have physical meanings, the minimum segment length is about L_u . We are to discuss that such segments individually reflect the large-scale flow.

Figure 5(a) shows $\langle v_R^2 \rangle_\varepsilon / \langle v^2 \rangle$ as a function of

$\langle \varepsilon_R \rangle_\varepsilon / \langle \varepsilon \rangle$. The former varies with the latter. When $R \geq 10^3 \eta \simeq L_u$, the microscale Reynolds number $\propto \langle v_R^2 \rangle_\varepsilon / \langle \varepsilon_R \rangle_\varepsilon^{1/2}$ is almost constant at the value for the whole record, $\text{Re}_\lambda = 409$. Thus, segments obey a correlation between v_R^2 and ε_R characterized by the Re_λ value. Since Re_λ is determined by the large-scale flow, it follows that the large-scale flow affects each of the segments. A consistent result is obtained for $\langle \text{Re}_{\lambda,R} \rangle_\varepsilon$ in Fig. 5(b).

The above tendency toward a constant microscale Reynolds number originates in energy-containing scales. In general, through an empirical relation $\langle \varepsilon \rangle \propto \langle v^2 \rangle^{3/2} / L_v$, Re_λ is related to the Reynolds number for energy-containing scales as $\text{Re}_\lambda \propto (\langle v^2 \rangle^{1/2} L_v / \nu)^{1/2}$. Then, Re_λ is constant if $L_v \propto \langle v^2 \rangle^{-1/2}$. Fig. 6 shows $\langle v(x+r)v(x)_R \rangle_\varepsilon / \langle v_R^2 \rangle_\varepsilon$, which extends to larger scales for smaller $\langle v_R^2 \rangle_\varepsilon$. The correlation length $\int_0^\infty \langle v(x+r)v(x)_R \rangle_\varepsilon dr / \langle v_R^2 \rangle_\varepsilon$ should be accordingly larger.

The process for each segment to reflect the large-scale flow could be related to the energy transfer. As noted before, the energy transfer couples all scales. The energy transfer itself is affected by the large-scale flow. This is because energy is transferred between two scales via an interaction with some other scale. When the interaction occurs with a large scale, the energy transfer is strong.^{10,15}

C. Unimportance of ε fluctuation

Obukhov¹¹ discussed that, through the fluctuation of the ε_R value, the large-scale flow affects small-scale statistics for the whole region such as $\langle \delta u_r^n \rangle / u_{K,R}^n$. The reason is that $\langle \delta u_r^n \rangle$ is obtained at a fixed r , regardless of the fluctuations of $u_{K,R}^n$ and η_R induced by the fluctuation of ε_R . We are to discuss that the fluctuation of the ε_R value is not important so far as $R \gtrsim L_u$. This condition on R is required for the ε_R value to correlate with and thus statistically represent the mean rate of energy transfer that determines $\delta u_{r,R}^n$.

Figure 7 compares $\langle \delta u_r^n \rangle / u_K^n$ to $\langle \delta u_{r,R}^n / u_{K,R}^n \rangle$, i.e., average of $\delta u_{r,R}^n / u_{K,R}^n$ at each r/η_R over all segments, with $R = 10^3 \eta \simeq L_u$. Even for $n = 6$ [Fig. 7(b)], they are not distinguishable [see also Fig. 4(b)]. Although ε_R and hence $u_{K,R}$ and η_R fluctuate among segments, these fluctuations are not large enough for $\langle \delta u_r^n \rangle / u_K^n$ to differ from $\langle \delta u_{r,R}^n / u_{K,R}^n \rangle$. This conclusion is general. In various flows,^{3,4,5} including an atmospheric boundary layer at $\text{Re}_\lambda \simeq 9000$, the standard deviation of $\varepsilon_R / \langle \varepsilon_R \rangle$ at $R \simeq L_u$ is close to the value obtained here [Fig. 2(a)]. Hence, through the fluctuation of the ε_R value, the large-scale flow does not affect small-scale statistics for the whole record. It was suggested that the large-scale flow does affect $\langle \delta u_r^n \rangle / u_K^n$, even in the scaling exponents.^{6,9,10} If this is the case, the effect is already inherent in the individual segments. They are unlikely to be “pure¹¹” or elementary.




FIG. 6: $\langle v(x+r)v(x)_R \rangle_\varepsilon / \langle v_R^2 \rangle_\varepsilon$ at $R = 10^3 \eta$ as a function of r/η_R (solid lines) and $\langle v(x+r)v(x) \rangle / \langle v^2 \rangle$ as a function of r/η (dotted line).




FIG. 7: $\langle \delta u_{r,R}^n / u_{K,R}^n \rangle$ at $R = 10^3 \eta$ as a function of r/η_R (solid line) and $\langle \delta u_r^n \rangle / u_K^n$ as a function of r/η (dotted line). (a) $n = 2$. (b) $n = 6$.

IV. CONCLUDING REMARKS

Using segments of a long record of velocity data obtained in grid turbulence, we have studied fluctuations of statistics such as $\delta u_{r,R}^2 / u_{K,R}^2$, ε_R , and v_R^2 . The fluctuations are significant and have lognormal distributions at least as a good approximation (Figs. 2 and 3). In each segment, the mean rate of energy transfer that determines $\delta u_{r,R}^2$ is not in equilibrium with the mean rate of energy dissipation ε_R that determines $u_{K,R}^2$ and η_R . These two rates still correlate among segments with $R \gtrsim L_u$ (Fig. 4), which tend toward equilibrium between

the two rates. Also between ε_R and v_R^2 , there is a correlation characterized by Re_λ for the whole record (Fig. 5). Thus, the large-scale flow affects each of the segments.

The observed fluctuations depend on L_u and Re_λ , which in turn depend on the configuration for turbulence production, e.g., boundaries such as the grid used in our experiment. Nevertheless, the significance of those fluctuations implies that they have been developed in turbulence itself. Their lognormal distributions are explained by a multiplicative stochastic process in turbulence, which is related with the energy transfer among scales. The correlations among the fluctuations are also explained in terms of the energy transfer.

Previous studies^{5,6,7,8,9,10} suggested that the large-scale flow affects some small-scale statistics, although this has to be confirmed at higher Re_λ where large and small scales are more distinct.¹⁹ If the effect really exists, it is inherent individually in the segments.

Our study was motivated by Obukhov's discussion.¹¹ It implies the presence of equilibrium between the mean rates of energy transfer and dissipation in an ensemble of segments with similar values of ε_R . This is the case at $R \gtrsim L_u$ (Fig. 4).²⁰ Also as discussed by Obukhov, the distribution of ε_R is lognormal at least as a good approximation (Figs. 2 and 3). However, although Obukhov discussed that the large-scale flow affects small-scale statistics through the fluctuation of the ε_R value, this is not the case (Fig. 7).

The lognormal distributions observed here have to be distinguished from those proposed by Kolmogorov.²¹ While he was interested in small-scale intermittency and studied ε_r and δu_r^n at small r to obtain their scaling laws, we are interested in large-scale fluctuations and have studied ε_R and $\delta u_{r,R}^n$ at small r but at large R . The

scaling laws of ε_R and $\delta u_{r,R}^n$ are not necessary. In addition, the lognormality has been attributed to a different process. Hence, our study is not necessarily concerned with the well-known problems of Kolmogorov's lognormal model, e.g., violation of Novikov's inequality²² for scaling exponents. Still exists a possibility that small-scale intermittency is affected by large-scale fluctuations. The study of this possibility is desirable.

There were no studies of statistics among segments with large R . Hence, we have focused on grid turbulence, which is simple and thus serves as a standard. For flows other than grid turbulence, the fluctuations of statistics among segments are expected to be significant as well. In fact, regardless of the flow configuration and the Reynolds number, the large-scale fluctuation of ε_R is significant.^{3,4,5} Those fluctuations are also expected to have lognormal distributions and mutual correlations as observed here because they are due to the energy transfer in turbulence itself. However, grid turbulence is free from shear. It was previously found that δu_r correlates with u in shear flows such as a boundary layer but not in shear-free flows.^{7,8,9} The fluctuations of statistics among segments might be somewhat different in a shear flow. To this and other flow configurations, it is desirable to apply our approach.

Acknowledgments

We are grateful to K. R. Sreenivasan for inspiring this study and for helpful comments and also to M. Tanahashi for helpful comments.

¹ A. N. Kolmogorov, "The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers," *Dokl. Akad. Nauk SSSR* **30**, 301 (1941).

² L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon, London, 1959), Chap. 3.

³ A. Praskovsky and S. Oncley, "Comprehensive measurements of the intermittency exponent in high Reynolds number turbulent flows," *Fluid Dyn. Res.* **21**, 331 (1997).

⁴ J. Cleve, M. Greiner, and K. R. Sreenivasan, "On the effects of surrogacy of energy dissipation in determining the intermittency exponent in fully developed turbulence," *Europhys. Lett.* **61**, 756 (2003).

⁵ H. Mouri, M. Takaoka, A. Hori, and Y. Kawashima, "On Landau's prediction for large-scale fluctuation of turbulence energy dissipation," *Phys. Fluids* **18**, 015103 (2006).

⁶ V. R. Kuznetsov, A. A. Praskovsky, and V. A. Sabelnikov, "Fine-scale turbulence structure of intermittent shear flows," *J. Fluid Mech.* **243**, 595 (1992).

⁷ A. A. Praskovsky, E. B. Gledzer, M. Y. Karyakin, and Y. Zhou, "The sweeping decorrelation hypothesis and energy-inertial scale interaction in high Reynolds number flows," *J. Fluid Mech.* **248**, 493 (1993).

⁸ K. R. Sreenivasan and G. Stolovitzky, "Statistical dependence of inertial range properties on large scales in a high-Reynolds-number shear flow," *Phys. Rev. Lett.* **77**, 2218 (1996).

⁹ K. R. Sreenivasan and B. Dhruva, "Is there scaling in high-Reynolds-number turbulence?" *Prog. Theor. Phys. Suppl.* **130**, 103 (1998).

¹⁰ P. D. Mininni, A. Alexakis, and A. Pouquet, "Large-scale flow effects, energy transfer, and self-similarity on turbulence," *Phys. Rev. E* **74**, 016303 (2006).

¹¹ A. M. Obukhov, "Some specific features of atmospheric turbulence," *J. Fluid Mech.* **13**, 77 (1962).

¹² If we had obtained snapshots of the velocity field in the wind tunnel, we should have observed various eddies. Their strengths should have been random but on average decayed along the streamwise direction. Over scales along the streamwise direction, statistics of velocity fluctuations should have suffered from the decay, which is not relevant to our study. Over scales along the spanwise direction, so far as they were not too large, statistics of velocity fluctuations should have been identical to those for $u(x)$ and $v(x)$.

- ¹³ The alternative interpretation is that such fluctuations correspond to spatial fluctuations of some virtual turbulence. Local values of $u(x)$ and $v(x)$ represent local regions of actual turbulence, and they are continuously connected up to largest scales. Throughout the scales, $u(x)$ and $v(x)$ obtained here are consistent with those measured at a certain time in spatially homogeneous but temporally decaying turbulence (see Appendix of Ref. 5).
- ¹⁴ The reader might consider that $\delta u_{r,R}^2 \propto r^{2/3}$ is absent in some segments because their $\text{Re}_{\lambda,R}$ values are not high enough. This is not the case. For example, the segment shown in Fig. 1(a) has $\text{Re}_{\lambda,R} \simeq 1000$.
- ¹⁵ K. Ohkitani and S. Kida, "Triad interactions in a forced turbulence," *Phys. Fluids A* **4**, 794 (1992).
- ¹⁶ K. Kajita and T. Gotoh, "Statistics of the energy dissipation rate in turbulence," in *Statistical Theories and Computational Approaches to Turbulence*, edited by Y. Kaneda and T. Gotoh (Springer, Tokyo, 2003), p. 260.
- ¹⁷ L. D. Landau and E. M. Lifshitz, *Statistical Physics*, 3rd ed. (Pergamon, Oxford, 1979), Part 1, Chap. 12.
- ¹⁸ For such a segment, statistics do not strictly have local isotropy, and hence $\varepsilon_R = 15\nu(\partial_x v)_R^2/2$ is not strictly exact. The associated uncertainty is nevertheless small and not serious to our study. In fact, among segments with $R = 10^2\eta$ and $10^3\eta$, respectively, the correlation coefficients are as high as 0.89 and 0.95 between $15\nu(\partial_x v)_R^2/2$ and $15\nu[(\partial_x u)_R^2 + (\partial_x v)_R^2]/2$. The value of the latter tends to be more close to the true ε_R value (see Ref. 4), although it is not used in our study because of possible spurious correlations with the values of δu_r over small r .
- ¹⁹ R. A. Antonia and P. Burattini, "Approach to the 4/5 law in homogeneous isotropic turbulence," *J. Fluid Mech.* **550**, 175 (2006).
- ²⁰ To examine Obukhov's discussion in a strict manner, we have to use $\langle \delta u_{r,R}^2 \rangle_\varepsilon / (\nu \langle \varepsilon_R \rangle_\varepsilon)^{1/2}$ instead of $\langle \delta u_{r,R}^2 / u_{K,R}^2 \rangle_\varepsilon$. Even in this case, the result is almost the same as in Fig. 4. An analogous situation is seen in Fig. 5, where the Reynolds number obtained from $\langle v_R^2 \rangle_\varepsilon$ and $\langle \varepsilon_R \rangle_\varepsilon$ is almost the same as $\langle \text{Re}_{\lambda,R} \rangle_\varepsilon$.
- ²¹ A. N. Kolmogorov, "A refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at high Reynolds number," *J. Fluid Mech.* **13**, 82 (1962).
- ²² E. A. Novikov, "Intermittency and scale similarity in the structure of a turbulent flow," *J. Appl. Math. Mech.* **35**, 231 (1971).