Two graph isomorphism polytopes

Shmuel Onn

January 11, 2008

Abstract

The convex hull $\psi_{n,n}$ of certain $(n!)^2$ tensors was considered recently in connection with graph isomorphism. We consider the convex hull ψ_n of the n! diagonals among these tensors. We show: 1. The polytope ψ_n is a face of $\psi_{n,n}$. 2. Deciding if a graph G has a subgraph isomorphic to Hreduces to optimization over ψ_n . 3. Optimization over ψ_n reduces to optimization over $\psi_{n,n}$. In particular, this implies that the subgraph isomorphism problem reduces to optimization over $\psi_{n,n}$.

AMS Subject Classification: 05A, 15A, 51M, 52A, 52B, 52C, 62H, 68Q, 68R, 68U, 68W, 90B, 90C

1 Introduction

Let \mathcal{P}_n be the set of $n \times n$ permutation matrices and consider the following two polytopes,

 $\psi_n := \operatorname{conv} \{ P \otimes P : P \in \mathcal{P}_n \}, \qquad \psi_{n,n} := \operatorname{conv} \{ P \otimes Q : P, Q \in \mathcal{P}_n \}.$

The polytope $\psi_{n,n}$ was considered recently in [1] in connection with the graph isomorphism problem. Note that ψ_n and $\psi_{n,n}$ have n! and $(n!)^2$ vertices respectively. In this short note we show:

- 1. The polytope ψ_n is a face of the polytope $\psi_{n,n}$.
- 2. Deciding if a graph G has a subgraph isomorphic to a graph H reduces to optimization over ψ_n .
- 3. Optimization over ψ_n reduces to optimization over $\psi_{n,n}$.

In particular, this implies a result of [1] that subgraph isomorphism reduces to optimization over $\psi_{n,n}$.

So if $P \neq NP$ then optimization and separation over ψ_n and hence over $\psi_{n,n}$ cannot be done in polynomial time and a compact inequality description of ψ_n and hence of $\psi_{n,n}$ cannot be determined.

Deciding if G has a subgraph that is isomorphic to H can also be reduced to optimization over a related polytope ϕ_n defined as follows. Each permutation σ of the vertices of the complete graph K_n naturally induces a permutation Σ of its edges by $\Sigma(\{i, j\}) := \{\sigma(i), \sigma(j)\}$. Then ϕ_n is defined as the convex hull of all $\binom{n}{2} \times \binom{n}{2}$ permutation matrices of induced permutations Σ . This polytope and a broader class of so-called *Young polytopes* have been studied in [2]. In particular, therein it was shown that the graph of ϕ_n is complete, so pivoting algorithms cannot be exploited for optimization over this polytope. It is an interesting question whether ψ_n and ϕ_n , having n! vertices each, are isomorphic.

2 Statements

Define bilinear forms on $\mathbb{R}^{n \times n}$ and on $\mathbb{R}^{n \times n} \otimes \mathbb{R}^{n \times n}$ (note the shuffled indexation on the right) by

$$\langle A, B \rangle := \sum_{i,j} A_{i,j} B_{i,j} , \qquad \langle X, Y \rangle := \sum_{i,j,s,t} X_{i,s,j,t} Y_{i,j,s,t} .$$

Let *I* be the $n \times n$ identity matrix and for a graph *G* let A_G be its adjacency matrix. We show: **Theorem 2.1** The polytope ψ_n is a face of $\psi_{n,n}$ given by

$$\psi_n = \psi_{n,n} \cap \{X : \langle I \otimes I, X \rangle = n\}$$

Theorem 2.2 Let G and H be two graphs on n vertices with m the number of edges of H. Then

$$\max\{\langle A_G \otimes A_H, X \rangle : X \in \psi_n\} \leq 2m$$

with equality if and only if G has a subgraph that is isomorphic to H.

Theorem 2.3 Let $W = (W_{i,s,j,t})$ be any tensor and let $w := 2n^2 \max |W_{i,s,j,t}|$. Then

$$\max\{\langle W, X \rangle : X \in \psi_n\} = \max\{\langle W + wI \otimes I, X \rangle : X \in \psi_{n,n}\} - nw$$

Combining Theorems 2.2 and 2.3 with $W = A_G \otimes A_H$ and $w = n^2$ (sufficing since $W \ge 0$, as is clear from the proof of Theorem 2.3 below), we get the following somewhat tighter form of a result of [1].

Corollary 2.4 Let G and H be two graphs on n vertices with m the number of edges of H. Then

$$\max\{\langle A_G \otimes A_H + nI \otimes nI, X \rangle : X \in \psi_{n,n}\} \leq 2m + n^3$$

with equality if and only if G has a subgraph that is isomorphic to H.

3 Proofs

We record the following statement that follows directly from the definitions of the bilinear forms above.

Proposition 3.1 For any two simple tensors $X = A \otimes B$ and $Y = P \otimes Q$ we have

$$\langle X, Y \rangle = \langle A \otimes B, P \otimes Q \rangle = \sum_{i,j,s,t} A_{i,s} B_{j,t} P_{i,j} Q_{s,t} = \langle P B Q^{^{\mathsf{T}}}, A \rangle .$$

Proof of Theorem 2.1. For every $P, Q \in \mathcal{P}_n$, the matrix PIQ^{T} is a permutation matrix, with $PIQ^{\mathsf{T}} = I$ if and only if P = Q. It follows that for every two distinct $P, Q \in \mathcal{P}_n$ we have

$$\langle I \otimes I, P \otimes Q \rangle = \langle PIQ^{\mathsf{T}}, I \rangle \leq n-1 < n = \langle PIP^{\mathsf{T}}, I \rangle = \langle I \otimes I, P \otimes P \rangle.$$
 (1)

Proof of Theorem 2.2. For any $P \in \mathcal{P}_n$, the matrix PA_HP^{T} is the adjacency matrix of the permutation of H by P. So $\langle PA_HP^{\mathsf{T}}, A_G \rangle \leq 2m$ with equality if and only if H is isomorphic via P to a subgraph of G. Since the maximum of a linear form over a polytope is attained at a vertex we get

$$\max\{\langle A_G \otimes A_H, X \rangle : X \in \psi_n\} = \max\{\langle A_G \otimes A_H, P \otimes P \rangle : P \in \mathcal{P}_n\} \\ = \max\{\langle PA_H P^{\mathsf{T}}, A_G \rangle : P \in \mathcal{P}_n\} \le 2m$$

with the last inequality holding with equality if and only if G has a subgraph isomorphic to H.

Proof of Theorem 2.3. For every $P, Q \in \mathcal{P}_n$, the tensor $P \otimes Q = (P_{i,j}Q_{s,t})$ has n^2 entries that are equal to 1 and all other entries equal to 0, and therefore $-\frac{1}{2}w \leq \langle W, P \otimes Q \rangle \leq \frac{1}{2}w$. Combining this with inequality (1) we see that for every two distinct $P, Q \in \mathcal{P}_n$ we have

Since the maximum of a linear form over a polytope is attained at a vertex we obtain the inequality

$$\max\{\langle W + wI \otimes I, X \rangle : X \in \psi_{n,n}\} = \max\{\langle W + wI \otimes I, P \otimes Q \rangle : P, Q \in \mathcal{P}_n\} \\ = \max\{\langle W + wI \otimes I, P \otimes P \rangle : P \in \mathcal{P}_n\} \\ = \max\{\langle W, P \otimes P \rangle : P \in \mathcal{P}_n\} + nw \\ = \max\{\langle W, X \rangle : X \in \psi_n\} + nw . \blacksquare$$

References

- [1] S. Friedland, On the graph isomorphism problem, e-print: arXiv:0801.0398
- [2] S. Onn, Geometry, complexity and combinatorics of permutation polytopes, Journal of Combinatorial Theory Series A 64:31-49 (1993)

Shmuel Onn

Technion - Israel Institute of Technology, 32000 Haifa, Israel email: onn@ie.technion.ac.il, http://ie.technion.ac.il/~onn