

On the metric character of the quantum Jensen-Shannon divergence

P.W. Lamberti¹, A.P. Majtey², A. Borras², M. Casas² and A. Plastino³

¹ *Facultad de Matemática, Astronomía y Física, Universidad Nacional de Córdoba, Ciudad Universitaria, 5000 Córdoba, and CONICET, Argentina*

² *Departament de Física and IFISC, Universitat de les Illes Balears, 07122 Palma de Mallorca, Spain*

³ *Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, Argentina*

(Dated: January 26, 2023)

In a recent paper, the generalization of the Jensen Shannon divergence (JSD) in the context of quantum theory has been studied (Phys. Rev. A **72**, 052310 (2005)). This distance between quantum states has shown to verify several of the properties required for a good distinguishability measure. Here we investigate the metric character of this distance. More precisely we show, formally for pure states and by means of simulations for mixed states, that its square root verifies the triangle inequality.

PACS numbers: 03.67.-a; 03.67.Mn; 03.65.-w

I. INTRODUCTION

Fundamental physical theories are formulated in terms of an abstract space. This is the case of Relativity Theory, Quantum Mechanics (QM), Yang-Mills like theories, and every proposal for Unified Field Theory. On each abstract space different structures can be defined. For example topological, differentiable, affine and metric structures are ubiquitous in space-time models. A prescription for measuring just how close two points of the concomitant space are is what we mean here by a metric structure. A more precise distinction between a distance and a metric will be given below.

In principle each one of above mentioned structures can be defined in an independent way. In Special Relativity theory the space-time is the standard \mathbb{R}^4 manifold provided with the (fixed, non-dynamical) Minkowskian metric. In General Relativity, instead, the space-time is a differentiable four dimensional manifold where the metric is given by the matter-energy distribution (throughout the Einstein's field equations). In both cases the metric is compatible with Lorentz's covariance. It is worth mentioning here (as known since the pioneering works of Gauss and Riemann) that the metric defines every geometrical property of a differentiable manifold.

In QM the corresponding abstract space is a (finite or infinite dimensional) Hilbert space \mathcal{H} . In its mathematical formalism the states of a physical system \mathcal{S} are represented by operators (density operators) acting on \mathcal{H} . More precisely the states of the system \mathcal{S} are represented by the elements of $\mathcal{B}(\mathcal{H})_1^+$, that is, the set of positive trace one operators on \mathcal{H} . The notion of a state as a unit vector of \mathcal{H} refers to the extremal elements of $\mathcal{B}(\mathcal{H})_1^+$ ($\rho \in \mathcal{B}(\mathcal{H})_1^+$ is extremal if and only if it is idempotent, $\rho^2 = \rho$). In this case ρ is of the form $|\varphi\rangle\langle\varphi|$ for some unit vector $|\varphi\rangle \in \mathcal{H}$, and is called a pure state.

In the case of a Hilbert space, the basic underlying structure is that of a vectorial space provided with an internal product $\langle | \rangle$ between elements of \mathcal{H} . From this inner product several ways of measuring "proximity" between two elements of \mathcal{H} can be defined. For example, the Wootters's distance

$$d_W(|\varphi\rangle, |\psi\rangle) \equiv d_W(|\varphi\rangle\langle\varphi|, |\psi\rangle\langle\psi|) = \arccos(|\langle\varphi|\psi\rangle|) \quad (1)$$

is a very important one. On one side (1) represents the angle between the (pure) states $|\varphi\rangle$ and $|\psi\rangle$; on the other, it has to do with the statistical fluctuations in the outcomes of measurements into the QM formalism [1]. Finally, (1) is invariant under unitary evolution. Therefore, we can think of (1) as a very natural distance between pure states in QM, in some sense imposed by the quantum theory itself. A generalization of this distance to mixed states have been studied by Braunstein and Caves [2].

Before going on let us remind the reader of a formal distance-definition. Let \mathbb{X} be an abstract set. A function

$$d : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$$

is a distance defined over the set \mathbb{X} , if for every $x, y \in \mathbb{X}$, it satisfies the following properties:

$$\begin{aligned} d(x, y) &> 0 \text{ for } x \neq y \text{ and } d(x, x) = 0 \\ d(x, y) &= d(y, x) \end{aligned} \quad (2)$$

If, for every $x, y, z \in \mathbb{X}$, the function d also verifies the triangle inequality:

$$d(x, y) + d(y, z) - d(x, z) \geq 0 \quad (3)$$

it is said that d is a **metric** for the space \mathbb{X} . Incidentally, we mention that the function given by (1) verifies the properties of a distance but is not a metric. As a matter of fact, only a few among all distances between quantum states historically introduced in the literature verify condition (3).

The definition of distance between mixed quantum states is a topic of permanent interest. This interest has been lately rekindled on account of problems emerging in information theory (QIT)[3, 4, 5, 6, 7]. In introducing distances between quantum states, different roads have been traversed. We have already mentioned the case of the Wootters's distance and its generalization, presented in [2]. Recently, a rather interesting approach has been advanced by Lee et al. in reference [5]. There these authors characterize the degree of closeness of two states with regards to the information that can be attained for each of them from a complete set of mutually complementary measurements plus an invariance criterium. The resulting distance-measure is equivalent to the Hilbert-Schmidt metric. Let us recall that this metric emerges from the primitive structure of the Hilbert space. Indeed, an inner product between bounded operators acting over the Hilbert space \mathcal{H} can be defined in the fashion

$$\langle A|B \rangle = Tr(A^\dagger B)$$

The Hilbert-Schmidt norm of the operator A is given by $\|A\|_{HS}^2 = \langle A|A \rangle$ and from this, the Hilbert-Schmidt metric between two operators A and B is defined as

$$d_{HS}(A, B) = \|A - B\|_{HS} \quad (4)$$

Another way of dealing with the problem of introducing distances between quantum states is generalizing the notions of distance defined in the space of classical probability distributions. This is the case of the relative entropy, which is a generalization of information theoretic Kullback-Leibler divergence. The relative entropy of an operator ρ with respect to an operator σ , both belonging to $\mathcal{B}(\mathcal{H})_1^+$, is

$$S(\rho, \sigma) = Tr[\rho(\log \rho - \log \sigma)], \quad (5)$$

where \log stands for logarithm in base two. The relative entropy is not a distance (and obviously is not a metric either) because it is not symmetric and does not verify the triangle inequality (3). Worst, it may even be unbounded. In particular, the relative entropy is well defined only when the support of σ is equal to or larger than that of ρ [3] (the support of an operator is the subspace spanned by the eigenvectors of the operator with nonzero eigenvalues). This is a strong restriction which is violated in some physically relevant situations, as for example when σ is a pure reference state.

To overcome such problems we have recently investigated an alternative to the relative entropy [8] that emerges as a natural extension of a symmetrized version of the Kullback-Leibler divergence to the realm of quantum theory. In the classical context this quantity is known as the Jensen-Shannon divergence (JSD) and was introduced by C. Rao [9] and, independently, by J. Lin [10]. It has been applied to a diversity of problems arising in statistics and physics [11, 12, 13, 14, 15]. Among its most significant properties one can include its boundedness and its metric character [17]. In reference [12] it is shown that the JSD can be taken as a unifying distance between probability distributions.

In our previous study of the quantum JSD we showed that it verifies all the properties required for a good measure of distinguishability between quantum states. In this paper we investigate the metric property of the quantum JSD (QJSD), that could be regarded as essential to check on the convergence of iterative algorithms in quantum computation [16].

The structure of this paper is as follows: next Section is devoted to the formal definition of the classical and QJSD. In Section III we investigate the metric character of the QJSD. In the first place we consider the pure states case and then we investigate the metric properties for arbitrary mixed states recourse to numerical simulations in different Hilbert spaces. Finally, some conclusions are drawn in Sect IV.

II. CLASSICAL JENSEN-SHANNON DIVERGENCE AND ITS QUANTUM EXTENSION

The classical JSD between two (discrete) probability distributions $P = (p_1, p_2, \dots, p_N)$ and $Q = (q_1, q_2, \dots, q_N)$, $\sum_i p_i = \sum_i q_i = 1$ is defined as

$$JS(P, Q) = \frac{1}{2} \left[S \left(P, \frac{P+Q}{2} \right) + S \left(Q, \frac{P+Q}{2} \right) \right] \quad (6)$$

where $S(P, Q) = \sum_i p_i \log \frac{p_i}{q_i}$ is the Kullback-Leibler divergence. $JS(P, Q)$ can be also expressed in the form

$$\begin{aligned} JS(P, Q) &= H\left(\frac{P+Q}{2}\right) - \frac{1}{2}H(P) - \frac{1}{2}H(Q) \\ &= \frac{1}{2} \left[\sum_i p_i \log \left(\frac{2p_i}{p_i+q_i}\right) + \sum_i q_i \log \left(\frac{2q_i}{p_i+q_i}\right) \right] \end{aligned} \quad (7)$$

where $H(P) = -\sum_i p_i \log p_i$ is the Shannon entropy. The classical JSD exhibits several interesting properties. Among them we recall the following ones

- $JS(P, Q)$ is symmetric and always well defined;
- it is bounded

$$0 \leq JS(P, Q) \leq 1,$$

and, as it was already stated,

- its square root,

$$d_{JS}(P, Q) \equiv \sqrt{JS(P, Q)} \quad (8)$$

verifies the triangle inequality Eq. (3) (but JS does not).

A proof of this last fact can be found in references [17, 18]. Alternatively, this can be proved by using some results of harmonic analysis due to I. Schoenberg [19]. The basic fact that makes Schoenberg's theorem applicable to the classical JSD resides in that it is a definite negative kernel, that is, for all finite collection of real numbers $(c_i)_{i \leq N}$, and for all corresponding probability distributions $(P_i)_{i \leq N}$, the implication

$$\sum_{i=1}^N c_i = 0 \Rightarrow \sum_{i,j} c_i c_j JS(P_i, P_j) \leq 0 \quad (9)$$

is valid. A corollary of Schoenberg's theorem allows one to assert that the probability distributions-space, with the metric (8), can be isometrically mapped into a subset of a Hilbert space [20].

The classical JSD can be used to distinguish two probability distributions and therefore can be used as well to do so for two quantum states described by their density operators, say, ρ and σ . Indeed, let us suppose we choose a positive operator value measure (POVM), $\sum_{i=1}^M \mathbb{E}_i = \mathbb{I}$, that generates two probability distributions via

$$\begin{aligned} p_i &= Tr(\mathbb{E}_i \rho) \\ q_i &= Tr(\mathbb{E}_i \sigma), \end{aligned}$$

for $i = 1, \dots, M$. Then we can use the JSD (6) to distinguish between these two distributions. In this procedure we have the freedom of choosing the POVM which most clearly distinguishes p_i from q_i , that is, which makes the value of $JS(p_i, q_i)$ the largest. This reasoning motivates to introduce the quantity

$$JS_1(\rho, \sigma) = \sup_{\{\mathbb{E}_i\}} JS(p_i, q_i), \quad (10)$$

where the supremum is taken over all POVM's. Physically JS_1 gives the best discrimination between the states ρ and σ that we can achieve by means of measurements.

By mimicking the extension of Kullback-Leibler divergence to the realm of quantum theory, we define the QJSD as [8]

$$JS(\rho, \sigma) = \frac{1}{2} \left[S\left(\rho, \frac{\rho+\sigma}{2}\right) + S\left(\sigma, \frac{\rho+\sigma}{2}\right) \right], \quad (11)$$

that can be recast in terms of the von Neumann entropy $H_N(\rho) = -Tr(\rho \log \rho)$ in the fashion

$$JS(\rho, \sigma) = H_N\left(\frac{\rho+\sigma}{2}\right) - \frac{1}{2}H_N(\rho) - \frac{1}{2}H_N(\sigma), \quad (12)$$

This quantity is always well defined, symmetric, positive definite and bounded ($0 \leq JS(\rho, \sigma) \leq 1$). By using the corresponding properties of the relative entropy [21] and expression (11) it can be shown that, for arbitrary ρ and σ , the following inequality

$$JS(\rho, \sigma) \geq JS_1(\rho, \sigma), \quad (13)$$

is valid. The equality is satisfied *if and only if* ρ and σ commute, that is, the upper bound in (13) is, in general, not attainable for any POVM.

To conclude this section we give the explicit expression for the QJSD in terms of the eigenvalues and eigenvectors of the operators involved in its expression.

$$JS(\rho, \sigma) = \frac{1}{2} \left[\sum_{k,i} |\langle t_k | r_i \rangle|^2 r_i \log \left(\frac{2r_i}{\tau_k} \right) + \sum_{k,j} |\langle t_k | s_j \rangle|^2 s_j \log \left(\frac{2s_j}{\tau_k} \right) \right], \quad (14)$$

where $\rho = \sum_i r_i |r_i\rangle\langle r_i|$, $\sigma = \sum_i s_i |s_i\rangle\langle s_i|$, $(\rho + \sigma) = \sum_i t_i |t_i\rangle\langle t_i|$ and $\tau_k = \sum_i r_i |\langle t_k | r_i \rangle|^2 + \sum_i s_i |\langle t_k | s_i \rangle|^2$

It should be noted that, when ρ and σ do not commute, the structure of (14) is quite different from that of (7).

III. THE METRIC CHARACTER OF THE QUANTUM \sqrt{JS}

In this section we investigate the putative metric character of the QJSD, that is we study if the square root of the QJSD,

$$d_{JS}(\rho, \sigma) = \sqrt{JS(\rho, \sigma)} \quad (15)$$

verifies the triangle inequality. The other three properties for a metric are obviously verified by (15). A formal proof of property (3) for $\sqrt{JS(\rho, \sigma)}$ has till now eluded us. Unfortunately there is no analog of Schoember's theorem when operators are involved. Still more, there is no direct way of verifying condition (9) for expression (14). No extension to the case of the QJSD of the proof given in [17] has been possible. Incidentally it should be observed that, if the upper bound in (13) could be attained for some POVM, the proof of the triangle inequality for (15) would be obvious (because $\sqrt{JS_1}$ verifies it).

The results to be presented here correspond to a separate analysis of the metric condition for (15) for the two cases: when (15) is restricted to pure states and when it acts on the complete set $\mathcal{B}(\mathcal{H})_1^\dagger$. In the first instance we were able to give a formal proof of inequality (3); in the second one, we checked it by means of simulations.

1. Pure states

For a pure state the von Neumann entropy vanishes. Then, for two pure states,

$$\rho = |\psi\rangle\langle\psi| \quad \text{and} \quad \sigma = |\varphi\rangle\langle\varphi|, \quad (16)$$

the QJSD (12), becomes

$$JS(\rho, \sigma) = H_N \left(\frac{\rho + \sigma}{2} \right) \quad (17)$$

After some algebra, we can rewrite (17) in terms of the inner product $\langle \psi | \varphi \rangle$:

$$\begin{aligned} JS(\rho, \sigma) &= \Phi(|\langle \psi | \varphi \rangle|) \equiv \\ &= - \left(\frac{1 - |\langle \psi | \varphi \rangle|}{2} \right) \log \left(\frac{1 - |\langle \psi | \varphi \rangle|}{2} \right) - \left(\frac{1 + |\langle \psi | \varphi \rangle|}{2} \right) \log \left(\frac{1 + |\langle \psi | \varphi \rangle|}{2} \right). \end{aligned} \quad (18)$$

The entropy of the average $\frac{1}{2}(|\psi\rangle\langle\psi| + |\varphi\rangle\langle\varphi|)$ can be interpreted to the light of quantum information theory. Indeed, let us suppose that Alice has a source of pure qubit signal states $|\psi\rangle$ and $|\varphi\rangle$. Each emission is chosen to be $|\psi\rangle$ or $|\varphi\rangle$ with an equal prior probability a half. Then the density matrix of the source is $\frac{1}{2}(|\psi\rangle\langle\psi| + |\varphi\rangle\langle\varphi|)$. Alice may communicate the sequence of states to Bob by transmitting one qubit per emitted state. But according to the quantum source coding theorem, (17) gives the lowest number of qubits per states that Alice needs to communicate the quantum information (with arbitrarily high fidelity) [22].

Let us take two fixed arbitrary pure states $\rho = |\psi\rangle\langle\psi|$ and $\sigma = |\varphi\rangle\langle\varphi|$ and an arbitrary third one, $\xi = |\chi\rangle\langle\chi|$. Let us denote the absolute value of the inner products $|\langle\psi|\varphi\rangle|$, $|\langle\psi|\chi\rangle|$ and $|\langle\chi|\varphi\rangle|$ by x, y and z respectively, and let us introduce the function

$$H(x, y, z) = \sqrt{\Phi(y)} + \sqrt{\Phi(z)} - \sqrt{\Phi(x)}.$$

In terms of these variables the triangle inequality for (15) reads:

$$0 \leq H(x, y, z) \quad (19)$$

We can decompose the vector $|\chi\rangle$ in a part belonging to the plane developed by $|\psi\rangle$ and $|\varphi\rangle$ and in an other one perpendicular to this plane:

$$|\chi\rangle = a|\psi\rangle + b|\varphi\rangle + |\chi_{\perp}\rangle$$

with $|a| \leq 1$ and $|b| \leq 1$. Then

$$y = |a + b\langle\psi|\varphi\rangle| \quad \text{and} \quad z = |a^*\langle\varphi|\psi\rangle + b^*|$$

As a function of a and b , for x fixed, H is a concave function on the circles $|a| \leq 1$ and $|b| \leq 1$ (in the sense that its second derivative is negative) and it vanishes for $y = x$ and $z = x$. This guarantees that inequality (19) is satisfied for arbitrary y and z .

2. Arbitrary mixed states

Here we attempt a numerical verification of the triangle inequality for (15) when arbitrary mixed states are involved. To do this we restrict ourselves to the Hilbert space corresponding to a bipartite system. More specifically, we evaluate numerically the inequality (3) by generating random bipartite states in a Hilbert space. The space of all (pure and mixed) such states can be regarded as a product space of the form [24]:

$$\mathcal{H} = \mathcal{P} \times \Delta,$$

where \mathcal{P} stands for the family of all complete sets of ortho-normal projectors $\{\hat{P}_i\}_i^N$, $\sum_i \hat{P}_i = \mathbb{I}$ (\mathbb{I} the identity matrix), and Δ is the set of all real N -tuples of the form $\{\lambda_1, \dots, \lambda_N\}$; $\lambda_i \in \mathbb{R}$; $\sum_i \lambda_i = 1$; $0 \leq \lambda_i \leq 1$. Any state in \mathcal{H} is of the form $\rho = \sum_i \lambda_i \hat{P}_i$.

In exploring exhaustively \mathcal{H} we need to introduce an appropriate measure μ on this space. Such a measure is required to compute volumes within \mathcal{H} , as well as to determine what is to be understood by a uniform distribution of states on \mathcal{H} . The measure that we adopt here is taken from the work of Zyczkowski *et al.* [25, 26].

An arbitrary (pure or mixed) state ρ of a quantum system described by an N -dimensional Hilbert space can always be expressed as a product of the form:

$$\rho = UD[\{\lambda_i\}]U^\dagger. \quad (20)$$

Here U is an $N \times N$ unitary matrix and $D[\{\lambda_i\}]$ is an $N \times N$ diagonal matrix whose diagonal elements are, precisely, our above defined $\{\lambda_1, \dots, \lambda_N\}$. The group of unitary matrices $U(N)$ is endowed with a unique, uniform measure, known as the Haar's measure, ν [27]. On the other hand, the N -simplex Δ , consisting of all the real N -uples $\{\lambda_1, \dots, \lambda_N\}$ appearing in (20), is a subset of a $(N - 1)$ -dimensional hyperplane of \mathbb{R}^N . Consequently, the standard normalized Lebesgue measure \mathcal{L}_{N-1} on \mathbb{R}^{N-1} provides a measure for Δ . The aforementioned measures on $U(N)$ and Δ lead to a measure μ on the set \mathcal{S} of all the states of our quantum system [25, 26, 27], namely,

$$\mu = \nu \times \mathcal{L}_{N-1}. \quad (21)$$

In our numerical computations we randomly generate mixed states according to the measure (21). In order to assess, for these randomly generated states, how the triangle inequality (3) is satisfied, we define the auxiliary quantity

$$\Delta d_{JS}(\rho, \xi, \sigma) = d_{JS}(\rho, \xi) + d_{JS}(\xi, \sigma) - d_{JS}(\rho, \sigma) \quad (22)$$

and evaluate it for 10^6 simulated states. This procedure is repeated for different dimensions of the Hilbert space. The metric character of the square root of the QJSD is given by the positivity of the quantity Δd_{JS} . In Fig. 1 we plot the




FIG. 1: Values of Δd_{JS} obtained for two qubits states. Here we plot only 10^4 points.




FIG. 2: Probability distribution for Δd_{JS} for different Hilbert space dimensions. The PDFs were constructed with 10^6 generated mixed random states. Inset: Δd_{JS}^{min} as a function of the Hilbert space dimension $N = N_1 \times N_2$.

values of Δd_{JS} for each one of the simulated terms corresponding to a two qubits space. This plot makes it evident the positivity of Δd_{JS} , as required.

Alternatively we investigate the positivity of Δd_{JS} by constructing the probability distributions for the values of Δd_{JS} . The corresponding histograms, for different dimensions of the Hilbert space, are depicted in Fig. 2. In the accompanying inset we plot the minimum value for this quantity as a function of the Hilbert space dimension N . This minimum, Δd_{JS}^{min} , is always positive and increases with N , giving further evidence that the inequality (3) will be satisfied, even for higher Hilbert space dimensions.

IV. CONCLUSIONS

The main purpose of this work was to investigate the metrical property of the QJSD. We were able to show that the square root of the QJSD verifies the triangle inequality, giving to this distance the character of a metric. Although we

have proved this claim (for mixed states) merely by recourse to simulations, we believe that the cases here analyzed are sufficiently representative so as to render credible the claim that metric properties are verified in general for the QJSD.

A second item deserves to be pointed out, which emerges from the following two facts:

- On the one hand, we have showed that, when restricted to pure states, the square root of the entropy of the average $\frac{1}{2}(|\psi\rangle\langle\psi| + |\varphi\rangle\langle\varphi|)$ is a true metric.
- On the other hand, a classical result from Uhlmann [4] asserts that the fidelity of states ρ and σ

$$F(\rho, \sigma) = \text{Tr} \sqrt{\rho^{\frac{1}{2}} \sigma \rho^{\frac{1}{2}}}$$

can be expressed in the form

$$F(\rho, \sigma) = \max_{|\psi\rangle, |\varphi\rangle} |\langle\psi|\varphi\rangle| \quad (23)$$

where the maximization is over all purifications $|\psi\rangle$ of ρ and all purifications $|\varphi\rangle$ of σ [28].

These two facts motivates us to introduce an alternative metric for arbitrary mixed states. Given two arbitrary mixed states ρ and σ we define

$$d_H(\rho, \sigma) = \min_{|\psi\rangle, |\varphi\rangle} \sqrt{H_N \left(\frac{|\psi\rangle\langle\psi| + |\varphi\rangle\langle\varphi|}{2} \right)} \quad (24)$$

where the minimum is taken over all purification $|\psi\rangle$ of ρ and all purifications $|\varphi\rangle$ of σ . In (24) we must look for the minimum, not for the maximum as in (23), due to the decreasing nature of Φ , eq. (18), as a function of $|\langle\psi|\varphi\rangle|$.

Obviously the basic properties required for a good distinguishability measure are inherited by (24) from those verified by the QJSD. Additionally, several interesting questions arise from this proposal. For example, what relations exist between (24) and (11); or, in general, how (24) relates to other quantum distances. A more detailed study of the properties of this quantity will be presented elsewhere.

Acknowledgments

This work was partially supported by the MEC grant FIS2005-02796 (Spain) and FEDER (EU) and by CONICET (Argentine Agency). AB and APM acknowledge support from MEC through FPU grant AP-2004-2962 and contract SB-2006-0165. PWL wants to thank SECyT-UNC (Argentina) and CONICET for financial support.

-
- [1] W.K. Wootters, Phys. Rev. D **23** 357 (1981).
[2] S.L. Braunstein and C.M. Caves, Phys. Rev. Lett. **72**, 3439 (1994).
[3] G. Lindblad, Commun. Math Phys. **33**, 305 (1973).
[4] R. Jozsa, J. Mod. Optics, **41**, 2315 (1994).
[5] J. Lee, M.S. Kim and C. Brukner, Phys. Rev. Lett. **91** 087902 (2003).
[6] S. Luo and Q. Zhang, Phys. Rev. A **69** 032106 (2004).
[7] D. Markham, J. Miszczak, Z. Puchala and K. Zyczkowski, arXiv:0711.4286v1 (2007).
[8] A.P. Majtey, P.W. Lamberti, and D.P. Prato, Phys. Rev. A **72**, 052310 (2005).
[9] C. Rao, IMS-Lect. Notes **10**, 217 (1987).
[10] J. Lin, IEEE Trans. Inf. Theory **37**, 145 (1991).
[11] I. Grosse, P. Bernaola-Galvan, P. Carpena, R. Roman-Roldan, J. Oliver and H.E. Stanley, Phys. Rev. E **65**, 041905 (2002).
[12] A. Majtey, P.W. Lamberti, M.T. Martin and A. Plastino, Eur. Phys. J. D **32**, 413 (2005).
[13] M. Pereyra, P.W. Lamberti and O.A. Rosso, Physica A **379**, 122 (2007).
[14] O. A. Rosso, H. A. Larrondo, M. T. Martin, A. Plastino, and M. A. Fuentes, Phys. Rev. Lett. **99**, 154102 (2007).
[15] G.E. Crooks, Phys. Rev. Lett. **99**, 100602 (2007).
[16] A. Galindo and M. A. Martín-Delgado, Rev. Mod. Phys. **74** 347 (2002).
[17] D. Endres and J. Schindelin, IEEE Trans. Inf. Theory, **49**, 7, 1858 (2003).
[18] F. Österreicher and I. Vajda, Ann. Inst. Stat. Math. **55**, pag. 639 (2003).

- [19] I.J. Schoenberg, *Trans. Am. Math. Soc.* **44**, 3 (1938).
- [20] B. Fuglede and F. Topsøe, *Proceedings of the International Symposium on Information Theory, ISIT 2004*, pag 31. Published by IEEE (2004).
- [21] M.J. Donald, *Comm. Math. Phys.* **105**, pag 13 (1986).
- [22] B. Schumacher, *Phys. Rev. A* **51**, 2738 (1995).
- [23] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge University Press, Cambridge, 2006).
- [24] J. Batle, M. Casas, A. R. Plastino, and A. Plastino, *Phys. Lett. A* **353**, 161 (2006).
- [25] K. Życzkowski, P. Horodecki, A. Sanpera and M. Lewenstein, *Phys. Rev. A* **58**, 883 (1998).
- [26] K. Życzkowski, *Phys. Rev. A* **60**, 3496 (1999).
- [27] M. Pozniak, K. Życzkowski and M. Kus, *J. Phys. A: Math. Gen.* **31**, 1059 (1998).
- [28] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, England, 2000.