

A Most General Edge Elimination Polynomial — Thickening of Edges

Christian Hoffmann

November 6, 2018

Abstract

We consider a graph polynomial $\xi(G; x, y, z)$ introduced by Averbouch, Godlin, and Makowsky (2007). This graph polynomial simultaneously generalizes the Tutte polynomial as well as a bivariate chromatic polynomial defined by Dohmen, Pönitz and Tittmann (2003). We derive an identity which relates the graph polynomial of a thicked graph (i.e. a graph with each edge replaced by k copies of it) to the graph polynomial of the original graph. As a consequence, we observe that at every point (x, y, z) , except for points lying within some set of dimension 2, evaluating ξ is $\#P$ -hard.

1 Introduction

We consider the following three-variable graph polynomial which has been introduced by I. Averbouch, B. Godlin, and J. A. Makowsky [AGM07]:

$$\xi(G; x, y, z) = \sum_{(A \sqcup B) \subseteq E} x^{k(A \cup B) - k_{cov}(B)} \cdot y^{|A| + |B| - k_{cov}(B)} \cdot z^{k_{cov}(B)}, \quad (1)$$

where $G = (V, E)$ is a graph with multiple edges and self loops allowed, $A \sqcup B$ denotes a *vertex-disjoint* union of edge sets A and B , $k(A \cup B)$ is the number of components of $(V, A \cup B)$, and $k_{cov}(B)$ is the number of components of $(V(B), B)$.

The polynomial ξ simultaneously generalizes two interesting graph polynomials: the Tutte polynomial and a bivariate chromatic polynomial $P(G; x, y)$ defined by K. Dohmen, A. Pönitz, and P. Tittmann [DPT03].

It is known that the Tutte polynomial of a graph with “thicked” edges evaluated at some point equals the Tutte polynomial of the original graph evaluated at another

point (parallel edge reduction). This property can be used to prove that at almost every point evaluating the Tutte polynomial is hard [JWV90, Sok05, BM06, BDM07].

In Section 2 of this note we observe that edge thickening has a similar effect on ξ as on the Tutte polynomial (Theorem 3). In Section 3 we conclude that for every point $(x, y, z) \in \mathbb{Q}^3$, except on a set of dimension at most 2, it is $\#\mathbf{P}$ -hard to compute $\xi(G; x, y, z)$ from G (Theorem 5). This supports a difficult point conjecture for graph polynomials [Mak07, Conjecture 1], [AGM07, Question 1].

2 A point-to-point reduction from thickening

In this section we apply Sokal's approach to ξ and obtain Lemma 2 (cf. [Sok05, Section 4.4]), the main technical contribution of this note.

We define the following auxiliary polynomial, which has a different y -variable for each edge of the graph, $\bar{y} = (y_e)_{e \in E(G)}$.

$$\psi(G; x, \bar{y}, z) = \sum_{(A \sqcup B) \subseteq E(G)} w(G; x, \bar{y}, z; A, B), \quad (2)$$

where

$$w(G; x, \bar{y}, z; A, B) = x^{k(A \cup B)} \left(\prod_{e \in (A \cup B)} y_e \right) z^{k_{\text{cov}}(B)}.$$

We write $\psi(G; x, y, z)$ for $\psi(G; x, \bar{y}, z)$ if for each $e \in E(G)$ we have $y_e = y$.

Lemma 1. *We have the polynomial identities $\psi(G; x, y, zx^{-1}y^{-1}) = \xi(G; x, y, z)$ and $\xi(G; x, y, zxy) = \psi(G; x, y, z)$.* \square

Let G be a graph and $e \in E(G)$ an edge. Let $E' := E \setminus \{e\}$ and G_{ee} be the graph G with e doubled, i.e. $G_{ee} = (V(G), E' \cup \{e_1, e_2\})$ with e_1, e_2 being new edges.

Lemma 2. *$\psi(G_{ee}; x, \bar{y}, z) = \psi(G; x, \bar{Y}, z)$ with $Y_e = (1 + y_{e_1})(1 + y_{e_2}) - 1$ and $Y_{\tilde{e}} = y_{\tilde{e}}$ for all $\tilde{e} \in E'$.*

Proof. Let $M(G) = \{(A, B) \mid A \sqcup B \subseteq E(G)\}$ and $M(G_{ee}) = \{(\tilde{A}, \tilde{B}) \mid \tilde{A} \sqcup \tilde{B} \subseteq E(G_{ee})\}$. We define a map $\tau : M(G) \rightarrow 2^{M(G_{ee})}$ in the following way. Consider $(A, B) \in M(G)$. If $e \notin A \cup B$, we set $\tau(A, B) = \{(A, B)\}$. If $e \in A$, we let $A' := A \setminus \{e\}$ and define $\tau(A, B) = \{(A' \cup \{e_1\}, B), (A' \cup \{e_2\}, B), (A' \cup \{e_1, e_2\}, B)\}$. (Note that in this case $e \notin B$, as A and B are vertex-disjoint.) If $e \in B$, we let $B' := B \setminus \{e\}$ and define $\tau(A, B) = \{(A, B' \cup \{e_1\}), (A, B' \cup \{e_2\}), (A, B' \cup \{e_1, e_2\})\}$. Observe that

$$M(G_{ee}) = \cup_{(A, B) \in M(G)} \tau(A, B), \quad (3)$$

and that this union is a union of *pairwise disjoint* sets.

Calculation yields

$$w(G; x, \bar{Y}, z; A, B) = \sum_{(\tilde{A}, \tilde{B}) \in \tau(A, B)} w(G_{ee}, x, \bar{y}, z; \tilde{A}, \tilde{B}) \quad (4)$$

for every $(A, B) \in M(G)$. Thus,

$$\begin{aligned} \psi(G_{ee}; x, \bar{y}, z) &= \sum_{(\tilde{A}, \tilde{B}) \in M(G_{ee})} w(G_{ee}; x, \bar{y}, z; \tilde{A}, \tilde{B}) \\ &= \sum_{(A, B) \in M(G)} \sum_{(\tilde{A}, \tilde{B}) \in \tau(A, B)} w(G_{ee}; x, \bar{y}, z; \tilde{A}, \tilde{B}) && \text{by (3)} \\ &= \sum_{(A, B) \in M(G)} w(G; x, \bar{Y}, z; A, B) && \text{by (4)} \\ &= \psi(G; x, \bar{Y}, z). \end{aligned}$$

□

Applying Lemma 2 repeatedly and Lemma 1 to convert between ψ and ξ we obtain

Theorem 3. *Let G_k be the k -thickening of G (i.e. the graph obtained out of G by replacing each edge by k copies of it). Then*

$$\psi(G_k; x, y, z) = \psi(G; x, (1+y)^k - 1, z), \quad (5)$$

$$\xi(G_k; x, y, z) = \xi\left(G; x, (1+y)^k - 1, z \frac{(1+y)^k - 1}{y}\right). \quad (6)$$

3 Hardness

The following theorem has been proven independently by I. Averbouch (J. A. Makowsky, personal communication, October 2007).

Theorem 4. *Let P denote the bivariate chromatic polynomial defined by K. Dohmen, A. Pönitz, and P. Tittmann [DPT03]. For every $(x, y) \in \mathbb{Q}$, $y \neq 0$, $(x, y) \notin \{(1, 1), (2, 2)\}$, it is #P-hard to compute $P(G; x, y)$ from G .*

Proof (Sketch). Given a graph $G = (V, E)$ let \tilde{G} denote the graph obtained out of G by inserting a new vertex \tilde{v} and connecting \tilde{v} to all vertices in V . Let $P(G; y)$ denote the chromatic polynomial [Rea68]. It is well known that

$$P(\tilde{G}; y) = yP(G; y - 1). \quad (7)$$

From this and [DPT03, Theorem 1] we can derive

$$P(\tilde{G}; x, y) = yP(G; x - 1, y - 1) + (x - y)P(G; x, y). \quad (8)$$

The proof of the theorem now works in the same fashion as a proof that $P(G; y)$ is $\#P$ -hard to evaluate almost everywhere using (7) would work: using (8) we reduce along the lines $x = y + d$, which eventually enables us to evaluate P at $(1 + d, 1)$ (if y is a positive integer, we reach $(1 + d, 1)$ directly; otherwise we obtain arbitrary many points on the line $x = y + d$, which enables us to interpolate the polynomial on this line). On the line $y = 1$ the polynomial P equals the independent set polynomial [DPT03, Corollary 2], which is $\#P$ -hard to evaluate almost everywhere [AM07, BH07]. \square

Theorem 5. *For every $(x, y, z) \in \mathbb{Q}$, $x \neq 0$, $z \neq -xy$, $(x, z) \notin \{(1, 0), (2, 0)\}$, $y \notin \{-2, -1, 0\}$, the following statement holds true: It is $\#P$ -hard to compute $\xi(G; x, y, z)$ from G .*

Proof (Sketch). For $x, y \in \mathbb{Q}$, $x, y \neq 0$ and $(x, y) \notin \{(1, 1), (2, 2)\}$ the following problem is $\#P$ -hard by Theorem 4: Given G , compute

$$P(G; x, y) = \xi(G; x, -1, x - y) = \psi\left(G; x, -1, \frac{y - x}{x}\right),$$

where the first equality is by [AGM07, Proposition 18] and the second by Lemma 1. We will argue that, for any fixed $\tilde{y} \in \mathbb{Q} \setminus \{-2, -1, 0\}$, this reduces to compute $\psi(G; x, \tilde{y}, \frac{y-x}{x})$ from G . We have

$$\psi\left(G; x, \tilde{y}, \frac{y - x}{x}\right) = \xi(G; x, \tilde{y}, (y - x)\tilde{y})$$

by Lemma 1. An easy calculation converts the conditions on x, \tilde{y}, y into conditions on x, y, z and yields the statement of the theorem.

Now assume that we are able to evaluate ψ at some fixed $(x, y, z) \in \mathbb{Q}^3$, i.e. given G we can compute $\psi(G; x, y, z)$. Then Theorem 3 allows us to evaluate ψ at (x, y', z) for infinitely many different $y' = (1 + y)^k - 1$ provided that $|1 + y| \neq 0$ and $|1 + y| \neq 1$. As ψ is a polynomial, this enables interpolation in y and eventually gives us the ability to evaluate ψ at (x, y', z) for *any* $y' \in \mathbb{Q}$. In particular, being able to evaluate ψ at $(x, \tilde{y}, \frac{y-x}{x})$, $\tilde{y} \in \mathbb{Q} \setminus \{-2, -1, 0\}$, implies the ability to evaluate it at $(x, -1, \frac{y-x}{x})$. \square

References

- [AGM07] Ilia Averbouch, Benny Godlin, and J. A. Makowsky. A most general edge elimination polynomial, 2007. Preprint, arXiv:0712.3112v1 (math.CO).
- [AM07] Ilia Averbouch and J. A. Makowsky. The complexity of multivariate matching polynomials, February 2007. Preprint.
- [BDM07] M. Bläser, H. Dell, and J.A. Makowsky. Complexity of the Bollobás-Riordan polynomial: Dichotomy results and uniform reductions, 2007. Preprint.
- [BH07] Markus Bläser and Christian Hoffmann. On the complexity of the interlace polynomial, 2007. Preprint, arXiv:cs.CC/0707.4565.
- [BM06] Markus Bläser and Johann Makowsky. Hip hip hooray for Sokal, 2006. Unpublished note.
- [DPT03] Klaus Dohmen, André Pönitz, and Peter Tittmann. A new two-variable generalization of the chromatic polynomial. *Discrete Mathematics & Theoretical Computer Science*, 6(1):69–90, 2003.
- [JVW90] F. Jaeger, D. L. Vertigan, and D. J. A. Welsh. On the computational complexity of the Jones and the Tutte polynomials. *Math. Proc. Cambridge Philos. Soc.*, 108:35–53, 1990.
- [Mak07] J. A. Makowsky. From a zoo to a zoology: Towards a general theory of graph polynomials. To appear in *Theory of Computing Systems*, 2007. doi:10.1007/s00224-007-9022-9.
- [Rea68] R. C. Read. An introduction to chromatic polynomials. *J. Combin. Theory*, 4:52–71, 1968.
- [Sok05] Alan D. Sokal. The multivariate Tutte polynomial (alias Potts model) for graphs and matroids. In Bridget S. Webb, editor, *Surveys in Combinatorics 2005*. Cambridge University Press, 2005.