

**THE GRAPHICAL TRAVELING SALESMAN POLYHEDRON IS THE  
INTERSECTION OF THE POSITIVE ORTHANT WITH THE MINKOWSKI  
SUM OF THE SYMMETRIC TRAVELING SALESMAN POLYTOPE AND THE  
POLAR OF THE METRIC CONE  
(SHORT COMMUNICATION)**

DIRK OLIVER THEIS

ABSTRACT. In this short communication, we observe that the Graphical Traveling Salesman Polyhedron is the intersection of the positive orthant with the Minkowski sum of the Symmetric Traveling Salesman Polytope and the polar of the metric cone. This follows trivially from known facts. There are two reasons why we find this observation worth communicating none-the-less: It is very surprising; it helps to understanding the relationship between these two important families of polyhedra.

1. INTRODUCTION

The *Symmetric Traveling Salesman Polytope* is the convex hull of all characteristic vectors edge sets of cycles (i.e., circuits) on the vertex set  $[n] := \{1, \dots, n\}$ . More precisely, denote by  $E_n$  the set of all two-element subsets of  $[n]$ . This is the set of all possible edges of a graph with vertex set  $[n]$ . We let

$$S_n := \text{conv} \left\{ \chi^{E(C)} \mid C \text{ is a cycle with vertex set } [n] \right\} \subset \mathbb{R}^{E_n}.$$

Here, for an edge set  $F$ ,  $\chi^F$  is characteristic vector in  $\mathbb{R}^{E_n}$  with  $\chi_e^F = 1$  if  $e \in F$ , and zero otherwise. The importance of the Symmetric Traveling Salesman Polytope comes mainly (but not exclusively) from its use in the solution of the Symmetric Traveling Salesman Problem.

The *Graphical Traveling Salesman Polyhedron* is the convex hull of all non-negative integer vectors describing the edge multi-sets of connected Eulerian multi-graphs on the same vertex set. More precisely, identifying multi-sets of elements of  $E_n$  with vectors in  $\mathbb{Z}_+^{E_n}$ , we denote

$$P_n := \text{conv} \left\{ x \mid x \text{ is the edge multi-set of a connected Eulerian multi-graph} \right. \\ \left. \text{with vertex set } [n] \right\} \subset \mathbb{R}^{E_n}.$$

Ever since the seminal work of Naddef & Rinaldi [4, 5] on the two polyhedra,  $P_n$  is considered to be an important tool for investigating the facets of  $S_n$ . Moreover, in works of Carr [2] and Applegate, Bixby, Chvátal & Cook [1],  $P_n$  has been used algorithmically in contributing to solution schemes for the Symmetric Traveling Salesman Problem.

---

*Date:* June 9, 2019.

2000 *Mathematics Subject Classification.* 52B12.

Research supported by *Deutsche Forschungsgemeinschaft (DFG)* as project RE 776/9-1. Author currently supported by *Communauté française de Belgique – Actions de Recherche Concertées*.

Numerous authors have expressed how close the connection between Graphical and Symmetric Traveling Salesman Polyhedra is, an opinion which finds its most basic witness in the fact that  $S_n$  is a face of  $P_n$  — consisting of all points  $x$  whose “degree” is two at every vertex:  $\sum_{v \neq u} x_{uv} = 2$  for all  $u \in [n]$ . However, the connections are far deeper (see [3] and the references therein). In this short communication, we contribute the following surprising geometric observation to the issue of the relationship between these two polyhedra:

**Theorem.**  $P_n$  is the intersection of the positive orthant with the Minkowski sum of  $S_n$  and the polar of the metric cone,  $C_n$ :

$$P_n = (S_n + C_n^\Delta) \cap \mathbb{R}_+^{E_n} \quad (1)$$

The metric cone consists of all  $a \in \mathbb{R}^{E_n}$  which satisfy the triangle inequality:

$$a_{uv} \leq a_{uw} + a_{wv} \quad (2)$$

for all pairwise distinct vertices  $u, v, w \in [n]$ . Consequently, its polar is generated as a cone by the vectors (we abbreviate  $\chi^{\{e\}}$  to  $\chi^e$ )

$$\chi^{uw} + \chi^{wv} - \chi^{uv} \quad (3)$$

## 2. PROOF

We start with showing that  $P_n \subset (S_n + C_n^\Delta) \cap \mathbb{R}_+^{E_n}$ . While  $P_n \subset \mathbb{R}_+^{E_n}$  holds trivially,  $P_n \subset S_n + C_n^\Delta$ , follows from an argument of [5], which we reproduce here for the sake of completeness.

Let  $x \in \mathbb{Z}_+^{E_n}$  define an edge multi-set of a connected Eulerian multi-graph  $G$  with vertex set  $[n]$ . We prove by induction on the number  $m$  of edges of  $G$ , that  $x$  can be written as a sum of a cycle and a number of vectors (3). If  $m = n$ , then there is nothing to prove. Let  $m \geq n + 1$ . There exists a vertex  $w$  of degree at least four in  $G$ . We distinguish two cases. If the graph  $G \setminus w$  has at least two connected components, let  $u$  and  $v$  be two neighbors of  $w$  in distinct components of  $G \setminus w$ . Then, replacing the edges  $uw$  and  $wv$  of  $G$  with the new edge  $uv$  gives a connected Eulerian multi-graph  $G'$  with fewer edges than  $G$ . The change in the vector  $x$  amounts to subtracting the expression(3):  $x' = x - (\chi^{uw} + \chi^{wv} - \chi^{uv})$ . By induction, we conclude that  $x'$ , and hence  $x$ , can be written as a sum of a cycle and a number of vectors (3).

We now prove  $P_n \supset (S_n + C_n^\Delta) \cap \mathbb{R}_+^{E_n}$ . For this, let  $a \cdot x \geq \alpha$  be a facet-defining inequality for  $P_n$  (we denote by “ $\cdot$ ” the inner product). We show that it is also valid for the right hand side. Again we invoke an argument from [5], which we again reproduce. Naddef & Rinaldi have shown that these inequalities fall into one of two categories: the non-negativity inequalities  $x_e \geq 0$  and inequalities with left-hand sides satisfying the triangle inequality (2). To see this, suppose that  $a \cdot x \geq \alpha$  is not dominated by a non-negativity inequality, and let  $u, v, w$  be three distinct vertices in  $[n]$ . Then there exists an  $x \in \mathbb{Z}_+^{E_n}$  defining the edge multi-set of a connected Eulerian multi-graph  $G$  which has an edge between  $u$  and  $v$ , such that  $a \cdot x = \alpha$ . If we replace the edge  $uv$  of  $G$  by the two edges  $uw$  and  $wv$ , then we obtain a connected Eulerian multi-graph, whose edge multi-set is given by  $x' := x + \chi^{uw} + \chi^{wv} - \chi^{uv}$ . Now  $a \cdot x' \geq \alpha$ , implies  $a_{uw} + a_{wv} - a_{uv} \geq 0$ , i.e., the triangle inequality.

To conclude the proof of the theorem, note that the non-negativity inequalities are clearly satisfied by the right hand side of (1). If  $a$  satisfies the triangle inequality, then using the argument we have used to show  $P_n \subset S_n + C_n^\Delta$ , it is easy to see that the minimum of  $x \mapsto a \cdot x$  over all  $x \in \mathbb{Z}_+^{E_n}$  defining edge multi-sets of a connected Eulerian multi-graphs is attained in a cycle (this is another argument of Naddef & Rinaldi which

we have reproduced). Hence,  $a \cdot x \geq \alpha$  is valid for  $S_n$ , and, since  $a$  satisfies the triangle inequality,  $a \in C_n = (C_n^\Delta)^\Delta$ , i.e.,  $a \cdot x \geq 0$  is valid for  $C_n$ . This concludes the proof of the theorem.

#### ACKNOWLEDGMENTS

Thanks are extended to the *Deutsche Forschungsgemeinschaft*, DFG, for funding this research, and to the *Communauté française de Belgique – Actions de Recherche Concertées* for supporting the author during the time the paper was written.

#### REFERENCES

- [1] D. Applegate, R. Bixby, V. Chvátal, and W. Cook. TSP cuts which do not conform to the template paradigm. In M. Jünger and D. Naddef, editors, *Computational Combinatorial Optimization*, pages 261–303. Springer-Verlag Berlin Heidelberg, 2001.
- [2] R. Carr. Separation algorithms for classes of STSP inequalities arising from a new STSP relaxation. *Math. Oper. Res.*, 29(1):80–91, 2004.
- [3] D. Naddef. Polyhedral Theory and Branch-and-Cut Algorithms for the Symmetric TSP. In Gregory Gutin and Abraham P. Punnen, editors, *The Traveling Salesman Problem and Its Variations*, pages 29–116. Kluwer Academic Publishers, 2002.
- [4] D. Naddef and G. Rinaldi. The Symmetric Traveling Salesman Polytope and its graphical relaxation: Composition of valid inequalities. *Math. Program.*, 51:359–400, 1991.
- [5] D. Naddef and G. Rinaldi. The graphical relaxation: A new framework for the Symmetric Traveling Salesman Polytope. *Math. Program.*, 58:53–88, 1993.

DIRK OLIVER THEIS, SERVICE DE GÉOMETRIE COMBINATOIRE ET THÉORIE DES GROUPES, DEPARTMENT OF MATHEMATICS, UNIVERSITÉ LIBRE DE BRUXELLES, BRUSSELS, BELGIUM  
*E-mail address:* theis@uni-heidelberg.de