Modified KdV hierarchy: Lax pair representation and bi-Hamiltonian structure

Amitava Choudhuri† , B. Talukdar‡ and U. Das∗

†Department of Physi
s, Visva-Bharati University, Santiniketan 731235, India E-mail: amitava_ch26@yahoo.com

‡Department of Physi
s, Visva-Bharati University, Santiniketan 731235, India E-mail: $\textit{binou123@bsnl.in}$

[∗]Abhedananda Mahavidyalaya, Sainthia 731234, India

Abstract

We consider equations in the modified KdV (mKdV) hierarchy and make use of the Miura transformation to construct expressions for their Lax pair. We derive a Lagrangian-based approach to study the bi-Hamiltonian structure of the mKdV equations. We observe that the complex mKdV equation has an additional analytic structure. The omplex equation is hara
terized by two alternative Lagrangians not onne
ted by a gauge term. We examine how the alternative Lagrangian description of the system affects the bi-Hamiltonian structure.

Key Words: Real and complex modified KdV equations; Lax pair representation; Bi-Hamiltonian stru
ture

PACS Number: 47.20.Ky, 42.81.Dp, 02.30.Jr

1. Introduction

Nearly forty years ago Lax $[1]$ showed that the Korteweg de Vries (KdV) initial value problem for $u = u(x, t)$ given by

$$
u_t = u_{3x} - 6uu_x \tag{1}
$$

with

$$
u(x,0) = \mathcal{V}(x) \tag{2}
$$

is but one of the infinite family of equations that leave the eigenvalue of the Schrödinger equation with the potential $\mathcal{V}(x)$ invariant in time. The subscripts of u in (1) denote differentiation with respect to the associated independent variables. The family of equations dis
overed by Lax often goes by the name KdV hierar
hy and is generated by making use of the re
ursion operator $[2]$

$$
\Lambda = \partial_x^2 - 4u - 2u_x \partial_x^{-1}, \partial_x = \frac{\partial}{\partial x} \tag{3}
$$

in the differential relation

$$
u_t = \Lambda^n u_x \ , \quad n = 0, 1, 2, 3, \dots \ .
$$
 (4)

The KdV equation (1) is recognized as the solvability condition for the system

$$
L\psi = \lambda\psi \tag{5a}
$$

and

$$
\partial_t \psi = A \psi \ , \partial_t = \frac{\partial}{\partial t} \tag{5b}
$$

with

$$
L = -\partial_x^2 + u \t{,} \t(6a)
$$

the so-called Schrödinger operator. Here A is a third-order linear operator written as

$$
A = 4\partial_x^3 - 3u\partial_x - 3\partial_x u \ . \tag{6b}
$$

The existence of the solution $\psi = \psi(\lambda, x, t)$ for every constant λ is equivalent to

$$
\partial_t L = AL - LA = [A, L] . \tag{7}
$$

The result in (7) is called the Lax equation and the operators L and A are called Lax pair $[1]$. The Lax pair representation holds good for all equations in the KdV hierarchy. In the context of Lax's method it is often said that L defines the original spectral problem while A represents an auxiliary spectral problem. As one goes along the hierar
hy, L remains un
hanged but the differential operator associated with the auxiliary spectral problems changes according to

$$
A_n = (4)^n \partial_x^{2n+1} + \sum_{j=1}^n \{a_j \partial_x^{2j-1} + \partial_x^{2j-1} a_j \}, \quad n = 0, 1, 2, 3, \dots \tag{8}
$$

The operator $A_0 = \partial_x$ and A_1 stands for A in (6b). It is not always easy to obtain results for other A_n 's which generate higher-order KdV equations. The coefficient a_j depends on the solution u and derivatives $u_n (= \frac{\partial^n u}{\partial x^n})$. From (8) it is clear that as j varies, the dimension of a_j changes. Thus a_j should be chosen as a linear combination of power and products of u and u_n 's such that the terms in the curly bracket have the right dimension of ∂_x^{2n+1} . The constructed expression for A_n will then generate the KdV hierarchy when used in the Lax equation $[3]$. On the other hand, one can postulate that for an evolution equation of the form $u_t = K[u]$ the terms in the Fréchet derivative of K[u] contribute additively with unequal weights to form the operator A_n such that L and A_n via (7) reproduces the equations in the hierarchy [4]. Of ourse, there should not be any in
onsisten
y in determining the values of the weight fa
tors.

Zakharov and Faddeev [5] developed the Hamiltonian approach to integrability of nonlinear evolution equations in one spatial and one temporal $(1+1)$ dimension and, in particular, Gardner [6] interpreted the KdV equation as a completely integrable Hamiltonian system with ∂_x as the relevant Hamiltonian operator. A significant development in the Hamiltonian theory is due to Magri $[7]$, who realized that integrable Hamiltonian systems have an additional structure. They are bi-Hamiltonian i.e. they are Hamiltonian with respect to two different compatible Hamiltonian operators ∂_x and $(\partial_x^3 - 4u\partial_x - 2u_x)$ such that

$$
u_t = \partial_x \left(\frac{\delta H_n}{\delta u} \right) = \left(\partial_x^3 - 4u \partial_x - 2u_x \right) \left(\frac{\delta H_{n-1}}{\delta u} \right) , \quad n = 1, 2, 3... \tag{9}
$$

Here $H_n = \int \mathcal{H}_n dx$ with $\mathcal{H}_n,$ the conserved densities for the equations in the KdV hierarchy. These conserved densities generate flows which commute with the KdV flow and as such give rise to an appropriate hierarchy. Traditionally, the expression for \mathcal{H}_n is constructed using a mathematical formulation that does not make expli
it referen
e to the Lagrangians of the equations in the hierarchy. However, a Lagrangian-based approach can be used to identify \mathcal{H}_n as the Hamiltonian density of the nth hierarchical equation [8].

The nonlinear transformation of Miura or the soalled Miura transformation $[9]$

$$
u = v_x + v^2 \ , \ v = v(x, t) \tag{10}
$$

onverts the KdV equation into a modied KdV (mKdV) equation

$$
v_t = v_{3x} - 6v^2 v_x \tag{11}
$$

This equation differs from the KdV equation only because of its cubic nonlinearity. It has many applicative relevance. For example, mKdV equation has been used to describe acoustic waves in anharmonic lattices and Alfvén waves in collisionless plasma. It is of interest to note that the recursion operator Λ for the mKdV equation $[10]$

$$
\Lambda_m = \partial_x^2 - 4v^2 - 4v_x \partial_x^{-1} . v \tag{12}
$$

can be identified from (4) and (10) . The equation of the mKdV hierarchy an be generated by using the relation

$$
v_t = \Lambda_m^n v_x \ , \quad n = 0, 1, 2, 3, \dots \ .
$$
 (13)

It is straightforward to obtain the equations in the mKdV hierarchy from those in the KdV hierar
hy by the use of Miura transformation. However, it is a nontrivial problem to derive the Lax representation and onstru
t the bi-Hamiltonian structure of the equations in mKdV hierarchy starting from corresponding results for the KdV equation [11]. In this work we shall mainly deal with these problems. Admittedly, we shall make use of a Lagrangianbased approach to derive the Magri structure. In this context it is important to note that the omplex mKdV (
mKdV) equation given by

$$
v_t = v_{3x} - 6|v|^2 v_x , \quad v = p + iq
$$
 (14)

follows from third-order nonlinear S
hrödinger equation via an appropriate variable transformation $[12]$. Unfortunately, the equation is nonLagrangian even in the potential function representation. If $|v|^2$ in (14) is replaced by v^2 the resultant mKdV equation has a Lax pair representation and follows from an action principle. In fact, we shall show that the cmKdV equation has two alternative Lagrangian representations with some interesting onsequen
es on its bi-Hamiltonian stru
ture.

In §2 we introduce the equations in the mKdV hierarchy and derive their Lax pair representation. We find that the system of equations follows from the action principle and as such can be obtained from appropriate Lagrangian densities via the so-called Euler-Lagrange equations. The corresponding Hamiltonian densities onstitute the onserved densities of the mKdV equation. We then study the bi-Hamiltonian structure of the mKdV equations. In $\S 3$ we envisage similar studies for a complex mKdV (cmKdV) equation. The mKdV equation has two alternative Lagrangian representations. The alternative Lagrangians yield the same equation of motion but are not related by a gauge term $[13]$. We examine how the presence of alternative Lagrangians affects the bi-Hamiltonian structure. Finally, in $\S 4$ we try to summarize our outlook on the present work.

2. mKdV hierar
hy

The equations of the mKdV hierarchy follow from (13) for $n =$ 0, 1, 2, 3... . We shall onstru
t Lax pair representations of these equations by taking recourse to the use of (10) in $(6a)$ and (8) . For these equations we shall use a Lagrangian-based method to obtain the conserved densities which are in involution and generate the so-called mKdV flow. We shall then try to realize the bi-Hamiltonian structure by an appropriate modification of (9) by the use of Miura transform.

(a) Lax pair representation

From $(6a)$, $(6b)$ and (10) we write

$$
L = -\partial_x^2 + v^2 + v_x \tag{15a}
$$

and

$$
A = 4\partial_x^3 - 3(v^2 + v_x)\partial_x - 3\partial_x.(v^2 + v_x) .
$$
 (15b)

Using (15) in (7) we get

$$
(\partial_x + 2v) (v_t - v_{3x} + 6v^2 v_x) = 0.
$$
 (16)

As with (11), (16) gives the mKdV equation. In view of this we shall denote the Lax pair in (15) by L^m and A^m just to indicate that these refer to the mKdV equation. We shall follow this onvention for all operators and functions related to the mKdV equation. In consistent with the notation of (8) $A^m (= A)$ stands for A_1^m . In close analogy with the case of higher KdV equations the original spectral problem for the mKdV equations characterized by the operator L^m remains unchanged as we go up the hierarchy but the differential operator A_n^m 's change with n . From (10) and the results given in refs 3 and 4 one can calculate the expressions for A_n^m , $n = 2, 3, 4, ...$. In the following we present some of our results.

$$
A_2^m = 16\partial_x^5 + (25v_x^2 + 30v^2v_x + 10vv_{2x} + 15v^4 + 5v_{3x})\partial_x - 20(v^2 + v_x)\partial_x^3 +
$$

\n
$$
\partial_x \cdot (25v_x^2 + 30v^2v_x + 10vv_{2x} + 15v^4 + 5v_{3x}) - 20\partial_x^3 \cdot (v^2 + v_x), \qquad (17a)
$$

\n
$$
A_3^m = 64\partial_x^7 - 140(v^4 + 3v_x^2 + 2v^2v_x + 2vv_{2x} + v_{3x})\partial_x^3 + 112(v^2 + v_x)\partial_x^5 -
$$

\n
$$
140\partial_x^3 \cdot (v^4 + 3v_x^2 + 2v^2v_x + 2vv_{2x} + v_{3x}) + 112\partial_x^5 \cdot (v^2 + v_x) +
$$

\n
$$
(70v^6 + 210v^4v_x + 1050v^2v_x^2 + 210v_x^3 + 140v^3v_{2x} + 840vv_xv_{2x} + 721v_{2x}^2 + 70v^2v_{3x})\partial_x +
$$

\n
$$
(798v_xv_{3x} + 182v_xv^4 + 91v_{5x})\partial_x + \partial_x \cdot (70v^6 + 210v^4v_x + 1050v^2v_x^2 + 210v_x^3)
$$

\n
$$
+ \partial_x \cdot (140v^3v_{2x} + 840vv_xv_{2x} + 721v_{2x}^2 + 70v^2v_{3x} + 798v_xv_{3x} + 182v_xv^4 + 91v_{5x})
$$

\n(17b)

$$
A_4^m = 256\partial_x^9 - 255v_{8x} - 1794v_{7x}\partial_x - 510vv_{7x} - 1152v_x\partial_x^7 - 1152v^2\partial_x^7 - 5628v_{6x}\partial_x^2 + 2100v_xv_{6x} - 3588vv_{6x}\partial_x + 5670v^2v_{6x} - 4032v_{2x}\partial_x^6 - 8064vv_x\partial_x^6 - 10248v_{5x}\partial_x^3 - 7224v_{2x}v_{5x} - 11256vv_{5x}\partial_x^2 - 15594v_xv_{5x}\partial_x + 18312vv_xv_{5x} + 5934v^2v_{5x}\partial_x + 11340v^3v_{5x} - 8736v_{3x}\partial_x^5 - 17472vv_{2x}\partial_x^5 - 15456v_x^2\partial_x^5 + 4032v^2v_x\partial_x^5 + 2016v^4\partial_x^5 - 11760v_{4x}\partial_x^4 - 11760v_{3x}v_{4x} - 20496vv_{4x}\partial_x^3 - 39204v_{2x}v_{4x}\partial_x + 19152vv_{2x}v_{4x} - 11760v_{4x}\partial_x^4 - 11760v_{4x}\partial_x^4 - 20496vv_{4x}\partial_x^3 - 39204v_{2x}v_{4x}\partial_x + 19152vv_{2x}v_{4x} - 11760v_{4x}\partial_x^4 - 1176
$$

$$
43680v_xv_{4x}\partial_x^2 + 12600v^2v_{4x}\partial_x^2 + 68670v_x^2v_{4x} + 41100vv_xv_{4x}\partial_x + 70224v^2v_xv_{4x} + 11868v^3v_{4x}\partial_x - 210v^4v_{4x} - 23520vv_{3x}\partial_x^4 - 60240v_xv_{2x}\partial_x^4 + 10320v^2v_{2x}\partial_x^4 + 20640v^3v_x\partial_x^4 - 25758v_{3x}^2\partial_x + 12180vv_{3x}^2 - 66864v_xv_{3x}\partial_x^3 + 15120v^2v_{3x}\partial_x^3 - 87360v_{2x}v_{3x}\partial_x^2 + 170268v_xv_{2x}v_{3x} + 69720v_xv_{2x}v_{3x}\partial_x + 130200v^2v_{2x}v_{3x} + 75600vv_xv_{3x}\partial_x^2 + 25200v^3v_{3x}\partial_x^2 + 81660v_x^2v_{3x}\partial_x + 64596vv_x^2v_{3x} + 93336v^2v_xv_{3x} + 75600vv_xv_{3x}\partial_x^2 + 25200v^3v_{3x}\partial_x^2 + 81660v_x^2v_{3x}\partial_x + 64596vv_x^2v_{3x} + 93336v^2v_xv_{3x}\partial_x - 15960v^3v_xv_{3x} - 6300v^4v_{3x}\partial_x - 420v^5v_{3x} - 50568v_{2x}^2\partial_x^3 + 73920vv_xv_{2x}\partial_x^3 + 28560v_x^3\partial_x^3 + 68880v^2v_x^2\partial_x^3 - 5040v^4v_x\partial_x^3 - 1680v^6\partial_x^3 + 19026v_{2x}^2 + 57960v_xv_{2x}\partial_x^2 + 114480v_x^2v_{2x}\partial_x + 88452vv_xv_{2x}^2 + 67272v^2v_{2x}^2\partial_x - 15120v^3v
$$

Using $(15a)$ and $(17a)$ in (7) we get

$$
(\partial_x + 2v)\{v_t - v_{5x} + 40vv_xv_{2x} + 10v^2v_{3x} + 10v_x^3 - 30v^4v_x\} = 0.
$$
 (18)

The expression inside the curly bracket represents the equation obtained from (13) with $n = 2$. Results similar to that in (18) hold good for any pair like $[A_n^m, L^m]$. This observation serves as a useful check on our results for A_n^m with arbitrary values of *n*.

(b) Bi-Hamiltonian stru
ture

Here we shall demonstrate that the bi-Hamiltonian structure of (11) and all higher-order equations obtained from (13) with $n = 2, 3, 4...$. We note that a single evolution equation $u_t = P[u]$, $u \in \mathbb{R}$ is never the Euler-Lagrange equation of a variational problem $[10]$. One common trick to put a single evolution equation into a variational form is to replace v by a potential function

$$
v = -w_x, \ w = w(x, t) \ . \tag{19}
$$

The function w is often called the Casimir potential. Our expressions for the Lagrangian densities will be written in terms of w and its appropriate derivatives. Hamiltonian densities obtained by use of Legendre map can. however, be expressed in terms of field variable $v(x, t)$ and its derivatives.

The linear equation obtained from (13) with $n = 0$ reads

$$
v_t = v_x \tag{20}
$$

From (19) and (20)

$$
w_{xt} = w_{xx} = P[w_x] \text{ (say)}.
$$
 (21)

Equivalently,

$$
w_t = w_x \tag{22}
$$

In writing (22) we have used the boundary condition limit $w(x,t) = 0$ as $x \to \pm \infty$. The self adjointness of $P[w_x]$ ensures the existance of a Lagrangian for (21) and (22) . In this case, the Lagrangian density can be constructed using the homotopy formula $[10]$

$$
\mathcal{L}[\xi] = \int_0^1 \xi P[\lambda \xi] d\lambda \tag{23}
$$

From (23) we get

$$
\mathcal{L}_0^m = \frac{1}{2} w_t w_x - \frac{1}{2} w_x^2 \,, \tag{24a}
$$

The subs
ript zero is self explanatory. The Hamiltonian density obtained from $(24a)$ is given by

$$
\mathcal{H}_0^m = \frac{1}{2} w_x^2 = \frac{1}{2} v^2 \ . \tag{25a}
$$

The Lagrangian and Hamiltonian densities for the mKdV $(n = 1)$ and the higher-order equations obtained from (13) for $n = 2$, 3 and 4 are given by

$$
\mathcal{L}_1^m = \frac{1}{2} w_t w_x - \frac{1}{2} w_x w_{3x} + \frac{1}{2} w_x^4 , \qquad (24b)
$$

$$
\mathcal{H}_1^m = \frac{1}{2} v v_{2x} - \frac{1}{2} v^4 , \qquad (25b)
$$

$$
\mathcal{L}_2^m = \frac{1}{2} w_t w_x - \frac{1}{2} w_{3x}^2 - w_x^6 - 5 w_x^2 w_{2x}^2 \,, \tag{24c}
$$

$$
\mathcal{H}_2^m = \frac{1}{2}v_{2x}^2 + v^6 + 5v^2v_x^2 \,,\tag{25c}
$$

$$
\mathcal{L}_3^m = \frac{1}{2} w_t w_x - \frac{1}{2} w_x w_{7x} + \frac{7}{2} w_x^3 w_{5x} + 14 w_x^2 w_{2x} w_{4x} + \frac{21}{2} w_x^2 w_{3x}^2 + \frac{35}{2} w_x w_{2x}^2 w_{3x} - \frac{35}{3} w_x^5 w_{3x} - \frac{70}{3} w_x^4 w_{2x}^2 + \frac{5}{2} w_x^8 ,
$$
\n(24d)

$$
\mathcal{H}_3^m = \frac{1}{2} v v_{6x} - \frac{7}{2} v^3 v_{4x} - 14 v^2 v_x v_{3x} - \frac{21}{2} v^2 v_{2x}^2 - \frac{35}{2} v v_x^2 v_{2x} + \frac{35}{3} v^5 v_{2x} - \frac{70}{3} v^4 v_x^2 - \frac{5}{2} v^8
$$
\n
$$
(25d)
$$

and

$$
\mathcal{L}_4^m = \frac{1}{2} w_t w_x - \frac{1}{2} w_x w_{9x} + \frac{9}{2} w_x^3 w_{7x} + 27 w_x^2 w_{2x} w_{6x} + 57 w_x^2 w_{3x} w_{5x} + \frac{105}{2} w_x w_{2x}^2 w_{5x}
$$

$$
-21 w_x^5 w_{5x} + \frac{69}{2} w_x^2 w_{4x}^2 + 189 w_x w_{2x} w_{3x} w_{4x} - 168 w_x^4 w_{2x} w_{4x} + \frac{91}{2} w_x w_{3x}^3 - 126 w_x^4 w_{3x}^2
$$

$$
-518w_x^3w_{2x}^2w_{3x} + \frac{105}{2}w_x^7w_{3x} - 133w_x^2w_{2x}^4 + \frac{315}{2}w_x^6w_{2x}^2 - 7w_x^{10}
$$
 (24*e*)

$$
\mathcal{H}_4^m = \frac{1}{2} v v_{8x} - \frac{9}{2} v^3 v_{6x} - 27 v^2 v_x v_{5x} - 57 v^2 v_{2x} v_{4x} - \frac{105}{2} v v_x^2 v_{4x} + 21 v^5 v_{4x} - \frac{69}{2} v^2 v_{3x}^2 \n- 189 v v_x v_{2x} v_{3x} + 168 v^4 v_x v_{3x} - \frac{91}{2} v v_{2x}^3 + 126 v^4 v_{2x}^2 + 518 v^3 v_x^2 v_{2x} - \frac{105}{2} v^7 v_{2x} + \n133 v^2 v_x^4 - \frac{315}{2} v^6 v_x^2 - 7 v^{10} .
$$
\n(25*e*)

Results of \mathcal{H}^{m}_{n} 's for still higher values of n can be obtained in a similar manner. As a useful he
k on our expressions, one an verify that our results are in exact agreement with those obtained by the application of Miura transformation on the well known onserved densities of the KdV equations.

2

The bi-Hamiltonian structure of equations in the mKdV hierarchy can easily be verified by using our Hamiltonian functionals in

$$
v_t = \partial_x \left(\frac{\delta H_n^m}{\delta v} \right) = \mathcal{E} \left(\frac{\delta H_{n-1}^m}{\delta v} \right) , \quad n = 1, 2, 3... , \qquad (26)
$$

where

$$
\mathcal{E} = \left(\partial_x^3 - 4v^2 \partial_x - 4v_x \partial_x^{-1} v \partial_x\right) . \tag{27}
$$

The first Hamiltonian operator ∂_x in (26) is the same as that in (9) while the second has been obtained from [10]

$$
\mathcal{E} = \Lambda_m \partial_x \ . \tag{28}
$$

The operators ∂_x and $\mathcal E$ are skew symmetric and satisfy the Jacobi-identity. Thus they constitute two compatible Hamiltonian operators such that all equations obtained from (13) are integrable in the Liouville's sense [7].

3. mKdV equation

For the omplex mKdV equation of our interest we write

$$
v(x,t) = p(x,t) + iq(x,t)
$$
\n
$$
(29)
$$

where p and q are real. From (11) and (29) we get the coupled nonlinear equations

$$
p_t = p_{3x} - 6(p^2 - q^2)p_x + 12pqq_x \tag{30a}
$$

and

$$
q_t = q_{3x} - 6(p^2 - q^2)q_x - 12pqp_x.
$$
 (30b)

We shall now study some interesting feature for the Lax pair representation and bi-Hamiltonian stru
ture of (30).

(a) Lax pair representation

The Lax pair for the complex field under consideration is easily obtained from (15) and (29) . Writing L and A in the form

$$
L = L_1 + iL_2 \tag{31a}
$$

and

$$
A = B_1 + iB_2 \tag{31b}
$$

we get

$$
L_1 = -\partial_x^2 + p_x + p^2 - q^2 \t\t(32a)
$$

$$
L_2 = q_x + 2pq \tag{32b}
$$

$$
B_1 = 4\partial_x^3 - 6(p_x + p^2 - q^2)\partial_x - 3(p_{2x} + 2pp_x - 2qq_x)
$$
 (33*a*)

and

$$
B_2 = -6(q_x + 2pq)\partial_x - 3(q_{2x} + 2pq_x + 2qp_x) . \tag{33b}
$$

It is interesting to see that both

$$
\frac{dL_1}{dt} = [B_1, L_1] - [B_2, L_2] \tag{34a}
$$

and

$$
\frac{dL_2}{dt} = [B_1, L_2] + [B_2, L_1] \tag{34b}
$$

yield the coupled equations in (30). Thus the complexly coupled mKdV equations appear to have alternative Lax pair representations.

(b) Bi-Hamiltonian stru
ture

To derive the bi-Hamiltonian structure for the coupled equation in (30) we first look for a possible variational formulation of these equations. It will be convenient to work with velocity potential corresponding to $p(x, t)$ and $q(x, t)$. We thus write

$$
p = -\psi_x
$$
 and $q = -\phi_x$, $\psi = \psi(x, t)$, $\phi = \phi(x, t)$. (35)

The potentials ψ and ϕ can be introduced into the problem by observing that $(30a)$ and $(30b)$ can be obtained from the one forms

$$
\alpha_1 = pdx + (p_{2x} - 2p^3 + 6pq^2)dt , \qquad (36a)
$$

$$
\alpha_2 = qdx + (q_{2x} + 2q^3 - 6p^2q)dt
$$
\n(36b)

and oditions for them to be losed

$$
d\alpha_1 = 0 \text{ and } d\alpha_2 = 0. \tag{37}
$$

From (37) , using poincare lemma [13], we write

$$
\alpha_1 = -d\psi \quad \text{and} \quad \alpha_2 = -d\phi \tag{38}
$$

where ψ and ϕ are zero form or scalars. Combining (36) and (38) we obtain

$$
-\psi_x = p \ , \quad -\psi_t = (p_{2x} - 2p^3 + 6pq^2) \tag{39a}
$$

and

$$
-\phi_x = q \ , \quad -\phi_t = (q_{2x} + 2q^3 - 6p^2q) \ . \tag{39b}
$$

From the integrability onditions of (39) we get

$$
\psi_t = \psi_{3x} - 2\psi_x^3 + 6\psi_x \phi_x^2 \tag{40a}
$$

and

$$
\phi_t = \phi_{3x} + 2\phi_x^3 + 6\phi_x\psi_x^2 \,. \tag{40b}
$$

We have found that these equations follow from a variational principle

$$
\delta I = 0 \ , \quad I = \int \mathcal{L} \, dx dt \tag{41}
$$

with two different Lagrangian densities given by

$$
\mathcal{L}^d = \frac{1}{2}(\psi_x \psi_t - \phi_x \phi_t) - \frac{1}{2}(\psi_x \psi_{3x} - \phi_x \phi_{3x}) + \frac{1}{2}\psi_x^4 + \frac{1}{2}\phi_x^4 - 3\psi_x^2 \phi_x^2 \quad (42a)
$$

and

$$
\mathcal{L}^a = \frac{1}{2}(\psi_x \phi_t + \phi_x \psi_t) - \frac{1}{2}(\psi_x \phi_{3x} + \phi_x \psi_{3x}) + 2\psi_x^3 \phi_x - 2\psi_x \phi_x^3. \tag{42b}
$$

The superscripts d and a have been used for the terms direct and alternative. We call \mathcal{L}^d , a direct Lagrangian because \mathcal{L}^d when substituted in the Euler-Lagrange equations for ψ and ϕ gives the equation of motion for ψ and ϕ in $(40a)$ and $(40b)$. On the other hand, the alternative Lagrangian does the opposite. Note that, these Lagrangians are not connected by a gauge term. The results in $(42a)$ and $(42b)$ could also be derived by the use of a symmetry-based approach for solving the inverse variational problem [14].

From (42) we find the Hamiltonian densities

$$
\mathcal{H}^{d} = \frac{1}{2}(pp_{2x} - qq_{2x}) - \frac{1}{2}p^{4} - \frac{1}{2}q^{4} + 3p^{2}q^{2}
$$
 (43a)

and

$$
\mathcal{H}^{a} = \frac{1}{2}(p_{2x}q + pq_{2x}) - 2p^{3}q + 2pq^{3}.
$$
 (43b)

It is an interesting uriosity to note that

$$
\mathcal{H} = \mathcal{H}^d + i\mathcal{H}^a \tag{44}
$$

represents the Hamiltonian density of the cmKdV equation such that via the anoni
al equation

$$
(p + iq)_t = \partial_x \frac{\delta \mathcal{H}}{\delta(p + iq)}\tag{45}
$$

gives the oupled equations in (30). To realize the bi-Hamiltonian stru
ture of the mKdV equation we onsider equations obtained from the omplex form of the linear equation in (13) with $n = 0$. For these equations the direct and alternative Hamiltonian densities

$$
\mathcal{H}^{d} = \frac{1}{2}(p^{2} - q^{2}) \text{ and } \mathcal{H}^{a} = pq .
$$
 (46)

give the total Hamiltonian densities

$$
\mathcal{H} = \frac{1}{2}(p^2 - q^2) + ipq \ . \tag{47}
$$

The result for H when used in

$$
(p+iq)_t = \left(\partial_x^3 - 4(p+iq)^2\partial_x - 4(p+iq)_x\partial_x^{-1} \cdot (p+iq)\partial_x\right) \frac{\delta \mathcal{H}}{\delta(p+iq)}\tag{48}
$$

on
e again gives the oupled equations in (30). From (45) and (48) we see that as with the mKdV equation the mKdV equation is also bi-Hamiltonian.

4. Con
lusion

The nonlinear transformation of Miura or the soalled Miura transformation is an aid to obtain the modified KdV (mKdV) equation from the KdV equation. We find that this transformation also provides an effective way to construct expressions for Lax pairs of all equations in the mKdV hierarchy. As with the KdV equations the bi-hamiltonian structure of the mKdV equations are traditionally studied using involutive set of onserved Hamiltonian densities without expli
it referen
e to their Lagrangians. We derive a Lagrangian-based approach to realize the bi-Hamiltonian structure. A similar approa
h when applied to orresponding omplex equations leads to an interesting curiosity. The complex equations have alternative Lax pairand Lagrangian representations.

A
knowledgements

This work is supported by the University Grants Commission, Government of India, through grant No. F.32-39/2006(SR).

[1] Lax P D 1986 Comm. Pure Appl. Math. $21\,467$

[2] Calogero F and Degasperis A 1982 Spectral Transform and Soliton (North-Holland Publising Company, New York)

[3] Chakraborti S, Pal J, Shamanna J and Talukdar B 2003 Czech. J. Phys. 52 853

[4] Choudhuri A, Talukdar B and Datta S B 2006 Z. Naturforsch. 61a 7

[5] Zakharov V E and Faddeev L D 1971 Funct. Anal. Phys. 5 18

[6] Gardner C S 1971 J. Math. Phys. 12 1548

[7] Magri F 1971 J. Math. Phys. **19** 1584

[8] Ghosh S, Talukdar B and Shamanna J 2003 Czech. J. Phys. 53 425

[9] Miura R M 1968 J. Math. Phys. 9 1202

[10] Olver P J 1993 Application of Lie Groups to Differential Equation. (Springer-Verlag, New York)

[11] Toda K 2002 Proc. Inst. Math. of NAS, Ukraine 43 377

[12] Yang J 2003 Phys. Rev. Lett. 91 143903

[13] Santilli R M 1978 Foundations of Theoretical Mechanics I (Springer-Verlag, New York)

[14] Ghosh S, Talukdar B, Sarkar P and Das U 2007 Acta Mechanica 190 73