

VECTOR-VALUED RIESZ POTENTIALS: CARTAN TYPE ESTIMATES AND RELATED CAPACITIES

V. EIDERMAN, F. NAZAROV, AND A. VOLBERG

ABSTRACT. Our aim is to give sharp upper bounds for the size of the set of points where the Riesz transform of a linear combination of N point masses is large. This size will be measured by the Hausdorff content with various gauge functions. Among other things, we shall characterize all gauge functions for which the estimates do not blow up as N tends to infinity (in this case a routine limiting argument will allow us to extend our bounds to all finite Borel measures). We also show how our techniques can be applied to estimates for certain capacities.

1. INTRODUCTION

The aim of this paper is to give sharp upper bounds for the size of the set of points where the singular s -Riesz transform of a linear combination of N point masses is large. This size will be measured by the Hausdorff content M_h with various gauge functions h . Among other things, we shall characterize all h for which the estimates do not blow up as $N \rightarrow +\infty$ (in this case a routine limiting argument will allow us to extend our bounds to all finite Borel measures).

The main motivation for this work came from [2], [7] where such bounds were obtained for the Cauchy transform in \mathbb{C} , which is a special case of s -Riesz transform in \mathbb{R}^d corresponding to $d = 2$, $s = 1$.

The main challenge in the generalizing the result to higher dimensions and different s is the absence of the Menger's curvature tool. The methods of the current paper allow us to cover the range $0 < s < \infty$.

2. DEFINITIONS AND NOTATION

Let ν be a finite Borel measure in \mathbb{R}^d (not necessarily positive). For $s > 0$, put

$$K^s(x) = \frac{x}{|x|^{s+1}}, \quad x \in \mathbb{R}^d \setminus \{0\}.$$

For $\varepsilon > 0$, define the ε -truncated s -Riesz transform of ν by

$$R_{\nu,\varepsilon}^s(x) = \int_{|y-x|>\varepsilon} K^s(y-x) d\nu(y).$$

If the limit $\lim_{\varepsilon \rightarrow 0+} R_{\nu,\varepsilon}^s(x)$ exists, we shall call it the s -Riesz transform of ν at x and denote it by $R_\nu^s(x)$. For $0 < s < d$, this is true almost everywhere in \mathbb{R}^d with respect to Lebesgue

1991 *Mathematics Subject Classification*. Primary: 42B20. Secondary: 28A78, 30C85, 31B15, 31C45, 28A75, 28A80.

Research of the authors was supported in part by NSF grants DMS-0501067 (Nazarov, Volberg). The first-named author is grateful to the Department of Mathematics, Michigan State University, for its hospitality.

measure but this observation will be rather useless to us because sets of Lebesgue measure 0 can easily have arbitrarily large Hausdorff content M_h when $\lim_{t \rightarrow 0^+} \frac{h(t)}{t^d} = +\infty$ (we remind the definition of M_h below). Of course, in the case when ν is a finite linear combination of point masses, $R_\nu^s(x)$ makes sense everywhere except finitely many points. Nevertheless, since we are aiming at extending our results to all finite Borel measures whenever possible, we shall introduce one more quantity that always makes sense, namely the so called maximal s -Riesz transform

$$R_{\nu,*}^s(x) = \sup_{\varepsilon > 0} |R_{\nu,\varepsilon}^s(x)|$$

(note that $R_{\nu,\varepsilon}^s(x)$ and $R_\nu^s(x)$ are vectors and $R_{\nu,*}^s(x)$ is a number).

Take $P > 0$. The sets whose size we will be interested in are

$$\mathcal{Z}(\nu, P) = \{x \in \mathbb{R}^d : R_\nu^s(x) \text{ exists and } |R_\nu^s(x)| > P\}$$

and

$$\mathcal{Z}^*(\nu, P) = \{x \in \mathbb{R}^d : R_{\nu,*}^s(x) > P\}.$$

Clearly, $\mathcal{Z}(\nu, P) \subset \mathcal{Z}^*(\nu, P)$.

By a *gauge* (or *measuring*) function, we shall understand any continuous strictly increasing function $h : [0, +\infty) \rightarrow [0, +\infty)$ such that $h(0) = 0$, $\lim_{r \rightarrow +\infty} h(r) = +\infty$ and $\frac{h(r)}{r^d}$ is non-increasing. The last condition, which may seem a regularity condition at the first glance, is actually not a restriction at all. As we shall see, for any increasing function h vanishing at the origin, there exists another function \tilde{h} satisfying this restriction and such that the Hausdorff contents M_h and $M_{\tilde{h}}$ coincide up to a constant factor depending only on the dimension d .

The *Hausdorff content* $M_h(G)$ of a set $G \subset \mathbb{R}^d$ is defined by

$$M_h(G) = \inf \sum_j h(r_j),$$

where the infimum is taken over all (at most countable) coverings of G by balls of radii r_j .

By c, C we denote various positive constants, and we set $B(x, r) := \{y \in \mathbb{R}^d : |y - x| < r\}$. To show that the assumption that $\frac{h(r)}{r^d}$ is non-increasing is not restrictive, we consider a continuous increasing function h satisfying $h(0) = 0$, without this property in general. Set

$$\tilde{h}(r) = r^d \inf_{0 < t \leq r} \frac{h(t)}{t^d}.$$

Clearly, $\tilde{h}(r) \leq h(r)$, and $\frac{\tilde{h}(r)}{r^d}$ is non-increasing. We are going to prove that

$$M_{\tilde{h}}(G) \leq M_h(G) \leq C_d M_{\tilde{h}}(G)$$

for any set G , with C_d depending only on d . The first inequality is obvious. Choose $\varepsilon > 0$. Let $\{B_j\}$, $B_j = B(x_j, r_j)$, be a covering of G by balls. For every ball B_j , we take t_j such that $0 < t_j \leq r_j$ and

$$\frac{\varepsilon}{2^j} + \tilde{h}(r_j) > r_j^d \frac{h(t_j)}{t_j^d}.$$

There is a constant C_d depending only on d such that B_j can be covered by at most $C_d \left(\frac{r_j}{t_j}\right)^d$ balls $B_{j,k}$ of radii $r_{j,k} = t_j$. For this new covering of G , we have

$$\sum_{j,k} h(r_{j,k}) = \sum_j C_d \left(\frac{r_j}{t_j}\right)^d h(t_j) < C_d \sum_j \tilde{h}(r_j) + C_d \varepsilon.$$

Thus, $M_h(G) \leq C_d M_{\tilde{h}}(G)$, and we are done.

For nonnegative Borel measure μ , we introduce the ε -truncated s -Riesz operator defined by

$$\mathfrak{R}_{\mu,\varepsilon}^s f(x) = \int_{|y-x|>\varepsilon} K^s(y-x) f(y) d\mu(y), \quad f \in L^2(\mu), \quad \varepsilon > 0.$$

For every $\varepsilon > 0$, the operator $\mathfrak{R}_{\mu,\varepsilon}^s$ is bounded on $L^2(\mu)$. We set

$$|\mathfrak{R}_\mu^s| := \sup_{\varepsilon>0} \|\mathfrak{R}_{\mu,\varepsilon}^s\|_{L^2(\mu) \rightarrow L^2(\mu)}.$$

3. FUNCTION $\mathfrak{M}_h(\varkappa, N)$

Let h be a measuring function, $\varkappa > 0$, $N \geq 2$, and let h^{-1} be inverse to h . Define $\mathfrak{M}_h(\varkappa, N)$ as the unique solution $M > 0$ of the equation

$$\varkappa^2 \int_{1/N}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t} = 1. \quad (3.1)$$

Note that the left hand side is a decreasing function of M that tends to $+\infty$ at 0 and to 0 at $+\infty$, so the definition makes sense. Moreover, it is clear that \mathfrak{M}_h is increasing in both \varkappa and N .

Lemma 3.1. (*Doubling property of \mathfrak{M}_h*). For each measuring function h and $0 < s < d$,

$$\mathfrak{M}_h(2\varkappa, 2N) \leq C(s, d) \mathfrak{M}_h(\varkappa, N).$$

Proof. It is enough to show that for sufficiently large C , one has

$$4 \int_{1/(2N)}^1 \left[\frac{t}{h^{-1}(CMt)^s} \right]^2 \frac{dt}{t} \leq \int_{1/N}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t}.$$

Assuming that $C > 2$, and making the change of variable $t \mapsto Ct$, we see that the left hand side does not exceed

$$\frac{4}{C^2} \int_{1/N}^C \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t}.$$

Now notice that the function

$$H(t) = \frac{t^{s/d}}{h^{-1}(Mt)^s}$$

is non-increasing. Therefore,

$$\int_1^C [t^{1-\frac{s}{d}} H(t)]^2 \frac{dt}{t} \leq H(1)^2 \int_1^C t^{2(1-\frac{s}{d})} \frac{dt}{t} \leq H(1)^2 \frac{1}{2(1-\frac{s}{d})} C^{2(1-\frac{s}{d})}.$$

Also,

$$\int_{1/N}^1 [t^{1-\frac{s}{d}} H(t)]^2 \frac{dt}{t} \geq H(1)^2 \int_{1/2}^1 t^{2(1-\frac{s}{d})} \frac{dt}{t} \geq H(1)^2 \cdot \frac{3}{8}.$$

Hence,

$$\frac{4}{C^2} \int_{1/N}^C \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t} \leq \frac{4}{C^2} \left[1 + \frac{4}{3(1-\frac{s}{d})} C^{2(1-\frac{s}{d})} \right] \int_{1/N}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t},$$

and it remains to note that the factor in front of the integral on the right hand side tends to 0 as $C \rightarrow \infty$. \square

Lemma 3.2. *Let $\varkappa > 0$, $N \geq 2$, $c \in (0, 1)$, and $0 < s < d$. If*

$$M \leq C\varkappa \left[\int_{h^{-1}(\frac{c}{N}M)}^{h^{-1}(M)} \left(\frac{h(y)}{y^s} \right)^2 \frac{dy}{y} \right]^{1/2},$$

then $M \leq C'(c, C, s, d) \mathfrak{M}_h(\varkappa, N)$.

Proof. Let n be the least positive integer satisfying $2^{-n} < c/N$. Note that

$$\begin{aligned} \int_{h^{-1}(\frac{c}{N}M)}^{h^{-1}(M)} \left[\frac{h(y)}{y^s} \right]^2 \frac{dy}{y} &\leq \int_{h^{-1}(2^{-n}M)}^{h^{-1}(M)} \left[\frac{h(y)}{y^s} \right]^2 \frac{dy}{y} \\ &= \sum_{j=1}^n \int_{h^{-1}(2^{-j}M)}^{h^{-1}(2^{-(j-1)}M)} \left[\frac{h(y)}{y^s} \right]^2 \frac{dy}{y} \leq C(s) \sum_{j=1}^n \left[\frac{2^{-j}M}{h^{-1}(2^{-j}M)^s} \right]^2 \\ &\leq C''(s) M^2 \sum_{j=1}^n \int_{2^{-(j+1)}}^{2^{-j}} \left[\frac{t}{h^{-1}(tM)^s} \right]^2 \frac{dt}{t} \leq C''(s) M^2 \int_{c/(4N)}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t}. \end{aligned}$$

Thus the inequality in the condition of the lemma implies that

$$\sqrt{C''(s)} C\varkappa \left[\int_{c/(4N)}^1 \left(\frac{t}{h^{-1}(Mt)^s} \right)^2 \frac{dt}{t} \right]^{1/2} \geq 1,$$

i.e.,

$$M \leq \mathfrak{M}_h(\sqrt{C''(s)} C\varkappa, \frac{4}{c}N) \leq C'(c, C, s, d) \mathfrak{M}_h(\varkappa, N).$$

□

Remark 3.3. *Let $c, C > 0$, $\varkappa > 0$, $N > 2C$, and $0 < s < d$. If*

$$M \geq c\varkappa \left[\int_{h^{-1}(\frac{c}{N}M)}^{h^{-1}(M)} \left(\frac{h(y)}{y^s} \right)^2 \frac{dy}{y} \right]^{1/2},$$

then $M \geq c'(c, C, s, d) \mathfrak{M}_h(\varkappa, N)$.

Proof. For $y = h^{-1}(Mt)$,

$$\begin{aligned} \int_{h^{-1}(\frac{c}{N}M)}^{h^{-1}(M)} \left[\frac{h(y)}{y^s} \right]^2 \frac{dy}{y} &= M^2 \int_{C/N}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dh^{-1}(Mt)}{h^{-1}(Mt)} \\ &\geq \frac{M^2}{d} \int_{C/N}^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t} \end{aligned} \tag{3.2}$$

because, since $t \mapsto [\log h^{-1}(Mt) - \frac{1}{d} \log(Mt)]$ is non-decreasing, we have

$$\frac{dh^{-1}(Mt)}{h^{-1}(Mt)} \geq \frac{1}{d} \frac{dt}{t}.$$

Thus, the condition in the remark implies

$$\frac{1}{\sqrt{d}} c\varkappa \left[\int_{C/N}^1 \left(\frac{t}{h^{-1}(Mt)^s} \right)^2 \frac{dt}{t} \right]^{1/2} \leq 1,$$

i.e.,

$$M \geq \mathfrak{M}_h\left(\frac{c}{\sqrt{d}}\varkappa, \frac{N}{C}\right) \geq c\mathfrak{M}_h(\varkappa, N).$$

□

Hence, the condition in Lemma 3.2 is essentially equivalent to $M \leq C\mathfrak{M}_h(\varkappa, N)$.

4. MAIN RESULTS

We formulate our results for $0 < s < d$ and for $s \geq d$ separately. The main and the most difficult case is $0 < s < d$.

Theorem 4.1. *Let $N \geq 2$, $P > 0$, $s \in (0, d)$. Let h be any measuring function.*

I) There exists $C = C(s, d) > 0$ such that for every measure ν that is a linear combination of N Dirac point masses, the inequality

$$M_h(\mathcal{Z}^*(\nu, P)) \leq C\mathfrak{M}_h\left(\frac{\|\nu\|}{P}, N\right) \quad (4.1)$$

holds.

II) There exists $c = c(s, d) > 0$ such that, for every $\eta > 0$, one can find a measure ν that is a linear combination of N Dirac point masses and such that $\|\nu\| = \eta$ and

$$M_h(\mathcal{Z}(\nu, P)) \geq c\mathfrak{M}_h\left(\frac{\|\nu\|}{P}, N\right). \quad (4.2)$$

Theorem 4.1 is a generalization of the corresponding results in [2], [7], [8]. In some important cases one can derive explicit estimates for $M_h(\mathcal{Z}^*(\nu, P))$ from (4.1).

Example 4.2. *For $h(t) = t^\beta$, $s < d$ and $\beta \leq d$, easy calculations yield*

$$M_h(\mathcal{Z}^*(\nu, P)) \leq \begin{cases} C\left(\frac{\|\nu\|}{P}\right)^{\beta/s} \left(\frac{N^{2(s-\beta)/\beta} - 1}{s - \beta}\right)^{\beta/(2s)}, & \beta \neq s, \\ C\frac{\|\nu\|}{P}(\ln N)^{1/2}, & \beta = s, N \geq 2, \end{cases} \quad (4.3)$$

where C depends only on d and s .

For $\beta > d$ the h -content of every set in \mathbb{R}^d is zero.

It is interesting to compare inequalities (4.3) with estimates for the h -content of the set

$$\mathcal{X}(|\nu|, P) := \left\{ x \in \mathbb{R}^d : \int |K^s(y-x)| d|\nu|(y) = \int \frac{1}{|y-x|^s} d|\nu|(y) > P \right\}.$$

Obviously, $\mathcal{Z}^*(\nu, P) \subset \mathcal{X}(|\nu|, P)$. Corollary 1.2 in [8] yields the following estimate: for $h(t) = t^\beta$ and $\beta \leq d$,

$$M_h(\mathcal{X}(|\nu|, P)) \leq \begin{cases} C\left(\frac{\|\nu\|}{P}\right)^{\beta/s} \left(\frac{N^{(s-\beta)/\beta} - 1}{s - \beta}\right)^{\beta/s}, & \beta \neq s, \\ C\frac{\|\nu\|}{P} \ln N, & \beta = s, N \geq 2, \end{cases} \quad (4.4)$$

with another constant C depending only on d and s .

If $s < \beta$, then the estimate

$$M_h(\mathcal{X}(|\nu|, P)) \leq C \left(\frac{\|\nu\|}{P} \cdot \frac{1}{\beta - s} \right)^{\beta/s}$$

(the limiting case of (4.4) as $N \rightarrow +\infty$) holds for every (not necessarily discrete) measure ν . The exponent $1/2$ in (4.3) reflects the mutual annihilation of terms in the passage from the sum of moduli to the modulus of the sum of the corresponding fractions.

Consider now the case when

$$\int_0^1 \left[\frac{h(t)}{t^s} \right]^2 \frac{dt}{t} < \infty. \quad (4.5)$$

Under this assumption we obtain estimates for the h -content of $M_h(\mathcal{Z}^*(\nu, P))$ not only for discrete measures but also for arbitrary finite Borel measures ν . (Note that for the function $h(t) = t^\beta$ with $0 < \beta \leq d$, the condition (4.5) holds iff $s < \beta \leq d$. This is exactly the case when the right-hand sides of (4.3) and (4.4) do not blow up as $N \rightarrow +\infty$.)

The condition (4.5) implies that for every $M > 0$,

$$\int_0^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t} < \infty$$

(see (3.2)). Since this integral is a decreasing function of M that tends to $+\infty$ at 0 and to 0 at $+\infty$, the equation

$$\varkappa^2 \int_0^1 \left[\frac{t}{h^{-1}(Mt)^s} \right]^2 \frac{dt}{t} = 1$$

has the unique solution $M > 0$, which we denote by $\mathfrak{M}_h(\varkappa, \infty)$.

Theorem 4.3. *Let ν be a Borel measure (generally, complex-valued) with finite total variation, and let h be a measuring function satisfying (4.5). Then for any $P > 0$,*

$$M_h(\mathcal{Z}^*(\nu, P)) \leq C \mathfrak{M}_h \left(\frac{\|\nu\|}{P}, \infty \right), \quad (4.6)$$

where C depends only on d and s .

Theorem 4.3 can be viewed as a limiting case of Theorem 4.1 as $N \rightarrow \infty$.

Let us consider the case $s \geq d$.

Theorem 4.4. *Let $N \geq 2$, $P > 0$, $s \geq d$, and let h be any measuring function. There exists $C = C(s, d) > 0$ such that for every measure ν that is a linear combination of N Dirac point masses, the inequality*

$$M_h(\mathcal{Z}^*(\nu, P)) \leq CNh \left(\left(\frac{\|\nu\|}{PN} \right)^{1/s} \right) \quad (4.7)$$

holds.

Thus, for $s > d$ cancelation plays no role and the sharp estimate for $M_h(\mathcal{Z}^*(\nu, P))$ is the same (up to a constant factor depending on s and d) as the estimate for the h -Hausdorff content of the set $\mathcal{X}(|\nu|, P)$ obtained in [8], which in this case reduces to (4.7). Note that the right hand side of this estimate always (when $s > d$) blows up as $N \rightarrow \infty$ (since

$h(t) \geq ct^d$, $t \rightarrow 0+$). So, no meaningful generalization to the case of arbitrary Borel measures ν is possible here.

The sharpness of the estimate (4.7) for $s \geq d$ is obvious: just consider N equal point masses located far away from each other.

Proofs of Theorems 4.1, 4.3 are based on the weak type 1 – 1 estimate for the maximal Calderón-Zygmund operator obtained by Nazarov, Treil and Volberg in [21], p. 483. We quote this general result for the case of the maximal Riesz transform $R_{\nu,*}^s(x)$.

Let Σ_s be the class of nonnegative Borel measures μ in \mathbb{R}^d such that

$$\mu(B(x, r)) \leq r^s \quad \text{for all } x \in \mathbb{R}^d \text{ and } r > 0. \quad (4.8)$$

Theorem 4.5. [21] *Suppose that $\mu \in \Sigma_s$ and $|\mathfrak{R}_\mu^s| < \infty$. Then for every complex-valued Radon measure ν one has*

$$\mu\{x \in \mathbb{R}^d : R_{\nu,*}^s(x) > t\} < \frac{C\|\nu\|}{t} \quad (4.9)$$

with C depending only on s and $|\mathfrak{R}_\mu^s|$.

To apply Theorem 4.5 we should be able to construct the auxiliary measure μ and to estimate $|\mathfrak{R}_\mu^s|$. This estimate has various applications (see for example Section 11 below) and as we believe is of independent interest.

Theorem 4.6. *For every nonnegative Borel measure μ and $0 < s < d$ we have*

$$|\mathfrak{R}_\mu^s|^2 \leq C \sup_{x \in \text{supp } \mu} \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r} \quad (4.10)$$

with C depending only on s and d .

As a byproduct of our calculations we obtain the following estimate for Calderón-Zygmund capacity $\gamma_{s,+}(E)$ (the corresponding definitions are given in Section 11).

Theorem 4.7. *For any compact set $E \subset \mathbb{R}^d$,*

$$\gamma_{s,+}(E) \geq c \sup \|\mu\|^{3/2} \left[\int_{\mathbb{R}^d} W^\mu(x) d\mu(x) \right]^{-1/2}, \quad W^\mu(x) := \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r}, \quad (4.11)$$

where the supremum is taken over all positive Radon measures supported by E , and c depends only on d, s .

We give two applications of Theorem 4.7. We use it to derive relations between Hausdorff content $M_h(E)$ and capacity $\gamma_{s,+}(E)$. Interestingly enough, there exists another completely different capacity $C_{\frac{2}{3}(d-s), \frac{3}{2}}(E)$ (see [1], [17], [11] and the multiple references therein), that can be characterized via the same potential $W^\mu(x)$. Theorem 4.7 immediately implies that $\gamma_{s,+}(E) \geq c \cdot C_{\frac{2}{3}(d-s), \frac{3}{2}}(E)$ (see Section 11).

We prove Theorem 4.6 in Section 5. The construction of the appropriate measure μ is given in Section 6. The first part of Theorem 4.1 is proved in Section 7 and the second part in Section 8. Section 9 contains the proof of Theorem 4.3. In Section 10 we consider the case $s \geq d$ and prove Theorem 4.4. In Section 11 we investigate metric properties of various capacities generated by vector-valued Riesz potentials. In particular, we obtain the Frostman type theorem on comparison of these capacities and Hausdorff content.

5. PROOF OF THEOREM 4.6

The main trick in [18] (which led to the use of Menger's curvature in non-homogeneous harmonic analysis) is to symmetrize an expression involving Cauchy kernel by using averaging over all permutations of coordinates. Amazingly this averaging a) is non-negative, b) is "considerably smaller" than the absolute value of the original expression, and c) is equal to a certain curvature. This observation is no longer true when one averages a similar expression involving vector Riesz kernels in \mathbb{R}^d , $d > 2$, see the paper of Hany Farag [9]. This is why we said (repeating the expression of Guy David) that the tool of curvature is "cruelly missing" for $d > 2$.

However, the following simple observation still holds for all dimensions. If we symmetrize the pertinent expression involving Riesz kernels we generally miss a) and c) above, but we still have b): the symmetrized expression has "considerably smaller" absolute value than the original one. This should be understood maybe not pointwise, but in average.

This observation saves our day, proves Theorem 4.6, and in general allows us to obtain very sharp estimates of various Calderón-Zygmund capacities $\gamma_{s,+}$ from below.

Lemma 5.1. *Let x, y, z be three distinct points in \mathbb{R}^d , $d \geq 1$, and let $|z-x| \leq |z-y| \leq |y-x|$. Then for $s > 0$,*

$$\begin{aligned} q_s(x, y, z) &:= \frac{x-z}{|x-z|^{s+1}} \cdot \frac{y-z}{|y-z|^{s+1}} + \frac{y-x}{|y-x|^{s+1}} \cdot \frac{z-x}{|z-x|^{s+1}} \\ &\leq \frac{2^{s+1}}{|y-x|^{s+1}} \cdot \frac{1}{|z-x|^{s-1}}. \end{aligned} \quad (5.1)$$

Proof. Let $a = |y-x|$, $b = |z-y|$, $c = |z-x|$, and let α, β, γ be the angles opposite to sides a, b, c respectively. Then

$$q_s(x, y, z) = (abc)^{-s} (a^s \cos \alpha + b^s \cos \beta).$$

Since

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}, \quad \cos \beta = \frac{a^2 + c^2 - b^2}{2ac},$$

we have with $u = b/a$, $v = c/a$

$$\begin{aligned} q_s(x, y, z) &= 2^{-1} (abc)^{-s-1} [a^{s+1}(b^2 + c^2 - a^2) + b^{s+1}(a^2 + c^2 - b^2)] \\ &= 2^{-1} (uv)^{-s-1} a^{-2s} [u^2 + v^2 - 1 + u^{s+1}(1 + v^2 - u^2)] \\ &= 2^{-1} (uv)^{-s-1} a^{-2s} [u^{s+1}(1 - u^2) + v^2(1 + u^{s+1}) - (1 - u^2)] \\ &= 2^{-1} (uv)^{-s-1} a^{-2s} [v^2(1 + u^{s+1}) - (1 - u^2)(1 - u^{s+1})] \\ &\leq u^{-s-1} v^{-s+1} a^{-2s} \leq 2^{s+1} a^{-s-1} c^{-s+1}, \end{aligned}$$

because $1/2 \leq u \leq 1$. □

Proof of Theorem 4.6. Without loss of generality we assume that

$$\mathbf{S} := \sup_{x \in \text{supp } \mu} \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r} < \infty. \quad (5.2)$$

Otherwise (4.10) becomes trivial. We consider the measure

$$\eta := (2s\mathbf{S})^{-1/2} \mu.$$

Since for every $x \in \text{supp } \mu$ and $r > 0$,

$$\mathbf{S} \geq \int_0^\infty \left[\frac{\mu(B(x, t))}{t^s} \right]^2 \frac{dt}{t} \geq [\mu(B(x, r))]^2 \int_r^\infty \frac{dt}{t^{2s+1}} = \frac{[\mu(B(x, t))]^2}{2sr^{2s}},$$

we see that $\eta \in \Sigma_s$.

We are going to give *two proofs* of the theorem. They have a lot in common but they are different. The first one will be based on the non-homogeneous $T1$ theorem.

The first approach. Let Q be any cube in \mathbb{R}^d . If for each $\varepsilon > 0$ we prove the inequality

$$\|\mathfrak{R}_{\eta, \varepsilon}^s \chi_Q\|_{L^2(\eta|_Q)}^2 \leq C\eta(Q), \quad C = C(d, s), \quad (5.3)$$

then the theorem follows. In fact, for $\eta \in \Sigma_s$ satisfying this condition, the norm of Calderón-Zygmund operator on a space of non-homogeneous type can be estimated by a constant depending only on C in (5.3) and the constants in Calderón-Zygmund kernel, see [20], [22], [28]. In the spaces of homogeneous type, it is the famous $T1$ theorem of David-Journé (see [5] for the Euclidean setting and [4] for homogeneous setting). Notice that a measure η does not have any doubling property in general. So we cannot use the homogeneous $T1$ theorem, and we can use neither [5] nor [4]. But the non-homogeneous $T1$ theorem [20], [22] works fine. Thus,

$$\|\mathfrak{R}_{\eta, \varepsilon}^s\|_{L^2(\eta) \rightarrow L^2(\eta)}^2 \leq C, \quad C = C(d, s),$$

and we have

$$|\mathfrak{R}_\mu^s|^2 = 2s\mathbf{S} |\mathfrak{R}_\eta^s|^2 \leq C\mathbf{S}, \quad C = C(d, s),$$

that is (4.10). So, it is enough to prove (5.3) or equivalently,

$$\|\mathfrak{R}_{\mu, \varepsilon}^s \chi_Q\|_{L^2(\mu|_Q)}^2 \leq C\mathbf{S}\mu(Q), \quad C = C(d, s). \quad (5.4)$$

We fix $\varepsilon > 0$ and set

$$\mathcal{U} = \{(x, y, z) \in Q^3 : |y - x| > \varepsilon, |z - x| > \varepsilon\},$$

$$\mathcal{U}_1 = \{(x, y, z) \in Q^3 : |y - x| \geq |z - x| > \varepsilon\},$$

$$\mathcal{U}_2 = \{(x, y, z) \in Q^3 : \varepsilon < |y - x| \leq |z - x|\},$$

$$\mathcal{U}_{1,1} = \{(x, y, z) \in Q^3 : |y - x| \geq |z - x| > \varepsilon, |y - z| \geq |z - x|\},$$

$$\mathcal{U}_{1,2} = \{(x, y, z) \in Q^3 : |y - x| \geq |z - x| > \varepsilon, |y - z| < |z - x|\}.$$

Then

$$\begin{aligned} \int_Q |\mathfrak{R}_{\mu, \varepsilon}^s \chi_Q(x)|^2 d\mu(x) &= \iiint_{\mathcal{U}} \frac{y-x}{|y-x|^{s+1}} \cdot \frac{z-x}{|z-x|^{s+1}} d\mu(z) d\mu(y) d\mu(x) \\ &\leq \iiint_{\mathcal{U}_1} + \iiint_{\mathcal{U}_2} =: A + B. \end{aligned}$$

It is enough to estimate A . We have

$$\begin{aligned} |A| &\leq \left| \iiint_{\mathcal{U}_{1,1}} \frac{y-x}{|y-x|^{s+1}} \cdot \frac{z-x}{|z-x|^{s+1}} d\mu(z) d\mu(y) d\mu(x) \right| \\ &\quad + \left| \iiint_{\mathcal{U}_{1,2}} \frac{y-x}{|y-x|^{s+1}} \cdot \frac{z-x}{|z-x|^{s+1}} d\mu(z) d\mu(y) d\mu(x) \right| =: A_1 + A_2. \end{aligned}$$

We put the absolute value in A_2 inside the integral. Since $|z - x| > \frac{1}{2}|y - x|$ in A_2 , we get

$$\begin{aligned} A_2 &\leq 2^s \int_Q \int_{|y-x|>0} \frac{1}{|y-x|^{2s}} \mu(B(x, |y-x|)) d\mu(y) d\mu(x) \\ &= 2^s \int_Q \int_0^\infty \frac{1}{r^{2s}} \mu(B(x, r)) d\mu(B(x, r)) d\mu(x) \\ &= 2^s \int_Q \int_0^\infty \frac{1}{r^{2s}} d\left[\frac{\mu(B(x, r))^2}{2}\right] d\mu(x). \end{aligned} \quad (5.5)$$

From (5.2) one can easily deduce that

$$\lim_{r \rightarrow 0} \frac{\mu(B(x, r))}{r^s} = 0 \quad (5.6)$$

(see, for example, [12], p. 219). Moreover,

$$\frac{\mu(B(x, r))}{r^s} \leq \frac{\mu(\mathbb{R}^d)}{r^s} \rightarrow 0 \text{ as } r \rightarrow \infty.$$

Integrating by parts in the last integral of (5.5) we get

$$A_2 \leq s2^s \int_Q \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s}\right]^2 \frac{dr}{r} d\mu(x) \leq s2^s \mathbf{S}\mu(Q).$$

Let us estimate A_1 . By the symmetry of $\mathcal{U}_{1,1}$ with respect to z, x we have

$$A_1 = \frac{1}{2} \left| \iiint_{\mathcal{U}_{1,1}} \left(\frac{y-x}{|y-x|^{s+1}} \cdot \frac{z-x}{|z-x|^{s+1}} + \frac{y-z}{|y-z|^{s+1}} \cdot \frac{x-z}{|x-z|^{s+1}} \right) d\mu(z) d\mu(y) d\mu(x) \right|.$$

Lemma 5.1 yields

$$A_1 \leq 2^s \iiint_{\mathcal{U}_{1,1}} \frac{1}{|y-x|^{s+1}} \cdot \frac{1}{|z-x|^{s-1}} d\mu(z) d\mu(y) d\mu(x).$$

Clearly,

$$\begin{aligned} A_1 &\leq 2^s \int_Q \int_{|y-x| \geq \varepsilon} \frac{1}{|y-x|^{s+1}} \left[\int_\varepsilon^{|y-x|} \frac{d\mu(B(x, t))}{t^{s-1}} \right] d\mu(y) d\mu(x) \\ &\leq 2^s \int_Q \int_0^\infty \frac{1}{r^{s+1}} \left[\int_0^r \frac{d\mu(B(x, t))}{t^{s-1}} \right] d\mu(B(x, r)) d\mu(x). \end{aligned}$$

Set

$$H_x(r) := \int_0^r \frac{d\mu(B(x, t))}{t^{s-1}}.$$

Then the last expression can be written in the form

$$2^s \int_Q \int_0^\infty \frac{1}{r^2} H_x(r) dH_x(r) d\mu(x) = 2^{s-1} \int_Q \int_0^\infty \frac{dH_x^2(r)}{r^2} d\mu(x). \quad (5.7)$$

Obviously,

$$H_x(r) = \frac{\mu(B(x, r))}{r^{s-1}} + (s-1) \int_0^r \frac{\mu(B(x, t))}{t^s} dt, \quad (5.8)$$

and

$$\lim_{r \rightarrow \infty} \frac{H_x(r)}{r} = 0, \quad \lim_{r \rightarrow 0} \frac{H_x(r)}{r} = 0$$

(the last equality follows from (5.6)). Thus,

$$\begin{aligned} \int_0^\infty \frac{dH_x^2(r)}{r^2} &= 2 \int_0^\infty \frac{H_x^2(r)}{r^3} dr \stackrel{(5.8)}{\leq} 4 \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r} \\ &\quad + 4(s-1)^2 \int_0^\infty \frac{1}{r^3} \left[\int_0^r \frac{\mu(B(x, t))}{t^s} dt \right]^2 dr. \end{aligned} \quad (5.9)$$

The first term in the right hand side of (5.9) is what we need. Let us estimate the second term. By the Cauchy–Bunyakovskii–Schwarz inequality,

$$\left[\int_0^r \frac{\mu(B(x, t))}{t^s} dt \right]^2 \leq \int_0^r \left[\frac{\mu(B(x, t))}{t^s} \right]^2 dt \cdot \int_0^r dt.$$

Hence, applying integration by parts, we obtain

$$\begin{aligned} \int_0^\infty \frac{1}{r^3} \left[\int_0^r \frac{\mu(B(x, t))}{t^s} dt \right]^2 dr &\leq \int_0^\infty \left[\int_0^r \left[\frac{\mu(B(x, t))}{t^s} \right]^2 dt \right] \frac{dr}{r^2} \\ &= \left(-\frac{1}{r} \int_0^r \left[\frac{\mu(B(x, t))}{t^s} \right]^2 dt \right) \Big|_0^\infty + \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r}. \end{aligned}$$

According to (5.6), the substitution of limits gives zero. Thus, (see (5.9))

$$\int_0^\infty \frac{dH_x^2(r)}{r^2} < C(s) \int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r}.$$

Now (5.7) yields (5.4), and Theorem 4.6 is proved.

The second approach. Notice that there was nothing specific in using cubes Q in (5.4). Verbatim the same proof gives more, namely, that for every measurable set E ,

$$\|\mathfrak{R}_{\mu, \varepsilon}^s \chi_E\|_{L^2(\mu|_E)}^2 \leq C \mathbf{S} \mu(E), \quad C = C(d, s). \quad (5.10)$$

But operators $\mathfrak{R}_{\mu, \varepsilon}^s$ have one drawback: they are not operators with Calderón–Zygmund kernels. So, instead of $\mathfrak{R}_{\mu, \varepsilon}^s$, we will use similar operators, but with Calderón–Zygmund kernels. Let us introduce them. Let ϕ be a C_0^∞ “bell-like” function, $\phi = 1$ on the unit ball and $\phi(x) = 0$, $|x| \geq 2$. Put $\psi = 1 - \phi$, and let $\psi_\varepsilon(\cdot) := \psi(\frac{\cdot}{\varepsilon})$. Set

$$\begin{aligned} \tilde{\mathfrak{R}}_{\mu, \varepsilon}^s f(x) &= \int \psi_\varepsilon(|x - y|) K^s(y - x) f(y) d\mu(y), \quad f \in L^2(\mu), \quad \varepsilon > 0, \\ \tilde{\mathfrak{R}}_{\mu, *}^s f(x) &= \sup_{\varepsilon > 0} |\tilde{\mathfrak{R}}_{\mu, \varepsilon}^s f(x)|. \end{aligned}$$

We denote by $\tilde{R}_{\nu, \varepsilon}^s$ the corresponding modified s -Riesz transform of a finite Borel measure ν (not necessarily positive):

$$\tilde{R}_{\nu, \varepsilon}^s(x) = \int \psi_\varepsilon(|x - y|) K^s(y - x) d\nu(y).$$

It is easy to see that if $\eta \in \Sigma_s$ then

$$|\mathfrak{R}_{\eta, \varepsilon}^s f(x) - \tilde{\mathfrak{R}}_{\eta, \varepsilon}^s f(x)| \leq \frac{C}{\varepsilon^s} \int_{B(x, 2\varepsilon)} |f| d\eta \leq C(d, s) \tilde{M} f(x), \quad (5.11)$$

where

$$\widetilde{M}f(x) := \sup_{r>0} \frac{1}{\eta(B(x, 3r))} \int_{B(x,r)} |f| d\eta. \quad (5.12)$$

Maximal operator \widetilde{M} is bounded in any $L^2(\eta)$ (see [21], Lemma 2.1). Hence, (5.11) shows that operators $\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s$ and $\mathfrak{R}_{\eta,\varepsilon}^s$ are bounded simultaneously, and that their norms differ at most by C . Thus, (5.10) yields the corresponding estimate for $\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s$:

$$\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_E\|_{L^2(\eta|_E)}^2 \leq C\eta(E), \quad C = C(d, s). \quad (5.13)$$

Suppose that we would have even more, namely that

$$\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_E\|_{L^2(\eta)}^2 \leq C\eta(E), \quad C = C(d, s). \quad (5.14)$$

Using Theorem 5.1 of [21] and (5.14), one can prove that operators

$$\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s : L^1(\eta) \rightarrow L^{1,\infty}(\eta)$$

are uniformly bounded in ε . Then Theorem 10.1 of [21] proves that norms of operators $\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s$ are uniformly bounded in $L^2(\eta)$ by the same number (up to a constant). This is what we need. The second proof of our theorem would be done.

Unfortunately, we do not have (5.14). And (5.13) alone seems (at the first glance) to be too weak to carry through the proofs of [21]. It is a pity because it would be nice to prove Theorem 4.6 without referring the reader to the proof of non-homogeneous $T1$ theorem from [20].

However, there is a way to use (5.13) and to avoid the non-homogeneous $T1$ theorem. To do that, let us first notice that (5.13) can be “strengthened” as follows: for any measurable sets E and F ,

$$\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_E\|_{L^2(\eta|_F)}^2 \leq C(\eta(E) + \eta(F)), \quad C = C(d, s). \quad (5.15)$$

Indeed, using (5.13) for various sets, we have

$$\begin{aligned} \|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_E\|_{L^2(\eta|_F)}^2 &= \|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{E \cup F} - \widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{F \setminus E}\|_{L^2(\eta|_F)}^2 \\ &\leq 2\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{E \cup F}\|_{L^2(\eta|_{E \cup F})}^2 + 2\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{F \setminus E}\|_{L^2(\eta|(F \setminus E) \cup (F \cap E))}^2 \\ &\leq 2C\eta(E \cup F) + 2C\eta(F \setminus E) + 2\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_F - \widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{F \cap E}\|_{L^2(\eta|_{F \cap E})}^2 \\ &\leq 2C(\eta(E \cup F) + \eta(F \setminus E)) + 4\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_F\|_{L^2(\eta|_F)}^2 + 4\|\widetilde{\mathfrak{R}}_{\eta,\varepsilon}^s \chi_{F \cap E}\|_{L^2(\eta|_{F \cap E})}^2 \\ &\leq 2C(\eta(E \cup F) + \eta(F \setminus E)) + 2\eta(F) + 2\eta(F \cap E) \leq C'(\eta(E) + \eta(F)). \end{aligned}$$

Lemma 5.2. *Let η be a finite measure on some set X , and let function f and constants $\tau, K \in (0, +\infty)$ be such that for any measurable set F ,*

$$\int_F |f|^2 d\eta \leq \tau + K\eta(F).$$

Then $f = f_1 + f_2$, where $\|f_2\|_\infty \leq 2K^{\frac{1}{2}}$, and $\|f_1\|_{L^2(\eta)}^2 \leq \frac{4}{3}\tau$.

Proof. Put $f_1 := f \chi_{\{|f| > 2K^{\frac{1}{2}}\}}$, $f_2 := f \chi_{\{|f| \leq 2K^{\frac{1}{2}}\}}$. Then

$$4K\eta(\{|f| > 2K^{\frac{1}{2}}\}) \leq \int_{\{|f| > 2K^{\frac{1}{2}}\}} |f|^2 d\eta \leq \tau + K\eta(\{|f| > 2K^{\frac{1}{2}}\}),$$

and therefore $\eta(\{|f| > 2K^{\frac{1}{2}}\}) \leq \frac{\tau}{3K}$. Hence,

$$\int |f_1|^2 d\eta = \int_{\{|f| > 2K^{\frac{1}{2}}\}} |f|^2 d\eta \leq \tau + K\eta(\{|f| > 2K^{\frac{1}{2}}\}) \leq \frac{4}{3}\tau.$$

The lemma is proved. \square

Next we need the following modification of Guy David's lemma [6] (or Lemma 4.1 from [21]):

Lemma 5.3. *For any operator $\tilde{\mathfrak{R}}$ satisfying (5.15) with Calderón-Zygmund kernel, for any Borel set E , and for any point $x \in \text{supp } \eta$,*

$$\tilde{\mathfrak{R}}_* \chi_E(x) \leq C \tilde{M} \tilde{\mathfrak{R}} \chi_E(x) + A, \quad C = C(d),$$

where A depends only on d and Calderón-Zygmund constants of the kernel.

Proof. We can try to repeat line by line the proof of Lemma 4.1 in [21], but we cannot use that $\tilde{\mathfrak{R}}$ is $L^2(\eta)$ -bounded. This is not given (and actually this is what we wish to prove), but instead we just use (5.15) to estimate the term on page 474 of [21] as follows

$$\left| \int \chi_{B(x,R)} \tilde{\mathfrak{R}} \chi_{E \cap B(x,3R)} d\eta \right| \leq C(\eta(B(x,R)))^{\frac{1}{2}} (\eta(B(x,R)) + \eta(B(x,3R)))^{\frac{1}{2}}.$$

This is enough to finish the proof exactly as in [21]. In fact, the right hand side is at most $C\eta(B(x,R))$ because $3R$ is a doubling radius (being chosen in the proof of Lemma 4.1 in [21]). \square

Notice a simple thing: the Calderón-Zygmund constants of the kernels of $\tilde{\mathfrak{R}}_{\eta,\varepsilon}^s$ do not depend on ε . This allows us to have uniform C_0 in the following lemma.

Lemma 5.4.

$$\|\tilde{\mathfrak{R}}_{\eta,\varepsilon}^s\|_{L^1(\eta) \rightarrow L^{1,\infty}(\eta)} \leq C_0, \quad (5.16)$$

where C_0 depends only on d and s (and does not depend on ε).

Proof. It is sufficient to prove that

$$\|\tilde{R}_{\nu,\varepsilon}^s\|_{L^{1,\infty}(\eta)} \leq C \|\nu\|, \quad (5.17)$$

where ν is a finite linear combination of unit point masses with positive coefficients. To show that, note first of all that (5.17) remains valid (with constant doubled) for ν with arbitrary real coefficients (just write ν as a difference of two measures with positive coefficients). Fix $f \in L^1(\eta)$, $\varepsilon > 0$, and a ball $B \subset \mathbb{R}^d$. Since the kernel of $\tilde{\mathfrak{R}}_{\nu,\varepsilon}^s$ forms an equicontinuous family of functions $k_x(y) := \psi_\varepsilon(|x-y|)K^s(y-x)$, we can approximate $f d\eta$ by a measure ν of the form $\nu = \sum_{j=1}^M \alpha_j \delta_{y_j}$, $M \in \mathbb{N}_+$, $\alpha_j \in \mathbb{R}$, in such a way that

$$(\{|\tilde{\mathfrak{R}}_{\eta,\varepsilon}^s f(x)| > t\} \cap B) \subset (\{|\tilde{R}_{\nu,\varepsilon}^s(x)| > \frac{1}{2}t\} \cap B).$$

With this end in view we cover a sufficiently large ball $B' \supset B$ by the net of small cubes Q_j , and take $\alpha_j = \int_{Q_j} f d\eta$. Using (5.17), we have

$$\eta(\{|\tilde{\mathfrak{R}}_{\eta,\varepsilon}^s f(x)| > t\} \cap B) \leq \eta(\{|\tilde{R}_{\nu,\varepsilon}^s(x)| > \frac{1}{2}t\}) \leq 2C \|\nu\| \leq 2C \|f\|_{L^1(\eta)}.$$

Since B is arbitrarily large, we get (5.16) with $C_0 = 2C$.

The proof of (5.17) is a modification of the proof of Theorem 5.1 of [21]. This latter can be repeated line by line till we come to p. 477 of [21]. There we need to estimate $(\tilde{\mathfrak{R}} := \tilde{\mathfrak{R}}_{\eta,\varepsilon}^s, \tilde{\mathfrak{R}}_* := \tilde{\mathfrak{R}}_{\eta,*}^s)$

$$\tilde{\mathfrak{R}}_* \chi_E \leq C \widetilde{M} \tilde{\mathfrak{R}} \chi_E + A,$$

where constants C, A depend only on s, d . Lemma 4.1 of [21] is not applicable here, but we have the replacement, namely, Lemma 5.3 above.

Another hitch is that we are required to estimate the following expression:

$$I := \left| \int \chi_F \widetilde{M} \tilde{\mathfrak{R}} \chi_E d\eta \right|.$$

Here

$$\eta(F) = \eta(E), \quad (5.18)$$

and one is tempted to use (5.15) to obtain $I \leq C\eta(E)$, which would be enough to finish the proof verbatim as in [21], p. 477. However, we cannot use here (5.15). In fact, (5.15) does not have any \widetilde{M} in it.

This is why here we need to be more subtle, and we need to use Lemma 5.2. Keep in mind (5.15) and apply Lemma 5.2 to $f := \tilde{\mathfrak{R}} \chi_E$, $K = C$, $\tau = C\eta(E)$. Then we get

$$\tilde{\mathfrak{R}} \chi_E = f_1 + f_2, \quad \|f_2\|_\infty \leq C, \quad \|f_1\|_2^2 \leq C\eta(E).$$

Hence,

$$\|\widetilde{M} f_2\|_\infty \leq C, \quad \|\widetilde{M} f_1\|_{L^2(\eta)}^2 \leq C' C \eta(E),$$

where $C' := \|\widetilde{M}\|_{L^2(\eta) \rightarrow L^2(\eta)}$. This constant is absolute. As $I \leq \int_F |\widetilde{M} f_1| d\eta + \int_F |\widetilde{M} f_2| d\eta$, it becomes obvious that

$$I \leq C'' \eta(E)^{\frac{1}{2}} \eta(F)^{\frac{1}{2}} + C\eta(F).$$

We already mentioned (5.17). So,

$$I \leq C\eta(E),$$

and the proof of Lemma 5.4 is finished exactly as that of Lemma 5.1 of [21]. \square

We started from (5.10) and we derived from it the uniform bound $C_0 \mathbf{S}^{\frac{1}{2}}$ for the norms $\|\tilde{\mathfrak{R}}_{\mu,\varepsilon}^s\|_{L^1(\mu) \rightarrow L^{1,\infty}(\mu)}$. Now we just use Theorem 10.1 from [21], which claims that for operator with Calderón-Zygmund kernel the boundedness of the latter norms implies its boundedness in $L^2(\mu)$ by $C \mathbf{S}^{\frac{1}{2}}$.

It is the time to recall that we have (5.11). Therefore, we obtained the uniform estimate for $\|\tilde{\mathfrak{R}}_{\mu,\varepsilon}^s\|_{L^2(\mu) \rightarrow L^2(\mu)}$. \square

From Theorem 4.6, we derive a useful corollary for Cantor sets. Let ℓ_0, \dots, ℓ_n and λ be such that

$$0 < \ell_{k+1} < \lambda \ell_k, \quad k = 0, \dots, n-1, \quad 0 < \lambda < 1/2.$$

For N of the form $N = 2^{nd}$ we consider N Cantor cubes Q_j^n , $j = 1, \dots, N$, built by the usual procedure from the cube Q_1^0 with edge length ℓ_0 by the corner construction, namely having 2^d corner cubes Q_j^1 with edge length ℓ_1 , 2^{2d} cubes Q_j^2 with edge length ℓ_2 , et cetera. Let $E_n = \bigcup_j Q_j^n$, and let m be the measure uniform on each n -cube and of mass 2^{-nd} on each Q_j^n . Set

$$\theta_{s,k} = \theta_k = \frac{2^{-kd}}{\ell_k^s}.$$

Corollary 5.5. *For the measure m defined above,*

$$|\mathfrak{R}_m^s|^2 \leq C \sum_{k=0}^n \theta_k^2, \quad C = C(d, s). \quad (5.19)$$

Proof. Denote by ρ the maximal density of m , that is $\rho = 2^{-nd}/\ell_n^d$. For every $x \in E_n$ we have

$$m(B(x, r)) \leq \begin{cases} \rho r^d, & 0 < r < \ell_n, \\ C 2^{-kd}, & \ell_k \leq r \leq \ell_{k-1}, \quad k = 1, \dots, n, \\ 1, & \ell_0 \leq r < \infty, \end{cases}$$

with the positive constant C depending only on d . Hence, for every $x \in \mathbb{R}^d$,

$$\begin{aligned} \int_0^\infty \left[\frac{m(B(x, r))}{r^s} \right]^2 \frac{dr}{r} &\leq \int_0^{\ell_n} \left(\frac{2^{-nd}}{\ell_n^d} \right)^2 r^{2d-2s-1} dr + C \sum_{k=1}^n 2^{-2kd} \int_{\ell_k}^{\ell_{k-1}} \frac{dr}{r^{2s+1}} + \int_{\ell_0}^\infty \frac{dr}{r^{2s+1}} \\ &\leq \left(\frac{2^{-nd}}{\ell_n^d} \right)^2 \frac{\ell_n^{2d-2s}}{2d-2s} + C \sum_{k=1}^n \frac{1}{2s} \frac{2^{-2kd}}{\ell_k^{2s}} + \frac{1}{2s} \frac{1}{\ell_0^{2s}} \\ &\leq C(d, s) \sum_{k=0}^n \left(\frac{2^{-kd}}{\ell_k^s} \right)^2. \end{aligned}$$

Now (5.19) follows immediately from (4.10). \square

It was proved in [13] that under the condition $\theta_{k+1} \leq \theta_k$

$$C^{-1} \sum_{k=0}^n \theta_k^2 \leq |\mathfrak{R}_m^s|^2 \leq C \sum_{k=0}^n \theta_k^2, \quad (5.20)$$

where C depends on λ , s and d . Thus, Theorem 4.6 is a generalization of the estimate from above in (5.20).

Due to X. Tolsa [27] we know now that the condition $\theta_{k+1} \leq \theta_k$ is superfluous in the estimate from below as well.

6. CONSTRUCTION OF THE AUXILIARY MEASURE

We start with one property of Hausdorff contents.

Lemma 6.1. *Let h be a measuring function. For given $t_1 > 0$ we set*

$$\bar{h}(t) = \begin{cases} t^d h(t_1) t_1^{-d}, & 0 \leq t < t_1, \\ h(t), & t \geq t_1. \end{cases} \quad (6.1)$$

If $\alpha \in (0, 1)$, and the sets $F, G \in \mathbb{R}^d$ are such that

$$B(x, \alpha t_1) \subset G \quad \text{for every } x \in F,$$

then

$$M_h(F) \leq C M_{\bar{h}}(G), \quad C = C(\alpha, d).$$

Proof. We cover \mathbb{R}^d by the net of cubes Q_l with edge length $\alpha t_1 / \sqrt{d}$. For fixed $\varepsilon > 0$ we take a covering of G by balls $B_j = B(x_j, r_j)$, such that

$$M_{\bar{h}}(G) + \varepsilon > \sum_j \bar{h}(r_j).$$

Let Q_l be a cube from our net for which $Q_l \cap F \neq \emptyset$. Then $Q_l \subset G \subset \bigcup_j B_j$. We replace each ball B_j with $r_j \geq t_1$ by the ball $B'_j = 2B_j$. We remark that $h(r_j) \geq 2^{-d}h(2r_j)$. If Q_l intersects a ball B_j with $r_j \geq t_1$, then $Q_l \subset 2B_j$. Suppose now that Q_l intersects only balls B_j with $r_j < t_1$. These balls cover Q_l , and in turn can be covered and substituted by a ball of radius $3t_1$. Since $\sum_{j: B_j \cap Q_l \neq \emptyset} r_j^d \geq c_d(\alpha t_1)^d$, we have

$$\sum_{j: B_j \cap Q_l \neq \emptyset} \bar{h}(r_j) \geq \frac{\bar{h}(3t_1)}{(3t_1)^d} \sum_{j: B_j \cap Q_l \neq \emptyset} r_j^d \geq c(d, \alpha) \bar{h}(3t_1).$$

We discard all (possibly) remaining balls B_j with $r_j < t_1$. Thus, we obtain a new covering of F by balls $B'_j = B(x'_j, r'_j)$ with $r'_j \geq t_1$. Since each B_j of $r_j < t_1$ intersects at most $C = C(d, \alpha)$ cubes Q_l , we have

$$\begin{aligned} M_{\bar{h}}(G) + \varepsilon &> \sum_j \bar{h}(r_j) > c(d, \alpha) \sum_j \bar{h}(r'_j) \\ &= c(d, \alpha) \sum_j h(r'_j) \geq c(d, \alpha) M_h(F). \end{aligned}$$

□

Lemma 6.2. *Let $P > 0$ be given, and let ν be a linear combination of N Dirac point masses. There is a constant C_1 depending only on d and s , with the following property. If*

$$\mathbf{M} := M_h(\mathcal{Z}^*(\nu, P)) > \frac{C_1 \|\nu\|}{P} \max_{t_1 \leq t \leq t_2} \frac{h(t)}{t^s}, \quad (6.2)$$

where $t_1 = h^{-1}(0.1\mathbf{M}/N)$, $t_2 = h^{-1}(\mathbf{M})$, then there exists a positive Borel measure μ such that

- 1) $\text{supp } \mu \subset \mathcal{Z}^*(\nu, 0.8P)$;
- 2) $c\mathbf{M} \leq \|\mu\| \leq \mathbf{M}$, $c = c(d)$;
- 3) for every ball $B(x, r) \subset \mathbb{R}^d$, one has

$$\mu(B(x, r)) \leq \bar{h}(r), \quad (6.3)$$

where $\bar{h}(r)$ is defined by (6.1).

For $d = 2$ a similar assertion was proved in [8] (see Lemma 5.1 in [8]). The proof given here essentially differs from the arguments in [8].

Proof of Lemma 6.2. Let $|\nu| = \sum_{j=1}^N |\nu_j| \delta_{y_j}$ be the variation of ν . First of all we exclude from \mathbb{R}^d the set with high density of $|\nu|$ and with comparatively “small” h -content.

We say that a point $x \in \mathbb{R}^d$ is *normal* (with respect to $|\nu|$ and h) if the inequality

$$|\nu|(B(x, r)) \leq C_2^{-1} P \rho^{-1} h(r), \quad \rho := \max_{t_1 \leq t \leq t_2} \frac{h(t)}{t^s}, \quad (6.4)$$

holds for all $r \geq 0$. Here $C_2 < C_1$ is the constant depending only on d and s , which will be specified later. Let G_1 be the set of non-normal points $x \in \mathbb{R}^d$. For each $x \in G_1$ there exists $r = r(x)$ such that

$$h(r) < C_2 P^{-1} \rho |\nu|(B(x, r)). \quad (6.5)$$

We obtain a covering of G_1 by balls. Since (see (6.2), (6.5))

$$h(t_2) = \mathbf{M} > \frac{C_1 \|\nu\|}{P} \rho \geq \frac{C_1 |\nu|(B(x, r))}{P} \rho > h(r),$$

the radii of these balls are bounded by t_2 . By Besicovitch's covering lemma (see for example [15], p. 30), there is a subcovering $\{B'_k\}$, $B'_k = B(w'_k, r'_k)$, of multiplicity not exceeding A_d (that is, every point $x \in G$ is covered by at most A_d balls B'_k). We set $\mathcal{Z}_1 = \bigcup_k B'_k$. Then

$$M_h(\mathcal{Z}_1) \leq \sum_k h(r'_k) \stackrel{(6.5)}{<} \frac{C_2 \rho}{P} \sum_k |\nu|(B'_k) \leq \frac{A_d C_2 \rho}{P} \|\nu\| \stackrel{(6.2)}{<} 0.3\mathbf{M},$$

if $A_d C_2 < 0.3C_1$. We denote by $\{y_1, \dots, y_N\}$ the support of ν . Set $\mathcal{Z}_2 = \bigcup_{k=1}^N B(y_k, t_1)$. Then

$$M_h(\mathcal{Z}_2) \leq N h(t_1) = 0.1\mathbf{M}.$$

We claim that

$$|\nu|(B(x, r)) < C_2^{-1} P r^s \quad \text{for all } x \in \mathbb{R}^d \setminus (\mathcal{Z}_1 \cup \mathcal{Z}_2) \text{ and } r > 0. \quad (6.6)$$

Indeed, $|\nu|(B(x, r)) = 0$ when $0 \leq r < t_1$. For $t_1 \leq r < t_2$, the inequality (6.6) follows from (6.4). Assume that $r \geq t_2$. By (6.2),

$$\mathbf{M} > \frac{C_1 \|\nu\|}{P} \frac{h(t_2)}{t_2^s} = \frac{C_1 \|\nu\|}{P} \frac{\mathbf{M}}{t_2^s}.$$

Hence,

$$|\nu|(B(x, r)) \leq \|\nu\| < C_1^{-1} P t_2^s \leq C_1^{-1} P r^s,$$

and we get (6.6). We set

$$F = \mathcal{Z}^*(\nu, P) \setminus (\mathcal{Z}_1 \cup \mathcal{Z}_2).$$

Clearly,

$$M_h(F) \geq \mathbf{M} - M_h(\mathcal{Z}_1) - M_h(\mathcal{Z}_2) > \mathbf{M} - 0.3\mathbf{M} - 0.1\mathbf{M} = 0.6\mathbf{M}. \quad (6.7)$$

We will prove that

$$B(x_0, 0.4t_1) \subset \mathcal{Z}^*(\nu, 0.8P) \quad \text{for every } x_0 \in F. \quad (6.8)$$

Fix $x_0 \in F$, and let x be such that $|x - x_0| < 0.4t_1$. Since $x_0 \in \mathcal{Z}^*(\nu, P)$, there is $\varepsilon > 0$, for which $|R_{\nu, \varepsilon}^s(x_0)| > P$. We may assume that $\varepsilon > 0.9t_1$. (Indeed, $|\nu|(B(x_0, r)) = 0$ as $0 \leq r < t_1$. Hence, all integrals $R_{\nu, \varepsilon}^s(x_0)$ are the same for $0 < \varepsilon < t_1$.) Obviously,

$$\begin{aligned} & |R_{\nu, \varepsilon}^s(x_0) - R_{\nu, \varepsilon - |x - x_0|}^s(x)| \\ & \leq \int_{|y - x_0| > \varepsilon} \left| \frac{y - x_0}{|y - x_0|^{s+1}} - \frac{y - x}{|y - x|^{s+1}} \right| d|\nu|(y) + \int_{\{|y - x| > \varepsilon - |x - x_0|\} \cap \{|y - x_0| \leq \varepsilon\}} \frac{1}{|y - x|^s} d|\nu|(y). \end{aligned} \quad (6.9)$$

Since $|y - x_0| > \varepsilon > 0.9t_1$ implies $|y - x| > |y - x_0| - 0.4t_1 > 0.5|y - x_0|$, we have

$$\left| \frac{y - x_0}{|y - x_0|^{s+1}} - \frac{y - x}{|y - x|^{s+1}} \right| \leq C(s) \frac{|x - x_0|}{|y - x_0|^{s+1}} < \frac{C(s)\varepsilon}{|y - x_0|^{s+1}}.$$

Hence, the first integral on the right hand side of (6.9) does not exceed

$$\begin{aligned} C(s)\varepsilon \int_{|y - x_0| > \varepsilon} \frac{d|\nu|(y)}{|y - x_0|^{s+1}} &= C(s)\varepsilon \int_{\varepsilon}^{\infty} \frac{d|\nu|(B(x_0, t))}{t^{s+1}} \\ &\leq C(s)\varepsilon \int_{\varepsilon}^{\infty} \frac{(s+1)|\nu|(B(x_0, t))}{t^{s+2}} dt \leq \frac{C(s)\varepsilon(s+1)P}{C_2} \int_{\varepsilon}^{\infty} \frac{dt}{t^2} < 0.1P, \end{aligned}$$

if C_1 and C_2 are big enough (we integrated by parts and used (6.6)). Using (6.6) again we see that the second integral on the right hand side of (6.9) is bounded by

$$\frac{|\nu|(\overline{B}(x_0, \varepsilon))}{(\varepsilon - |x - x_0|)^s} < \frac{P\varepsilon^s}{C_2(0.5\varepsilon)^s} < 0.1P,$$

if $C_2 > 10 \cdot 2^s$. Thus,

$$|R_{\nu, \varepsilon}^s(x_0) - R_{\nu, \varepsilon - |x - x_0|}^s(x)| < 0.2P \quad \text{for all } x \in B(x_0, 0.4t_1).$$

Since $|R_{\nu, \varepsilon}^s(x_0)| > P$ and x_0 is any point in F , we get (6.8).

We consider the compact set

$$E = \overline{\bigcup_{x \in F} B(x, 0.2t_1)}.$$

By (6.8), $E \subset \mathcal{Z}^*(\nu, 0.8P)$. By Frostman's theorem (see for example [3], p. 7) there is a measure μ supported by E such that

$$\|\mu\| \geq a_d M_{\overline{h}}(E) \quad \text{and} \quad \mu(B(x, r)) \leq \overline{h}(r) \quad \text{for every } B(x, r) \subset \mathbb{R}^d.$$

Lemma 6.1 yields

$$\|\mu\| \geq a_d M_{\overline{h}}(E) \geq c(d) M_h(F) \stackrel{(6.7)}{>} c'(d) \mathbf{M}.$$

It remains to consider the inequality $\|\mu\| \leq \mathbf{M}$. If $\|\mu\| > \mathbf{M}$, we multiply μ by the constant $\mathbf{M}/\|\mu\| < 1$, fulfilling in this way all the requirements of Lemma 6.2. \square

7. PROOF OF THE FIRST PART OF THEOREM 4.1

As above, we set $\mathbf{M} = M_h(\mathcal{Z}^*(\nu, P))$. It is enough to prove the inequality

$$\mathbf{M} \leq C(s, d) \frac{\|\nu\|}{P} \left[\int_{t_1}^{t_2} \left(\frac{h(t)}{t^s} \right)^2 \frac{dt}{t} \right]^{1/2}, \quad (7.1)$$

where $t_1 = h^{-1}(0.1 \mathbf{M}/N)$, $t_2 = h^{-1}(\mathbf{M})$. Then Lemma 3.2 will imply the desired assertion.

We remark that

$$\max_{t_1 \leq t \leq t_2} \frac{h(t)}{t^s} =: \frac{h(t_0)}{t_0^s} < C_3 \left[\int_{t_1}^{t_2} \left(\frac{h(t)}{t^s} \right)^2 \frac{dt}{t} \right]^{1/2}, \quad C_3 = C_3(d, s). \quad (7.2)$$

Indeed, we have

$$0.05 > \frac{0.1}{N} = \frac{h(t_1)}{h(t_2)} = \frac{h(t_1)}{t_1^d} \frac{t_2^d}{h(t_2)} \left(\frac{t_1}{t_2} \right)^d \geq \left(\frac{t_1}{t_2} \right)^d.$$

Hence, $t_1 \leq (0.05)^{1/d} t_2$. There is the interval $[\alpha, \beta] \subset [t_1, t_2]$, containing t_0 and such that $\alpha = c(d)\beta$, $0 < c(d) < 1$. Since $h(t) > c'(d)h(t_0)$ as $t \in [\alpha, \beta]$, we get

$$\frac{h(t_0)}{t_0^s} < C(d, s) \left[\int_{\alpha}^{\beta} \left(\frac{h(t)}{t^s} \right)^2 \frac{dt}{t} \right]^{1/2},$$

that implies (7.2). So, if

$$\mathbf{M} \leq \frac{C_1 \|\nu\|}{P} \max_{t_1 \leq t \leq t_2} \frac{h(t)}{t^s},$$

then (7.1) holds, and the first part of Theorem 4.1 is proved. Thus, we may assume that (6.2) is fulfilled. Let μ be the measure in Lemma 6.2. Relations (6.3) and $\|\mu\| \leq \mathbf{M} = h(t_2)$ imply the estimate

$$\frac{\mu(B(x, r))}{r^s} \leq \max_{t_1 \leq t \leq t_2} \frac{h(t)}{t^s} \quad \text{for every ball } B(x, r) \subset \mathbb{R}^d.$$

Set

$$a_h := C_3 \left[\int_{t_1}^{t_2} \left(\frac{h(t)}{t^s} \right)^2 \frac{dt}{t} \right]^{1/2}, \quad \eta := a_h^{-1} \mu.$$

Then $\eta \in \Sigma_s$ (see (7.2) and (4.8)). Moreover, by (6.1) for every $x \in \mathbb{R}^d$, we have

$$\int_0^\infty \left[\frac{\mu(B(x, r))}{r^s} \right]^2 \frac{dr}{r} \leq \frac{h^2(t_1)}{t_1^{2d}} \int_0^{t_1} t^{2d-2s-1} dt + \int_{t_1}^{t_2} \left[\frac{h(t)}{t^s} \right]^2 \frac{dt}{t} + \int_{t_2}^\infty \frac{\|\mu\|^2}{t^{2s+1}} dt \leq C(d, s) a_h^2$$

(we remind the reader that $\|\mu\| \leq \mathbf{M} = h(t_2)$). Theorem 4.6 yields

$$|\mathfrak{R}_\eta^s|^2 = a_h^{-2} |\mathfrak{R}_\mu^s|^2 \leq C, \quad C = C(d, s). \quad (7.3)$$

We apply Theorem 4.5 with $t = 0.8P$ and η instead of μ . By (7.3), the constant C in (4.9) depends only on d and s . Since $\text{supp } \eta \subset \mathcal{Z}^*(\nu, 0.8P)$, (4.9) and the properties 1), 2) in Lemma 6.2 imply

$$\frac{C\|\nu\|}{0.8P} > \eta(\mathcal{Z}^*(\nu, 0.8P)) = \|\eta\| = a_h^{-1} \|\mu\| \geq c(d) a_h^{-1} \mathbf{M},$$

that is equivalent to (7.1). \square

8. PROOF OF THE SECOND PART OF THEOREM 4.1

Without loss of generality we may assume that $N = 2^{nd}$. Fix $\mathbf{M} > 0$, and let

$$\ell_j = h^{-1}(2^{-dj} \mathbf{M}), \quad j = 0, \dots, n-1, \quad \ell_n = \frac{1}{5} h^{-1}(2^{-dn} \mathbf{M}).$$

Since $t^{-d}h(t)$ is non-increasing, $\ell_{j+1} \leq \frac{1}{2}\ell_j$. Choose the set $J \subset \{0, \dots, n\}$ inductively as follows: $0 \in J$; if $j \in J$, then the least $k > j$ such that $\ell_k \leq \frac{1}{5}2^{j-k}\ell_j$, also belongs to J . If j_0, j_1, \dots, j_m are the elements of J listed in the increasing order, then $j_0 = 0$, $j_m = n$, and for every $k = 0, \dots, m-1$ we have


$$\ell_{j_k} \geq 2\ell_{j_{k+1}} \geq 4\ell_{j_{k+2}} \geq \dots \geq 2^{j_{k+1}-j_k-1} \ell_{j_{k+1}-1} \geq \frac{1}{5} \ell_{j_k} \geq 2^{j_{k+1}-j_k} \ell_{j_{k+1}}.$$

Construct the random set E recursively as follows: E_m is just the cube with edge length ℓ_{j_m} centered at the origin. Suppose that E_{k+1} is already defined as a random set. Let Q be the cube with edge length $\frac{1}{5}\ell_{j_k}$ centered at the origin. Partition it into $D_k := 2^{d(j_{k+1}-j_k-1)}$ equal subcubes. Let x_1, \dots, x_{D_k} be the centers of those subcubes. Take D_k independent copies of E_{k+1} and define F_k to be the union of those copies shifted by x_1, \dots, x_{D_k} . Now take 2^d independent copies of F_k and shift them by

$$\frac{\ell_{j_k}}{5}(\varepsilon + v_\varepsilon),$$

where ε runs over all 2^d vectors in \mathbb{R}^d whose coordinates are ± 1 , and v_ε are independent random vectors uniformly distributed over the cube with edge length $\frac{1}{10}$ centered at the origin and also independent of all F_k . The resulting random set is E_k .

It is easy to show by induction that each E_k is contained in the cube with edge length ℓ_{j_k} centered at the origin and, for $k < n$, the 2^d shifted copies of F_k whose union is E_k are separated by $\frac{1}{10}\ell_{j_k}$. Note also that $E = E_0$ consists of 2^{nd} randomly located cubes with edge length ℓ_n , which we will call the base cubes.



Denote by \tilde{E} the sure set constructed exactly in the same way as E , but with $v_\varepsilon = 0$.

Define the random measures μ and ν supported by E as follows. For each base cube Q , we put

$$\mu(Q) = 2^{-nd}, \quad \nu(Q) = 2^{-nd}\eta.$$

The measure μ will be proportional to Lebesgue measure on each base cube Q , and the measure ν will be a multiple of the Dirac point mass located at the center of Q .

Lemma 8.1. *For every ball $B_r \subset \mathbb{R}^d$ of radius $r > 0$, we have*

$$\mathbf{M}\mu(B_r) \leq Ch(r) \quad \text{with some } C = C(d, s) > 0.$$

Proof. Note first of all that the base cubes are disjoint. Thus, the density of μ with respect to the Lebesgue measure is not greater than

$$\frac{2^{-nd}}{\ell_n^d} = \frac{5^d \mathbf{M}^{-1} h(5\ell_n)}{(5\ell_n)^d}.$$

Hence,

$$\mu(B_r) \leq C \mathbf{M}^{-1} r^d \frac{h(r)}{r^d} = C \mathbf{M}^{-1} h(r),$$

if $r < 5\ell_n$.

Suppose now that $\ell_0 \geq r \geq 5\ell_n$. Then $r \in (\ell_{j_{k+1}}, \ell_{j_k}]$ for some k . Note that B_r can intersect only a bounded number of random copies of E_k constituting E because each such copy lies in its own cube with edge length ℓ_{j_k} .

Now E_k consists of 2^d blocks F_k . In each block the μ -measure of every cube of edge length

$$\frac{\ell_{j_k}}{5} 2^{j_{k+1}-j_k} \geq 2\ell_{j_{k+1}}$$

is $2^{-j_{k+1}d}$. Since B_r can intersect at most

$$\left[\frac{5r}{\ell_{j_k}} 2^{j_{k+1}-j_k} + 1 \right]^d \leq C_d \left[\left(\frac{5r}{\ell_{j_k}} 2^{j_{k+1}-j_k} \right)^d + 1 \right]$$

such cubes, we conclude that

$$\begin{aligned} \mu(B_r) &\leq C r^d \frac{2^{-j_k d}}{\ell_{j_k}^d} + C 2^{-j_{k+1}d} \leq C \mathbf{M}^{-1} r^d \frac{h(\ell_{j_k})}{\ell_{j_k}^d} + C \mathbf{M}^{-1} h(5\ell_{j_{k+1}}) \\ &\leq C \mathbf{M}^{-1} r^d \frac{h(r)}{r^d} + C' \mathbf{M}^{-1} h(\ell_{j_{k+1}}) < C'' \mathbf{M}^{-1} h(r). \end{aligned}$$

Finally, if $r \geq \ell_0$, we have

$$\mu(B_r) \leq \mu(E) = 1 = \mathbf{M}^{-1} h(\ell_0) \leq \mathbf{M}^{-1} h(r).$$

□

Corollary 8.2. *For every Borel set $G \subset \mathbb{R}^d$, we have*

$$M_h(G) \geq C^{-1} \mathbf{M} \mu(G), \quad C = C(d, s) > 0.$$

Lemma 8.3. *Let $\theta_k > 0$, $k = 0, \dots, m-1$. Let ξ_k be \mathbb{R}^d -valued independent random variables satisfying*

$$|\xi_k| \leq C\theta_k, \quad \sum_k \text{Var } \xi_k \geq c \sum_k \theta_k^2.$$

Then there exists $\delta = \delta(C, c, d) > 0$ such that

$$\mathcal{P} \left\{ \left| \sum_{k=0}^{m-1} \xi_k + a \right| \geq \delta \left(\sum_{k=0}^{m-1} \theta_k^2 \right)^{1/2} \right\} \geq \delta \text{ for all } a \in \mathbb{R}^d.$$

Proof. Denote

$$\sigma = \sum_{k=0}^{m-1} \xi_k + a, \quad \zeta_k = \xi_k - \mathcal{E} \xi_k.$$

It is enough to consider the case $d = 1$. Indeed, since $\int_{|e|=1} \text{Var}\langle \xi_k, e \rangle dm(e) = \frac{1}{d} \text{Var} \xi_k$, we can find a unit vector e such that $\sum_k \text{Var}\langle \xi_k, e \rangle \geq \frac{c}{d} \sum_k \theta_k^2$. But, obviously, $\langle \xi_k, e \rangle \leq C\theta_k$ and $|\sigma| \geq |\sum_k \langle \xi_k, e \rangle + \langle a, e \rangle|$.

Take $\lambda > 0$ and consider

$$|\mathcal{E} e^{i\lambda\sigma}| = \left| e^{i\lambda a} \prod_{k=0}^{m-1} \mathcal{E} e^{i\lambda\xi_k} \right| = \prod_{k=0}^{m-1} |\mathcal{E} e^{i\lambda\xi_k}| = \prod_{k=0}^{m-1} |\mathcal{E} e^{i\lambda\zeta_k}|.$$

Note now that

$$e^{i\lambda\zeta_k} = 1 + i\lambda\zeta_k - \frac{\lambda^2}{2}\zeta_k^2 + O(\lambda^3\zeta_k^3).$$

Hence,

$$|\mathcal{E} e^{i\lambda\zeta_k}| = \left| 1 - \frac{\lambda^2}{2} \text{Var} \xi_k + C\lambda^3\theta_k^3 \right| \leq \exp\left(-\frac{\lambda^2}{2} \text{Var} \xi_k + C\lambda^3\theta_k^3\right),$$

and

$$\begin{aligned} \prod_{k=0}^{m-1} |\mathcal{E} e^{i\lambda\zeta_k}| &\leq \exp\left(-c\lambda^2 \sum \theta_k^2 + C\lambda^3 \sum \theta_k^3\right) \\ &\leq \exp\left(-c' \left[\lambda \left(\sum \theta_k^2\right)^{1/2}\right]^2 + C' \left[\lambda \left(\sum \theta_k^2\right)^{1/2}\right]^3\right). \end{aligned}$$

Now choose

$$\lambda = \frac{c'}{2C'} \left(\sum \theta_k^2\right)^{-1/2}.$$

Then

$$|\mathcal{E} e^{i\lambda\sigma}| \leq \exp\left(-\frac{(c')^3}{8(C')^2}\right).$$

On the other hand, for every $\delta > 0$, one has

$$\begin{aligned} |\mathcal{E} e^{i\lambda\sigma} - 1| &\leq \lambda\delta \left(\sum \theta_k^2\right)^{1/2} + 2\mathcal{P}\left\{|\sigma| > \delta \left(\sum \theta_k^2\right)^{1/2}\right\} \\ &\leq \frac{c'}{2C'}\delta + 2\mathcal{P}\left\{|\sigma| > \delta \left(\sum \theta_k^2\right)^{1/2}\right\}. \end{aligned}$$

Hence,

$$\mathcal{P}\left\{|\sigma| > \delta \left(\sum \theta_k^2\right)^{1/2}\right\} \geq \frac{1}{2} \left[1 - \exp\left(-\frac{(c')^3}{8(C')^2}\right) - \frac{c'}{2C'}\delta\right] > \delta,$$

if δ is chosen small enough. \square

Lemma 8.4. *Let ν be a positive measure supported by the cube Q with edge length ℓ centered at the origin. Let $x \in [\frac{5}{4}\ell, \frac{11}{4}\ell]^d$. Let v be the random vector uniformly distributed over the cube with edge length $\frac{1}{2}\ell$ centered at the origin. Then the random variable $\xi = R_\nu^s(x + v)$ satisfies*

$$|\xi| \leq C\theta, \quad \text{Var} \xi \geq c\theta^2, \quad \text{where } \theta = \frac{\|\nu\|}{\ell^s}.$$

Proof. Since for every $j = 1, \dots, d$, and every $y \in Q$, we have

$$(x + v - y)_j \geq \frac{5\ell}{4} - \frac{\ell}{2} - \frac{\ell}{4} \geq \frac{\ell}{2},$$

we conclude that

$$|R_\nu^s(x+v)| \leq \frac{2^s}{\ell^s} \|\nu\| = 2^s \theta.$$

Now we have to consider two cases.

Case 1: $d = 1$.

Then

$$\begin{aligned} \Delta &= |\mathcal{E}\{R_\nu^s(x+v)|v < 0\} - \mathcal{E}\{R_\nu^s(x+v)|v > 0\}| \\ &= \frac{4}{\ell} \int_{[-\ell/4, 0]} \left[R_\nu^s(x+v) - R_\nu^s\left(x+v + \frac{\ell}{4}\right) \right] dv \\ &= \frac{4}{\ell} \int_{[-\ell/4, 0]} dv \int_Q \left[\frac{1}{(x+v-y)^s} - \frac{1}{(x+v + \frac{1}{4}\ell - y)^s} \right] d\nu(y). \end{aligned}$$

Note now that

$$\frac{\partial}{\partial t} \frac{1}{t^s} = -\frac{s}{t^{s+1}} \leq -\frac{s}{(\frac{7}{2}\ell)^{s+1}}, \quad \text{whenever } t \in \left[x+v-y, x+v + \frac{\ell}{4} - y \right] \subset \left[\frac{\ell}{2}, \frac{7\ell}{2} \right].$$

Thus, the difference of the conditional expectations is at least

$$s \left(\frac{2}{7} \right)^{s+1} \frac{1}{4} \frac{\|\nu\|}{\ell^s} = c\theta.$$

Since $\text{Var } \xi \geq \Delta^2/4$, we are done in this case.

Case 2: $d \geq 2$.

Let now

$$\Delta = |\mathcal{E}\{[R_\nu^s(x+v)]_1 | v_2 < 0\} - \mathcal{E}\{[R_\nu^s(x+v)]_1 | v_2 > 0\}|.$$

Since for $t \in [\frac{1}{2}\ell, \frac{7}{2}\ell]^d$, we have

$$\frac{\partial}{\partial t_2} \frac{t_1}{|t|^{s+1}} = -(s+1) \frac{t_1 t_2}{|t|^{s+3}} \leq -\frac{c}{\ell^{s+1}}.$$

Denote $\xi^{(1)} := (\xi, e_1)$. Then the arguments analogous to that for the one-dimensional case allows us to conclude that

$$\text{Var } \xi \geq \text{Var } \xi^{(1)} \geq \frac{\Delta^2}{4} \geq c\theta^2.$$

□

Fix $x \in \tilde{E}$ and consider the random variable $\sigma = R_\nu^s(\Phi(x))$, where $\Phi : \tilde{E} \mapsto E$ is the (random) canonical measure preserving mapping between \tilde{E} and E defined above. Namely, Φ shifts each base cube of \tilde{E} onto the corresponding perturbed base cube of E .

Now fix the random shifts responsible for the positioning of the copies of the blocks F_k ($k = 0, \dots, m-1$), that contain $\Phi(x)$.

Suppose now that some copy of the block F_k contains $\Phi(x)$. Then fix all shifts responsible for the positioning of the base cubes within the copies F_k not containing $\Phi(x)$. That will leave free $2^d - 1$ shifts of the copies of the block F_k at each of the levels $k = 0, \dots, m-1$. Out of these shifts, fix all except the one responsible for the positioning of the copy of F_k “opposite” in each coordinate to the copy containing $\Phi(x)$ in the copy of E_k containing $\Phi(x)$.

Now notice that $R_\nu^s(\Phi(x))$ is the sum of a fixed part, that we shall call a , and m independent random parts ξ_0, \dots, ξ_{m-1} , where ξ_k is the potential of the part of the measure μ supported by the copy of F_k in the copy of E_k containing $\Phi(x)$ that is “opposite” to the copy of F_k containing $\Phi(x)$.

Applying Lemma 8.4 to ξ_k with $\ell = \frac{1}{5}\ell_{j_k}$ (in Lemma 8.4 we moved the point and fixed the measure while here we are moving the measure and fixing the point, but for convolution kernels it is the same; also the obvious change of directions of the coordinate axes should be performed before we find ourselves in the conditions of Lemma 8.4), we conclude that

$$|\xi_k| \leq C \frac{\eta 2^{-j_k d}}{\ell_{j_k}^s}, \quad \text{and} \quad \text{Var } \xi_k \geq C \left(\frac{\eta 2^{-j_k d}}{\ell_{j_k}^s} \right)^2, \quad k = 0, 1, \dots, m-1.$$

Hence, Lemma 8.3 is applicable and we get

$$\mathcal{P} \left\{ |R_\nu^s(\Phi(x))| \geq \delta \left(\sum_{k=0}^{m-1} \theta_k^2 \right)^{1/2} \right\} \geq \delta \quad \text{with} \quad \theta_k = \frac{\eta 2^{-j_k d}}{\ell_{j_k}^s}.$$

This is a conditional probability, of course, conditioned on freezing a lot of shifts. But, since δ and θ_k do not depend on the values of the frozen shifts, we get the same inequality for the full probability.

The immediate conclusion is that there exists a set of shifts such that

$$\mu(G) \geq \delta, \quad \text{where} \quad G = \left\{ y \in E : |R_\nu^s(y)| \geq \delta \left(\sum_{k=0}^{m-1} \theta_k^2 \right)^{1/2} \right\},$$

and, by Corollary 8.2, $M_h(G) \geq c\mathbf{M}$.

Now note that

$$\begin{aligned} \int_{C/N}^1 \left[\frac{t}{h^{-1}(\mathbf{M}t)^s} \right]^2 \frac{dt}{t} &\leq \sum_{j=0}^{n-2} \int_{2^{-(j+1)d}}^{2^{-jd}} \left[\frac{t}{h^{-1}(\mathbf{M}t)^s} \right]^2 \frac{dt}{t} \\ &\leq C \sum_{j=0}^{n-2} \left[\frac{2^{-jd}}{h^{-1}(\mathbf{M}2^{-(j+1)d})^s} \right]^2 = C \sum_{j=1}^{n-1} \left[\frac{2^{-jd}}{\ell_j^s} \right]^2 \\ &\leq C \sum_{k=0}^{m-1} \sum_{j_k \leq j < j_{k+1}} \left[\frac{2^{-jd}}{\ell_j^s} \right]^2 \leq C \sum_{k=0}^{m-1} \left[\frac{2^{-j_k d}}{\ell_{j_k}^s} \right]^2 \sum_{j \geq j_k} 2^{-2(j-j_k)(d-s)} \\ &\leq C \sum_{k=0}^{m-1} \left[\frac{2^{-j_k d}}{\ell_{j_k}^s} \right]^2 = C \eta^{-2} \sum_{k=0}^{m-1} \theta_k^2. \end{aligned}$$

Thus, we constructed a linear combination of N Dirac point masses ν such that $\|\nu\| = \eta$ and

$$M_h \left(\mathcal{Z} \left(\nu, c\eta \left\{ \int_{C/N}^1 \left[\frac{t}{h^{-1}(\mathbf{M}t)^s} \right]^2 \frac{dt}{t} \right\}^{1/2} \right) \right) \geq c\mathbf{M}.$$

Now, given $P > 0$, put

$$\mathbf{M} = \mathfrak{M}_h \left(\frac{c\eta}{P}, \frac{N}{C} \right).$$

Then

$$c\eta \left\{ \int_{C/N}^1 \left[\frac{t}{h^{-1}(\mathbf{M}t)^s} \right]^2 \frac{dt}{t} \right\}^{1/2} = P.$$

It remains to note that

$$\mathfrak{M}_h \left(\frac{c\eta}{P}, \frac{N}{C} \right) \geq c' \mathfrak{M}_h \left(\frac{\eta}{P}, N \right).$$

This proves the theorem for $N \geq 2C$.

For $2 \leq N < 2C$, just put all N masses to one point. Then

$$M_h(\mathcal{Z}(\nu, P)) = h\left(\left(\frac{\eta}{P}\right)^{1/s}\right).$$

Note now that

$$\varkappa^2 \int_{1/2}^1 \left[\frac{t}{h^{-1}(2h(\varkappa^{1/s})t)^s} \right]^2 \frac{dt}{t} \leq \varkappa^2 \int_{1/2}^1 \left[\frac{t}{\varkappa} \right]^2 \frac{dt}{t} < 1,$$

so $h(\varkappa^{1/s}) \geq \frac{1}{2}\mathfrak{M}_h(\varkappa, 2)$. Thus,

$$h\left(\left(\frac{\eta}{P}\right)^{1/s}\right) \geq \frac{1}{2}\mathfrak{M}_h\left(\frac{\eta}{P}, 2\right) \geq c\mathfrak{M}_h\left(\frac{\eta}{P}, 2C\right) \geq c\mathfrak{M}_h\left(\frac{\eta}{P}, N\right).$$

9. PROOF OF THEOREM 4.3

At the beginning we consider a measure ν with compact support. Without loss of generality we assume that

$$\mathbf{M} := M_h(\mathcal{Z}^*(\nu, P)) > \frac{C_1\|\nu\|}{P}\rho_0, \quad \text{where } \rho_0 := \max_{0 < t \leq t_2} \frac{h(t)}{t^s}, \quad h(t_2) = \mathbf{M}.$$

Otherwise the same arguments as in Section 7 (with sufficiently big N) yield (4.6). In the same way as in the proof of Lemma 6.2 we define the set \mathcal{Z}_1 (we use the notation in Lemma 6.2), taking ρ_0 instead of ρ . Repeating the arguments in the proof of Lemma 6.2 and choosing $C_2 > 10$, we see that

$$M_h(\mathcal{Z}_1) < 0.3\mathbf{M}, \quad (9.1)$$

$$|\nu|(B(x, r)) < 0.1Pr^s \quad \text{for any } x \in \mathbb{R}^d \setminus \mathcal{Z}_1 \text{ and } r > 0. \quad (9.2)$$

For fixed $\delta > 0$ we set

$$\mathcal{Z}_\delta^*(\nu, P) = \{x \in \mathbb{R}^d : \sup_{\varepsilon > \delta} |R_{\nu, \varepsilon}^s(x)| > P\}.$$

There exists a measure ν' consisting of finitely many point charges such that

$$\|\nu'\| = \|\nu\|, \quad |R_{\nu, \varepsilon}^s(x) - R_{\nu', \varepsilon}^s(x)| < 0.2P \quad \text{for any } x \in \mathbb{R}^d \setminus \mathcal{Z}_1 \text{ and } \varepsilon \geq \delta. \quad (9.3)$$

For the construction of ν' we cover \mathbb{R}^d by a sufficiently fine net of cubes and put at the center of each cube a charge equal to the ν -measure of this cube (the parts of the measure concentrated on sides of adjacent cubes can be ascribed to any of them). The difference of the integrals over cubes lying in the domain $\{y \in \mathbb{R}^d : |y - x| > \varepsilon \geq \delta\}$ can be made arbitrarily small, while the difference of the integrals over those pieces of cubes intersecting the sphere $|y - x| = \varepsilon$ which lie in this domain, can be made not exceeding $0.15P$ (we use here (9.2)). For $\delta = 1/n$ we denote this measure ν' by ν_n . By (9.3),

$$\mathcal{Z}_{1/n}^*(\nu, P) \setminus \mathcal{Z}_1 \subset \mathcal{Z}_{1/n}^*(\nu_n, 0.8P) \setminus \mathcal{Z}_1 \subset \mathcal{Z}^*(\nu_n, 0.8P), \quad n \in \mathbb{Z}_+.$$

Obviously,

$$\mathcal{Z}_{1/n}^*(\nu, P) \setminus \mathcal{Z}_1 \nearrow \mathcal{Z}^*(\nu, P) \setminus \mathcal{Z}_1.$$

Then

$$c(d)M_h(\mathcal{Z}^*(\nu, P) \setminus \mathcal{Z}_1) \leq \lim_{n \rightarrow \infty} M_h(\mathcal{Z}_{1/n}^*(\nu, P) \setminus \mathcal{Z}_1).$$

This inequality follows from arguments given by Carleson in [3], p. 9–11 (see also [8], Lemma 7.1). By (9.1), $0.7\mathbf{M} \leq M_h(\mathcal{Z}^*(\nu, P) \setminus \mathcal{Z}_1)$. Applying Theorem 4.1 for $\nu = \nu_n$ and Lemma 3.1, for sufficiently big n we get

$$\mathbf{M} \leq CM_h(\mathcal{Z}^*(\nu_n, 0.8P)) \leq C' \mathfrak{M}_h\left(\frac{\|\nu\|}{P}, \infty\right),$$

that is (4.6).

Suppose that $\text{supp } \nu$ is not bounded.

Fix $R > 0$ and take $R_1 > R$. Let ν_0 be a measure such that $\nu = \nu_0$ in $B(0, R_1)$, and $\nu_0 = 0$ outside of $B(0, R_1)$. For sufficiently big R_1 ,

$$\{\mathcal{Z}^*(\nu, P) \cap B(0, R)\} \subset \{\mathcal{Z}^*(\nu_0, 0.9P) \cap B(0, R)\} \subset \mathcal{Z}^*(\nu_0, 0.9P).$$

Hence,

$$M_h(\mathcal{Z}^*(\nu, P) \cap B(0, R)) \leq C(d, s) \mathfrak{M}_h\left(\frac{\|\nu_0\|}{0.9P}, \infty\right) \leq C'(d, s) \mathfrak{M}_h\left(\frac{\|\nu\|}{P}, \infty\right) \quad \text{for all } R > 0 \quad (9.4)$$

(it is clear that Lemma 3.1 is correct for $N = \infty$ as well). We will prove that this inequality implies (4.6).

Choose $R > 0$ such that $h(\frac{1}{2}R) - 1$ is greater then the right hand side of (9.4), and set

$$G_1 = \mathcal{Z}^*(\nu, P) \cap B(0, R), \quad G_k = \mathcal{Z}^*(\nu, P) \cap \{y : (k-1)R \leq |y| < kR\}.$$

Fix $\varepsilon \in (0, 1)$ and $K \in \mathbb{N}$, and consider a covering of $\mathcal{Z}^*(\nu, P) \cap B(0, KR)$ by balls $B_j = B(x_j, r_j)$ such that

$$\sum_j h(r_j) < M_h(\mathcal{Z}^*(\nu, P) \cap B(0, KR)) + \varepsilon \leq C'(d, s) \mathfrak{M}_h\left(\frac{\|\nu\|}{P}, \infty\right) + \varepsilon.$$

Every ball B_j intersects at most two sets G_k , $k = 1, \dots, K$. Hence,

$$2C'(d, s) \mathfrak{M}_h\left(\frac{\|\nu\|}{P}, \infty\right) + 2\varepsilon > 2 \sum_j h(r_j) \geq \sum_{k=1}^K \sum_{j: B_j \cap G_k \neq \emptyset} h(r_j) \geq \sum_{k=1}^K M_h(G_k),$$

and we get the estimate

$$M_h(\mathcal{Z}^*(\nu, P)) \leq \sum_{k=1}^{\infty} M_h(G_k) \leq 2C'(d, s) \mathfrak{M}_h\left(\frac{\|\nu\|}{P}, \infty\right).$$

□

10. THE CASE $s \geq d$: PROOF OF THEOREM 4.4

Case $s > d$. Let $K(t)$, $t \in (0, +\infty)$, be a non-increasing continuous function such that $K(t) \rightarrow +\infty$ as $t \rightarrow +0$. By [8], Theorem 1.1,

$$M_h\left(\left\{x \in \mathbb{R}^d : \int K(|x-y|) d|\nu|(y) > P\right\}\right) < Ch(r_0), \quad C = C(d),$$

where r_0 is a solution of the equation

$$h(r_0) = \frac{\|\nu\|}{P} \left(h(r'_0)K(r'_0) + \int_{r'_0}^{r_0} K(t) dh(t) \right) \quad \text{with } h(r'_0) = \frac{h(r_0)}{N}.$$

We apply this result for $K(t) = t^{-s}$, $s > d$. Integrating by parts, we see that the right hand side does not exceed

$$\frac{\|\nu\|}{P} \left(\frac{h(r_0)}{r_0^s} + s \int_{r'_0}^{r_0} \frac{h(t)}{t^{s+1}} dt \right).$$

Since $h(t) \leq h(r'_0)(r'_0)^{-d}t^d$ for $t \geq r'_0$, we have

$$h(r_0) \leq \frac{\|\nu\|}{P} \left(\frac{h(r_0)}{r_0^s} + \frac{sh(r'_0)}{(r'_0)^d} \frac{(r'_0)^{d-s}}{s-d} \right) \leq C(s, d) \frac{\|\nu\|}{P} \frac{h(r'_0)}{(r'_0)^s}.$$

Hence,

$$r'_0 \leq \left(C(s, d) \frac{\|\nu\|}{NP} \right)^{1/s}.$$

Therefore,

$$\begin{aligned} M_h(\mathcal{Z}^*(\nu, P)) &\leq M_h(\mathcal{X}(|\nu|, P)) \leq Ch(r_0) = CNh(r'_0) \\ &< C'Nh \left(\left(\frac{\|\nu\|}{NP} \right)^{1/s} \right), \quad C' = C'(s, d) \end{aligned}$$

(we used the inequality $h(2t) \leq 2^d h(t)$).

Case $s = d$. Without loss of generality we assume that

$$h(t_2) = \mathbf{M} := M_h(\mathcal{Z}^*(\nu, P)) > C_1Nh \left(\left(\frac{\|\nu\|}{NP} \right)^{1/d} \right)$$

with sufficiently big C_1 . Set

$$t_1 = \left(\frac{\|\nu\|}{NP} \right)^{1/d}, \quad \rho = \frac{h(t_1)}{t_1}.$$

As in the proof of Lemma 6.2, we construct the sets \mathcal{Z}_1 , \mathcal{Z}_2 and F such that

$$\begin{aligned} F &= \mathcal{Z}^*(\nu, P) \setminus (\mathcal{Z}_1 \cup \mathcal{Z}_2), \quad M_h(F) \geq 0.6\mathbf{M}, \\ B(x, 0.4t_1) &\subset \mathcal{Z}^*(\nu, 0.8P) \text{ for every } x \in F. \end{aligned}$$

Let μ be the d -dimensional Lebesgue measure, and let

$$G = \bigcup_{x \in F} B(x, 0.4t_1).$$

Obviously, $G \subset \mathcal{Z}^*(\nu, 0.8P)$. By the classical Calderón-Zygmund result,

$$\frac{C\|\nu\|}{P} > \mu(\mathcal{Z}^*(\nu, 0.8P)) \geq \mu(G), \quad C = C(d).$$

The set G can be covered by balls B_j with the same radii t_1 in such a way that

$$\mu(G) \geq c \sum_j t_1^d, \quad c = c(d).$$

Obviously,

$$\begin{aligned} \mu(G) &\geq c \sum_j t_1^d = \frac{c t_1^d}{h(t_1)} \sum_j h(t_1) \geq \frac{c t_1^d}{h(t_1)} M_h(G) \\ &\geq \frac{c t_1^d}{h(t_1)} M_h(F) \geq \frac{c' t_1^d}{h(t_1)} \mathbf{M}, \quad c' = c'(d). \end{aligned}$$

Thus,

$$\mathbf{M} \leq \frac{C\|\nu\|}{P} \frac{h(t_1)}{t_1^d} = \frac{C\|\nu\|}{P} \frac{NP}{\|\nu\|} h(t_1) = CNh(t_1).$$

Theorem 4.4 is proved. \square

11. HAUSDORFF CONTENT AND CAPACITY

The main object of this section is the capacity $\gamma_{s,+}(E)$ of a compact set $E \subset \mathbb{R}^d$ defined by the equality

$$\gamma_{s,+}(E) := \sup\{\|\mu\| : \mu \in M_+(E), \|R_\mu^s(x)\|_\infty \leq 1\},$$

where $M_+(E)$ is the class of positive Radon measures supported by E .

Remark. In [28], p. 46, the capacity $\gamma_{s,+}(E)$ is defined in the following way:

$$\gamma_{s,+}(E) := \sup\{\|\mu\| : \mu \in \Sigma_s, \text{supp } \mu \subset E, \|R_\mu^s(x)\|_\infty \leq 1\}.$$

It is shown in [12], p. 217, that if $\|R_\mu^s(x)\|_\infty \leq 1$, then

$$\mu(B(x, r)) \leq Cr^s, \quad x \in \mathbb{R}^d, \quad r > 0$$

for every measure $\mu \in M_+(E)$. Arguments in this part of the proof of Lemma 4.1 in [12] valid not only for $0 < s < 1$, but for $0 < s < d$ as well. (We note that the reference [P], Lemma 11 in [12] should be replaced by [P], Lemma 3.1.) For $s = d - 1$, this fact is also noted in [28], p. 46. Therefore, these two definitions of $\gamma_{s,+}$ are equivalent.

This capacity is connected with various problems in analysis.

For $d = 2$, $s = 1$,

$$\gamma_{1,+}(E) \asymp \gamma(E),$$

where $\gamma(E)$ is analytic capacity (see [26] and [28]). Here $A \asymp B$ means that $C^{-1}A \leq B \leq CA$ with C depending (possibly) only on d and s .

For $s = d - 1$, $d \geq 2$,

$$\gamma_{s,+}(E) \asymp \kappa(E), \tag{11.1}$$

where $\kappa(E)$ is the Lipschitz harmonic capacity

$$\kappa(E) := \sup\{|\langle \Delta f, 1 \rangle| : f \in \text{Lip}_{\text{loc}}^1(\mathbb{R}^d), \text{supp}(\Delta f) \subset E, \|\nabla f\|_\infty \leq 1, \nabla f(\infty) = 0\},$$

introduced by Paramonov [23] in connection with problems of approximation by harmonic functions. Here (as usual) $\langle T, \varphi \rangle$ means the action of a distribution T with compact support on a smooth test function. It was noticed in [23] that $\kappa(E) \leq 2\pi\gamma(E)$ for $d = 2$. The relation (11.1) was proved in [26] for $d = 2$ and in [28] for $d > 2$ (see [28], Theorem 2.1 and Lemma 5.15). The null-sets for the capacity κ are the same as the removable sets for Lipschitz harmonic functions, see [23], [16]. In these papers Mattila and Paramonov established important geometrical properties of the capacity κ .

Moreover, $\gamma_{s,+}(E)$ is related to the Riesz capacity $C_{\alpha,p}$ in non-linear potential theory. We discuss this relation near the end of the paper.

Main results of this section concern the connections between Hausdorff content and the capacity $\gamma_{s,+}$. We need the following important characterization of $\gamma_{s,+}$ obtained in [28], Chapter 5:

$$\gamma_{s,+}(E) \asymp \gamma_{op}(E) := \sup\{\|\mu\| : \mu \in \Sigma_s, \text{supp } \mu \subset E, |\mathfrak{R}_\mu^s| \leq 1\}, \quad 0 < s < d. \tag{11.2}$$

Proof of Theorem 4.7. Choose $\mu \in M_+(E)$. We may assume that $\int_{\mathbb{R}^d} W^\mu(x) d\mu(x) < \infty$. Set

$$G := \left\{ x \in \mathbb{R}^d : W^\mu(x) > \frac{2}{\|\mu\|} \int_{\mathbb{R}^d} W^\mu(x) d\mu(x) \right\}.$$

It is easy to see that G is open and

$$\mu(G) \leq \frac{1}{2} \|\mu\|.$$

Let

$$\mu^* = \mu|_{(\mathbb{R}^d \setminus G)}, \quad \mathbf{S} = \sup_{x \in \text{supp } \mu^*} W^{\mu^*}(x).$$

We claim that

$$W^{\mu^*}(x) \leq 2^{2s+1} \mathbf{S} \quad \text{for all } x \in \mathbb{R}^d. \quad (11.3)$$

It is enough to consider x with $\delta := \text{dist}(x, \text{supp } \mu^*) > 0$. Let x' be such that $x' \in \text{supp } \mu^*$ and $|x - x'| = \delta$. Then

$$\begin{aligned} W^{\mu^*}(x) &= \int_{\delta}^{\infty} \left[\frac{\mu^*(B(x, r))}{r^s} \right]^2 \frac{dr}{r} \leq \int_{\delta}^{\infty} \left[\frac{\mu^*(B(x', r + \delta))}{r^s} \right]^2 \frac{dr}{r} \\ &= \int_{2\delta}^{\infty} \left[\frac{\mu^*(B(x', t))}{(t - \delta)^s} \right]^2 \frac{dt}{t - \delta} < 2^{2s+1} \int_{2\delta}^{\infty} \left[\frac{\mu^*(B(x', t))}{t^s} \right]^2 \frac{dt}{t} \leq 2^{2s+1} \mathbf{S}, \end{aligned}$$

and we get (11.3).

Let $\eta = (2^{2s+2} s \mathbf{S})^{-1/2} \mu^*$. Since for each ball $B(x, r)$

$$2^{2s+1} \mathbf{S} \geq \int_0^{\infty} \left[\frac{\mu^*(B(x, t))}{t^s} \right]^2 \frac{dt}{t} \geq \int_r^{\infty} \left[\frac{\mu^*(B(x, t))}{t^s} \right]^2 \frac{dt}{t} \geq \frac{[\mu^*(B(x, r))]^2}{2sr^{2s}},$$

we see that $\eta \in \Sigma_s$. Moreover, (4.10) implies

$$|\mathfrak{R}_\eta^s|^2 \leq C(2^{2s+2} s \mathbf{S})^{-1} \mathbf{S} = C'.$$

Relations (11.2) and $\|\mu^*\| \geq \frac{1}{2} \|\mu\|$ yield

$$\gamma_{s,+}(E) \geq C\eta(E) \geq C' \|\mu\| \mathbf{S}^{-1/2}.$$

Since

$$\mathbf{S} \leq \sup_{x \in \text{supp } \mu^*} W^\mu(x) \leq \frac{2}{\|\mu\|} \int_{\mathbb{R}^d} W^\mu(x) d\mu(x),$$

we have

$$\gamma_{s,+}(E) \geq C \|\mu\|^{3/2} \left[\int_{\mathbb{R}^d} W^\mu(x) d\mu(x) \right]^{-1/2},$$

that implies (4.11). □

Theorem 11.1. *Under assumption (4.5), for each compact set $E \subset \mathbb{R}^d$,*

$$\gamma_{s,+}(E) \geq CM_h(E) \left[\int_0^{t_2} \left(\frac{h(t)}{t^s} \right)^2 \frac{dt}{t} \right]^{-1/2}, \quad 0 < s < d, \quad (11.4)$$

where C depends only on d, s , and t_2 is defined by the equality $h(t_2) = M_h(E)$.

Proof. By Frostman's theorem (see [3], p. 7) there is a positive measure μ such that

$$\begin{aligned} \text{supp } \mu &\subset E, \\ \mu(B(x, r)) &\leq h(r) \text{ for each ball } B(x, r) \subset \mathbb{R}^d, \\ \mu(E) &\geq CM_h(E) \text{ with } C \text{ depending only on } d. \end{aligned}$$

Without loss of generality we can assume that $\|\mu\| \leq M_h(E)$ (otherwise we divide μ by the constant $\|\mu\|/M_h(E) > 1$). Then

$$W^\mu(x) \leq \int_0^{t_2} \left[\frac{h(t)}{t^s} \right]^2 \frac{dt}{t} + \int_{t_2}^\infty \left[\frac{h(t_2)}{t^s} \right]^2 \frac{dt}{t} \leq C \int_0^{t_2} \left[\frac{h(t)}{t^s} \right]^2 \frac{dt}{t},$$

since by the doubling property of h ,

$$\left[\frac{h(t_2)}{t_2^s} \right]^2 \leq C \int_0^{t_2} \left[\frac{h(t)}{t^s} \right]^2 \frac{dt}{t}.$$

Inequality (11.4) follows directly from (4.11). \square

For $h(t) = t^\beta$ easy calculations give the following result.

Corollary 11.2. *For each compact set $E \subset \mathbb{R}^d$,*

$$\gamma_{s,+}(E) \geq C(\beta - s)^{1/2} [M_h(E)]^{s/\beta}, \quad \text{where } 0 < s < d, \quad h(t) = t^\beta, \quad \beta > s,$$

and C depends only on d and s .

The next statement can be viewed as a counterpart of the classical Frostman's theorem on connections between capacities generated by potentials with positive kernels and Hausdorff measure $\Lambda_h(E)$ (see, for example, [3], Section IV, Theorem 1).

Corollary 11.3. *For each compact set $E \subset \mathbb{R}^d$, the condition $\gamma_{s,+}(E) > 0$ implies $\Lambda_h(E) > 0$ for $h(t) = t^s$.*

On the other hand, if $\Lambda_h(E) > 0$ for a measuring function h satisfying (4.5), then $\gamma_{s,+}(E) > 0$.

Proof. The first part of Corollary 11.3 is a direct consequence of the following result by Prat [24], p. 946: for $0 < s < d$

$$C_\varepsilon [M_{t^{s+\varepsilon}}(E)]^{s/(s+\varepsilon)} \leq \gamma_s(E) \leq CM_{t^s}(E)$$

(we need the second inequality). Indeed, by definition $\gamma_{s,+}(E) \leq \gamma_s(E)$, and $M_h(E)$, $\Lambda_h(E)$ vanish simultaneously. (We remark that for $0 < s < 1$, Prat [24] has obtained the following essentially stronger result: if $\gamma_s(E) > 0$ then $\Lambda_h(E) = \infty$.)

The second part is an immediate consequence of (11.4). \square

Obviously, there is a gap between the assumptions about h in the first and the second parts of Corollary 11.3. We claim that this gap cannot be reduced, that is, both parts are sharp. Concerning the first part it means that if $\liminf_{t \rightarrow 0} h(t)t^{-s} = 0$, then there is a compact set E for which $\gamma_{s,+}(E) > 0$ but $\Lambda_h(E) = 0$. This assertion follows from the more general and strong result [3], p. 34, Theorem 4: for any positive decreasing kernel $K(r)$ and any measuring function $h(r)$ such that

$$\liminf_{r \rightarrow 0} h(r) \overline{K}(r) = 0,$$

there is a Cantor type set E with $C_K(E) > 0$ and $\Lambda_h(E) = 0$. Here

$$\overline{K}(r) = \frac{1}{r^d} \int_0^r K(t)t^{d-1} dt,$$

$$C_K(E) := \sup \left\{ \|\mu\| : \mu \in M_+(E), \int_{\mathbb{R}^d} K(|x-y|) d\mu(y) \leq 1 \text{ on } E \right\}.$$

For $K(r) = r^{-s}$ we have $\overline{K}(r) = \frac{1}{d-s}r^{-s}$. By the maximum principle, $\gamma_{s,+}(E) \geq C \cdot C_K(E)$, and we get the needed assertion.

The second part of Corollary 11.3 is also precise: if the integral in (4.5) is divergent, then there exists a compact set E for which $\Lambda_h(E) > 0$ but $\gamma_{s,+}(E) = 0$. The industrious reader can obtain this claim from Section 8. The reader who does not care about conditions of regularity of h , can derive this statement under the additional condition $\frac{h(t)}{t^s} \nearrow$, using the estimate for the capacity γ_s of Cantor sets given at the end of [13].

The results of this section mentioned above generalize the corresponding results in [8], Section 12.

Remark. In Carleson's book [3] (see also [8], Sections 1,2 and the references therein) another condition similar to (4.5) plays an important part, namely

$$\int_0^\infty \frac{h(t)}{t^s} \frac{dt}{t} < \infty. \quad (11.5)$$

It is interesting that the difference between these two conditions is explained by the fact that we are dealing here with capacities with *signed* kernels, and (11.5) is pertinent to the classical capacities with the *positive* kernel $K(|x|) = |x|^{-s}$.

In conclusion we consider the relation between $\gamma_{s,+}(E)$ and the Riesz capacity $C_{\alpha,p}$ in non-linear potential theory. One of a number of equivalent definitions is the following equality (see [1], p. 34, Theorem 2.5.1):

$$C_{\alpha,p}(E) = \sup_{\mu \in M_+(E)} \left(\frac{\mu(E)}{\|I_\alpha * \mu\|_{p'}} \right)^p, \quad I_\alpha(x) = \frac{A_{d,\alpha}}{|x|^{d-\alpha}}, \quad \frac{1}{p'} + \frac{1}{p} = 1,$$

where $1 < p < \infty$, $0 < \alpha p \leq d$, $\|\cdot\|_{p'}$ is the $L^{p'}$ -norm with respect to the Lebesgue measure in \mathbb{R}^d , and $A_{d,\alpha}$ is the certain constant depending on d and α . It was proved in [12] that

$$\gamma_{s,+}(E) \asymp \gamma_s(E) \asymp C_{\frac{2}{3}(d-s), \frac{3}{2}}(E), \quad 0 < s < 1, \quad (11.6)$$

where

$$\gamma_s(E) := \sup |\langle T, 1 \rangle|,$$

and the supremum is taken over all distributions T supported by E such that $T * \frac{x}{|x|^{s+1}}$ is a

function in $L^\infty(\mathbb{R}^d)$ with $\left\| T * \frac{x}{|x|^{s+1}} \right\|_\infty \leq 1$. Theorem 4.7 implies that one of the inequalities between $\gamma_{s,+}(E)$ and $C_{\frac{2}{3}(d-s), \frac{3}{2}}(E)$ holds for wider range of s .

Corollary 11.4. For $0 < s < d$, one has

$$\gamma_{s,+}(E) \geq c \cdot C_{\frac{2}{3}(d-s), \frac{3}{2}}(E). \quad (11.7)$$

Proof. We may assume that $C_{\frac{2}{3}(d-s), \frac{3}{2}}(E) > 0$. As in [12], our proof is based on the following Wolff's equality [1], p. 110, Theorem 4.5.4: for any $\mu \in M_+(\mathbb{R}^d)$ and $1 < p < \infty$, $0 < \alpha p \leq d$,

$$\int_{\mathbb{R}^d} W_{\alpha,p}^\mu(x) d\mu(x) \asymp \|I_\alpha * \mu\|_{p'}^{p'}, \quad W_{\alpha,p}^\mu(x) := \int_0^\infty \left[\frac{\mu(B(x,r))}{r^{d-\alpha p}} \right]^{p'-1} \frac{dr}{r}. \quad (11.8)$$

Take $\alpha = \frac{2}{3}(d-s)$, $p = \frac{3}{2}$. Then $p' = 3$, $d - \alpha p = s$, and

$$W_{\alpha,p}^\mu(x) = \int_0^\infty \left[\frac{\mu(B(x,r))}{r^s} \right]^2 \frac{dr}{r} =: W^\mu(x).$$

Choose $\mu \in M_+(E)$ for which

$$C_{\alpha,p}(E) < 2\|\mu\|^p \|I_\alpha * \mu\|_{p'}^{-p}. \quad (11.9)$$

By (4.11) and (11.8) we have

$$\gamma_{s,+}(E) \geq c \|\mu\|^{3/2} \|I_\alpha * \mu\|_{p'}^{-p'/2} \stackrel{(11.9)}{>} \frac{c}{2} C_{\frac{2}{3}(d-s), \frac{3}{2}}(E),$$

and we get (11.7). \square

For integer $s \in (0, d)$ the opposite inequality $\gamma_{s,+}(E) \leq C \cdot C_{\frac{2}{3}(d-s), \frac{3}{2}}(E)$ is false. In fact, for a smooth s -dimensional manifold E in \mathbb{R}^d we have $\gamma_{s,+}(E) > 0$ by the obvious reason that natural Lebesgue measure on it gives bounded Riesz transform operator (this is from the classical Calderón-Zygmund theory). On the other hand, it has been noticed (for example in [12]) that any measure μ with finite Wolff's energy should have $\mu(B(x,r)) = o(r^s)$ for μ a. e. x . On a smooth s -dimensional manifold it can be only zero measure, so $C_{\frac{2}{3}(d-s), \frac{3}{2}}(E) = 0$. The question about validity of the inequality $\gamma_{s,+}(E) \leq C \cdot C_{\frac{2}{3}(d-s), \frac{3}{2}}(E)$ for all non-integer $s \in (0, d)$ remains open. We believe that this is the case.

REFERENCES

- [1] D. R. Adams, L. I. Hedberg, *Function spaces and potential theory*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **314**. Springer-Verlag, Berlin, 1996.
- [2] J. M. Anderson, V. Ya. Eiderman, *Cauchy transforms of point masses: the logarithmic derivative of the polynomial*, Ann. of Math., **163** (2005), No. 3, 1057–1076.
- [3] L. Carleson, *Selected problems on exceptional sets*, Van Nostrand, Princeton, NJ-Toronto, ON-London, 1967.
- [4] M. Christ, *A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral*, Colloquium Math., LX/LXI (1990), 601–628.
- [5] G. David, J. L. Journé, *A boundedness criterion for generalized Calderón-Zygmund operators*, Ann. of Math., **120** (1984), 371–397.
- [6] G. David, *Completely unrectifiable 1-sets on the plane have vanishing analytic capacity*, Revista Mat. Iberoamericana, **14** (1998), 369–479.
- [7] V. Ya. Eiderman, *Cartan-type estimates for the Cauchy Potential*, Dokl. Math., **73** (2006), No. 2, 273–276.
- [8] V. Ya. Eiderman, *Cartan type estimates for potentials with Cauchy kernel and with real-valued kernels*, Mat. Sb. **198**, no. 8 (2007), 115–160 (Russian). English transl. in Sb. Math. **198**(8) (2007), 1175–1220.
- [9] H. Farag, *The Riesz kernels do not give rise to higher-dimensional analogues of the Menger-Melnikov curvature*, Publ. Mat. **43**, no. 1 (1999), 251–260.
- [10] J. Garnett, L. Prat, X. Tolsa, *Lipschitz harmonic capacity and bilipschitz images of Cantor sets*, Math. Res. Lett., **13** (2006), No. 6, 865–884.
- [11] L. I. Hedberg, Th. H. Wolff, *Thin sets in nonlinear potential theory*, Ann. Inst. Fourier (Grenoble) **33** (1983), no. 4, 161–187.

- [12] J. Mateu, L. Prat, J. Verdera, *The capacity associated to signed Riesz kernels, and Wolff potentials*, J. Reine Angew. Math. **578** (2005), 201–223.
- [13] J. Mateu, X. Tolsa, *Riesz transforms and harmonic Lip_1 -capacity in Cantor sets*, Proc. London Math. Soc., **89** (2004), no. 3, 676–696.
- [14] J. Mateu, X. Tolsa, J. Verdera, *The planar Cantor sets of zero analytic capacity and local $T(b)$ theorem*, J. Amer. Math. Soc., **16** (2003), 19–28.
- [15] P. Mattila, *Geometry of sets and measures in Euclidean spaces*, Cambridge Stud. Adv. Math., vol. 44, Cambridge Univ. Press, Cambridge, 1995.
- [16] P. Mattila, P. V. Paramonov, *On geometric properties of harmonic Lip_1 -capacity*, Pacific J. Math., **171** (1995), no. 2, 469–491.
- [17] V. G. Maz'ya, V. P. Havin, *A nonlinear potential theory*, Uspehi Mat. Nauk, **27** (1972), no. 6, 67–138 (Russian). English transl.: Russian Math. Surveys, **27** (1972), no. 6, 71–148 (1974).
- [18] M. Melnikov, *Analytic capacity: discrete approach and curvature of the measure*, Mat. Sb. **186** (1995), 827–846.
- [19] M. Melnikov, J. Verdera, *A geometric proof of the L^2 boundedness of the Cauchy integral on Lipschitz graphs*, Intern. Math. Research Notices **IMRN**, 1995, No. 7, 325–331.
- [20] F. Nazarov, S. Treil, A. Volberg, *Cauchy integral and Calderon-Zygmund operators on non-homogeneous spaces*, Internat. Math. Res. Notices **IMRN**, 1997, No. 15, 703–726.
- [21] F. Nazarov, S. Treil, A. Volberg, *Weak type estimates and Cotlar inequalities for Calderón-Zygmund operators on non-homogeneous spaces*, Intern. Math. Research Notices **IMRN**, 1998, No. 9, 463–486.
- [22] F. Nazarov, S. Treil, A. Volberg, *The Tb -theorem on non-homogeneous spaces*, Acta Math., **190** (2003), no. 2, 151–239.
- [23] P. V. Paramonov, *On harmonic approximation in the C^1 -norm*, Math. Sb. **181** (1990), no. 10, 1341–1365 (in Russian); translation in Math. USSR-Sb. **71** (1992), no. 1, 183–207.
- [24] L. Prat, *Potential theory of signed Riesz kernels: capacity and Hausdorff measure*, Intern. Math. Research Notices **IMRN**, 2004, No. 19, 937–981.
- [25] X. Tolsa, *On the analytic capacity γ_+* , Indiana Univ. Math. J., **51** (2002), No. 2, 317–343.
- [26] X. Tolsa, *Painlevé's problem and the semiadditivity of analytic capacity*, Acta Math., **190** (2003), No. 1, 105–149.
- [27] X. Tolsa, *Riesz Transforms on Cantor Sets*, Manuscript.
- [28] A. Volberg, *Calderón-Zygmund capacities and operators on non-homogeneous spaces*. CBMS Regional Conference Series in Mathematics, **100**. AMS, Providence, RI, 2003.

VLADIMIR EIDERMAN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI
E-mail address: eiderman@ms.uky.edu

FEDOR NAZAROV, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI
E-mail address: nazarov@math.wisc.edu

ALEXANDER VOLBERG, DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY AND THE UNIVERSITY OF EDINBURGH
E-mail address: volberg@math.msu.edu, a.volberg@ed.ac.uk