Active Quantum Memory Using Oscillating Dark States

Jun He,* Yong-Sheng Zhang,[†] Xiang-Fa Zhou, Qun-Feng Chen, and Guang-Can Guo

Key Laboratory of Quantum Information, University of Science and Technology of China(CAS), Hefei, Anhui 230026, China

(Dated: January 26, 2023)

An active method for long time storage of quantum superposition state in atomic system using the Oscillating Dark States (ODS) is presented. Quantum state of a three-level Λ configuration atomic system oscillates periodically between two ground levels, when two pairs of classical detuning laser fields driving the system into the ODS under evolving adiabatic conditions. When considering another uploading/unloading adiabatic conditions and applying the oscillation of the ODS to quantum state storage, surprisingly, we can obtain the greatly suppressed decoherence of the system and high fidelity of the retrieved state, even if decay factor of coherence term of the system density matrix γ_{21} . $t\gg1$. The storage time is not limited by coherence decay time of the atomic system any longer, and can be thousands of times longer than that in those passive schemes without additional laser fields.

PACS numbers: 42.50.Gy, 32.80.Qk, 03.67.-a

Dark state or coherent population trapping (CPT), whose essential feature is linear superposition of distinct quantum states, has given rise to growing interests for past few decades, since its first discovery in 1976 [1]. If a three-level Λ atomic system is coherently trapped in dark states, there is no absorption of incident laser fields even in the presence of resonant transitions, due to destructive quantum interference between two transition pathways. Well-known dark state or CPT [2, 3, 4, 5, 6, 7, 8, 9] involves wide applications in many fields such as electromagnetically induced transparency (EIT) [10], lasing without inversion [11, 12, 13], the enhancement of the refractive index [14, 15], adiabatic population transfer [16, 17], subrecoil laser cooling [18], and atom interferometry [19]. Quantum information science, a new field that combines ideas from quantum mechanics and information theory, has led to many intriguing new concepts such as quantum teleportation, quantum cryptography, and quantum computation [20]. Coherent manipulation of a number of coupled quantum systems, i.e., quantum network, which consists of quantum memory elements storing quantum states information and interconnecting channels communicating with each other, is a generally difficult task, when we implement quantum information and computation process practically. One of the major difficulties is how to realize long time quantum superposition state storage, when scalable quantum network is required. The physics of 'slow light'propagation [8], and Duan, Lukin, Cirac, and Zoller (DLCZ) scheme [21] for quantum memory elements in an optically thick atomic ensembles are two main development aspects in quantum optical systems, which provide photons for ideal carriers of quantum information and individual atoms or atomic ensembles for reliable quantum memory elements. However, storage time is mainly limited by decoherence time of the system in those passive schemes [22].

Here we propose an active method for long time storage of quantum superposition state in a three-level Λ configuration atomic system, using the Oscillating Dark States (ODS), where probability amplitudes of the system state oscillate periodically between two ground levels. When applying the oscillation feature to quantum memory elements for quantum state storage, and considering evolving adiabatic conditions and uploading/unloading adiabatic conditions, decoherence of the system can be greatly suppressed, which may be a new way for decoherence control, and therefore long time storage of quantum state is realizable, in contrary to those passive schemes without additional laser fields.

The three-level Λ configuration atomic system is shown in Fig. 1. The ODS is implemented with two pairs of classical detuning laser fields coupling two lower ground levels $|1\rangle$, $|2\rangle$ to a single upper level $|3\rangle$, respectively. We assume that the $|1\rangle$ - $|2\rangle$ transition is always dipole-forbidden. The time-dependent interaction Hamiltonian in the interaction picture that describes the atom-laser coupling within the dipole and rotating wave approximation (RWA) in the rotating frame reads (\hbar =1),

$$H_{int} = \begin{bmatrix} 0 & 0 & P^* \\ 0 & \Delta' & Q^* \\ P & Q & \Delta \end{bmatrix}, \qquad (1)$$

where $P=-ie^{-i\phi_{12}}\Omega_{12}\sin(\frac{\Delta_1-\Delta_2}{2}t)$, $Q=-e^{-i\phi_{34}}\Omega_{34}\cos(\frac{\Delta_3-\Delta_4}{2}t)$, $\Delta=\frac{1}{2}(\Delta_1 + \Delta_2)$, and $\Delta'=\frac{1}{2}[(\Delta_1 + \Delta_2) - (\Delta_3 + \Delta_4)]$. $\Delta_i=\omega_{31}-\omega_i(i=1,2)$, and $\Delta_j=\omega_{32}-\omega_j(j=3,4)$ are the detunings of two pairs of laser angular frequencies $\omega_i(i=1,2)$ and $\omega_j(j=3,4)$ from the corresponding atomic transitions ω_{31} and ω_{32} . Real Ω_i and ϕ_i (*i*=1, 2; 3, 4) are corresponding Rabi frequencies and phases of two pairs of laser fields, and we assume $\Omega_1=\Omega_2=\Omega_{12}$, $\Omega_3=\Omega_4=\Omega_{34}$, and $\phi_1=\phi_2+\pi=\phi_{12}$, $\phi_3=\phi_4=\phi_{34}$, with the difference $\Delta\phi=\phi_{12}-\phi_{34}$. The eigenstates of the interaction Hamiltonian (1), which can be generally described in terms of the "mixing angles" θ and φ , are given as

$$|a_{+}\rangle = -ie^{i\Delta\phi}\sin\theta\sin\varphi|1\rangle + \cos\theta\sin\varphi|2\rangle - \cos\varphi e^{-i\phi_{34}}|3\rangle, \quad (2)$$

$$|a_0\rangle = \cos\theta |1\rangle - i\sin\theta e^{-i\Delta\phi} |2\rangle, \qquad (3)$$

$$|a_{-}\rangle = -ie^{i\Delta\phi}\sin\theta\cos\varphi|1\rangle + \cos\theta\cos\varphi|2\rangle + \sin\varphi e^{-i\phi_{34}}|3\rangle,$$
(4)



FIG. 1: Two pairs of classical laser fields Ω_1 , Ω_2 and Ω_3 , Ω_4 couple two lower ground levels $|1\rangle$, $|2\rangle$ to a single upper level $|3\rangle$, respectively, with their corresponding detunings Δ_1 , Δ_2 and Δ_3 , Δ_4 , and their fixed phases: $\phi_1 = \phi_2 + \pi$, $\phi_3 = \phi_4$, in a three-level Λ configuration atomic system.

where the mixing angles that are dependent upon time *t* and Rabi frequencies are written, when we meet the condition $\Delta' = 0$, i.e. $(\Delta_1 + \Delta_2) = (\Delta_3 + \Delta_4)$, as

$$\tan \theta = \frac{\Omega_{12} \sin(\frac{\Delta_1 - \Delta_2}{2}t)}{\Omega_{34} \cos(\frac{\Delta_3 - \Delta_4}{2}t)},$$
(5)

$$\tan 2\varphi = \frac{2\sqrt{|P|^2 + |Q|^2}}{\Delta} . \tag{6}$$

The eigenstate $|a_0\rangle$ without the contribution from $|3\rangle$, the ODS, corresponds the null eigenvalue of interaction Hamiltonian (1). If a system is in state $|a_0\rangle$, there is no possibility of excitation to $|3\rangle$ and subsequent spontaneous emission, thus the atom is effectively decoupled from laser fields. The eigenvalues of the pair of states $|a_{\pm}\rangle$ which contain a component of all three bare atomic states, are shifted up and down by an amount λ_{\pm} ,

$$\lambda_{\pm} = \frac{1}{2} \left[\Delta \pm \sqrt{\Delta^2 + 4|P|^2 + 4|Q|^2} \right] \,. \tag{7}$$

The probability amplitudes of the system state can oscillate periodically between two ground levels $|1\rangle$ and $|2\rangle$ upon time *t*, when the system is driven into the ODS by laser fields with relatively fixed detunings and phases of four laser fields.

Evolving adiabatic conditions, under which the system avoids diabatic coupling to bright states and is always in some dark states if it is initially in a certain one, is considered, since the ODS follows the evolution of a time-dependent Hamiltonian (1). The change rate of the mixing angle θ must be small compared with the separation of the corresponding eigenvalues [23, 24],

$$\left|\frac{d}{dt}\theta\right|^2 \ll \left|\lambda_0 - \lambda_{\pm}\right|^2\,,\tag{8}$$

Let $\Omega_{12}=\Omega_{34}=\Omega$ (fixing Rabi frequencies when the system evolves) and $\Delta_1-\Delta_2=\Delta_3-\Delta_4=2\delta$, thus we can obtain evolving adiabatic conditions according to (8) as

$$|\delta|^2 \ll \frac{1}{4} |\Delta \pm \sqrt{\Delta^2 + 4\Omega^2}|^2 . \tag{9}$$

which change into $|\delta| \ll \Omega$, when $\Delta = 0$, i.e. $\Delta_1 = -\Delta_2$ and $\Delta_3 = -\Delta_4$. Evolving adiabatic conditions for the ODS ensure that the values of all parameters can be established in advance, because the evolution process is automated by laser-atomic system itself, in contrast to ordinary dark states [5, 17], where we change relative values of two Rabi frequencies to attain different dark states, and adiabatic evolution can be controlled by experimental instruments.

A process of quantum superposition state storage using the ODS is as follows: first, an arbitrary initial state (3) with a fixed $\Delta\phi$, using two bare ground states of the atomic system as two orthogonal basis states, which constitutes initial condition of the system evolution, is prepared. Then we concurrently and rapidly upload two pairs of laser fields as shown in Fig. 1, and the system periodically evolves subsequently. *N* (*N*=1, 2, 3, ..., natural number) oscillation periods later, rapid unloading all laser fields simultaneously retrieves final system state that is almost the same as the initial one. The total storage time *t* depends on the product of oscillation times *N* and its periods $T(= 2\pi/\delta)$.

When uploading/unloading two pairs of laser fields, uploading/unloading adiabatic conditions are necessary besides evolving adiabatic conditions (9), in order to keep the least diabatic coupling between bright states and dark states of the system. We assume a concurrent increase/decrease of four laser fields in a short time $\tau(\tau \ll T)$, i.e. $\Omega_1 = \Omega_2 = \Omega_3 = \Omega_4 = \Omega$ and $\frac{d}{dt}\Omega_1 = \frac{d}{dt}\Omega_2 = \frac{d}{dt}\Omega_3 = \frac{d}{dt}\Omega_4$. Thus, we can obtain the same result as evolving adiabatic conditions in (9) for uploading/unloading adiabatic conditions, according to (8).

Time-dependent interaction Hamiltonian in interaction picture that describes atom-laser coupling can be written as

$$H'_{int} = -\frac{1}{2} [\Omega_1 e^{-i\phi_1} e^{i\Delta_1 t} \sigma_{31} + \Omega_2 e^{-i\phi_2} e^{i\Delta_2 t} \sigma_{31} + \Omega_3 e^{-i\phi_3} e^{i\Delta_3 t} \sigma_{32} + \Omega_4 e^{-i\phi_4} e^{i\Delta_4 t} \sigma_{32} + H.c.], \qquad (10)$$

where $\sigma_{ij} = |i\rangle\langle j|$ (*i*, *j*=1, 2, 3) are atomic projection operators. A master equation [9, 25], using density matrix ρ ,

$$\begin{aligned} \frac{d\rho}{dt} &= -i[H_{int}',\rho] + \frac{\Gamma_{31}}{2} [2\sigma_{13}\rho\sigma_{31} - \sigma_{33}\rho - \rho\sigma_{33}] \\ &+ \frac{\Gamma_{32}}{2} [2\sigma_{23}\rho\sigma_{32} - \sigma_{33}\rho - \rho\sigma_{33}] \\ &+ \frac{\gamma_{3deph}}{2} [2\sigma_{33}\rho\sigma_{33} - \sigma_{33}\rho - \rho\sigma_{33}] \\ &+ \frac{\gamma_{2deph}}{2} [2\sigma_{22}\rho\sigma_{22} - \sigma_{22}\rho - \rho\sigma_{22}] \\ &+ \frac{\Gamma_{21}}{2} [2\sigma_{12}\rho\sigma_{21} - \sigma_{22}\rho - \rho\sigma_{22}] , \end{aligned}$$
(11)

is required, because the decoherence of a practical atomic system must be considered, when using the ODS for long time state storage. Γ_{31} and Γ_{32} indicate rates of spontaneous emission from level $|3\rangle$ to $|1\rangle$ and $|2\rangle$, respectively, and rates γ_{2deph} and γ_{3deph} describe energy-conserving dephasing processes. Γ_{21} that is longitudinal relaxation from level $|2\rangle$ to $|1\rangle$, is mainly due to higher-order transition, and is much small



FIG. 2: Probability amplitudes evolution of an arbitrary initial superposition state finds expression in the time-dependent populations of three levels ρ_{11} , ρ_{22} , ρ_{33} , and coherence term between two ground levels $|\rho_{21}|^2$ with $\Delta_1 = \Delta_3 = 0.3$, $\Delta_2 = \Delta_4 = 0.2$, $\gamma_{2deph} = 0.02$, $\Gamma_{21} = 0.002$, for different case: (a). $\Omega = 2$, (b). $\Omega = 0.2$, (c). $\Omega = 0.08$, (d). $\Omega = 0$, respectively. All parameters are in the units of γ_{31} , and total evolution time t=4T is marked.

compared with γ_{2deph} . We can define coherence decay rates as $\gamma_{31}=\Gamma_{31}+\Gamma_{32}+\gamma_{3deph}$, and $\gamma_{21}=\Gamma_{21}+\gamma_{2deph}$, and estimate Γ_{21} at 1/10 of γ_{2deph} in the following analysis.

The evolution stage in storage process is stimulated numerically as shown in Fig. 2. We give an arbitrary initial superposition state (3) for storage (random $\Delta \phi$, and t_0 for initial θ , but in fact $\Delta \phi$ should be predicted in practical experiments), and analyze its probability amplitudes evolution with different Rabi frequencies Ω . The populations of two ground lower levels, ρ_{11} and ρ_{22} , oscillate periodically after applying interaction Hamiltonian (10), and the population of upper level ρ_{33} stays almost zero in total evolution stage, when evolving adiabatic conditions (9) are well satisfied for case (a) in Fig. 2. Especially and surprisingly, the same periodical oscillation of the coherence term $|\rho_{21}|^2$ between two ground levels as ρ_{11} and ρ_{22} , and $|\rho_{21}|^2 \approx \rho_{11} \cdot \rho_{22}$, imply a coherent evolution of quantum superposition states, i.e. the ODS. Any arbitrary initial superposition state in more than thousands of oscillation periods gives the same periodical evolution process as case (a) under evolving adiabatic conditions. For case (b), Rabi frequency Ω is decreased, and evolving adiabatic conditions are not well met, which resulting in a slight occupation of population ρ_{33} , but the oscillations of population ρ_{11} , ρ_{22} and of coherence term $|\rho_{21}|^2$ still dominate whole evolution process. When we decrease Ω further, laser fields are too weak to make upper level $|3\rangle$ occupied, and a decoherence process appears for case (c), despite the violation of evolving adiabatic conditions. A complete decoherence process of the system emerges when Ω becomes to be 0 for case (d), which is obvious without laser fields, and the initial superposition state becomes a mixed one on the time scale $\sim 1/\gamma_{21}$, since $|\rho_{21}|^2$ decays to 0 absolutely.

From case (d) to (a), the value of Rabi frequencie Ω of laser fields increased little by little, it can be seen that coherent reconstruction mechanism of laser fields becomes superior gradually to the decoherence effect of the the system for case (d), and entirely dominates periodical oscillation of the system for case (a). Under evolving adiabatic conditions, it is unreasonable that, the system state does not follow the evolution of time-dependent hamiltonian if decoherence effect changes the state to the one that is not the instantaneous eigenstate of the system. Long time state storage using a coherent evolution process of the ODS for case (a), where upper level $|3\rangle$ population is little and the oscillation of probability amplitudes of two ground levels $|1\rangle$ and $|2\rangle$ is periodical, is reliable. Total state storage time t as long as t is an integral multiple of oscillation period T, after which we can retrieve almost the same initial system state, is largely increased compared with that without laser fields for case (d), since the decoherence between two ground levels $|1\rangle$ and $|2\rangle$ can be greatly suppressed, owing to coherent reconstruction contribution from laser fields.

Without contribution of applied laser fields, an initial superposition state will exponentially decay to a mixed one with a factor $\gamma_{21} \cdot t$, expressed in coherence term of the system density matrix $|\rho_{21}|^2$. When applying two pairs of laser fields, the dependence of average fidelity of initial storage and final retrieved state on total evolution time t is analyzed in Fig. 3, with ten arbitrary initial states chosen randomly as before. γ_{21} is fixed at 0.022 in the units of γ_{31} (reference to D_1 line of ${}^{87}Rubidium$ with $\gamma_{31} \sim 36.10 \times 10^6 s^{-1}$ and estimated $\gamma_{2deph}=0.02\gamma_{31}$), and different total evolution times t that should be integral multiples of oscillation period T, is given. It can be seen that, high average fidelity above 0.95 of ten arbitrary initial states is surprisingly available when total evolution time t is up to thousands of multiples of oscillation period T (N~1000), where $\gamma_{21} \cdot t \gg 1$, as well as each one of ten. Fidelity F decreases slightly in the first one T, during which a certain complicated amplitude decay of the ODS experiences, and stays unchangeably after it, which also can be found in Fig. 2. The long time stability of F implies a pure periodical evolution of the system. In those passive schemes without additional laser fields [8, 21, 22], the storage time is mainly limited by coherence decay time of atomic system scaled $\sim 1/\gamma_{21}$, within which coherence of photons can be controlled and preserved in atomic system. Using the ODS, the storage time is not limited by coherence decay time of the atomic system any longer, and can be thousands of times longer than that in those passive schemes, since coherence of the atomic system can be preserved in laser fields, which is significant for quantum state storage. High fidelity F mainly depends on relatively large value of Ω and always can be obtained by controlling laser fields.

The stability of frequency difference 2δ , and π phase difference and $\Delta\phi$ of each pair of laser fields determine mainly the precision of oscillation process. A concurrent uploading/unloading time $\tau \sim 10^{-8} s$ (a reasonable 0.01*T* in our anal-



FIG. 3: Dependence of average fidelity of ten arbitrary initial storage and their respective final retrieved states on total evolution time of the system, with $\Omega = 2$, $\Delta_1 = \Delta_3 = 0.3$, $\Delta_2 = \Delta_4 = 0.2$, $\gamma_{21} = 0.022$, and independent variable *t* (integral multiples of oscillation period *T*, up to a maximum 1000*T* here). After the first one *T* (see inset), High fidelity above 0.95 is available and keep stable at all times, even if $\gamma_{21} \cdot t \gg 1$.

ysis which is short enough to avoid a certain decay of ρ at this stage) can be carried out using an optical switch. Laser fields may not be monochromatic, but the stability of relative phases of two pairs of laser fields can also guarantee almost the same result, despite a slow collective phase drift. A chosen δ whose value is larger than the linewidth of laser fields can avoid undesirable interference of laser fields and optical pumping. The state storage using the ODS is immune to Doppler frequency shift and concurrent fluctuation of Rabi frequencies Ω_i (*i*=1, 2; 3, 4). Well developed laser techniques can meet all above demands to obtain a perfect long time quantum state storage. It is interesting to compare this scheme that can be regarded as a certain temporal storage of atomic state by laser fields with the spatial storage of light in atomic system in those passive ones. The ODS can be implemented in a three-level atomic system by using at least three laser fields in an appropriate configuration, but we use two pairs of laser fields in a symmetric scheme, in order to obtain complete oscillation of probability amplitudes between two ground levels. The ODS is similar to the famous Rabi resonance in a two-level atomic system driven by one resonant laser field [5], where we can not find the same interesting result.

In conclusion, we propose an active method using the ODS for long time quantum superposition storage, where coherence of the atomic system can be preserved temporally by laser fields, compared with those passive ones without additional laser fields, where preserved coherence of photons keeps spatially in atomic system within its coherence time, and decoherence time of the system limits storage time. High fidelity, greatly suppressed decoherence, and long storage time, make the ODS a potential method for quantum information storage in the fields of quantum information and quantum computation. The ODS can also involves many other significant applications to nonlinear optics, quantum optics, optics and laser spectroscopy, and atomic physics. This work was funded by the National Fundamental Research Program (Grant No. 2006CB921907), the National Natural Science Foundation of China (Grant No. 10674127 and No. 60621064), the Innovation Funds from the Chinese Academy of Sciences, Program for NCET and International cooperation program from CAS.

- * Electronic address: jhe@mail.ustc.edu.cn
- [†] Electronic address: yshzhang@ustc.edu.cn
- G. Alzetta, A. Gozzini, L. Moi, and G. Orriols, Nuovo Cimento B 36, 5 (1976).
- [2] E. Arimondo and G. Orriols, Nuovo Cimento Lett. 17, 333 (1976).
- [3] H. R. Gray, R. M. Whitley, and C. R. Stroud, Jr., Opt. Lett. 3, 218 (1978).
- [4] E. Arimondo, Progress in Optics XXXV, (edited by E. Wolf, North-Holland, Amsterdam, 1996), pp.259-354.
- [5] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 1997).
- [6] K. Bergmann, H. Theuer, and B. W. Shore, Rev. Mod. Phys. 70, 1003 (1998).
- [7] M. D. Lukin, S. F. Yelin, M. Fleischhauer, and M. O. Scully, Phys. Rev. A 60, 3225 (1999).
- [8] M. Fleischhauer and M. D. Lukin, Phys. Rev. Lett. 84, 5094 (2000).
- [9] M. Fleischhauer, A. Imamoglu, and J. P. Marangos, Rev. Mod. Phys. 77, 633 (2005).
- [10] S. E. Harris, Phys. Today **50**(7), 36 (1997).
- [11] S. E. Harris, Phys. Rev. Lett. 62, 1033 (1989).
- [12] M. O. Scully, S. -Y. Zhu, and A. Gavrielides, Phys. Rev. Lett. 62, 2813 (1989).
- [13] A. S. Zibrov, M. D. Lukin, D. E. Nikonov, L. Hollberg, M. O. Scully, V. L. Velichansky, and H. G. Robinson, Phys. Rev. Lett. 75, 1499 (1995).
- [14] M. O. Scully, Phys. Rev. Lett. 67, 1855 (1991).
- [15] A. S. Zibrov, M. D. Lukin, L. Hollberg, D. E. Nikonov, M. O. Scully, H. G. Robinson, and V. L. Velichansky, Phys. Rev. Lett. 76, 3935 (1996).
- [16] J. Oreg, F. T. Hioe, and J. H. Eberly, Phys. Rev. A 29, 690 (1984).
- [17] J. R. Kuklinski, U. Gaubatz, F. T. Hioe, and K. Bergmann, Phys. Rev. A 40, 6741(R) (1989).
- [18] F. Bardou, J. P. Bouchaud, O. Emile, A. Aspect, and C. Cohen-Tannoudji, Phys. Rev. Lett. 72, 203 (1994).
- [19] B. Young, M. Kasevich, and S. Chu, *Atom Interferometry*, (edited by P. Berman, Academic Press, New York, 1997), pp.363.
- [20] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [21] L. -M. Duan, M. D. Lukin, J. I. Cirac, and P. Zoller, Nature 414, 413 (2001).
- [22] T. Chanelière, D. N. Matsukevich, S. D. Jenkins, S.-Y. Lan, T. A. B. Kennedy, and A. Kuzmich, Nature 438, 833 (2005).
- [23] A. Messiah, *Quantum Mechanics* (North-Holland, Amsterdam, 1962), Vol.II, pp.744.
- [24] U. Gaubatz, P. Rudecki, S. Schiemann, and K. Bergmann, J. Chem. Phys. 92, 5363 (1990).
- [25] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon Interactions* (Wiely, New York, 1992).