

FINITENESS OF MAPPING DEGREES AND $\widetilde{\text{PSL}}(2, \mathbf{R})$ -VOLUME ON GRAPH MANIFOLDS

PIERRE DERBEZ AND SHICHENG WANG

ABSTRACT. For given closed orientable 3-manifolds M and N let $\mathcal{D}(M, N)$ be the set of mapping degrees from M to N . We address the problem: For which N , $\mathcal{D}(M, N)$ is finite for all M ? The answer is known in Thurston's picture of closed orientable irreducible 3-manifolds unless the target is a non-trivial graph manifold. We prove that for each closed non-trivial graph manifold N , $\mathcal{D}(M, N)$ is finite for all graph manifold M .

The proof uses a recently developed standard forms of maps between graph manifolds and the estimation of the $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume for certain class of graph manifolds.

CONTENTS

1. Introduction	2
2. Notations and known results	3
2.1. Coordinated graph manifolds and gluing matrices	4
2.2. Canonical framings and canonical submanifold	4
2.3. Standard forms of nonzero degree maps	5
2.4. $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume, absolute volume, and absolute Euler number	5
3. Reduction of complexity via coverings	6
3.1. Two general statements	6
3.2. Separable and characteristic coverings	7
3.3. Lifting of coordinates and gluing matrices	8
3.4. The absolute volume and the absolute Euler number	9
4. $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume of graph manifolds	11
5. Proof of Theorem 1.1	16
5.1. Simplifications	16
5.2. Proof of Theorem 1.1 when $ e (N) \neq 0$.	16
5.3. Proof of Theorem 1.1 when $ e (N) = 0$.	17
References	18

Date: November 16, 2018.

1991 Mathematics Subject Classification. 57M50, 51H20.

Key words and phrases. Graph manifold, non-zero degree maps, Volume of a representation.

1. INTRODUCTION

Let M and N be two closed oriented 3-dimensional manifolds. Let $\mathcal{D}(M, N)$ be the set of degree of maps from M to N , that is

$$\mathcal{D}(M, N) = \{d \in \mathbf{Z} \mid f: M \rightarrow N, \deg(f) = d\}.$$

According to [CT], M. Gromov think it is a fundamental problem in topology to determine the set $\mathcal{D}(M, N)$. A basic property of $\mathcal{D}(M, N)$ is reflected in the following

Question 1. (see also [Re2, Problem A] and [W2, Question 1.3]): For which closed orientable 3-manifold N , the set $\mathcal{D}(M, N)$ is finite for all closed orientable 3-manifolds M ?

This answer is known in Thurston's picture of closed orientable irreducible 3-manifolds unless the target is a non-trivial graph manifold. We are going to explain this.

Thurston's geometrization conjecture states that a closed orientable irreducible 3-manifold N has a unique family of JSJ-tori \mathcal{T}_N such that $N \setminus \mathcal{T}_N$ consists of hyperbolic pieces and Seifert pieces ([Th2], [JS], and [Jo]). A closed orientable, irreducible 3-manifold is called *geometrizable* if it satisfies Thurston's geometrization conjecture. Call a geometrizable 3-manifold N a graph manifold if all components in $N \setminus \mathcal{T}_N$ are Seifert pieces. Call a graph manifold N is *non-trivial* if it is not covered by either a Seifert manifold or a torus bundle.

Let ω be a non-zero and non-negative 3-manifold invariant. Say ω has *degree property* or simply *Property D*, if for any map $f: M \rightarrow N$, then $\omega(M) \geq |\deg(f)|\omega(N)$. Say ω has *covering property* or simply *Property C*, if for any covering $p: M \rightarrow N$, $\omega(M) = |\deg(p)|\omega(N)$. The invariants with Property *D* are important to study Question 1 due to the following fact (see Lemma 3.1):

Fact (*) *If ω has property D and if N admits a finite covering \tilde{N} such that $\omega(\tilde{N}) \neq 0$ then the set $\mathcal{D}(M, N)$ is finite for all M .*

Two 3-manifold invariants having Property *D* are known.

(1) Simplicial volume $\| \! \| \! \|$ defined in [G]: For geometrizable 3-manifolds, $\|N\|$ is proportional to the sum of the hyperbolic volumes of the hyperbolic pieces in $N \setminus \mathcal{T}_N$, and moreover $\| \! \| \! \|$ has Property C. ([G], [Th1], [So]).

(2) $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume SV introduced in [BG1] and [BG2]: It is non-zero for Seifert manifolds supporting $\widetilde{\text{PSL}}(2, \mathbf{R})$ geometry.

With those two invariants, it follows that for any geometrizable 3-manifold N if either $N \setminus \mathcal{T}_N$ has a hyperbolic piece or N supports the $\widetilde{\text{PSL}}(2, \mathbf{R})$ geometry, then $\mathcal{D}(M, N)$ is finite for any M . On the other hand, suppose N is either covered by a torus bundle, or a Seifert manifold not supporting the $\widetilde{\text{PSL}}(2, \mathbf{R})$ geometry, it is known that $\mathcal{D}(N, N)$ is infinite. Hence for the geometrizable 3-manifolds, Question 1 is reduced to

Question 2. Suppose N is a non-trivial graph manifold, is $\mathcal{D}(M, N)$ finite for all closed orientable 3-manifolds M ?

It is clear that the simplicial volume is vanishing on graph manifolds. Based on Fact (*), it is natural to ask

Question 3. Let N be a closed orientable non-trivial graph manifold. Does $SV(\tilde{N}) \neq 0$ for some finite covering \tilde{N} of N ?

The $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume is rather strange and was very little known. It can be either zero or non-zero for hyperbolic 3-manifolds [BG1]; if it has Property C is still unclear, and it was not addressed for 3-manifolds with non-trivial JSJ-decomposition since it was introduced more than 20 years ago.

One main result of this paper is a partial answer of Question 3: we verify that for non-trivial graph manifolds N with "simplest" gluing, they do have finite cover \tilde{N} with $SV(\tilde{N}) \neq 0$ (Proposition 4.1), and it follows that Question 2 has positive answer for each non-trivial closed graph manifold whose dual graph has two vertices and one edge. Such a partial answer, combined with the standard form of nonzero degree maps developed in [D1], enable us to solve Question 2 when we restrict on graph manifolds.

Theorem 1.1. *For any given closed non-trivial graph manifold N , $\mathcal{D}(M, N)$ is finite for any graph manifold M .*

Remark 1.2. Some facts related to Theorem 1.1 are known before: $\mathcal{D}(N, N)$ is finite for any non-trivial graph manifold N ([W1], see also [D2]). The covering degrees is uniquely determined by the graph manifolds involved [YW].

This paper is organized as follows.

In Section 2 we define the objects which will be used in the paper: For graph manifolds, we will define their coordinates and gluing matrices, canonical framings, the standard forms of nonzero degree maps, the absolute Euler number and the absolute volume. We also recall $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume and its basic properties.

In Section 3 we state and prove some results on coverings of graph manifolds which will be used in the paper.

Section 4 is devoted to the proof of Proposition 4.1. The strategy is to use a finite sequence of coverings to get a very "large" and "symmetric" covering space which allows some free action of a finite cyclic group so that the quotient can be sent onto a 3-manifold supporting $\widetilde{\text{PSL}}(2, \mathbf{R})$ geometry via a nonzero degree map.

In Section 5 we prove Theorem 1.1. The strategy is to use the standard form of nonzero degree maps between graph manifolds to show that one can reduce the problem to the case where the target is a graph manifold satisfying the hypothesis of Proposition 4.1.

2. NOTATIONS AND KNOWN RESULTS

From now on all 3-manifolds are irreducible and oriented, and all graph manifolds are non-trivial.

Suppose F (resp. P) is a properly embedded surface (resp. an embedded 3-manifold) in a 3-manifold M . We use $M \setminus F$ (resp. $M \setminus P$) to denote the resulting manifold obtained by splitting M along F (resp. removing $\text{int}P$, the interior of P).

2.1. Coordinated graph manifolds and gluing matrices. Let N be a graph manifold. Denote by \mathcal{T}_N the family of JSJ tori of N , by N^* the set $N \setminus \mathcal{T}_N = \{\Sigma_1, \dots, \Sigma_n\}$ of the JSJ pieces of N , by $\tau: \partial N^* \rightarrow \partial N^*$ the associated sewing involution defined in [JS].

A *dual graph* of N , denoted by Γ_N , is given as follows: each vertex represents a JSJ piece of N ; each edge represents a JSJ torus of N ; an edge e connects two vertices v_1 and v_2 (may be $v_1 = v_2$) if and only if the corresponding JSJ torus is shared by the corresponding JSJ pieces.

Call a dual graph Γ_N *directed* if each edge of Γ_N is directed, in other words, is endowed with an arrow. Once Γ_N is directed, the sewing involution τ becomes a well defined map, still denoted by $\tau: \partial N^* \rightarrow \partial N^*$.

Suppose N^* contains no pieces homeomorphic to $I(K)$, the twisted I -bundle over the Klein bottle.

Let Σ be an oriented Seifert manifold which admits a unique Seifert fibration, up to isotopy, and $\partial\Sigma \neq \emptyset$. Denote by h the homotopy class of the regular fiber of Σ , by \mathcal{O} the base 2-orbifold of Σ and by Σ^0 the space obtained from Σ after removing the singular fibers of Σ . Then Σ^0 is a \mathbf{S}^1 -bundle over a surface \mathcal{O}^0 obtained from \mathcal{O} after removing the exceptional points. Then there exists a cross section $s: \mathcal{O}^0 \rightarrow \Sigma^0$. Call Σ is *coordinated*, if

- (1) such a section $s: \mathcal{O}^0 \rightarrow \Sigma^0$ is chosen,
- (2) both $\partial\mathcal{O}^0$ and h are oriented so that their product orientation is matched with the orientation of $\partial\Sigma$ induced by that of Σ .

Once Σ is coordinated, then the orientation on $\partial\mathcal{O}^0$ and the oriented fiber h gives a basis of $H_1(T; \mathbf{Z})$ for each component T of $\partial\Sigma$. We also say that Σ is endowed with a (s, h) -basis.

Since N^* has no $I(K)$ -components then each component Σ_i of N^* admits a unique Seifert fibration, up to isotopy. Moreover each component Σ_i has the orientation induced from N . Call N is *coordinated*, if each component Σ_i of N^* is coordinated and Γ_N is directed.

Once N is coordinated, then each torus T in \mathcal{T}_N is associated with a unique 2×2 -matrix A_T provided by the gluing map $\tau|: T_-(s_-, h_-) \rightarrow T_+(s_+, h_+)$: where T_-, T_+ are two torus components in ∂N^* provided by T , with basis (s_-, h_-) and (s_+, h_+) respectively, and

$$\tau(s_-, h_-) = (s_+, h_+)A_T.$$

Call $\{A_T, T \in \mathcal{T}\}$ the *gluing matrices*.

2.2. Canonical framings and canonical submanifold. Let Σ denote an orientable Seifert manifold with regular fiber h . A *framing* α of Σ is to assign a simple closed essential curve not homotopic to the regular fiber of Σ , for each component T of $\partial\Sigma$. Denote by $\Sigma(\alpha)$ the closed Seifert 3-manifold obtained from Σ after Dehn fillings along the family α and denote by $\pi_\Sigma: \Sigma \rightarrow \Sigma(\alpha)$ the natural quotient map. Let $p: \tilde{\Sigma} \rightarrow \Sigma$ be a finite covering. Assume that Σ and $\tilde{\Sigma}$ are endowed with a framing α and $\tilde{\alpha}$. Then we say that $(\tilde{\Sigma}, \tilde{\alpha})$ covers (Σ, α) if each component of $\tilde{\alpha}$ is a component of $p^{-1}(\alpha)$. In this case, the map $p: (\tilde{\Sigma}, \tilde{\alpha}) \rightarrow (\Sigma, \alpha)$ extends

to a map $\hat{p}: \hat{\Sigma}(\hat{\alpha}) \rightarrow \Sigma(\alpha)$ and the Euler number of $\Sigma(\alpha)$ is nonzero iff the Euler number of $\hat{\Sigma}(\hat{\alpha})$ is nonzero by [LW]. When N contains no $I(K)$ -component in its JSJ-decomposition, each Seifert piece Σ of N^* is endowed with a *canonical framing* α_Σ given by the regular fiber of the Seifert pieces of N^* adjacent to Σ . Denote by $\hat{\Sigma}$ the space $\Sigma(\alpha_\Sigma)$. By minimality of JSJ decomposition, $\hat{\Sigma}$ admits a unique Seifert fibration extending that of Σ .

Call a submanifold L of a graph manifold N is *canonical* if L is a union of some components of $N \setminus \mathcal{T}$, where \mathcal{T} is subfamily of \mathcal{T}_N . Similarly call $\alpha_L = \{t_U \subset U\}$ where t_U is the regular fiber of the Seifert piece adjacent to L along the component U , when U runs over the components of ∂L , the canonical framing of L , and denote by \hat{L} the closed graph 3-manifold obtained from L after Dehn fillings along the family α_L . From the definition we have

Lemma 2.1. *For a given closed graph manifold M , there are only finitely many canonical framed canonical submanifolds (L, α_L) , and thus only finitely many \hat{L} .*

2.3. Standard forms of nonzero degree maps. We recall here two results which are proved in [D1] in a more general case. The first result is related to the standard forms of nonzero degree maps.

Proposition 2.2. ([D1, Lemma 3.4]) *For a given closed graph manifold M , there is a finite set $\mathcal{H} = \{M_1, \dots, M_k\}$ of closed graph manifolds satisfying the following property: for any nonzero degree map $g: M \rightarrow N$ into a closed non-trivial graph manifold N without $I(K)$ piece in N^* , there exists some M_i in \mathcal{H} and a map $f: M_i \rightarrow N$ such that:*

- (i) $\deg(f) = \deg(g)$,
- (ii) for each piece Q in N^* , $f^{-1}(Q)$ is a canonical submanifold of M .

The following technical "mapping lemma" will be also useful:

Lemma 2.3. ([D1, Lemma 4.3]) *Suppose $f: M \rightarrow N$ is a map between closed graph manifolds and N^* contains no $I(K)$ piece. Let S and S' be two components of M^* which are adjacent in M along a subfamily \mathcal{T} of \mathcal{T}_M and satisfy:*

- (i) $f(S') \subset \text{int}(\Sigma')$ for some piece Σ' of N^* ,
- (ii) $f_*([h_S]) \neq 1$, where t_S is the regular fiber of S .

Then there exists a piece Σ of N^ and a homotopy of f supported in a regular neighborhood of S such that $f(S) \subset \text{int}(\Sigma)$. Moreover if $f(h_S)$ is not homotopic to a non-trivial power of the regular fiber of Σ , then one can choose $\Sigma = \Sigma'$.*

[D1, Lemma 4.3] was stated for Haken manifolds. Since here we consider only non-trivial graph manifolds instead of Haken manifolds, then we can state [D1, Lemma 4.3] in term of the JSJ-pieces of N instead of in term of the characteristic Seifert pair of N .

2.4. $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume, absolute volume, and absolute Euler number. $\widetilde{\text{PSL}}(2, \mathbf{R})$ -volume SV is introduced in [BG1] and [BG2]. (It is also considered as a special case of volumes of representations, see [Re1], [Re2] and [WZ]). Two basic properties of SV are reflected in the following

Lemma 2.4. (i) *SV has Property D [BG1],*

(ii) *If N supports $\widetilde{\mathrm{PSL}}(2, \mathbf{R})$ geometry, i.e., N admits a Seifert fibration with nonzero Euler number $e(N)$ and whose base 2-orbifold \mathcal{O}_N has a negative Euler characteristic, then [BG2]*

$$SV(N) = \left| \frac{\chi^2(\mathcal{O}_N)}{e(N)} \right|.$$

When N is a closed graph manifold with no $I(K)$ piece in N^* , using the notations introduced in Section 2.2, one can define the so-called *absolute volume* $|SV|$ by setting

$$|SV|(N) = \sum_{\Sigma \in N^*} SV(\hat{\Sigma}).$$

In the same way one can define the *absolute Euler number* of N by setting

$$|e|(N) = \sum_{\Sigma \in N^*} |e(\hat{\Sigma})|.$$

In Section 3.3 we will study the relations between $|e|(N)$ and $|SV|(N)$ (see Lemma 3.6).

3. REDUCTION OF COMPLEXITY VIA COVERINGS

In this section we state some results on finite coverings of surfaces and 3-manifolds which will be used in the proofs of Proposition 4.1 and Theorem 1.1.

3.1. Two general statements. The first result says that to prove the finiteness of the set $\mathcal{D}(M, N)$ one can replace N by a finite covering of it.

Lemma 3.1. (1) *Let M and N denote two closed oriented manifolds of the same dimension and let $p: N' \rightarrow N$ be a finite covering of N . If $\mathcal{D}(P, N')$ is finite for any closed manifold P then the set $\mathcal{D}(M, N)$ is finite.*

(2) *If all manifolds involved in (1) are graph manifolds, then the conclusion of (1) is still hold.*

Proof. (1) For each nonzero degree map $f: M \rightarrow N$, let $M(f)$ be the connected covering space of M corresponding to the subgroup $f_*^{-1}(p_*(\pi_1 N'))$ of $\pi_1 M$ which we denote by $r: M(f) \rightarrow M$. Let $f': M(f) \rightarrow N'$ be a lift of f , then $p \circ f' = f \circ r$. We claim that the set

$$\mathcal{C} = \{M(f), \text{ when } f \text{ runs over the nonzero degree maps from } M \text{ to } N\}$$

is finite. To see this, first note that the index of $f_*^{-1}(p_*(\pi_1 N'))$ in $\pi_1 M$ is bounded by the index of $p_*(\pi_1 N')$ in $\pi_1 N$. Indeed, the homomorphism $f_*: \pi_1 M \rightarrow \pi_1 N$ descends through an injective map

$$\bar{f}_*: \frac{\pi_1 M}{f_*^{-1}(p_*(\pi_1 N'))} \rightarrow \frac{\pi_1 N}{p_*(\pi_1 N')}$$

Since $\pi_1 M$ contains at most finitely many subgroup of a bounded index, it follows that $M(f)$ has only finitely many choices which proves that the set \mathcal{C} is finite. By the construction we have

$$\deg(f) = \frac{\deg(p)}{\deg(r)} \deg(f').$$

By the finiteness of the set \mathcal{C} and assumption on N' , the set of $\{\deg(f') | f : M' \rightarrow N', M' \in \mathcal{C}\}$ is finite. Clearly $\deg(r)$ have only finitely many choice, so the lemma is proved.

(2) If M and N are graph manifolds, then all manifolds $M(f)$, N' in the proof of (1) are graph manifolds. Clearly (2) follows. \square

Lemma 3.2. *Let N be a closed 3-manifold with non-trivial JSJ-decomposition. Then there exists a 2-fold covering \tilde{N} of N such that each JSJ-torus of \tilde{N} is shared by two different pieces of \tilde{N}^* .*

Proof. Let $\{T_1, \dots, T_k\}$ be the union JSJ-tori of N with each T_i is shared by the same piece of N^* . Let e_1, \dots, e_k be the corresponding edges in Γ_N . Then e_1, \dots, e_k are the edges of Γ_N with the two ends of each e_i being at the same vertex. Clearly $H_1(\Gamma_N; \mathbf{Z}) = \langle e_1 \rangle \oplus \dots \oplus \langle e_k \rangle \oplus G$.




FIGURE 1.

Let $r : N \rightarrow \Gamma_N$ be the retraction. Consider the following epimorphism

$$\phi : H_1(N, \mathbf{Z}) \xrightarrow{r_*} H_1(\Gamma_N; \mathbf{Z}) \xrightarrow{q} \langle e_1 \rangle \oplus \dots \oplus \langle e_k \rangle \xrightarrow{\lambda} \mathbf{Z}/2\mathbf{Z}$$

where r_* is induced by r , q is the projection, and λ is defined by $\lambda([e_i]) = \bar{1}$ for $i = 1, \dots, k$. Then the double covering \tilde{N} of N corresponding to ϕ satisfies the conclusion of the lemma, since the double covering of Γ_N corresponding to $\lambda \circ q$, which is the dual graph of \tilde{N} , contains no edge with two ends being at the same vertex. See Figure 1 for the local picture. \square

3.2. Separable and characteristic coverings. Let N be a closed graph manifold without $I(K)$ JSJ-piece. Let \mathcal{T} be a union of tori and let m be a positive integer. Call a covering $p : \tilde{\mathcal{T}} \rightarrow \mathcal{T}$ m -characteristic if for each component T of \mathcal{T} and

for each component \tilde{T} of $\tilde{\mathcal{T}}$ over T , the restriction $p| : \tilde{T} \rightarrow T$ is the covering map associated to the characteristic subgroup of index $m \times m$ in $\pi_1 T$. Call a finite covering $\tilde{N} \rightarrow N$ of a graph manifold N m -characteristic if its restriction to $\mathcal{T}_{\tilde{N}} \rightarrow \mathcal{T}_N$ is m -characteristic.

Next we define the *separable coverings*. Let Σ be a component of N^* with base 2-orbifold \mathcal{O} . Let Σ^0 , \mathcal{O}^0 , and $s : \mathcal{O}^0 \rightarrow \Sigma^0$ are given as in Section 2.1. Let $p : \tilde{\Sigma} \rightarrow \Sigma$ denote a finite covering. Recall that p is a fiber preserving map.

Recall that the *vertical degree* of p is the integer d_v such that $p_*(\tilde{h}) = h^{d_v}$, where h and \tilde{h} denote the homotopy class of the regular fiber in Σ and $\tilde{\Sigma}$, and the *horizontal degree* d_h is the degree of the induced branched covering $\bar{p} : \tilde{\mathcal{O}} \rightarrow \mathcal{O}$, where $\tilde{\mathcal{O}}$ denotes the base of the bundle $\tilde{\Sigma}$. We have $\deg(p) = d_v \times d_h$.

On the other hand, p induces a finite covering $p| : \tilde{\Sigma}^0 = p^{-1}(\Sigma^0) \rightarrow \Sigma^0$ and a covering $p|_b : \tilde{\mathcal{F}}^0 \rightarrow \mathcal{O}^0$, with $\tilde{\mathcal{F}}^0$ connected. More precisely, $p|_b$ corresponds to the subgroup $s_*^{-1}((p|)_*(\pi_1 \tilde{\Sigma}^0))$. Note that p and $p|$ have the same degree, same vertical degree and same horizontal degree. If $\deg(p|_b) = d_h$, then we say that the covering p is *separable*. The following result provides two classes of separable coverings which will be used later.

Lemma 3.3. *Let $p : \tilde{\Sigma} \rightarrow \Sigma$ be an oriented Seifert manifold finite covering.*

(i) *If p has fiber degree one, then p is a separable covering.*

(ii) *If $\Sigma = F \times \mathbf{S}^1$ and p is a regular covering corresponding to an epimorphism $\phi : \pi_1 N = \pi_1 F \times \mathbf{Z} \rightarrow G = G_1 \times G_2$ satisfying $\phi(\pi_1 F \times \{1\}) = G_1$ and $\phi(\{1\} \times \mathbf{Z}) = G_2$ then p is separable.*

Proof. Using the same notations as above it is easy to see that the map $p|_b : \tilde{\mathcal{F}}^0 \rightarrow \mathcal{O}^0$ factors through covering maps $q : \tilde{\mathcal{F}}^0 \rightarrow \tilde{\mathcal{O}}^0$ and $\bar{p} : \tilde{\mathcal{O}}^0 \rightarrow \mathcal{O}^0$ where $\tilde{\mathcal{O}}^0$ denote the base of the bundle $\tilde{\Sigma}^0$. Then we get

$$\deg(p|_b) = d_h \times \deg(q) = \deg(p) \times \deg(q)$$

since p has vertical degree one. On the other hand, since $\deg(p|_b) \leq \deg(p)$ then $\deg(q) = 1$. This proves (i).

If Σ is homeomorphic to a product $F \times \mathbf{S}^1$ then we have the following commutative diagram

$$\begin{array}{ccc} \tilde{F} & \xrightarrow{\tilde{s}} & \tilde{\Sigma} \\ p_b \downarrow & & \downarrow p \\ F & \xrightarrow{s} & \Sigma \end{array}$$

where \tilde{F} is connected. Since $\phi(\pi_1 F \times \{1\}) = G_1$ then $p^{-1}(s(F))$ has $|G_2|$ components and thus $\deg(p) = \deg(p_b) \times |G_2|$. Since $\deg(p) = |G_1| \times |G_2|$ then $\deg(p_b) = |G_1| = d_h$. This proves (ii). \square

3.3. Lifting of coordinates and gluing matrices. From now on we assume the graph manifold N is coordinated. Let $p : \tilde{N} \rightarrow N$ be a finite covering of graph

manifolds. Then obviously $\Gamma_{\tilde{N}}$ can be directed in a unique way such that the induced map $p_{\#} : \Gamma_{\tilde{N}} \rightarrow \Gamma_N$ preserves the directions of the edges. Below we also assume that $\Gamma_{\tilde{N}}$ is directed in such a way.

Let $p: \tilde{N} \rightarrow N$ be a finite covering of graph manifolds. Call p is *separable* if the restriction $p|: \tilde{\Sigma} \rightarrow \Sigma$ on connected Seifert pieces is separable for all possible $\tilde{\Sigma}$ and Σ . Call a coordinate on \tilde{N} is a *lift* of the coordinate of N , if for each possible covering $p|: \tilde{\Sigma} \rightarrow \Sigma$ on connected Seifert pieces, the (s, h) -basis of $\tilde{\Sigma}$ is lifted from the (s, h) -basis of Σ .

Lemma 3.4. (i) *Let $p: \tilde{N} \rightarrow N$ be a separable finite covering of graph manifolds. Then the coordinate of N can be lifted on \tilde{N} .*

(ii) *Moreover, if the covering p is characteristic, then for each component T of \mathcal{T}_N and for each component \tilde{T} over T we have $A_T = A_{\tilde{T}}$, where the coordinate of \tilde{N} is lifted from N .*

Proof. To prove (i), one need only to show that for a separable finite covering $p: \tilde{\Sigma} \rightarrow \Sigma$ of connected Seifert piece, then any (s, h) -basis of Σ lifts to a (s, h) -basis of $\tilde{\Sigma}$.

Using the same notation as in the proof of Lemma 3.3 we have

$$\deg(p|_b) = d_h \times \deg(q)$$

Since we assume $\deg(p|_b) = d_h$, then $\deg(q) = 1$ and thus \tilde{s} is a cross section. This proves (i).

Once N is coordinated, then each torus T in \mathcal{T}_N is associated with a unique 2×2 -matrix A_T provided by the gluing map $\tau|: T_-(s_-, h_-) \rightarrow T_+(s_+, h_+)$ such that $\tau(s_-, h_-) = (s_+, h_+)A_T$.

Similarly with lifted coordinate on \tilde{N} we have $\tilde{\tau}|: \tilde{T}_-(\tilde{s}_-, \tilde{h}_-) \rightarrow \tilde{T}_+(\tilde{s}_+, \tilde{h}_+)$ and $\tilde{\tau}(\tilde{s}_-, \tilde{h}_-) = (\tilde{s}_+, \tilde{h}_+)\tilde{A}_T$.

Since the coordinate of \tilde{N} are lifted from N , and p is m -characteristic for some m , we have the following commutative diagram

$$\begin{array}{ccc} (\tilde{s}_-, \tilde{h}_-) & \xrightarrow{\tilde{A}_T} & (\tilde{s}_+, \tilde{h}_+) \\ \times m \downarrow & & \downarrow \times m \\ (s_-, h_-) & \xrightarrow{A_T} & (s_+, h_+) \end{array}$$

Then one verifies directly that $\tilde{A}_T = A_T$. This proves (ii). \square

3.4. The absolute volume and the absolute Euler number. We end this section with a result (see Lemma 3.6) which states the relation between the absolute volume and the absolute Euler number of a graph manifold. First we begin with a technical result.

Lemma 3.5. *Suppose N is a closed graph manifolds without $I(K)$ JSJ-piece.*

(i) *For any finite covering $\tilde{N} \rightarrow N$, $|e|(\tilde{N}) = 0$ if and only if $|e|(N) = 0$.*

(ii) *There is a finite covering $p : \tilde{N} \rightarrow N$ which is separable and characteristic, and each Seifert piece of \tilde{N} is the product of a surface of genus at least 2 and the circle. Moreover \tilde{N} may be chosen so that $\Gamma_{\tilde{N}}$ has two vertices if Γ_N has two vertices.*

Proof. (i) follows from the definition and [LW, Proposition 2.3].

(ii) It has been proved in [LW, Proposition 4.4], that there is a characteristic finite covering $p : \tilde{N} \rightarrow N$ whose each piece is the product of a surface and the circle. By checking the proof, it is easy to see that the condition "genus at least 2" can be satisfied; moreover the restriction $p| : \tilde{\Sigma} \rightarrow \Sigma$ on connected JSJ-pieces is a composition of separable coverings described in Lemma 3.3, which is still separable.

If moreover Γ_N has exactly two vertices Σ_1 and Σ_2 , then for $i = 1, 2$, denote by $p_i : \hat{\Sigma}_i \rightarrow \Sigma_i$ the m -characteristic separable finite covering such that $\hat{\Sigma}_i$ is the product of a surface of genus at least 2 and the circle. There exists a 1-characteristic finite covering $q_i : \tilde{\Sigma}_i \rightarrow \hat{\Sigma}_i$ such that $\partial\tilde{\Sigma}_1$ and $\partial\tilde{\Sigma}_2$ have the same number of components. Next one can glue $\tilde{\Sigma}_1$ and $\tilde{\Sigma}_2$ by the lift of the sewing involution of N to get a characteristic and separable finite covering $p : \tilde{N} \rightarrow N$ whose dual graph has two vertices. This completes the proof of the lemma. \square

Lemma 3.6. *Let N be a closed graph manifold without $I(K)$ JSJ-pieces.*

(i) *If $|e|(N) \neq 0$ then N admits a finite covering \tilde{N} with $|SV|(\tilde{N}) \neq 0$.*

(ii) *If $|e|(N) = 0$ then N admits a finite covering \tilde{N} which can be coordinated such that each gluing matrix is in the form $\pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.*

Proof. By Lemma 3.5 (ii) let $p : \tilde{N} \rightarrow N$ be a finite covering which is separable and characteristic and each piece of \tilde{N}^* is a product $F \times S^1$ with $g(F) \geq 2$.

By Lemma 3.5 (i) $|e|(N) \neq 0$ implies $|e|(\tilde{N}) \neq 0$. By definition of $|e|$, $e(\hat{\Sigma}) \neq 0$ for some $\Sigma = F \times S^1 \in \tilde{N}^*$. Since $g(F) \geq 2$, $SV(\hat{\Sigma}) \neq 0$, and hence $|SV|(\tilde{N}) \neq 0$ by definition in Section 2.3. This proves (i).

Denote by $\Sigma_1, \dots, \Sigma_n$ the components of N^* . For each $i = 1, \dots, n$, denote by (Σ_i, α_i) the Seifert piece Σ_i of N^* endowed with its canonical framing. Since $e(N) = 0$ then $e(\Sigma_i(\alpha_i)) = e(\hat{\Sigma}_i) = 0$ and thus there exists a finite covering of $\hat{\Sigma}_i$, with fiber degree one, homeomorphic to a product. By pulling back this covering via the quotient map $\pi : \Sigma_i \rightarrow \hat{\Sigma}_i$ we get a covering $\tilde{\Sigma}_i$ of Σ_i such that the framing $(\tilde{\Sigma}_i, \tilde{\alpha}_i)$ satisfies the following condition: there exists a properly embedded incompressible surface F_i in $\tilde{\Sigma}_i$ such that $\tilde{\Sigma}_i \simeq F_i \times \mathbf{S}^1$ and $\partial F_i = \tilde{\alpha}_i$.

Suppose T is a component of $\partial\Sigma_i$ and T' is a component of $\partial\Sigma_j$ such that T is identified to T' then the sewing involution $\tau|T : T \rightarrow T'$ lifts to a sewing involution $\tilde{\tau} : \tilde{T} \rightarrow \tilde{T}'$, where \tilde{T} , resp. \tilde{T}' , denotes a component of $\partial\tilde{\Sigma}_i$, resp. a component of $\partial\tilde{\Sigma}_j$, over T , resp. T' . Indeed by our construction the induced coverings $\tilde{T} \rightarrow T$ and $\tilde{T}' \rightarrow T'$ correspond exactly to the subgroup of $\pi_1 T$, resp. of $\pi_1 T'$, generated by h and h' , where h is the fiber of Σ_i represented in T and h' is the fiber of Σ_j represented in T' , hence the gluing map lifts by the lifting criterion.

Corollary 4.2. *Suppose N is a closed graph manifold whose dual graph has two vertices and one edge. Then $|D(M, N)|$ is finite for any 3-manifold M .*

Proof. We may suppose that N contains no $I(K)$ piece. Otherwise N is doubly covered by a non-trivial graph manifold which contains no $I(K)$ piece and whose dual graph still has two vertices and one edge (since we assume that N is non-trivial graph manifold). In any case N has a finite cover \tilde{N} such that $SV(\tilde{N}) \neq 0$ by Proposition 4.1. Then by Lemmas 2.4 and 3.1, $|D(M, N)|$ is finite for any 3-manifold M . \square

The proof of Proposition 4.1 follows from the a sequence of lemmas below.

Let N be a n -multiple edges coordinated graph manifold. We say that N satisfies the *Property I* if

- (1) The JSJ-piece Σ_i is homeomorphic to a product $F_i \times \mathbf{S}^1$ where F_i is an oriented surface with genus ≥ 2 , for $i = 1, 2$;
- (2) $A_1 = A_2 = \dots = A_n$.

Lemma 4.3. *Let N be a n -multiple edges graph manifold satisfying the assumption of Proposition 4.1. Then there exist separable and characteristic finite coverings $p_1: N_1 \rightarrow N$ and $p_2: N_1 \rightarrow N_2$ such that N_2 satisfies Property I.*

Proof. By Lemma 3.5 (ii) and Lemma 3.4, we may assume that N is a n -multiple edges graph manifold satisfying the assumption of Proposition 4.1, and moreover Σ_i is homeomorphic to a product $F_i \times \mathbf{S}^1$ where F_i is an oriented surface with genus ≥ 2 .

May assume that $A_1 = \dots = A_k = -A$ and $A_{k+1} = \dots = A_n = A$, $0 < k < n$, shown as in the left of Figure 3.

Denote by $c_{i,j}$ the loops of Γ_N corresponding to the "composition" $T_i \cdot (-T_j)$, note that here T_i represents an oriented edge. Then $c_{i,k+1}$ for $i = 1, \dots, k$ and $c_{j,n}$ for $j = k+1, \dots, n-1$ form a basis of $H_1(\Gamma_N)$ and we have

$$H_1(\Gamma_N) = (\oplus_{i=1}^k \langle c_{i,k+1} \rangle) \oplus (\oplus_{j=k+1}^{n-1} \langle c_{j,n} \rangle)$$

Next we define an epimorphism


$$\phi: H_1(N, \mathbf{Z}) \xrightarrow{r_*} H_1(\Gamma_N; \mathbf{Z}) \xrightarrow{q} \oplus_{i=1}^k \langle c_{i,k+1} \rangle \xrightarrow{\lambda} \mathbf{Z}/2\mathbf{Z}$$

where r_* is induced by the retraction $r: N \rightarrow \Gamma_N$, q is the projection and λ is defined by $\lambda(c_{i,k+1}) = \bar{1}$ for any $i \in \{1, \dots, k\}$. Denote by $p_1: N_1 \rightarrow N$ the 2-fold covering corresponding to ϕ , and by μ the deck transformation of this covering.

It is easy to see this covering is separable and 1-characteristic. Moreover with the lifted coordinates of N , the directed graph Γ_{N_1} with gluing matrices $\pm A$, as well as the two lifts Σ_i^1 and Σ_i^2 of Σ_i , $i = 1, 2$, are shown in the right of Figure 3.

Let $\Sigma_i^j = F_i^j \times S^1$. It is not difficult to see that there is an orientation preserving involution η_i^j on Σ_i^j satisfying the following

- (1) for η_i^j reverses both the orientation of F_i^j and S^1 ,

FIGURE 4. Fixed point free action of $\mathbf{Z}/d\mathbf{Z}$

Keeping the coordinate on Σ'_1 and re-coordinating Σ'_2 by $(-s_2, -h'_2)$ if needed, we may assume that $A = \begin{pmatrix} 1 & b \\ 0 & -1 \end{pmatrix}$ or $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Suppose first $A = \begin{pmatrix} 1 & b \\ 0 & -1 \end{pmatrix}$. Denote $F'_1 \simeq F'_2$ by F . Let denote by

$$\pi: F'_1 \times h'_1 \amalg F'_2 \times h'_2 \rightarrow \Sigma = F \times h$$

the trivial 2-fold covering map, where $\pi(h'_i) = h$ and $\pi(s_i) = s$. Denote by $\rho: \Sigma \rightarrow \hat{\Sigma}$ the quotient map associated with the Dehn filling on $\partial\Sigma$ along the curve $\frac{b}{(2,b)}s - \frac{2}{(2,b)}h$, where $(2, b)$ denotes the greatest common divisor of 2 and b .

Note that, since $b \neq 0$, then $\hat{\Sigma}$ is a $\widehat{\text{PSL}}(2, \mathbf{R})$ -manifold.

One can verify routinely that in the π_1 level the relations provided by gluing Σ'_1 and Σ'_2 via τ' are sent to the relation provided by Dehn filling on Σ via $\frac{b}{(2,b)}s - \frac{2}{(2,b)}h$ under ρ , hence the map $\rho \circ \pi: \Sigma'_1 \amalg \Sigma'_2 \rightarrow \hat{\Sigma}$ factors through $\frac{\Sigma'_1 \amalg \Sigma'_2}{\tau'} \simeq N'$

which is sent into $\hat{\Sigma}$ by a degree 2 map, since the sewing involution τ' is orientation reversing so that N' inherits compatible orientations from the pieces Σ'_1 and Σ'_2 .

In the case of $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, we can perform the same construction as above, just replace the filling curve $\frac{b}{(2,b)}s - \frac{2}{(2,b)}h$ by the curve $s - h$. This proves Lemma 4.5. \square

By Lemmas 4.3, 4.4 and 4.5 and their proofs, we have the following diagram:

$$\begin{array}{ccccc} & N_1 & & N_3 & \\ & \downarrow p_1 & \searrow p_2 & \downarrow p_3 & \searrow p_4 \\ & N & & N_2 & & N_4 \end{array}$$

where p_1 and p_2 are coverings provided by Lemma 4.3, p_3 is the coverings provided by Lemma 4.4, and the non-zero degree map p_4 is provided by Lemma 4.5, where $SV(N_4) \neq 0$. Since SV has property D , $SV(N_3) \neq 0$.

Consider the covering \tilde{N} corresponding to the finite index subgroup $p_{2*}(\pi_1 N_1) \cap p_{3*}(\pi_1 N_3)$ in $\pi_1 N_2$. Then \tilde{N} covers both N_1 (and thus N) and N_3 , and $SV(\tilde{N}) \neq 0$. Then the proof of Proposition 4.1 is complete.

5. PROOF OF THEOREM 1.1

5.1. Simplifications. Let N be a closed non-trivial graph manifold. We are going to show that $|D(M, N)|$ is finite for any given graph manifold M .

(1) First we simplify N : By Lemma 3.2 and Lemma 3.5, there is a finite covering \tilde{N} of N satisfying the condition (*): each JSJ piece of \tilde{N} is a product of an oriented surface with genus ≥ 2 and the circle, and each JSJ torus is shared by two different JSJ pieces.

By Lemma 3.1, if $|D(M, N)|$ is not finite for some graph manifold M , then $|D(P, \tilde{N})|$ is not finite for some graph manifold P . So we may assume N already satisfies the condition (*).

(2) Then we simplify M : For given M , let $\mathcal{H} = \{M_1, \dots, M_k\}$ be the finite set of graph manifolds provided by Proposition 2.2. By Proposition 2.2 (i), if $|D(M, N)|$ is not finite, then $|D(M_i, N)|$ is not finite for some $M_i \in \mathcal{H}$. So may assume that (**) $M = M_i \in \mathcal{H}$ for some $i \in \{1, \dots, k\}$.

5.2. Proof of Theorem 1.1 when $|e|(N) \neq 0$. Suppose $|e|(N) \neq 0$. By Lemma 3.6 and (*) in 5.1, we may assume that $|SV|(N) \neq 0$. Then there exists a Seifert piece Q of N^* such that $SV(\hat{Q}) \neq 0$. By (**) in 5.1 and Propostion 2.2 (ii), we may assume that $L_Q(f) = f^{-1}(Q)$ is a canonical submanifold of M . Below we denote $L_Q(f)$ as L_Q for short.

Lemma 5.1. *L_Q can be chosen so that any component T of ∂L_Q is shared by two distinct Seifert pieces of M : one in L_Q and another in $M \setminus L_Q$.*

Proof. Indeed if not, then there exists two distinct components T and T' of ∂L_Q which are identified by the sewing involution τ_M of M and such that T and T' are sent by f into the same component of ∂Q . Denote by \bar{L}_Q the canonical submanifold of M obtained by identifying T and T' via τ_M . Since each component of ∂Q is shared by two distinct Seifert pieces of N by the assumption in 5.1, f induces a proper map $\bar{f}: \bar{L}_Q \rightarrow Q$. After finitely many such operations, we reach a new L_Q satisfying the requirement of Lemma 5.1 \square

Below we assume that L_Q satisfies the requirement of Lemma 5.1. Now we choose L_Q to be maximal in the sense that for any Seifert piece S in $M \setminus L_Q$ adjacent to L_Q , S is not able to be added into L_Q by homotopy on f . Then $f(S) \subset B_S$, where B_S is a Seifert piece of N , distinct from Q and adjacent to Q .

Since L_Q is maximal, by Lemma 2.3, we deduce that for any Seifert piece S adjacent to L_Q along a component of ∂L_Q , $f|_S: S \rightarrow B_S$ is fiber preserving. Hence the proper map $f|_{L_Q}: L_Q \rightarrow Q$ preserves the canonical framings, and it induces a map $\hat{f}: \hat{L}_Q \rightarrow \hat{Q}$ between the closed manifolds obtained after Dehn filling along the canonical framings. By Lemma 2.4 we have

$$SV(\hat{L}_Q) \geq |\deg(\hat{f})|SV(\hat{Q}).$$

Since $\deg(f) = \deg(f|_{L_Q}) = \deg(\hat{f})$, we get

$$|\deg(f)| \leq \frac{SV(\hat{L}_Q)}{SV(\hat{Q})}.$$

Therefore

$$|\deg(f)| \leq \max \left\{ \frac{SV(\hat{L})}{SV(\hat{Q})} \mid Q \in N^*, SV(\hat{Q}) \neq 0; L \text{ is canonical in } M \right\}$$

By Lemma 2.1 there are only finitely many \hat{Q} and only finitely many \hat{L} . So the right side of the above inequality is finite. This completes the proof of Theorem 1.1 when $|e|(N) \neq 0$.

5.3. Proof of Theorem 1.1 when $|e|(N) = 0$. By Lemma 3.1 and Lemma 3.6 we can assume each gluing matrix of N is equal to $\pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Choose two distinct adjacent Seifert pieces S_1 and S_2 in N , denote by $\mathcal{T} = \partial S_1 \cap \partial S_2$ and by Q the connected graph manifold $S_1 \cup_{\mathcal{T}} S_2$ (such Seifert pieces exist by 5.1). By Proposition 2.2, we may assume that $f^{-1}(Q) = L_Q$ is a canonical submanifold of M .

Since each JSJ-torus of N is shared by two different JSJ-pieces, by the same arguments as in 5.2, we may assume that each component of ∂L_Q is shared by two distinct Seifert pieces of M one in L_Q and another in $M \setminus L_Q$. Furthermore we can arrange L_Q to be maximal in the sense of 5.2, then by Lemma 2.3, we deduce that any Seifert piece S' of M adjacent to L_Q is sent by f to a Seifert piece B' adjacent to Q such that $f|_{S'}: S' \rightarrow B'$ is fiber preserving.

As in 5.2, it follows that the proper map $f|L_Q: L_Q \rightarrow Q$ induces a map $\hat{f}: \hat{L}_Q \rightarrow \hat{Q}$ between closed graph manifolds obtained by Dehn filling along the canonical framings. Moreover, as in 5.2 we have $\deg(f) = \deg(f|L_Q) = \deg(\hat{f})$ and thus

$$|\mathcal{D}(M, N)| \leq \max \left\{ |\mathcal{D}(\hat{L}, \hat{Q})| \mid L \text{ is canonical in } M \right\}$$

Note that $\hat{Q} = \hat{S}_1 \cup_{\mathcal{T}} \hat{S}_2$, where \hat{S}_i is obtained by Dehn filling along the canonical framings on $\partial S_i \setminus \mathcal{T}$, $i = 1, 2$. It follows that \hat{Q} satisfies the hypothesis of Proposition 4.1. Then \hat{Q} has a finite covering \tilde{Q} with $SV(\tilde{Q}) \neq 0$ by Proposition 4.1. Hence by Lemma 3.1, the set $|\mathcal{D}(\hat{L}, \hat{Q})|$ is finite for any \hat{L} . Since by Lemma 2.1 there are only finitely many \hat{L} , this completes the proof of Theorem 1.1 when $|e|(N) \neq 0$. Hence Theorem 1.1 is proved.

REFERENCES

- [BG1] R. BROOKS, W. GOLDMAN, *Volumes in Seifert space*, Duke Math. J. 51 (1984), no. 3, 529–545.
- [BG2] R. BROOKS, G. GOLDMAN, The Godbillon-Vey invariant of a transversely homogeneous foliation. Trans. Amer. Math. Soc. 286 (1984), no. 2, 651–664.
- [CT] J. CARLSON, D. TOLEDO, *Harmonic mappings of Kähler manifolds to locally symmetric spaces*. Inst. Hautes Études Sci. Publ. Math. No. 69 (1989), 173–201.
- [D1] P. DERBEZ, *Non-zero degree maps between closed orientable three-manifolds*, Trans. Amer. Math. Soc. 359 (2007), 3887–3911.
- [D2] P. DERBEZ, *Topological rigidity and Gromov simplicial volume*. math.GT/0607766.
- [G] M. GROMOV, *Volume and bounded cohomology*, Inst. Hautes Études Sci. Publ. Math. No. 56, (1982), 5–99.
- [JS] W. JACO, P.B. SHALEN, *Seifert fibered space in 3-manifolds*, Mem. Amer. Math. Soc. 21 (1979).
- [Jo] K. JOHANNSON, *Homotopy equivalences of 3-manifolds with boundaries*, Lecture Notes in Mathematics, 761. Springer, Berlin, 1979.
- [LW] J. LUECKE, Y. WU, *Relative Euler number and finite covers of graph manifolds* Geometric topology (Athens, GA, 1993), 80–103, AMS/IP Stud. Adv. Math., 2.1, Amer. Math. Soc., Providence, RI, 1997.
- [N] W. D. NEUMANN, *A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves* Trans. Amer. Math. Soc. 268 (1981), no. 2, 299–344.
- [Re1] A. REZNIKOV, *Rationality of secondary classes*, J. Differential Geom. 43 (1996), no. 3, 674–692.
- [Re2] A. REZNIKOV, *Volumes of discrete groups and topological complexity of homology spheres*, Math. Ann. 306 (1996), no. 3, 547–554.
- [Ro] Y. Y. RONG, *Degree one maps of Seifert manifolds and a note on Seifert volume* Topology Appl. 64 (1995), no. 2, 191–200.
- [So] T. Soma, *The Gromov invariant of links*. Invent. Math. **64** 1981, 445–454.
- [Th1] W. THURSTON, *The geometry and topology of 3-manifolds*, Lectures Notes, Princeton Univ., 1979.
- [Th2] W. THURSTON, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*. Bull. Am. Math. Soc. 6, 357–381 (1982)
- [W1] S.C. WANG, *The π_1 -injectivity of self-maps of nonzero degree on 3-manifolds*, Math. Ann. 297 (1993), no. 1, 171–189.

- [W2] S.C. WANG, *Non-zero degree maps between 3-manifolds*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 457–468, Higher Ed. Press, Beijing, 2002.
- [WZ] S.C. WANG, Q. ZHOU, *Any 3-manifold 1-dominates at most finitely many geometric 3-manifolds*, Math. Ann. 322 (2002), no. 3, 525–535.
- [Wa] F. WALDHAUSEN, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. 87 (1968), 56–88.
- [YW] F. YU; S.C. WANG, *Covering degrees are determined by graph manifolds involved*. Comment. Math. Helv. 74 (1999), no. 2, 238–247.

C.M.I, TECHNOPÔLE CHÂTEAU-GOMBERT, 39, RUE JOLIOT CURIE, 13453 MARSEILLE
CEDEX 13, FRANCE

E-mail address: derbez@cmi.univ-mrs.fr

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY, BEIJING, CHINA

E-mail address: wangsc@math.pku.edu.cn