## Comment on "Direct space-time observation of pulse tunneling in an electromagnetic band gap"

G. Nimtz<sup>\*</sup> and A.A. Stahlhofen<sup>†</sup>

Institut für Integrierte Naturwissenschaft, Universität Koblenz Universiätsstrasse 1, 56070 Koblenz, Germany (Dated: October 24, 2018)

The investigation presented by Doiron, Hache, and Winful [Phys. Rev. A 76, 023823 (2007)] is not valid for the tunneling process as claimed in the paper.

PACS numbers: 42.25.Bs,03.65.Ta,42.70.Qs,73.40.Gk

The title of a recent article by Doiron et al. [1] is misleading. The authors have investigated a dielectric mirror but not a tunneling barrier, see also Refs.[2]. The measured and discussed superluminal group velocity is similar to that studied on a Lorentz-Lorenz oscillator by Sommerfeld and Brillouin a hundred years ago [3]. A dielectric mirror is based on a periodical quarter wavelength structure of two refractive indices in which destructive interference causes a standing wave pattern decaying exponentially with mirror thickness as shown in Figs. 2(a - f) in Ref.[1]. There are nodes in the intensity every half wavelength.

Tunneling, however, is understood and performed by electromagnetic evanescent modes or by tunneling solutions of the Schrödinger equation, which have purely imaginary wave numbers. The latter includes a purely imaginary refractive index. Signals with purely evanescent frequency components may travel at a superluminal velocity [4, 5]. Inside the barrier there is no standing wave and tunneling proceeds even instantaneously, it represents a process well described by virtual photons [6].

Recently, the transmission of dielectric mirrors and equivalent structures has often been misinterpreted as tunneling, see for instance Refs. [4, 7].

Actually, in the paper there are some errors: Fig.3 \*Also at II.Physikalisches Institut, Universität zu Köln †Electronic address: G.Nimtz@uni-koeln.de shows the vacuum light velocity and in section (II,D) the dwell time is not directly measured but it is derived from an approximately integrated stored energy and from the measured input power. This indirect measuring procedure is indirect like the light measurement from direct frequency and wavelength measurements, see for information NBS and NIST, for instance.

In addition it is claimed to have measured a resonator decay time, but detectors measure the traversal time of a pulse through a black box independent of the content of the box. The authors are asking whether an identifiable pulse peak actually propagates through the barrier? According to their Fig.1 not only the peak but also the pulse half width (for instance representing a digital signal) propagated faster than light and were correctly detected. Figure 2(a-f) displays the standing wave patterns at various times. Figure 2(d) shows the peak of the pulse from the mirrors entrance to the end of the mirror. Taking the x = 0 values from Figs.2(a) to 2(f) we see the peak in 2(d) instantaneously propagating from the entrance to the exit with the exponential decay of the standing wave.

Describing the traversal time of a mirror as a decay time of a cavity does not represent a contribution to the understanding of the tunneling time as claimed by Doiron et al. [1].

- S. Doiron, A. Hache, and H. Winful, Phys. Rev. A 76, 023823 (2007)
- [2] H. Winful, Phys. Rev. Lett. 90, 023901 (2003); Phys. Rev. Lett. 91, 260401 (2003); Phys. Rep. 436, 1 (2006); Opt. Express 10, 1491 (2002); Nature 424, 638 (2003); Phys. Rev E 68, 016615 (2003); Phys. Rev. E 72, 046608 (2005)
- [3] L. Brillouin, Wave Propagation and Group Velocity, Academic Press, New York (1960)
- [4] G. Nimtz and W. Heitmann, Progr. Quantum Electron.,

**21**, 81 (1997)

- [5] G. Nimtz, Lecture Notes In Physics, Eds. J. Ehlers and C. Lämmerzahl, **702**, 506 (2006)
- [6] A.A.Stahlhofen and G. Nimtz, Euro. Phys. Lett. 76, 189 (2006) and arXiv:0708.0681v1
- [7] A. M. Steinberg, P. G. Kwiat, and R. Y. Chiao, Phys. Rev. Lett. **71**, 708 (1993)