

On the oscillatory integration of some ordinary differential equations

Octavian G. Mustafa*
*Faculty of Mathematics, D.A.L.,
 University of Craiova, Romania*
e-mail: octaviangenghiz@yahoo.com

Abstract Conditions are given for a class of nonlinear ordinary differential equations $x'' + a(t)w(x) = 0$, $t \geq t_0 \geq 1$, which includes the linear equation to possess solutions $x(t)$ with prescribed oblique asymptote that have an oscillatory pseudo-wronskian $x'(t) - \frac{x(t)}{t}$.

2000 Mathematics Subject Classification: 34A30 (Primary), 34E05, 34K25

Keywords: Ordinary differential equation; Asymptotic integration; Prescribed asymptote; Non-oscillation of solutions

1 Introduction

A certain interest has been shown recently in studying the existence of bounded and positive solutions to a large class of elliptic partial differential equations which can be displayed as

$$\Delta u + f(x, u) + g(|x|)x \cdot \nabla u = 0, \quad x \in G_R, \quad (1)$$

where $G_R = \{x \in \mathbb{R}^n : |x| > R\}$ for any $R \geq 0$ and $n \geq 2$. We would like to mention the contributions [1], [3], [8] – [11], [13, 14], [18] and their references in this respect.

It has been established, see [8, 9], that it is sufficient for the functions f , g to be Hölder continuous, respectively continuously differentiable in order to analyze the asymptotic behavior of the solutions to (1) by the comparison

*Correspondence address: Str. Tudor Vladimirescu, Nr. 26, 200534 Craiova, Dolj, Romania

method [15]. In fact, given $\zeta > 0$, let us assume that there exist a continuous function $A : [R, +\infty) \rightarrow [0, +\infty)$ and a nondecreasing, continuously differentiable function $W : [0, \zeta] \rightarrow [0, +\infty)$ such that

$$0 \leq f(x, u) \leq A(|x|)W(u) \quad \text{for all } x \in G_R, u \in [0, \zeta]$$

and $W(u) > 0$ when $u > 0$. Then we are interested in the positive solutions $U = U(|x|)$ of the elliptic partial differential equation

$$\Delta U + A(|x|)W(U) = 0, \quad x \in G_R,$$

for the rôle of super-solutions to (1).

M. Ehrnström [13] noticed that, by imposing the restriction

$$x \cdot \nabla U(x) \leq 0, \quad x \in G_R,$$

upon the super-solutions U , an improvement of the conclusions from the literature is achieved for the special subclass of equations (1) where g takes only nonnegative values. Further developments of Ehrnström's idea are given in [1, 3, 11, 14].

Translated into the language of ordinary differential equations, the research about U reads as follows: given $c_1, c_2 \geq 0$, find (if any) a positive solution $x(t)$ of the nonlinear differential equation

$$x'' + a(t)w(x) = 0, \quad t \geq t_0 \geq 1, \quad (2)$$

where the coefficient $a : [t_0, +\infty) \rightarrow \mathbb{R}$ and the nonlinearity $w : \mathbb{R} \rightarrow \mathbb{R}$ are continuous and given by means of A, W , such that

$$x(t) = c_1 t + c_2 + o(1) \quad \text{when } t \rightarrow +\infty \quad (3)$$

and

$$\mathcal{W}(x, t) = \frac{1}{t} \begin{vmatrix} x'(t) & 1 \\ x(t) & t \end{vmatrix} = x'(t) - \frac{x(t)}{t} < 0, \quad t > t_0. \quad (4)$$

The symbol $o(f)$ for a given functional quantity f has here its standard meaning. In particular, by $o(1)$ we refer to a function of t that decreases to 0 as t increases to $+\infty$.

The papers [2, 3, 20, 22, 23] present various properties of the functional quantity \mathcal{W} , which shall be called *pseudo-wronskian* in the sequel. Our aim in this note is to complete their conclusions by giving some sufficient conditions upon a and w which lead to the existence of a solution x to (2) that verifies (3) while having an *oscillatory pseudo-wronskian* (this means that there exist the unbounded from above sequences $(t_n^\pm)_{n \geq 1}$ and $(t_n^0)_{n \geq 1}$ such that $t_{2n-1}^0 < t_n^+ < t_{2n}^0 < t_n^- < t_{2n+1}^0$ and $\mathcal{W}(t_n^+) > \mathcal{W}(t_n^0) = 0 > \mathcal{W}(t_n^-)$ for all $n \geq 1$). We answer thus to a question raised in [3, p. 371], see also the comment in [2, pp. 46–47].

2 The sign of \mathcal{W}

Let us start the discussion with a simple condition to settle the sign issue of the pseudo-wronskian.

Lemma 1 *Given $x \in C^2([t_0, +\infty), \mathbb{R})$, suppose that $x''(t) \leq 0$ for all $t \geq t_0$. Then $\mathcal{W}(x, \cdot)$ can change from being nonnegative-valued to being negative-valued at most once in $[t_0, +\infty)$. In fact, its set of zeros is an interval (possibly degenerate).*

Proof. Notice that

$$\frac{d^2}{dt^2}[x(t)] = \frac{1}{t} \cdot \frac{d}{dt}[t\mathcal{W}(x, t)], \quad t \geq t_0.$$

The function $t \mapsto t\mathcal{W}(x, t)$ being nonincreasing, it is clear that, if it has zeros, it has either a unique zero or an interval of zeros. \square

The result has an obvious counterpart.

Lemma 2 *Given $x \in C^2([t_0, +\infty), \mathbb{R})$, suppose that $x''(t) \geq 0$ for all $t \geq t_0$. Then, $\mathcal{W}(x, \cdot)$ can change from being nonpositive-valued to being positive-valued at most once in $[t_0, +\infty)$. Again, its set of zeros is an interval (possibly reduced to one point).*

Consider that x is a positive solution of equation (2) in the case where $a(t) \geq 0$ in $[t_0, +\infty)$ and $w(u) > 0$ for all $u > 0$. Then, we have

$$\frac{d\mathcal{W}}{dt} = -\frac{\mathcal{W}}{t} - a(t)w(x(t)), \quad t \geq t_0,$$

which leads to

$$\mathcal{W}(x, t) = \frac{1}{t} \left[t_0\mathcal{W}_0 - \int_{t_0}^t sa(s)w(x(s))ds \right], \quad \mathcal{W}_0 = \mathcal{W}(x, t_0), \quad (5)$$

throughout $[t_0, +\infty)$ by means of Lagrange's variation of constants formula.

The integrand in (5) being nonnegative-valued, we regain the conclusion of Lemma 1. In fact, if $T \in [t_0, +\infty)$ is a zero of $\mathcal{W}(x, \cdot)$ then it is a solution of the equation

$$t_0\mathcal{W}_0 = \int_{t_0}^T sa(s)w(x(s))ds. \quad (6)$$

On the other hand, if the pseudo-wronskian of x is positive-valued throughout $[t_0, +\infty)$ then it is necessary to have

$$(t_0\mathcal{W}_0 \geq) \quad \int_{t_0}^{+\infty} sa(s)w(x(s))ds < +\infty. \quad (7)$$

It has become clear at this point that whenever the equation (2) has a positive solution x such that $\mathcal{W}_0 \leq 0$, the functional coefficient a is nonnegative-valued and has at most isolated zeros and $w(u) > 0$ for all $u > 0$, the pseudo-wronskian \mathcal{W} satisfies the restriction (4). Now, returning to the problem stated in the Introduction, we can evaluate the main difficulty of the investigation: *if the positive solution x has prescribed asymptotic behavior, see formula (3) or a similar development, then we cannot decide upfront whether or not $\mathcal{W}_0 \leq 0$.* The formula (6) shows that there are also certain difficulties to estimate the zeros of the pseudo-wronskian.

3 The behavior of \mathcal{W}

Let us survey in this section some of the recent results regarding the pseudo-wronskian.

It has been established that its presence in the structure of a nonlinear differential equation

$$x'' + f(t, x, x') = 0, \quad t \geq t_0 \geq 1, \quad (8)$$

where the nonlinearity $f : [t_0, +\infty) \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous, allows for a remarkable flexibility of the hypotheses when searching for solutions with the asymptotic development (3) (or similar).

Theorem 1 ([20, p. 177]) *Assume that there exist the nonnegative-valued, continuous functions $a(t)$ and $g(s)$ such that $g(s) > 0$ for all $s > 0$ and $xg(s) \leq g(x^{1-\alpha}s)$, where $x \geq t_0$ and $s \geq 0$, for a certain $\alpha \in (0, 1)$. Suppose further that*

$$|f(t, x, x')| \leq a(t)g\left(\left|x' - \frac{x}{t}\right|\right) \quad \text{and} \quad \int_{t_0}^{+\infty} \frac{a(s)}{s^\alpha} ds < \int_{c+|\mathcal{W}_0|t_0^{1-\alpha}}^{+\infty} \frac{du}{g(u)}.$$

Then the solution of equation (8) given by (5) exists throughout $[t_0, +\infty)$ and has the asymptotic behavior

$$x(t) = c \cdot t + o(t), \quad x'(t) = c + o(1) \quad \text{when } t \rightarrow +\infty \quad (9)$$

for some $c = c(x) \in \mathbb{R}$.

To compare this result with the standard conditions in asymptotic integration theory regarding the development (9), see the papers [2, 3, 24] and the monograph [19].

Another result is concerned with the presence of the pseudo-wronskian in the function space $L^1((t_0, +\infty), \mathbb{R})$.

Theorem 2 ([3, p. 371]) *Assume that f does not depend explicitly of x' and there exists the continuous function $F : [t_0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$, which is nondecreasing with respect to the second variable, such that*

$$|f(t, x)| \leq F\left(t, \frac{|x|}{t}\right) \quad \text{and} \quad \int_{t_0}^{+\infty} t \left[1 + \ln\left(\frac{t}{t_0}\right)\right] F\left(t, |c| + \frac{\varepsilon}{t_0}\right) dt < \varepsilon$$

for certain numbers $c \neq 0$ and $\varepsilon > 0$. Then there exists a solution $x(t)$ of equation (8) defined in $[t_0, +\infty)$ such that

$$x(t) = c \cdot t + o(1) \quad \text{when } t \rightarrow +\infty \quad \text{and} \quad \mathcal{W}(x, \cdot) \in L^1.$$

The effect of perturbations upon the pseudo-wronskian is investigated in the papers [2, 20, 22].

Theorem 3 ([20, p. 183]) *Consider the nonlinear differential equation*

$$x'' + f(t, x, x') = p(t), \quad t \geq t_0 \geq 1, \quad (10)$$

where the functions $f : [t_0, +\infty) \times \mathbb{R}^2 \rightarrow \mathbb{R}$ and $p : [t_0, +\infty) \rightarrow \mathbb{R}$ are continuous and verify the hypotheses

$$|f(t, x, x')| \leq a(t) \left|x' - \frac{x}{t}\right|, \quad \int_{t_0}^{+\infty} ta(t)dt < +\infty$$

and

$$\lim_{t \rightarrow +\infty} \frac{1}{t} \int_{t_0}^t sp(s)ds = C \in \mathbb{R} - \{0\}.$$

Then, given $x_0 \in \mathbb{R}$, there exists a solution $x(t)$ of equation (10) defined in $[t_0, +\infty)$ such that

$$x(t_0) = x_0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} \mathcal{W}(x, t) = C.$$

In particular,

$$\lim_{t \rightarrow +\infty} \frac{x(t)}{t \ln t} = C.$$

A slight modification of the discussion in [22, Remark 3], see [2, p. 47], leads to the next result.

Theorem 4 *Assume that f in (10) does not depend explicitly of x' and there exists the continuous function $F : [t_0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$, which is nondecreasing with respect to the second variable, such that*

$$|f(t, x)| \leq F(t, |x|) \text{ and } \int_t^{+\infty} sF\left(s, |P(s)| + \sup_{\tau \geq s} \{q(\tau)\}\right) ds \leq q(t), \quad t \geq t_0,$$

for a certain positive-valued, continuous function $q(t)$ possibly decaying to 0 as $t \rightarrow +\infty$. Here, P is the twice continuously differentiable antiderivative of p , that is $P''(t) = p(t)$ for all $t \geq t_0$. Suppose further that

$$\limsup_{t \rightarrow +\infty} \left[t \frac{\mathcal{W}(P, t)}{q(t)} \right] > 1 \quad \text{and} \quad \liminf_{t \rightarrow +\infty} \left[t \frac{\mathcal{W}(P, t)}{q(t)} \right] < -1.$$

Then equation (10) has a solution $x(t)$ throughout $[t_0, +\infty)$ such that

$$x(t) = P(t) + o(1) \quad \text{when } t \rightarrow +\infty$$

and $\mathcal{W}(x, \cdot)$ oscillates.

Finally, the presence of the pseudo-wronskian in the structure of a nonlinear differential equation can lead to multiplicity when searching for solutions with the asymptotic development (3).

Theorem 5 ([23, Theorem 1]) *Given the numbers $x_0, x_1, c \in \mathbb{R}$, with $c \neq 0$, and $t_0 \geq 1$ such that $t_0 x_1 - x_0 = c$, consider the Cauchy problem*

$$\begin{cases} x'' = \frac{1}{t}g(tx' - x), & t \geq t_0 \geq 1, \\ x(t_0) = x_0, & x'(t_0) = x_1, \end{cases} \quad (11)$$

where the function $g : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous, $g(c) = g(3c) = 0$ and $g(u) > 0$ for all $u \neq c$. Assume further that

$$\int_{c+}^{2c} \frac{du}{g(u)} < +\infty \quad \text{and} \quad \int_{2c}^{(3c)^-} \frac{du}{g(u)} = +\infty.$$

Then problem (11) has an infinity of solutions $x(t)$ defined in $[t_0, +\infty)$ and developable as

$$x(t) = c_1 t + c_2 + o(1) \quad \text{when } t \rightarrow +\infty$$

for some $c_1 = c_1(x)$ and $c_2 = c_2(x) \in \mathbb{R}$.

The asymptotic analysis of certain functional quantities attached to the solutions of equations (2), (8) and (10), as in our case the pseudo-wronskian, might lead to some surprising consequences. Among the functional quantities that gave the impetus to spectacular developments in the qualitative theory of linear/nonlinear ordinary differential equations we would like to refer to

$$\mathcal{K}(x)(t) = x(t)x'(t), \quad t \geq t_0,$$

employed in the theory of *Kneser-solutions*, see the papers [6, 7] for the linear and respectively the nonlinear case and the monograph [19], and

$$\mathcal{HW}(x) = \int_{t_0}^{+\infty} x(s)w(x(s))ds.$$

The latter quantity is the core of the nonlinear version of *Hermann Weyl's limit-point/limit-circle classification* designed for equation (2), see the well-documented monograph [5] and the paper [21].

4 The negative values of \mathcal{W}

We shall assume in the sequel that the nonlinearity w of equation (2) verifies some of the hypotheses listed below:

$$|w(x) - w(y)| \leq k|x - y|, \quad \text{where } k > 0, \quad (12)$$

and

$$w(0) = 0, \quad w(x) > 0 \text{ when } x > 0, \quad |w(xy)| \leq w(|x|)w(|y|) \quad (13)$$

for all $x, y \in \mathbb{R}$. We notice that restriction (13) implies the existence of a majorizing function F , as in Theorem 2, given by the estimates

$$|f(t, x)| = |a(t)w(x)| \leq |a(t)| \cdot w(t)w\left(\frac{|x|}{t}\right) = F\left(t, \frac{|x|}{t}\right).$$

We can now use the paper [24] to recall the main conclusions of an asymptotic integration of equation (2). It has been established that whenever $\int_{t_0}^{+\infty} tw(t)|a(t)|dt < +\infty$, all the solutions of (2) have asymptotes (3) and their first derivatives are developable as

$$x'(t) = c_1 + o(t^{-1}) \quad \text{when } t \rightarrow +\infty. \quad (14)$$

Consequently, $\mathcal{W}(x, t) = -c_2t^{-1} + o(t^{-1})$ for all large t 's. In this case (the functional coefficient a has varying sign), when dealing with the sign of the

pseudo-wronskian, of interest would be the subcase where $c_2 = 0$. Here, the asymptotic development does not even ensure that \mathcal{W} is eventually negative. Enlarging the family of coefficients to the ones subjected to the restriction $\int_{t_0}^{+\infty} t^\varepsilon w(t)|a(t)|dt < +\infty$, where $\varepsilon \in [0, 1)$, the developments (3), (14) become

$$x(t) = ct + o(t^{1-\varepsilon}), \quad x'(t) = c + o(t^{-\varepsilon}), \quad c \in \mathbb{R}, \quad (15)$$

yielding the less precise estimate $\mathcal{W}(x, t) = o(t^{-\varepsilon})$ when $t \rightarrow +\infty$. We have again a lack of precision in the asymptotic development of $\mathcal{W}(x, \cdot)$ with respect to the sign issue. We also deduce on the basis of (3), (15) that some of the coefficients a in these classes verify (7), a fact that complicates the discussion.

The next result establishes the existence of a positive solution to (2) subjected to (4), (15) for the largest class of functional coefficients: $\varepsilon = 0$. By taking into account Lemmas 1, 2 and the non-oscillatory character of equation (2) when the nonlinearity w verifies (13), we conclude that *for an investigation within this class of coefficients a of the solutions with oscillatory pseudo-wronskian it is necessary that a itself oscillates*. Also, when a is non-negative valued we recall that the condition

$$\int_{t_0}^{+\infty} a(t)dt < +\infty$$

is necessary for the linear case of equation (2) to be non-oscillatory, see [16], while in the case given by $w(x) = x^\lambda$, $x \in \mathbb{R}$, with $\lambda > 1$ (such an equation is usually called an *Emden-Fowler equation*, see the monograph [19]) the condition

$$\int_{t_0}^{+\infty} ta(t)dt = +\infty \quad (16)$$

is necessary and sufficient for oscillation, see [4]. In the case of Emden-Fowler equations with $\lambda \in (0, 1)$ and a continuously differentiable coefficient a such that $a(t) \geq 0$ and $a'(t) \leq 0$ throughout $[t_0, +\infty)$, another result establishes that equation (2) has no oscillatory solutions provided that condition (16) fails, see [17].

Regardless of the oscillation of a , it is known [3, p. 360] that the linear case of equation (2) has bounded and positive solutions with eventually negative pseudo-wronskian.

Theorem 6 *Assume that the nonlinearity w verifies hypothesis (13) and is nondecreasing. Given $c, d > 0$, suppose that the functional coefficient a is*

nonnegative-valued, with eventual isolated zeros, and

$$\int_{t_0}^{+\infty} w(t)a(t)dt \leq \frac{d}{w(c+d)}.$$

Then, the equation (2) has a solution x such that $\mathcal{W}_0 = 0$,

$$c - d \leq x'(t) < \frac{x(t)}{t} \leq c + d \quad \text{for all } t > t_0 \quad (17)$$

and

$$\lim_{t \rightarrow +\infty} x'(t) = \lim_{t \rightarrow +\infty} \frac{x(t)}{t} = c. \quad (18)$$

Proof. We introduce the set D given by

$$D = \{u \in C([t_0, +\infty), \mathbb{R}) : ct \leq u(t) \leq (c+d)t \text{ for every } t \geq t_0\}.$$

A partial order on D is provided by the usual pointwise order " \leq ", that is, we say that $v_1 \leq v_2$ if and only if $v_1(t) \leq v_2(t)$ for all $t \geq t_1$, where $v_1, v_2 \in D$. It is not hard to see that (D, \leq) is a complete lattice.

For the operator $V : D \rightarrow C([t_0, +\infty), \mathbb{R})$ with the formula

$$V(u)(t) = t \left\{ c + \int_t^{+\infty} \frac{1}{s^2} \int_{t_0}^s \tau a(\tau) w(u(\tau)) d\tau ds \right\}, \quad u \in D, t \geq t_0,$$

the next estimates hold

$$\begin{aligned} c &\leq \frac{V(u)(t)}{t} = c + \int_t^{+\infty} \frac{1}{s^2} \int_{t_0}^s \tau a(\tau) \cdot w(\tau) w\left(\frac{u(\tau)}{\tau}\right) d\tau ds \\ &\leq c + \sup_{\xi \in [0, c+d]} \{w(\xi)\} \cdot \int_t^{+\infty} \frac{1}{s^2} \int_{t_0}^s \tau w(\tau) a(\tau) d\tau ds \\ &= c + w(c+d) \left[\frac{1}{t} \int_{t_0}^t \tau w(\tau) a(\tau) d\tau + \int_t^{+\infty} w(\tau) a(\tau) d\tau \right] \\ &\leq c + w(c+d) \int_{t_0}^{+\infty} w(\tau) a(\tau) d\tau \leq c + d \end{aligned}$$

by means of (13). These imply that $V(D) \subseteq D$.

Since $c \cdot t \leq V(c \cdot t)$ for all $t \geq t_0$, by applying the Knaster-Tarski fixed point theorem [12, p. 14], we deduce that the operator V has a fixed point u_0 in D . This is the pointwise limit of the sequence of functions $(V^n(c \cdot \text{Id}_I))_{n \geq 1}$, where $V^1 = V$, $V^{n+1} = V^n \circ V$ and $I = [t_0, +\infty)$.

We deduce that

$$u_0'(t) = [V(u_0)]'(t) = \frac{u_0(t)}{t} - \frac{1}{t} \int_{t_0}^t \tau a(\tau) w(u_0(\tau)) d\tau < \frac{u_0(t)}{t},$$

when $t > t_0$, and thus (17), (18) hold true.

The proof is complete. \square

5 The oscillatory integration of equation (2)

Let the continuous functional coefficient a with varying sign satisfy the restriction

$$\int_{t_0}^{+\infty} t^2 |a(t)| dt < +\infty.$$

We call the problem studied in the sequel *an oscillatory (asymptotic) integration* of equation (2).

Theorem 7 *Assume that w verifies (12), $w(0) = 0$ and there exists $c > 0$ such that*

$$L_+^c > 0 > L_-^c, \quad (19)$$

where

$$L_+^c = \limsup_{t \rightarrow +\infty} \frac{t \int_t^{+\infty} sw(cs)a(s)ds}{\int_t^{+\infty} s^2 |a(s)| ds}, \quad L_-^c = \liminf_{t \rightarrow +\infty} \frac{t \int_t^{+\infty} sw(cs)a(s)ds}{\int_t^{+\infty} s^2 |a(s)| ds}.$$

Then the equation (2) has a solution $x(t)$ with oscillatory pseudo-wronskian such that

$$x(t) = c \cdot t + o(1) \quad \text{when } t \rightarrow +\infty. \quad (20)$$

Proof. There exist $\eta > 0$ such that $L_+^c > \eta$, $L_-^c < -\eta$ and two increasing, unbounded from above sequences $(t_n)_{n \geq 1}$, $(t^n)_{n \geq 1}$ of numbers from $(t_0, +\infty)$ such that $t^n \in (t_n, t_{n+1})$ and

$$t_n \int_{t_n}^{+\infty} sw(cs)a(s)ds + k\eta \int_{t_n}^{+\infty} s^2 |a(s)| ds < 0 \quad (21)$$

and

$$t^n \int_{t^n}^{+\infty} sw(cs)a(s)ds - k\eta \int_{t^n}^{+\infty} s^2 |a(s)| ds > 0 \quad (22)$$

for all $n \geq 1$.

Assume further that

$$\int_{t_0}^{+\infty} \tau^2 |a(\tau)| d\tau \leq \frac{\eta}{k(c + \eta)}$$

and introduce the complete metric space $S = (D, \delta)$ given by

$$D = \{y \in C([t_0, +\infty), \mathbb{R}) : t|y(t)| \leq \eta \text{ for every } t \geq t_0\}$$

and

$$\delta(y_1, y_2) = \sup_{t \geq t_0} \{t|y_1(t) - y_2(t)|\}, \quad y_1, y_2 \in D.$$

For the operator $V : D \rightarrow C([t_0, +\infty), \mathbb{R})$ with the formula

$$V(y)(t) = \frac{1}{t} \int_t^{+\infty} sa(s)w \left(s \left[c - \int_s^{+\infty} \frac{y(\tau)}{\tau} d\tau \right] \right) ds, \quad y \in D, t \geq t_0,$$

the next estimates hold (notice that $|w(x)| \leq k|x|$ for all $x \in \mathbb{R}$)

$$t|V(y)(t)| \leq k \int_t^{+\infty} s^2|a(s)| \left[c + \eta \int_s^{+\infty} \frac{d\tau}{\tau^2} \right] ds \leq \eta \quad (23)$$

and

$$\begin{aligned} t|V(y_2)(t) - V(y_1)(t)| &\leq k \int_t^{+\infty} s^2|a(s)| \left(\int_s^{+\infty} \frac{d\tau}{\tau^2} \right) ds \cdot \delta(y_1, y_2) \\ &\leq \frac{k}{t_0} \int_t^{+\infty} s^2|a(s)| ds \leq \frac{\eta}{c + \eta} \cdot \delta(y_1, y_2). \end{aligned}$$

These imply that $V(D) \subseteq D$ and thus $V : S \rightarrow S$ is a contraction.

From the formula of operator V we notice also that

$$\lim_{t \rightarrow +\infty} tV(y)(t) = 0 \quad \text{for all } y \in D. \quad (24)$$

Given $y_0 \in D$ the unique fixed point of V , one of the solutions to (2) has the formula $x_0(t) = t \left[c - \int_t^{+\infty} \frac{y_0(s)}{s} ds \right]$ for all $t \geq t_0$. Via (24) and L'Hospital's rule, we provide also an asymptotic development for this solution, namely

$$\begin{aligned} \lim_{t \rightarrow +\infty} [x_0(t) - c \cdot t] &= - \lim_{t \rightarrow +\infty} t \int_t^{+\infty} \frac{y_0(s)}{s} ds = - \lim_{t \rightarrow +\infty} ty_0(t) \\ &= - \lim_{t \rightarrow +\infty} tV(y_0)(t) = 0. \end{aligned}$$

The estimate

$$\begin{aligned} \left| ty_0(t) - \int_t^{+\infty} sw(cs)a(s)ds \right| &\leq k \int_t^{+\infty} s^2|a(s)| \left[\int_s^{+\infty} \frac{|y_0(\tau)|}{\tau} d\tau \right] ds \\ &\leq k\eta \cdot \frac{1}{t} \int_t^{+\infty} s^2|a(s)| ds, \quad t \geq t_0, \end{aligned}$$

accompanied by (21), (22), leads to

$$y_0(t_n) = \mathcal{W}(x_0, t_n) < 0 \quad \text{and} \quad y_0(t^n) = \mathcal{W}(x_0, t^n) > 0. \quad (25)$$

The proof is complete. \square

Remark 1 When equation (2) is linear, that is $w(x) = x$ for all $x \in \mathbb{R}$, the formula (19) can be recast as

$$L_+ = \limsup_{t \rightarrow +\infty} \frac{t \int_t^{+\infty} s^2 a(s) ds}{\int_t^{+\infty} s^2 |a(s)| ds} > 0 > \liminf_{t \rightarrow +\infty} \frac{t \int_t^{+\infty} s^2 a(s) ds}{\int_t^{+\infty} s^2 |a(s)| ds} = L_-.$$

We claim that *for all $c \neq 0$ there exists a solution $x(t)$ with oscillatory pseudo-wronskian which verifies (20)*. In fact, replace c with c_0 in the formulas (21), (22) for a certain c_0 subjected to the inequality $\min\{L_+, -L_-\} > \frac{\eta}{c_0}$. It is obvious that, when $L_+ = -L_- = +\infty$, formulas (21), (22) hold for all c_0 , $\eta > 0$. Given $c \in \mathbb{R} - \{0\}$, there exists $\lambda \neq 0$ such that $c = \lambda c_0$. The solution of equation (2) that we are looking for has the formula $x = \lambda \cdot x_0$, where $x_0(t) = t \left[c_0 - \int_t^{+\infty} \frac{y_0(s)}{s} ds \right]$ for all $t \geq t_0$ and y_0 is the fixed point of operator V in D . Its pseudo-wronskian oscillates as a consequence of the obvious identity

$$\lambda \cdot \mathcal{W}(x_0, t) = \mathcal{W}(x, t), \quad t \geq t_0.$$

Example 1 An immediate example of functional coefficient a for the problem of linear oscillatory integration is given by $a(t) = t^{-2} e^{-t} \cos t$, where $t \geq 1$.

We have

$$\int_t^{+\infty} s^2 a(s) ds = \frac{1}{\sqrt{2}} \cos\left(t + \frac{\pi}{4}\right) e^{-t} \quad \text{and} \quad \int_t^{+\infty} s^2 |a(s)| ds \leq e^{-t}$$

throughout $[1, +\infty)$ which yields $L_+ = +\infty$, $L_- = -\infty$.

Sufficient conditions are provided now for an oscillatory pseudo-wronskian to be in $L^p((t_0, +\infty), \mathbb{R})$, where $p > 0$. Since $\lim_{t \rightarrow +\infty} \mathcal{W}(x, t) = 0$ for any solution $x(t)$ of equation (2) with the asymptotic development (20), (14), we are interested in the case $p \in (0, 1)$.

Theorem 8 *Assume that, in the hypotheses of Theorem 7, the coefficient a verifies the condition*

$$\int_{t_0}^{+\infty} \left[\frac{t}{\int_t^{+\infty} s^2 |a(s)| ds} \right]^{1-p} t^2 |a(t)| dt < +\infty \quad \text{for some } p \in (0, 1). \quad (26)$$

Then the equation (2) has a solution $x(t)$ with an oscillatory pseudo-wronskian in L^p and the asymptotic expansion (20).

Proof. Recall that y_0 is the fixed point of operator V . Then, formula (23) implies that

$$|y_0(t)| \leq k(c + \eta) \cdot \frac{1}{t} \int_t^{+\infty} s^2 |a(s)| ds, \quad t \geq t_0.$$

Via an integration by parts, we have

$$\begin{aligned} & \frac{1}{[k(c + \eta)]^p} \int_t^T |y_0(s)|^p ds \\ & \leq \frac{T^{1-p}}{1-p} \left[\int_T^{+\infty} s^2 |a(s)| ds \right]^p + \frac{p}{1-p} \int_t^T \left[\frac{s}{\int_s^{+\infty} \tau^2 |a(\tau)| d\tau} \right]^{1-p} s^2 |a(s)| ds \end{aligned}$$

for all $T \geq t \geq t_0$.

The estimates

$$\begin{aligned} & \frac{T^{1-p}}{1-p} \left[\int_T^{+\infty} s^2 |a(s)| ds \right]^p = \frac{T^{1-p}}{1-p} \int_T^{+\infty} \left[\frac{1}{\int_T^{+\infty} \tau^2 |a(\tau)| d\tau} \right]^{1-p} s^2 |a(s)| ds \\ & \leq \frac{1}{1-p} \int_T^{+\infty} \left[\frac{s}{\int_s^{+\infty} \tau^2 |a(\tau)| d\tau} \right]^{1-p} s^2 |a(s)| ds \end{aligned}$$

allow us to establish that

$$\frac{1}{[k(c + \eta)]^p} \int_t^T |y_0(s)|^p ds \leq \frac{1+p}{1-p} \int_t^{+\infty} \left[\frac{s}{\int_s^{+\infty} \tau^2 |a(\tau)| d\tau} \right]^{1-p} s^2 |a(s)| ds.$$

The conclusion follows by letting $T \rightarrow +\infty$.


The proof is complete. \square

Example 2 An example of functional coefficient a in the linear case that verifies the hypotheses of Theorem 8 is given by the formula

$$t^2 a(t) = b(t) = \begin{cases} a_k(t - 9k), & t \in [9k, 9k + 1], \\ a_k(9k + 2 - t), & t \in [9k + 1, 9k + 3], \\ a_k(t - 9k - 4), & t \in [9k + 3, 9k + 4], \\ a_k(9k + 4 - t), & t \in [9k + 4, 9k + 5], \\ a_k(t - 9k - 6), & t \in [9k + 5, 9k + 7], \\ a_k(9k + 8 - t), & t \in [9k + 7, 9k + 8], \\ 0, & t \in [9k + 8, 9(k + 1)], \end{cases} \quad k \geq 1.$$

Here, we take $a_k = k^{-\alpha} - (k + 1)^{-\alpha}$ for a certain integer $\alpha > \frac{2-p}{p}$.

To help the computations, the k -th "cell" of the function b can be visualized next.



It is easy to observe that

$$\int_{9k}^{9k+4} b(t)dt = \int_{9k+4}^{9k+8} b(t)dt = 0 \quad \text{for all } k \geq 1.$$

We have

$$\int_{9k+2}^{+\infty} b(t)dt = \int_{9k+2}^{9k+4} b(t)dt = -a_k, \quad \int_{9k+6}^{+\infty} b(t)dt = \int_{9k+6}^{9k+8} b(t)dt = a_k$$

and respectively

$$\int_{9k+2}^{+\infty} |b(t)|dt = 3a_k + 4 \sum_{m=k+1}^{+\infty} a_m, \quad \int_{9k+6}^{+\infty} |b(t)|dt = a_k + 4 \sum_{m=k+1}^{+\infty} a_m.$$

By noticing that

$$L_+ = \lim_{k \rightarrow +\infty} \frac{(9k+6) \int_{9k+6}^{+\infty} b(t)dt}{\int_{9k+6}^{+\infty} |b(t)|dt}, \quad L_- = \lim_{k \rightarrow +\infty} \frac{(9k+2) \int_{9k+2}^{+\infty} b(t)dt}{\int_{9k+2}^{+\infty} |b(t)|dt},$$

we obtain $L_+ = \frac{9\alpha}{4}$ and $L_- = -\frac{9\alpha}{4}$.

To verify the condition (26), notice first that

$$\begin{aligned} I_k &= \int_{9k}^{9(k+1)} \left[\frac{t}{\int_t^{+\infty} |b(s)|ds} \right]^{1-p} t^2 |a(t)|dt \\ &\leq \int_{9k}^{9(k+1)} \left[\frac{9(k+1)}{\int_{9(k+1)}^{+\infty} |b(s)|ds} \right]^{1-p} a_k dt, \quad k \geq 1. \end{aligned}$$

The elementary inequality $a_k \leq (2^\alpha - 1)(k+1)^{-\alpha}$ implies that

$$I_k \leq \frac{c_\alpha}{(k+1)^{(1+\alpha)p-1}}, \quad \text{where } c_\alpha = 9 \left(\frac{9}{4} \right)^{1-p} (2^\alpha - 1),$$

and the conclusion follows from the convergence of the series $\sum_{k \geq 1} (k+1)^{1-(1+\alpha)p}$.

Acknowledgement The author is indebted to Professor Ondrej Došly and to a referee for valuable comments leading to an improvement of the initial version of the manuscript. The author was financed during this research by the Romanian AT Grant 97GR/25.05.2007 with the CNCSIS code 100.

References

- [1] R.P. Agarwal, O.G. Mustafa, *A Riccatian approach to the decay of solutions of certain semi-linear PDE's*, Appl. Math. Lett. **20** (2007), 1206–1210.
- [2] R.P. Agarwal, S. Djebali, T. Moussaoui, O.G. Mustafa, Yu.V. Rogovchenko, *On the asymptotic behavior of solutions to nonlinear ordinary differential equations*, Asympt. Anal. **54** (2007), 1–50.
- [3] R.P. Agarwal, S. Djebali, T. Moussaoui, O.G. Mustafa, *On the asymptotic integration of nonlinear differential equations*, J. Comput. Appl. Math. **202** (2007), 352–376.
- [4] F.V. Atkinson, *On second order nonlinear oscillation*, Pacific J. Math. **5** (1955), 643–647.
- [5] M. Bartušek, Z. Došlá, J.R. Graef, *The nonlinear limit-point/limit-circle problem*, Birkhäuser, Boston, 2004.
- [6] M. Cecchi, M. Marini, Gab. Villari, *Integral criteria for a classification of solutions of linear differential equations*, J. Differential Equations **99** (1992), 381–397.
- [7] M. Cecchi, M. Marini, Gab. Villari, *Comparison results for oscillation of nonlinear differential equations*, Nonlinear Differ. Equ. Appl. (NoDEA) **6** (1999), 173–190.
- [8] A. Constantin, *Existence of positive solutions of quasilinear elliptic equations*, Bull. Austral. Math. Soc. **54** (1996), 147–154.
- [9] A. Constantin, *Positive solutions of quasilinear elliptic equations*, J. Math. Anal. Appl. **213** (1997), 334–339.
- [10] J. Deng, *Bounded positive solutions of semilinear elliptic equations*, J. Math. Anal. Appl. **336** (2007), 1395–1405.
- [11] S. Djebali, T. Moussaoui, O.G. Mustafa, *Positive evanescent solutions of nonlinear elliptic equations*, J. Math. Anal. Appl. **333** (2007), 863–870.

- [12] J. Dugundji, A. Granas, *Fixed point theory I*, Polish Sci. Publ., Warszawa, 1982.
- [13] M. Ehrnström, *Positive solutions for second-order nonlinear differential equations*, Nonlinear Anal. TMA **64** (2006), 1608–1620.
- [14] M. Ehrnström, O.G. Mustafa, *On positive solutions of a class of nonlinear elliptic equations*, Nonlinear Anal. TMA **67** (2007), 1147–1154.
- [15] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, Berlin, 2001.
- [16] P. Hartman, *On non-oscillatory linear differential equations of second order*, Amer. J. Math. **74** (1952), 389–400.
- [17] J.W. Heidel, *A nonoscillation theorem for a nonlinear second order differential equation*, Proc. Amer. Math. Soc. **22** (1969), 485–488.
- [18] M. Hesaaraki, A. Moradifam, *On the existence of bounded positive solutions of Schrödinger equations in two-dimensional exterior domains*, Appl. Math. Lett. **20** (2007), 1227–1231.
- [19] I.T. Kiguradze, T.A. Chanturia, *Asymptotic properties of solutions of nonautonomous ordinary differential equations*, Kluwer, Dordrecht, 1993.
- [20] O.G. Mustafa, Yu.V. Rogovchenko, *Global existence and asymptotic behavior of solutions of nonlinear differential equations*, Funkc. Ekvac. **47** (2004), 167–186.
- [21] O.G. Mustafa, Yu.V. Rogovchenko, *Limit-point type solutions of nonlinear differential equations*, J. Math. Anal. Appl. **294** (2004), 548–559.
- [22] O.G. Mustafa, *On the existence of solutions with prescribed asymptotic behaviour for perturbed nonlinear differential equations of second order*, Glasgow Math. J. **47** (2005), 177–185.
- [23] O.G. Mustafa, *Initial value problem with infinitely many linear-like solutions for a second order differential equation*, Appl. Math. Lett. **18** (2005), 931–934.
- [24] O.G. Mustafa, Yu.V. Rogovchenko, *Asymptotic integration of a class of nonlinear differential equations*, Appl. Math. Lett. **19** (2006), 849–853.