

Classification of Irreducible Weight Modules with a Finite-dimensional Weight Space over the Twisted Schrödinger-Virasoro Lie algebra

Junbo Li

Department of Mathematics, Shanghai Jiao Tong University Shanghai 200240, P. R. China;
Department of Mathematics, Changshu Institute of Technology, Changshu 215500, P. R. China
Email: sd_junbo@163.com

Yucai Su¹

Department of Mathematics, University of Science and Technology of China
Hefei 230026, P. R. China
Email: ycsu@ustc.edu.cn

Abstract. It is shown that the support of an irreducible weight module over the Schrödinger-Virasoro Lie algebra with an infinite-dimensional weight space, coincides with the weight lattice and that all nontrivial weight spaces of such a module are infinite-dimensional. As a side-product, it is obtained that every simple weight module over the Schrödinger-Virasoro Lie algebra with a nontrivial finite-dimensional weight space, is a Harish-Chandra module.

Key Words: The Schrödinger-Virasoro Lie algebra, weight modules, irreducible modules
Mathematics Subject Classification (2000): 17B10, 17B65, 17B68.

1. Introduction

The *twisted Schrödinger-Virasoro Lie algebra* is the infinite-dimensional Lie algebra \mathcal{L} with \mathbb{C} -basis $\{L_n, Y_m, M_p \mid m, n, p \in \mathbb{Z}\}$ and the following relations

$$[L_n, L_p] = (p - n)L_{n+p},$$

$$[L_n, Y_m] = (m - \frac{n}{2})Y_{n+m},$$

$$[L_n, M_p] = pM_{n+p},$$

$$[Y_m, Y_{m'}] = (m' - m)M_{m+m'},$$

$$[Y_m, M_p] = [M_n, M_p] = 0.$$

where $m, m', n, p \in \mathbb{Z}$. This infinite-dimensional Lie algebra is the twisted deformation of the Schrödinger-Virasoro Lie algebra, which was originally introduced in [3] in the context of non-equilibrium statistical physics, as a by-product of the computation of n -point functions that are covariant under the action of the Schrödinger group, containing as subalgebras both the Lie algebra of invariance of the free Schrödinger equation and the central charge-free Virasoro algebra. Both original and twisted Schrödinger-Virasoro Lie algebras are closely

¹Supported by NSF grants 10471091 of China and “One Hundred Talents Program” from University of Science and Technology of China

related to the Schrödinger Lie algebras and the Virasoro Lie algebra (cf. [2,5-7,10,11,14]). They should consequently play a role akin to that of the Virasoro Lie algebra in two-dimensional equilibrium statistical physics.

One attempt of introducing two-dimensional conformal field theory is to understand the universal behaviour of two-dimensional statistical systems at equilibrium and at the critical temperature, where they undergo a second-order phase transition. A systematic investigation of the theory of representations of the Virasoro algebra in the 80es led to introduce a class of physical models (called unitary minimal models), corresponding to the unitary highest weight representations of the Virasoro algebra with central charge less than one. Miraculously, covariance alone is enough to allow the computation of the statistic correlators (or so-called ‘ n -point functions’) for these highly constrained models.

Motivated by the search for deformations and central extensions of both original and twisted Schrödinger-Virasoro Lie algebras, Roger and Unterberger presented the sets of generators provided by the cohomology classes of the cocycles in the fifth part of [9]. In [12], Unterberger constructed vertex algebra representations of the Schrödinger-Virasoro Lie algebras out of a charged symplectic boson and a free boson and its associated vertex operators.

A *weight module* \mathcal{M} over \mathcal{L} is an \mathfrak{H} -diagonalizable module, where

$$\mathfrak{H} := \text{Span}_{\mathbb{C}}\{L_0, M_0\}$$

is the *Cartan subalgebra* of \mathcal{L} . If, in addition, \mathcal{M} is irreducible and all weight spaces \mathcal{M}_λ (cf. (1.1)) are finite-dimensional, the module is called a *Harish-Chandra module*.

If \mathcal{M} is an irreducible weight \mathcal{L} -module, then M_0 must act as some complex number $h_{\mathcal{M}}$ on \mathcal{M} . Furthermore, \mathcal{M} has the weight space decomposition

$$\mathcal{M} = \bigoplus_{\lambda \in \mathbb{C}} \mathcal{M}_\lambda, \quad \text{where } \mathcal{M}_\lambda = \{v \in \mathcal{M} \mid L_0 v = \lambda v\}, \quad (1.1)$$

where \mathcal{M}_λ is called a *weight space* with weight λ . Denote the set of *weights* λ of \mathcal{M} by

$$\text{Supp}(\mathcal{M}) := \{\lambda \in \mathbb{C} \mid \mathcal{M}_\lambda \neq 0\},$$

called the *support* of \mathcal{M} . Obviously, if \mathcal{M} is an irreducible weight \mathcal{L} -module, then there exists $\lambda \in \mathbb{C}$ such that $\text{Supp}(\mathcal{M}) \subset \lambda + \mathbb{Z}$.

An irreducible weight module \mathcal{V} is called a *pointed module* if there exists a weight $\lambda \in \mathbb{C}$ such that $\dim \mathcal{V}_\lambda = 1$. Xu posted the following natural problem in [13]:

Problem 1.1 *Is any irreducible pointed module over the Virasoro algebra a Harish-chandra module?*

An irreducible weight module \mathcal{V} is called a *mixed module* if there exist $\lambda \in \mathbb{C}$ and $i \in \mathbb{Z}$ such that $\dim \mathcal{V}_\lambda = \infty$ and $\dim \mathcal{V}_{\lambda+i} < \infty$. The following conjecture was posted in [7]:

Conjecture 1.2 *There are no irreducible mixed modules over the Virasoro algebra.*

Mazorchuk and Zhao [8] gave the positive answers to the above question and conjecture. In [10], the authors gave positive answers to the above question and conjecture for the twisted Heisenberg-Virasoro algebra. In this note, we also give the positive answers to the above question and conjecture for the Schrödinger-Virasoro Lie algebra. Our main result is the following:

Theorem 1.3 *Let \mathcal{M} be an irreducible weight \mathcal{L} -module. Assume that there exists $\lambda \in \mathbb{C}$ such that $\dim \mathcal{M}_\lambda = \infty$. Then $\text{Supp}(\mathcal{M}) = \lambda + \mathbb{Z}$ and for every $k \in \mathbb{Z}$, we have $\dim \mathcal{M}_{\lambda+k} = \infty$.*

Theorem 1.3 also implies the following classification of all irreducible weight \mathcal{L} -modules which admit a nontrivial finite dimensional weight space:

Corollary 1.4 *Let \mathcal{M} be an irreducible weight \mathcal{L} -module. Assume that there exists $\lambda \in \mathbb{C}$ such that $0 < \dim \mathcal{M}_\lambda < \infty$. Then \mathcal{M} is a Harish-Chandra module.*

2. Proof of Theorem 1.3

We first recall a main result about the weight Virasoro-modules in [8].

Theorem 2.1 *Let \mathcal{V} be an irreducible weight Virasoro-module. Assume that there exists $\lambda \in \mathbb{C}$, such that $\dim \mathcal{V}_\lambda = \infty$. Then $\text{Supp}(\mathcal{V}) = \lambda + \mathbb{Z}$ and for every $k \in \mathbb{Z}$, we have $\dim \mathcal{V}_{\lambda+k} = \infty$.*

Lemma 2.2 *Assume that there exists $\mu \in \mathbb{C}$ and a non-zero element $v \in \mathcal{M}_\mu$, such that*

$$\begin{aligned} L_1 v = L_2 v = Y_1 v = M_1 v = 0 \quad \text{or} \\ L_{-1} v = L_{-2} v = Y_{-1} v = M_{-1} v = 0. \end{aligned}$$

Then \mathcal{M} is a Harish-Chandra module.

Proof. Indeed, under these conditions, v is either a highest or a lowest weight vector and hence \mathcal{M} is a Harish-Chandra module (see, e.g. [6]). □

Assume now that \mathcal{M} is an irreducible weight \mathcal{L} -module such that there exists $\lambda \in \mathbb{C}$ satisfying $\dim \mathcal{M}_\lambda = \infty$.

Lemma 2.3 *There exists at most one $i \in \mathbb{Z}$ such that $\dim \mathcal{M}_{\lambda+i} < \infty$.*

Proof. Assume that $\dim \mathcal{M}_{\lambda+i} < \infty$ and $\dim \mathcal{M}_{\lambda+j} < \infty$ for different $i, j \in \mathbb{Z}$. Without loss of generality we may assume $i = 1$ and $j > 1$. Let \mathcal{V} be the following \mathcal{L} -module:

$$\begin{aligned} & \text{Ker}(L_1 : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+1}) \cap \text{Ker}(Y_1 : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+1}) \cap \text{Ker}(M_1 : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+1}) \\ & \cap \text{Ker}(L_j : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+j}) \cap \text{Ker}(Y_j : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+j}) \cap \text{Ker}(M_j : \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda+j}). \end{aligned} \quad (2.1)$$

Since

$$\dim \mathcal{M}_\lambda = \infty, \quad \dim \mathcal{M}_{\lambda+1} < \infty \quad \text{and} \quad \dim \mathcal{M}_{\lambda+j} < \infty,$$

all cokernels of the maps appearing in (2.1) are finite-dimensional. Thus

$$\dim \mathcal{V} = \infty. \tag{2.2}$$

Since

$$\begin{aligned} [L_1, L_p] &= (p-1)L_{p+1} \neq 0, \quad [L_1, M_r] = rM_{r+1} \neq 0 \quad \text{and} \\ [L_1, Y_q] &= \left(q - \frac{1}{2}\right)Y_{q+1} \neq 0 \end{aligned}$$

for $1 \neq p \in \mathbb{Z}$, $q \in \mathbb{Z}$, $1 \leq r \in \mathbb{Z}$, we get

$$\begin{aligned} L_p \mathcal{V} &= 0, \quad p = 1, j, j+1, j+2, \dots, \quad \text{and} \\ Y_q \mathcal{V} &= M_r \mathcal{V} = 0, \quad q, r = 1, 2, \dots. \end{aligned} \tag{2.3}$$

If there would exist $0 \neq v \in \mathcal{V}$ such that $L_2 v = 0$, then $Y_1 v = M_1 v = L_1 v = L_2 v = 0$ and \mathcal{M} would be a Harish-Chandra module by Lemma 2.2, a contradiction. Hence $L_2 v \neq 0$ for all $v \in \mathcal{V}$. In particular,

$$\dim L_2 \mathcal{V} = \infty.$$

Since $\dim M_{\lambda+1} < \infty$, there exists $0 \neq w \in L_2 \mathcal{V}$ such that $Y_{-1} w = M_{-1} w = L_{-1} w = 0$ (as in the proof of (2.2)). Let $w = L_2 u$ for some $u \in \mathcal{V}$. For all $k \geq j$, using (2.3), we have

$$L_k w = L_k L_2 u = L_2 L_k u + (2-k)L_{k+2} u = 0.$$

Hence $L_k w = 0$ for all $k = 1, j, j+1, j+2, \dots$. Since

$$\begin{aligned} [L_{-1}, L_l] &= (l+1)L_{l-1} \neq 0, \quad [L_l, M_{-1}] = M_{l-1} \neq 0 \quad \text{and} \\ [Y_{-1}, L_l] &= \left(1 + \frac{l}{2}\right)Y_{l-1} \neq 0 \end{aligned}$$

for all $l > 1$, we get inductively $L_k w = Y_k w = M_k w = 0$ for all $k = 1, 2, \dots$. Hence \mathcal{M} is a Harish-Chandra module by Lemma 2.2, again a contradiction. The lemma follows. \square

Because of Lemma 2.3, we can now fix the following notation: \mathcal{M} is an irreducible weight \mathcal{L} -module, $\mu \in \mathbb{C}$ is such that $\dim \mathcal{M}_\mu < \infty$ and $\dim \mathcal{M}_{\mu+i} = \infty$ for all $i \in \mathbb{Z} \setminus \{0\}$.

Lemma 2.4 *Let $0 \neq v \in \mathcal{M}_{\mu-1}$ be such that $Y_1 v = M_1 v = L_1 v = 0$. Then*

- (1) $(L_1^3 - 6L_2 L_1 + 6L_3)L_2 v = 0$,
- (2) $Y_k v = M_k v = 0$,
- (3) $Y_k L_2 v = M_k L_2 v = 0$,

where $k = 1, 2, \dots$.

Proof. (1) is the conclusion of Lemma 4 in [8].

(2) Since $Y_1v = M_1v = L_1v = 0$ and $[L_1, Y_l] = (l - \frac{1}{2})Y_{l+1} \neq 0$, $[L_1, M_l] = lM_{l+1} \neq 0$ for $l \geq 1$, we inductively get (2).

(3) follows from $Y_kL_2 = L_2Y_k - (k-1)Y_{k+2}$, $M_kL_2 = L_2M_k - kM_{k+2}$ and (2). \square

Proof of Theorem 1.3. Denote

$$\mathcal{V} := \text{Ker}\{L_1 : \mathcal{M}_{\mu-1} \rightarrow \mathcal{M}_\mu\} \cap \text{Ker}\{Y_1 : \mathcal{M}_{\mu-1} \rightarrow \mathcal{M}_\mu\} \cap \text{Ker}\{M_1 : \mathcal{M}_{\mu-1} \rightarrow \mathcal{M}_\mu\}.$$

Since $\dim \mathcal{M}_{\mu-1} = \infty$ and $\dim \mathcal{M}_\mu < \infty$, similar with the discussion of (2.2), we have $\dim \mathcal{V} = \infty$. For any $0 \neq v \in \mathcal{V}$, consider the element L_2v . By Lemma 2.2, $L_2v = 0$ would imply that \mathcal{M} is a Harish-Chandra module, a contradiction. Hence $L_2v \neq 0$, in particular, $\dim L_2\mathcal{V} = \infty$. This implies that there exists $w \in L_2\mathcal{V}$ such that $w \neq 0$ and $Y_{-1}w = M_{-1}w = L_{-1}w = 0$. Using

$$\begin{aligned} [L_{-1}, M_{-k+1}] &= (1-k)M_{-k} \quad \text{and} \\ [L_{-1}, Y_{-k+1}] &= \left(\frac{3}{2} - k\right)Y_{-k}, \end{aligned}$$

we obtain

$$Y_{-k}w = M_{-k}w = 0, \quad k = 1, 2, \dots.$$

From Lemma 2.4, we have $(L_1^3 - 6L_2L_1 + 6L_3)w = 0$, and $Y_kw = M_kw = 0$, $k = 1, 2, \dots$. Therefore,

$$\begin{aligned} Y_0w &= \frac{2}{3}[L_{-1}, Y_1]w = 0 \quad \text{and} \\ M_0w &= [L_{-1}, M_1]w = 0. \end{aligned}$$

The above relations mean that Y_k, M_k act trivially on \mathcal{M} for all $k \in \mathbb{Z}$. Therefore, \mathcal{M} is simply a module over the Virasoro algebra \mathcal{Vir} . Thus, Theorem 1.3 follows from Theorem 2.1. \square

Proof of Corollary 1.4. Assume that \mathcal{M} is not a Harish-Chandra module. Then there should exist $i \in \mathbb{Z}$ such that $\dim \mathcal{M}_{\lambda+i} = \infty$. In this case, Theorem 1.3 implies $\dim \mathcal{M}_\lambda = \infty$, a contradiction. Hence \mathcal{M} is a Harish-Chandra module, and the rest of the statement follows from [6]. \square

References

- [1] Arbarello, E., De Concini, C., Kac, V. G., Procesi, C.: Moduli spaces of curves and representation theory, *Comm. Math. Phys.*, **117**(1988), 1-36.
- [2] Dokovic, D., Zhao, K.: Derivations, isomorphisms and second cohomology of generalized Block algebras, *Algebra Colloquium*, **3**(1996), 245-272.

- [3] Henkel, M.: Schrödinger invariance and strongly anisotropic critical systems, *J. Stat. Phys.*, **75**(1994), 1023-1029.
- [4] Henkel, M. Phenomenology of local scale invariance: from conformal invariance to dynamical scaling, *Nucl. Phys. B*, **641**(2002), 405-410.
- [5] Henkel, M.; Unterberger, J.: Schrödinger invariance and space-time symmetries, *Nucl. Phys. B*, **660**(2003), 407-412.
- [6] Lu, R., Zhao, K.: Classification of irreducible weight modules over the twisted Heisenberg-Virasoro algebra, preprint (arXiv:math.ST/ 0510194).
- [7] Mazorchuk, V.: On simple mixed modules over the Virasoro algebra, *Mat. Stud.*, **22**(2004), 121-128.
- [8] Mazorchuk, V. and Zhao, K.: Classification of simple weight Virasoro modules with a finite-dimensional weight space, *J. Algebra*, **307**(2007), 209-214.
- [9] Roger, C., and Unterberger, J.: The Schrödinger-Virasoro Lie group and algebra: from geometry to representation theory, preprint (arXiv:cond-mat/0601050).
- [10] Shen, R., Su, Y.: Classification of irreducible weight modules with a finite-dimensional weight space over twisted Heisenberg-Virasoro algebra, *Acta Mathematica Sinica, English Series*, **23**(2007), 189-192.
- [11] Su, Y.: Simple modules over the higher rank super-Virasoro algebras, *Lett. Math. Phys.*, **53**(2000), 263-272.
- [12] Unterberger, J.: On vertex algebra representations of the Schrödinger-Virasoro algebra, preprint (arXiv:cond-mat/0703214).
- [13] Xu, X.: Pointed representations of Virasoro algebra, A Chinese summary appears in *Acta Math. Sinica, Chinese Series*, **40**(3), 479; *Acta Math. Sinica, New Series*, **13**(1997), 161-168.
- [14] Zhao, K.: Automorphisms of the algebra of q-difference operators, *Acta Mathematica Sinica, English Series*, **15**(2)(1999), 145-152.