

A ZARISKI-TYPE THEOREM IN RIEMANNIAN GEOMETRY

FUQUAN FANG AND SÉRGIO MENDONÇA

ABSTRACT. In this paper we study the topology of complement of totally geodesic submanifold in a positively curved manifold. We prove that if $K \subset (S^5, g)$ is a totally geodesic 3-submanifold where (S^5, g) has positive sectional curvature. Then K is a simple knot uniquely determined up to isotopy by the equivalence class of its Seifert matrices. Moreover, we obtain the following Zariski type connectivity theorem (compare [Fr2]):

Let M be an m -dimensional closed manifold of positive sectional curvature, and let H, N be closed embedded totally geodesic submanifolds of M which intersect transversely. Then the homomorphism induced by the inclusion

$$i_* : \pi_1(N - (N \cap H)) \rightarrow \pi_1(M - H)$$

is surjective, provided that $n \geq \frac{m}{2}$ and $m \geq 5$.

1. INTRODUCTION

A classical theorem of Frankel ([Fr1]) says that if M is a closed manifold with positive sectional curvature, and $f_i : N_i \rightarrow M, i = 1, 2$ are totally geodesic immersions of closed manifolds, with dimensions satisfying $\dim(N_1) + \dim(N_2) \geq \dim(M)$, then their images must intersect. This result was extended by Kenmotsu and Xia ([KX]) for the case of positive k -Ricci curvature if $\dim(N_1) + \dim(N_2) \geq \dim(M) + k - 1$. Another classical Frankel theorem is the following:

Theorem A. (Frankel [Fr2]) *Consider an m -dimensional closed manifold M of positive sectional curvature. Let N be an n -dimensional closed manifold and $f : N \rightarrow M$ be a totally geodesic immersion with $n \geq m/2$. Then the homomorphism induced by the inclusion $i_* : \pi_1(f(N)) \rightarrow \pi_1(M)$ is surjective.*

Theorem A was generalized by Kenmotsu and Xia ([KX]) for positive k -Ricci curvature if $n \geq (m + k - 1)/2$.

2000 *Mathematics Subject Classification.* Primary 53C42; Secondary 53C22.

Key words and phrases. fundamental group, positive curvature, totally geodesic.

The second named author was supported by NSF Grant of China #10671097 and the Capital Normal University.

Those results are the counterpart in Riemannian geometry of similar results in algebraic geometry, which suggests that several results in algebraic geometry could have similar results in Riemannian geometry. In these correspondences, the complex projective space is replaced by some kind of positivity of curvature, and algebraic subvarieties are replaced by totally geodesic immersions or by immersions with large asymptotic index (cf. [FMR]).

In [W] Wilking obtained in Riemannian geometry the corresponding Barth-Lefschetz-type theorem (cf. [Ba], [Bt], [L]). In [FMR] the authors generalized the Wilking Theorem and the Frankel Theorems cited above to a connectedness principle. As a consequence, it was obtained a regularity theorem, which says that a totally geodesic immersion of a closed n -dimensional manifold into a closed m -dimensional manifold of positive sectional curvature must be an embedding if $n > m/2$. This result is the counterpart of a theorem of Fulton-Hansen ([FH]). Another consequence says that the intersection of totally geodesic closed immersions of dimensions n_1 and n_2 in a closed m -dimensional manifold of positive sectional curvature must be connected if $n_1 + n_2 \geq m + 1$, which is a counterpart of the classical Bertini Theorem (cf. [Fu]).

In [FM] the authors extended the results in [FMR] to complex geometry of biholomorphic k -Ricci positive curvature. In [Fa1] the author extended the result in [FMR] to quaternionic Kähler manifolds with geometric applications. Another related result was [BR].

However, the classical Zariski Theorem relative to the topology of the complement of a codimension one subvariety in \mathbb{P}^m still doesn't have a Riemannian counterpart. We believe that the following conjecture should be true, which may be viewed as a counterpart of the classical Zariski theorem in projective algebraic geometry (compare [Wi], [FMR]).

Conjecture 1.1. *Let M be an m -dimensional Riemannian manifold of positive sectional curvature. If N, H are totally geodesic closed submanifolds of dimension n and h which intersects transversely, then the homomorphism*

$$j_* : \pi_i(N - N \cap H) \rightarrow \pi_i(M - H)$$

induced by inclusion is an isomorphism for $i \leq 2n - m$ and an epimorphism for $i = 2n - m + 1$.

The difficulty here is that the energy of curves on the complement of a submanifold is not anymore a proper function, which does not allow us to use the Morse theory method as in [SW], [W], [FMR], [FM], [Fa1] and [BR]. It seems that a use of the stratified Morse theory

developed in [GM] could be used to prove such a version, though we do not have a proper approach right now. Nevertheless, with respect to the fundamental group we obtained a counterpart, which extends Theorem A to complement of totally geodesic submanifold.

Theorem 1.2. *Let M be an m -dimensional Riemannian manifold of positive sectional curvature. If N, H are totally geodesic closed submanifolds of dimension n and h which intersect transversely, then the homomorphism induced by the inclusion*

$$i_* : \pi_1(N - (N \cap H)) \rightarrow \pi_1(M - H)$$

is surjective if $2n \geq m$ and $m \geq 5$.

We remark that the theorem is only interesting when $h = m - 2$, since for $h < m - 2$, the fundamental group $\pi_1(N - (N \cap H))$ (resp. $\pi_1(M - H)$) is isomorphic to $\pi_1(N)$ (resp. $\pi_1(M)$).

We say that a Riemannian manifold M has positive k -Ricci curvature if

$$\sum_{i=1}^k K(v, e_i) > 0,$$

where $v \in T_p M$ is any unit tangent vector, $K(v, e_i)$ is the sectional curvature associated to the plane generated by v and e_i , and v, e_1, \dots, e_k are orthonormal vectors. Now we can state the following generalization of Theorem 1.2, which holds with both definitions of positive k -Ricci curvature. Its proof is an easy adaptation of the proof of Theorem 1.2 in a way similar as [FMR], for example.

Corollary 1.3. *Let M be an m -dimensional Riemannian manifold of positive k -Ricci curvature. If N, H are totally geodesic closed submanifolds of dimension n and h which intersect transversely, then the homomorphism induced by the inclusion*

$$i_* : \pi_1(N - (N \cap H)) \rightarrow \pi_1(M - H)$$

is surjective if $n \geq \frac{m}{2} + k \geq 4 + k$ and $m \geq k + 4$.

As a typical application we will address to the following problem, which is the original motivation of the present paper:

Let $i : N \rightarrow (S^n, g)$ be a codimension 2 totally geodesic submanifold, where (S^n, g) is a sphere with positive sectional curvature. Could $i(N)$ be a nontrivial knot?

The problem is already very interesting for $n = 3$. In [Re] certain torus type knots were realized as simple geodesics in some (S^3, g) with positive sectional curvature.

The following result shows that totally geodesic knot in sphere with positive sectional curvature is rather special.

Theorem 1.4. *Let (S^{2n+1}, g) be a Riemannian manifold with positive sectional curvature. Assume $n \geq 2$. If $K \subset (S^{2n+1}, g)$ is a totally geodesic submanifold of codimension 2, then the knot complement $S^{2n+1} - K$ has fundamental group \mathbb{Z} .*

Theorem 1.4 shows that 3-knot in a positively curved 5-manifold homeomorphic to S^5 is a simple knot (c.f. [Le]). Hence the result of [Le] implies readily that

Corollary 1.5. *Let $K \subset (S^5, g)$ be a totally geodesic 3-submanifold where (S^5, g) has positive sectional curvature. Then K is a simple knot uniquely determined up to isotopy by the equivalence class of its Seifert matrices.*

We believe that Corollary 1.5 should hold for higher all dimensional knots in a positively curved sphere.

The fixed point sets of isometric group actions arise naturally as totally geodesic submanifolds. The famous Smith conjecture, confirmed by S.T. Yau, implies that the fixed point set of a periodic transformation on S^3 is indeed unknotted. In higher dimension the Smith conjecture is false (cf. Gordan [Go]). It seems natural to ask:

Conjecture 1.6. *Let $K \subset (S^{2n+1}, g)$ be a codimension 2 fixed point set of a cyclic isometry of order p . If (S^{2n+1}, g) has positive sectional curvature, then K is unknotted.*

2. PRELIMINARIES

Lemma 2.1. *Let M be a closed manifold of positive sectional curvature and let H be a totally geodesic embedded submanifold of M . Let m, h , respectively, be the dimensions of M, H . Let V be the ϵ -tubular open neighborhood of H for some small $\epsilon > 0$, with boundary ∂V and closure \bar{V} . Let $\gamma : [0, 1] \rightarrow M$ be a geodesic with $\gamma(t) \in M - \bar{V}$ for $t \in (0, 1)$ and $\gamma(0), \gamma(1) \in \partial V$. Assume that $\gamma'(0), \gamma'(1) \perp \partial V$ and that $h \geq m/2$. Then there exists a smooth variation γ_s of $\gamma = \gamma_0$ with $\gamma_s(0), \gamma_s(1) \in \partial V$ and $\gamma_s(t) \in M - \bar{V}$ for all $t \in (0, 1)$ and all s , such that the length $L(\gamma_s) < L(\gamma)$ if $s \neq 0$.*

Proof. For ϵ sufficiently small ∂V is contained in the 2ϵ -tubular neighborhood U of H . We recall that, for all $x \in U$, the gradient of the distance function ρ from H satisfies $\nabla \rho(x) = \sigma'(0)$, where $\sigma :$

$[0, \rho(x)] \rightarrow U$ is a minimal geodesic from H to x with $\sigma'(0) \perp H$ and $|\sigma'(\rho(x))| = 1$. Since $\gamma'(0), \gamma'(1) \perp \partial V$ and $\partial V = \rho^{-1}(\{\epsilon\})$, we have that $\gamma'(0) = \lambda_1 \nabla \rho(\gamma(0))$ and $\gamma'(1) = \lambda_2 \nabla \rho(\gamma(1))$. Thus the geodesic γ can be extended to a geodesic $\tilde{\gamma} : [-\epsilon, 1 + \epsilon] \rightarrow M$ with $p = \tilde{\gamma}(-\epsilon) \in H, q = \tilde{\gamma}(1 + \epsilon) \in H$ and $\tilde{\gamma}'(-\epsilon), \tilde{\gamma}'(1 + \epsilon) \perp H$. The parallel transport along $\tilde{\gamma}$ defines an isometric linear imbedding $P : T_p H \rightarrow T_q M$. Since $h \geq m/2$ and $(P(T_p H) + T_q H) \subset \{\tilde{\gamma}'(1 + \epsilon)\}^\perp$ we have

$$\dim(P(T_p H) \cap T_q H) \geq \dim(P(T_p H)) + \dim(T_q H) - \dim(P(T_p H) + T_q H) \geq 1,$$

hence there exists a nontrivial parallel vector field $v(t)$ along $\tilde{\gamma}$ with $v(-\epsilon) \in T_p H, v(1 + \epsilon) \in T_q H$. Thus the positivity of sectional curvature, the fact that H is totally geodesic and the second variation formula imply together that there exists a variation $\tilde{\gamma}_s$ of $\tilde{\gamma}$ with $\tilde{\gamma}_0 = \tilde{\gamma}$, $\tilde{\gamma}_s(-\epsilon), \tilde{\gamma}_s(1 + \epsilon) \in H$ and $L(\tilde{\gamma}_s) < L(\tilde{\gamma})$ if $s \neq 0$. There exist $-\epsilon < t_s < u_s < 1 + \epsilon$ such that $\tilde{\gamma}_s(t_s), \tilde{\gamma}_s(u_s) \in \partial V$ and the image of the restriction $\tilde{\gamma}_s|_{(t_s, u_s)}$ is contained in $M - \bar{V}$. From the transversality of the intersection between γ and ∂V we obtain that t_s, u_s depend smoothly on s . For $s \neq 0$ we have:

$$\begin{aligned} 2\epsilon + L(\gamma) &= L(\tilde{\gamma}) > L(\tilde{\gamma}_s) = L(\tilde{\gamma}_s|_{[-\epsilon, t_s]}) + L(\tilde{\gamma}_s|_{[t_s, u_s]}) + L(\tilde{\gamma}_s|_{[u_s, 1 + \epsilon]}) \\ &\geq 2\epsilon + L(\tilde{\gamma}_s|_{[t_s, u_s]}). \end{aligned}$$

Thus we have $L(\gamma) > L(\tilde{\gamma}_s|_{[t_s, u_s]})$. We define $\gamma_s = \tilde{\gamma}_s|_{[t_s, u_s]} \circ \sigma_s$ for some smooth change of parameters $\sigma_s : [0, 1] \rightarrow [t_s, u_s]$. This concludes the proof. \square

Lemma 2.2. *Let M, H, V, ϵ, h, m as in Lemma 2.1 with $h \geq m/2$. Then we have $\pi_1(M - V, \partial V) = 0$.*

Proof. A curve τ will be called a V -deformation of $\gamma : (I, \partial I) \rightarrow (M - V, \partial V)$ if there exists a homotopy $\tau_s : (I, \partial I) \rightarrow (M - V, \partial V)$ for all $s \in [0, 1]$ with $\tau_0 = \gamma$ and $\tau_1 = \tau$, where $I = [0, 1]$.

We assume by contradiction that there exists a nonempty set S of continuous curves γ as above so that each γ does not admit any V -deformation τ such that $\tau([0, 1]) \subset \partial V$. Let $a \geq 0$ be the infimum of the length $L(\gamma)$ for $\gamma \in S$.

Let U be a δ -tubular neighborhood of ∂V so that U strongly deformation retracts to ∂V for some small $\delta > 0$. Note that $a \geq 2\delta > 0$. Now consider a sequence of curves $\sigma_k \in S$ satisfying $L(\sigma_k) \rightarrow a$. Since $L(\sigma_k)$ is uniformly bounded, the curves σ_k form an equicontinuous sequence. So the Ascoli-Arzelà Theorem implies that, considering a subsequence

if necessary, we can assume that σ_k converges uniformly to a continuous curve $\sigma : [0, 1] \rightarrow M - V$. From the uniform convergence we have $L(\sigma) \leq a$.

Now we will show that $\sigma \in S$. By the compactness of $[0, 1]$ there exist normal open r_i -balls B_1, \dots, B_j with $r_i < \delta$ for all i and a partition $t_0 = 0 < t_1 < \dots < t_j = 1$ such that $\sigma([t_{i-1}, t_i]) \subset B_i$, for all i . For k sufficiently large we have $\sigma_k([t_{i-1}, t_i]) \subset B_i$, for all i . Thus there exists a homotopy σ^u between σ_k and σ with $\sigma^u([0, 1]) \subset \Omega = (M - V) \cup U$. Let $\eta : \Omega \rightarrow M - V$ be the continuous map which is the identity on $M - V$ and the standard projection $V \cap U \rightarrow \partial V$ on $V \cap U$. Thus we obtain the homotopy $\eta \circ \sigma^u : [0, 1] \rightarrow M - V$, hence σ is a V -deformation of σ_k and then $\sigma \in S$.

As a consequence we have $L(\sigma) = a$. By the minimality of $L(\sigma)$ and the first variation formula we know that σ is a geodesic satisfying $\sigma'(0), \sigma'(1) \perp \partial V$. So, by using Lemma 2.1 we obtain a V -deformation τ of σ with $L(\tau) < a$. This contradiction concludes the proof of Lemma 2.2. \square

3. PROOFS OF THEOREMS 1.2 AND 1.4

Proof of Theorem 1.2. If $h < m - 2$, it is well-known (by transversality theory) that $\pi_1(M - H) \cong \pi_1(M)$ (resp. $\pi_1(N - N \cap H) \cong \pi_1(N)$), the desired epimorphism follows from Theorem A.

If $h = m - 1$, by [W][FMR] one gets easily that $\pi_1(M) = 0$ or \mathbb{Z}_2 , and the universal covering of M (resp. H) is homeomorphic to a sphere. In the former case, $M - H$ is the union of two open discs, and in the latter case, $M - H$ is contractible. The desired result is trivial.

In the remainder of the proof let us assume $h = m - 2$. Let V be the open ε -tubular neighborhood defined in Lemma 1.1. By Lemma 2.2 $\pi_1(\partial V) \rightarrow \pi_1(M - V)$ is surjective. Note that ∂V is a S^1 -bundle over H , whose pullback on $N \cap H$ is isomorphic to the normal circle bundle of $N \cap H$ in N , by the transversality of N and H . By Wilking's theorem Theorem 0.6 of [FMR] we know that $N \cap H \subset H$ induces an epimorphism on the fundamental groups, provided $n \geq 3$ (since $m \geq 4$). Therefore, the exact sequence for the circle bundle $S^1 \rightarrow \partial V \cap N \rightarrow N \cap H$ (resp. $S^1 \rightarrow \partial V \rightarrow H$) implies easily that $\pi_1(\partial V \cap N) \rightarrow \pi_1(\partial V)$ is surjective. Hence, the inclusion $\partial V \cap N \rightarrow N - N \cap V \rightarrow M - V$ induces an epimorphism on the fundamental groups. The desired result follows. \square

Proof of Theorem 1.4. By the Wilking's theorem 0.6 of [FMR] we know easily that K is a homotopy sphere and hence homeomorphic to S^{2n-1} by Smale's theorem for $n \geq 3$ and Poincare-Perelman's theorem for

$n = 3$. Note that the normal circle bundle of K in S^{2n+1} is trivial whose fundamental group is isomorphic to \mathbb{Z} . By Lemma 2.2 it follows that $\pi_1(S^{2n+1} - K)$ is cyclic. By duality we conclude that

$$\pi_1(S^{2n+1} - K) \cong H_1(S^{2n+1} - K) \cong H^{2n}(S^{2n+1}, K) \cong \mathbb{Z}.$$

□

REFERENCES

- [Ba] Barth, W., *Transplanting cohomology classes in complex projective space*, Amer. J. Math. 92 (1970), 951-967.
- [Bt] Bott, R., *On a theorem of Lefschetz*, Mich. Math. J. 6 (1959), 211-216.
- [BR] Borisenko, A., Rovenski, V., *About topology of saddle submanifolds*, Diff. Geom. and its Applications, 25 (2007), 220-233.
- [BRT] A. Borisenko, M. L. Rabelo, K. Tenenblat, *On saddle submanifolds of Riemannian manifolds*, Geom. Dedicata, 67 (1997), 233-243.
- [Fa1] Fang, F., *Positive quaternionic Kähler manifolds and symmetry rank*, J. für die reine und angew. Math. 576 (2004), 149-165.
- [Fu] Fulton, W. *On the topology of algebraic varieties*, Proc. Symp. in Pure Math. 46 (1987), 15-46.
- [FH] Fulton, W., Hansen, J., *A connectedness theorem for projective varieties, with applications to intersections and singularities of mappings*, Ann. of Math. 110 (1979), 159-166.
- [FL] Fulton, W., Lazarsfeld R., *Connectivity and its applications in algebraic geometry*, Lect. Notes in Math., 862, Springer-Verlag, 26-92.
- [FM] Fang, F., Mendonça, S., *Complex immersions in Kähler manifolds of positive holomorphic k -Ricci curvature*, Trans. Amer. Math. Soc. 357 (2005), no. 9, 3725-3738.
- [GM] Goresky, M., MacPherson, R., *Stratified Morse theory*, Springer-Verlag, Berlin, Heidelberg, New York, 1988.
- [L] Lefschetz, S. *L'analysis situs et la géométrie algébrique*, Gauthier-Villars, Paris (1924).
- [Fr1] Frankel, T., *Manifolds of positive curvature*, Pacific J. Math., 11 (1961), 165-174.
- [Fr2] Frankel, T., *On the fundamental group of a compact minimal submanifold*, Ann. Math., 83 (1966), 68-73.
- [FMR] Fang, F., Mendonça, S., Rong, X., *A connectedness principle in the geometry of positive curvature*, Comm. Anal. Geom., 13 (2005), no. 4, 671-695.
- [Go] Gordon, C. McA., *On the higher-dimensional Smith conjecture*, Proc. London Math. Soc. 29 (1974), 98-110
- [KX] Kenmotsu K., Xia C., *Hadamard-Frankel type theorems for manifolds with partially positive curvature*, Pacific J. Math., 176 (1996), no. 4, 129-139.
- [Le] Levine, J., *An algebraic classification of some knots of codimension two*, Comment. Math. Helv. 45 (1970), 185-198
- [SW] Schoen, R., Wolfson J., *Theorems of Barth-Lefschetz types and Morse theory on the spaces of paths*, Math. Zeit. 229 (1998), 77-89.
- [Re] Reznikov, A., *Knotted totally geodesic submanifolds in positively curved spheres*, Mat. Fiz. Anal. Geom. 7 (2000), 458-463.

- [W] Wilking, B., *Torus actions on manifolds of positive sectional curvature*, Acta Math. 191 (2003), 259-297.

DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, BEIJING,
P.R. CHINA

E-mail address: fuquan_fang@yahoo.com

DEPARTAMENTO DE ANÁLISE, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE
FEDERAL FLUMINENSE, NITERÓI, RJ, CEP 24020-140, BRASIL

E-mail address: mendonca@mat.uff.br, sergiomendoncario@yahoo.com.br