

LIE ALGEBRAS WITH S_3 OR S_4 -ACTION, AND GENERALIZED MALCEV ALGEBRAS

ALBERTO ELDUQUE* AND SUSUMU OKUBO*

ABSTRACT. Lie algebras endowed with an action by automorphisms of any of the symmetric groups S_3 or S_4 are considered, and their decomposition into a direct sum of irreducible modules for the given action is studied.

In case of S_3 -symmetry, the Lie algebras are coordinatized by some nonassociative systems, which are termed *generalized Malcev algebras*, as they extend the classical Malcev algebras. These systems are endowed with a binary and a ternary products, and include both the Malcev algebras and the Jordan triple systems.

INTRODUCTION

The Tetrahedron algebra is an infinite dimensional Lie algebra which is endowed with a natural action by automorphisms of S_4 , the symmetric group of degree 4 (see [Eld07]). Lie algebras with such an action have been investigated by the authors in [EO07], where it is shown that these Lie algebras are coordinatized by a class of nonassociative algebras that include, as a very important case, the structurable algebras introduced by Allison [All78]. These algebras are very useful in providing models of simple Lie algebras, models which reflect a S_4 -symmetry (see [EO07] and the references there in).

In [IT07, Problem 20.10] the authors posed the question of how the Tetrahedron algebra decomposes into a direct sum of irreducible modules for S_4 . The first aim of this paper is to study the decomposition into direct sums of irreducible S_4 -modules of any Lie algebra endowed with an action of S_4 by automorphisms. This will be done in Section 1. The answer for the Tetrahedron algebra is particularly simple, as only the two irreducible three dimensional modules for S_4 live inside the Lie algebra. This will be checked in Section 2.

On the other hand, given any Lie algebra endowed with an action of S_4 by automorphisms, Klein's 4-group induces a grading of the Lie algebra over $\mathbb{Z}_2 \times \mathbb{Z}_2$. The $(\bar{0}, \bar{0})$ -component of this grading, that is, the subspace of elements fixed by any element in Klein's 4-group, is naturally endowed with an action of the symmetric group S_3 by automorphisms. Lie algebras with such an action are studied in Section 3, where the *generalized Malcev algebras* are introduced.

Date: January 15, 2008.

* Supported by the Spanish Ministerio de Educación y Ciencia and FEDER (MTM 2007-67884-C04-02) and by the Diputación General de Aragón (Grupo de Investigación de Álgebra).

* Supported in part by U.S. Department of Energy Grant No. DE-FG02-91ER40685.

As already noticed by Mikheev [Mik92] and Grishkov [Gri03], Malcev algebras appear as coordinate algebras of Lie algebras with an action of S_3 by automorphisms satisfying an extra condition, which in our terms translates into the fact that, as a module for S_3 , there is no submodule isomorphic to the alternating one dimensional module for S_3 . This will be the subject of Section 4, where the examples of Malcev algebras and Jordan triple systems are discussed. Different examples of Lie algebras with an action of S_3 by automorphisms and of their coordinate algebras will be given in the last Section 5.

Throughout the paper, all the vector spaces considered will be defined over a ground field k of characteristic $\neq 2, 3$. Unadorned tensor products will be defined over k .

Recall that the symmetric group S_4 is the semidirect product of Klein's 4-group $V_4 = \langle \tau_1, \tau_2 \rangle$ and the symmetric group $S_3 = \langle \varphi, \tau \rangle$. Here

$$\begin{aligned}\tau_1 &= (12)(34), \\ \tau_2 &= (23)(14), \\ \varphi &= (123) : 1 \mapsto 2 \mapsto 3 \mapsto 1 \\ \tau &= (12).\end{aligned}$$

The symmetric group S_3 appears both as a subgroup and as a quotient S_4/V_4 of the symmetric group S_4 .

There are exactly five non-isomorphic irreducible modules for the symmetric group S_4 (see, for instance, [FH91]):

- $U = ku$, the *trivial module*: $\sigma(u) = u$ for any $\sigma \in S_4$.
- $U' = ku'$, the *alternating module*: $\sigma(u') = (-1)^\sigma u'$ for any $\sigma \in S_4$, where $(-1)^\sigma$ denotes the signature of the permutation σ .
- $W = \{(\alpha_1, \alpha_2, \alpha_3) \in k^3 : \alpha_1 + \alpha_2 + \alpha_3 = 0\}$, the two dimensional irreducible module. This is a natural module for S_3 ($\sigma((\alpha_1, \alpha_2, \alpha_3)) = (\alpha_{\sigma^{-1}(1)}, \alpha_{\sigma^{-1}(2)}, \alpha_{\sigma^{-1}(3)})$) and hence a module for S_4 in which the elements of V_4 act trivially.
- $V = \{(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \in k^4 : \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = 0\}$, the *standard module*.
- $V' = U' \otimes V$.

The modules U , U' and W form a family of representatives of the isomorphism classes of the irreducible modules for S_3 .

1. LIE ALGEBRAS WITH S_4 -ACTION

Let \mathfrak{g} be a Lie algebra over our ground field k endowed with a group homomorphism

$$S_4 \rightarrow \text{Aut}(\mathfrak{g}).$$

As in [EO07], the action of Klein's 4-group V_4 gives a $\mathbb{Z}_2 \times \mathbb{Z}_2$ -grading on \mathfrak{g} :

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2, \tag{1.1}$$

where

$$\begin{aligned} \mathfrak{t} &= \{x \in \mathfrak{g} : \tau_1(x) = x, \tau_2(x) = x\} (= \mathfrak{g}_{(\bar{0}, \bar{0})}), \\ \mathfrak{g}_0 &= \{x \in \mathfrak{g} : \tau_1(x) = x, \tau_2(x) = -x\} (= \mathfrak{g}_{(\bar{1}, \bar{0})}), \\ \mathfrak{g}_1 &= \{x \in \mathfrak{g} : \tau_1(x) = -x, \tau_2(x) = x\} (= \mathfrak{g}_{(\bar{0}, \bar{1})}), \\ \mathfrak{g}_2 &= \{x \in \mathfrak{g} : \tau_1(x) = -x, \tau_2(x) = -x\} (= \mathfrak{g}_{(\bar{1}, \bar{1})}). \end{aligned} \quad (1.2)$$

(Here, the subindices 0, 1, 2 must be considered modulo 3.)

Assume that the action of V_4 is not trivial, as otherwise the S_4 -action is just an action of $S_3 \simeq S_4/V_4$, and let then A denote the subspace \mathfrak{g}_0 . For $x \in \mathfrak{g}_0$, define

$$\iota_0(x) = x \in \mathfrak{g}_0, \quad \iota_1(x) = \varphi(\iota_0(x)) \in \mathfrak{g}_1, \quad \iota_2(x) = \varphi^2(\iota_0(x)) \in \mathfrak{g}_2. \quad (1.3)$$

Thus

$$\mathfrak{g} = \mathfrak{t} \oplus \left(\bigoplus_{i=0}^2 \iota_i(A) \right).$$

As shown in [EO07], A becomes an algebra with involution $(A, \cdot, -)$ where:

- The involution (involutive antiautomorphism) is given by

$$\iota_0(\bar{x}) = -\tau(\iota_0(x))$$

for any $x \in A$. Note that $\tau(\mathfrak{g}_0) = \mathfrak{g}_0$ and that this gives immediately, since $\varphi\tau = \tau\varphi^2$, the following actions:

$$\tau(\iota_0(x)) = -\iota_0(\bar{x}), \quad \tau(\iota_1(x)) = -\iota_2(\bar{x}), \quad \tau(\iota_2(x)) = -\iota_1(\bar{x}). \quad (1.4)$$

- The multiplication is given by

$$\iota_0(\overline{x \cdot y}) = [\iota_1(x), \iota_2(y)], \quad (1.5)$$

for any $x, y \in A$.

Moreover, consider the space of *Lie related triples* (see [AF93]):

$$\mathfrak{ltt}(A, \cdot, -) = \{(d_0, d_1, d_2) \in \mathfrak{gl}(A)^3 :$$

$$\bar{d}_i(x \cdot y) = d_{i+1}(x) \cdot y + x \cdot d_{i+2}(y), \quad \forall x, y \in A, \quad \forall i = 0, 1, 2\},$$

where

$$\bar{d}(x) = \overline{d(\bar{x})}.$$

This space $\mathfrak{ltt}(A, \cdot, -)$ is a Lie algebra under componentwise bracket, and it comes endowed with the action of S_3 by automorphisms given by:

$$\begin{aligned} \varphi((d_0, d_1, d_2)) &= (d_2, d_0, d_1), \\ \tau((d_0, d_1, d_2)) &= (\bar{d}_0, \bar{d}_2, \bar{d}_1), \end{aligned} \quad (1.6)$$

which is compatible with the action of S_3 on \mathfrak{t} , that is,

$$\rho(\sigma(d)) = \sigma(\rho(d)),$$

for any $\sigma \in S_3$ and $d \in \mathfrak{t}$ (see [EO07, eqs. (1.8) and (2.3)]), where ρ is the linear map:

$$\begin{aligned} \rho : \mathfrak{t} &\longrightarrow \mathfrak{gl}(A)^3 \\ d &\mapsto (\rho_0(d), \rho_1(d), \rho_2(d)), \end{aligned}$$

with

$$\iota_i(\rho_i(d)(x)) = [d, \iota_i(x)]$$

for any $d \in \mathfrak{t}$ and $x \in A$. It turns out that ρ is a Lie algebra homomorphism and that $\rho(\mathfrak{t})$ is contained in $\mathfrak{ltt}(A, \cdot, -)$. Moreover, there appears a skew-symmetric bilinear map

$$\begin{aligned} \delta : A \times A &\longrightarrow \mathfrak{ltt}(A, \cdot, -) \\ (x, y) &\mapsto \rho([\iota_0(x), \iota_0(y)]) = (\delta_0(x, y), \delta_1(x, y), \delta_2(x, y)). \end{aligned}$$

Theorem 1.7. ([EO07, Theorem 2.4]) *For any $a, b, x, y, z \in A$ and $i, j = 0, 1, 2$:*

- (i) $[\delta_i(a, b), \delta_j(x, y)] = \delta_j(\delta_{i-j}(a, b)(x), y) + \delta_j(x, \delta_{i-j}(a, b)(y)),$
- (ii) $\delta_0(\bar{x}, y \cdot z) + \delta_1(\bar{y}, z \cdot x) + \delta_2(\bar{z}, x \cdot y) = 0,$
- (iii) $\delta_0(x, y)(z) + \delta_0(y, z)(x) + \delta_0(z, x)(y) = 0,$
- (iv) $\delta_1(x, y) = L_{\bar{y}}L_x - L_{\bar{x}}L_y,$
- (v) $\delta_2(x, y) = R_{\bar{y}}R_x - R_{\bar{x}}R_y,$
- (vi) $\overline{\delta_i(x, y)} = \delta_{-i}(\bar{x}, \bar{y})$ (or $\tau(\delta(x, y)) = \delta(\bar{x}, \bar{y})$),

where L_x and R_x denote, respectively, the left and right multiplication by x in the algebra (A, \cdot) .

The 4-tuple $(A, \cdot, -, \delta)$ is then called a *normal Lie related triple algebra* (see [Oku05]), or normal LRTA for short.

Conversely, given a normal LRTA $(A, \cdot, -, \delta)$, consider three copies of A : $\iota_i(A)$, $i = 0, 1, 2$, and the Lie subalgebra of *inner Lie related triples*:

$$\mathfrak{inltt}(A, \cdot, -, \delta) = \sum_{i=0}^2 \varphi^i(\delta(A, A)).$$

Then the vector space

$$\mathfrak{g}(A, \cdot, -, \delta) = \mathfrak{inltt}(A, \cdot, -, \delta) \oplus \left(\bigoplus_{i=0}^2 \iota_i(A)\right)$$

is a Lie algebra [EO07, Theorem 2.6] with bracket determined by:

- $\mathfrak{inltt}(A, \cdot, -, \delta)$ is a Lie subalgebra,
 - $[(d_0, d_1, d_2), \iota_i(x)] = \iota_i(d_i(x)), \forall (d_0, d_1, d_2) \in \mathfrak{inltt}(A, \cdot, -, \delta), \forall x \in A,$
 - $[\iota_i(x), \iota_{i+1}(y)] = \iota_{i+2}(\overline{x \cdot y}) \forall x, y \in A, \forall i = 0, 1, 2,$
 - $[\iota_i(x), \iota_i(y)] = \varphi^i(\delta(x, y)), \forall x, y \in A, \forall i = 0, 1, 2.$
- (1.8)

Moreover, the symmetric group S_4 acts naturally by automorphisms on $\mathfrak{g}(A, \cdot, -, \delta)$ by means of:

- V_4 acts trivially on $\mathfrak{inltt}(A, \cdot, -, \delta)$ and φ and τ act by (1.6),
 - $\varphi(\iota_i(x)) = \iota_{i+1}(x), \forall i = 0, 1, 2,$
 - $\tau(\iota_0(x)) = -\iota_0(\bar{x}), \tau(\iota_1(x)) = -\iota_2(\bar{x}), \tau(\iota_2(x)) = -\iota_1(\bar{x}),$
 - $\tau_1(\iota_0(x)) = \iota_0(x), \tau_1(\iota_i(x)) = -\iota_i(x)$ for $i = 1, 2,$
 - $\tau_2(\iota_1(x)) = \iota_1(x), \tau_2(\iota_i(x)) = -\iota_i(x)$ for $i = 0, 2,$
- (1.9)

for any $x \in A$.

One can also consider the larger Lie algebra

$$\tilde{\mathfrak{g}}(A, \cdot, -, \delta) = \mathfrak{ltt}(A, \cdot, -, \delta) \oplus \left(\bigoplus_{i=0}^2 \iota_i(A)\right) \quad (1.10)$$

with the same bracket given in (1.8), which again is endowed with an action of S_4 by automorphisms given by (1.9). Besides, if \mathfrak{g} is a Lie algebra with an action of S_4 and $(A, \cdot, -, \delta)$ is the associated normal LRTA, then the Lie algebra homomorphism $\rho : \mathfrak{t} \rightarrow \mathfrak{ltt}(A, \cdot, -)$ extends to a Lie algebra homomorphism

$$\tilde{\rho} : \mathfrak{g} \longrightarrow \tilde{\mathfrak{g}}(A, \cdot, -, \delta)$$

compatible with the action of S_4 , and such that $\tilde{\rho}(\iota_i(x)) = \iota_i(x)$ for any $x \in A$ and $i = 0, 1, 2$.

Proposition 1.11. *Let \mathfrak{g} be a Lie algebra with S_4 -action and let $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$ be the associated $\mathbb{Z}_2 \times \mathbb{Z}_2$ -grading as in (1.1). Assume that $\mathfrak{g}_0 \neq 0$ and that \mathfrak{g} has no proper ideals invariant under the action of S_4 (this happens, in particular, if \mathfrak{g} is simple). Then the homomorphism $\tilde{\rho}$ is one-to-one and its image is $\mathfrak{g}(A, \cdot, -, \delta)$.*

Proof. Since $\tilde{\rho}$ is a homomorphism of Lie algebras with S_4 -action, its kernel $\ker \tilde{\rho}$ is an ideal invariant under the action of S_4 , so it is trivial. Hence $\tilde{\rho}$ is one-to-one. On the other hand, the subalgebra of \mathfrak{g} generated by $\mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$ is an ideal invariant under S_4 , so it is the whole \mathfrak{g} , and hence $\tilde{\rho}(\mathfrak{g})$ is the subalgebra generated by $\bigoplus_{i=0}^2 \iota_i(A)$, which is precisely $\mathfrak{g}(A, \cdot, -, \delta)$. \square

Now we are ready to decompose the subspace $\mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$ in (1.1) as a direct sum of irreducible modules for the action of S_4 .

Theorem 1.12. *Let \mathfrak{g} be a Lie algebra endowed with an action of S_4 by automorphisms, and let $0 \neq x \in A = \mathfrak{g}_0$. Then:*

- (i) *If $\bar{x} = -x$, then $\sum_{i=0}^2 k\iota_i(x)$ is a S_4 -module isomorphic to the standard module V .*
- (ii) *If $\bar{x} = x$, then $\sum_{i=0}^2 k\iota_i(x)$ is a S_4 -module isomorphic to $V' = U' \otimes V$.*

Proof. Equations (1.2), (1.3) and (1.4) show that in both cases the subspace $\sum_{i=0}^2 k\iota_i(x)$ is invariant under the action of S_4 . In the first case the assignment

$$\iota_0(x) \mapsto (1, 1, -1, -1), \quad \iota_1(x) \mapsto (-1, 1, 1, -1), \quad \iota_2(x) \mapsto (1, -1, 1, -1),$$

provides the required isomorphism, while in the second case, the assignment $\iota_0(x) \mapsto u' \otimes (1, 1, -1, -1), \dots$, gives the result. \square

Corollary 1.13. *Let \mathfrak{g} be a Lie algebra endowed with an action of S_4 by automorphisms. Then $\bigoplus_{i=0}^2 \iota_i(A) = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$ is a direct sum of copies of the two irreducible three-dimensional modules for S_4 .*

Let us turn now our attention to the action of S_4 (actually of $S_3 \simeq S_4/V_4$) on the Lie algebra $\mathfrak{ltt}(A, \cdot, -, \delta)$.

On the two one-dimensional irreducible modules (the trivial one U and the alternating one U' for S_4 (or S_3)), the cycle φ acts trivially, while φ acts with minimal polynomial $X^2 + X + 1$ on the two-dimensional irreducible module. But for $(A, \cdot, -)$ an algebra with involution:

$$\begin{aligned} & \{(d_0, d_1, d_2) \in \mathfrak{ltt}(A, \cdot, -) : \varphi((d_0, d_1, d_2)) = (d_0, d_1, d_2)\} \\ &= \{(d, d, d) \in \mathfrak{gl}(A)^3 : \bar{d}(x \cdot y) = d(x) \cdot y + x \cdot d(y) \quad \forall x, y \in A\} \\ &= \{(d, d, d) \in \mathfrak{gl}(A)^3 : d(x * y) = d(x) * y + x * d(y) \quad \forall x, y \in A\} \end{aligned}$$

where $x * y = \overline{x \cdot y}$ for any $x, y \in A$. Hence, the subspace of fixed elements by φ in $\mathfrak{ltr}(A, \cdot, -)$ is naturally isomorphic to the Lie algebra of derivations of the algebra $(A, *)$:

$$\text{Fix}_{\mathfrak{ltr}(A, \cdot, -)}(\varphi) \simeq \mathfrak{der}(A, *).$$

Given an algebra with involution $(A, \cdot, -)$, $\mathfrak{der}(A, \cdot, -)$ will denote the Lie algebra:

$$\mathfrak{der}(A, \cdot, -) = \{d \in \mathfrak{der}(A, \cdot) : \bar{d} = d\}.$$

Proposition 1.14. *Let $(A, \cdot, -)$ be an algebra with involution. Then:*

- (i) *The direct sum of the trivial S_4 -submodules of $\mathfrak{ltr}(A, \cdot, -)$ is the subspace $\{(d, d, d) : d \in \mathfrak{der}(A, \cdot, -)\}$.*
- (ii) *The direct sum of the S_4 -submodules of $\mathfrak{ltr}(A, \cdot, -)$ isomorphic to the alternating module is $\{(d, d, d) : d \in \mathfrak{sder}(A, \cdot, -)\}$, where $\mathfrak{sder}(A, \cdot, -)$ is the subspace of skew-derivations of $(A, \cdot, -)$:*

$$\mathfrak{sder}(A, \cdot, -) = \{d \in \mathfrak{gl}(A) : d(x \cdot y) = -d(x) \cdot y - x \cdot d(y) \text{ and } \bar{d} = -d\}.$$

- (iii) *The direct sum of the irreducible S_4 -submodules of $\mathfrak{ltr}(A, \cdot, -)$ isomorphic to the two-dimensional irreducible module W is*

$$\{(d_0, d_1, d_2) \in \mathfrak{ltr}(A, \cdot, -) : d_0 + d_1 + d_2 = 0\}.$$

Proof. The direct sum of the trivial submodules is

$$\begin{aligned} \{(d_0, d_1, d_2) \in \mathfrak{ltr}(A, \cdot, -) : \varphi((d_0, d_1, d_2)) = (d_0, d_1, d_2) = \tau((d_0, d_1, d_2))\} \\ = \{(d_0, d_1, d_2) \in \mathfrak{ltr}(A, \cdot, -) : d_0 = d_1 = d_2 = \bar{d}_0\} \end{aligned}$$

because of (1.6), and this gives (i).

Also, the direct sum of the alternating submodules is

$$\begin{aligned} \{(d_0, d_1, d_2) \in \mathfrak{ltr}(A, \cdot, -) : \varphi((d_0, d_1, d_2)) = (d_0, d_1, d_2) = -\tau((d_0, d_1, d_2))\} \\ = \{(d_0, d_1, d_2) \in \mathfrak{ltr}(A, \cdot, -) : d_0 = d_1 = d_2 = -\bar{d}_0\}, \end{aligned}$$

and this proves (ii).

Finally, the direct sum of the S_4 -submodules of $\mathfrak{ltr}(A, \cdot, -)$ which are isomorphic to the two-dimensional irreducible module W is the kernel of $1 + \varphi + \varphi^2$, and this gives the result in (iii). \square

Let us pause to give an example of a normal LRTA $(A, \cdot, -, \delta)$ where $\mathfrak{der}(A, \cdot, -)$ is strictly contained in the Lie algebra of derivations $\mathfrak{der}(A, \cdot)$.

Example 1.15. Let \mathfrak{g} be any $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie algebra:

$$\mathfrak{g} = \mathfrak{g}_{(\bar{0}, \bar{0})} \oplus \mathfrak{g}_{(\bar{1}, \bar{0})} \oplus \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})},$$

with $\mathfrak{g}_{(\bar{0}, \bar{1})} \neq 0 \neq \mathfrak{g}_{(\bar{1}, \bar{1})}$, and let ν and μ the order two automorphisms of \mathfrak{g} given by:

$$\begin{aligned} \nu &= \begin{cases} id & \text{on } \mathfrak{g}_{(\bar{0}, \bar{0})} \oplus \mathfrak{g}_{(\bar{1}, \bar{0})}, \\ -id & \text{on } \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}, \end{cases} \\ \mu &= \begin{cases} id & \text{on } \mathfrak{g}_{(\bar{0}, \bar{0})} \oplus \mathfrak{g}_{(\bar{0}, \bar{1})}, \\ -id & \text{on } \mathfrak{g}_{(\bar{1}, \bar{0})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}. \end{cases} \end{aligned}$$

The symmetric group S_4 acts by automorphisms on \mathfrak{g}^3 as follows:

$$\begin{aligned}\tau_1((x, y, z)) &= (x, \nu(y), \nu(z)), \\ \tau_2((x, y, z)) &= (\nu(x), y, \nu(z)), \\ \varphi((x, y, z)) &= (z, x, y), \\ \tau((x, y, z)) &= (\mu(x), \mu(z), \mu(y)).\end{aligned}$$

(Compare to [EO07, Example 3.5].) Then the decomposition in (1.1) is given by:

$$\begin{aligned}\mathfrak{t} &= (\mathfrak{g}_{(\bar{0}, \bar{0})} \oplus \mathfrak{g}_{(\bar{1}, \bar{0})})^3, \\ \mathfrak{g}_0 &= \{(x, 0, 0) : x \in \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}\}, \\ \mathfrak{g}_1 &= \{(0, x, 0) : x \in \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}\}, \\ \mathfrak{g}_2 &= \{(0, 0, x) : x \in \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}\},\end{aligned}$$

so that $A = \mathfrak{g}_{(\bar{0}, \bar{1})} \oplus \mathfrak{g}_{(\bar{1}, \bar{1})}$ is a normal LRTA with

$$x \cdot y = 0, \quad \bar{x} = -\mu(x),$$

for any $x \in A$.

The Lie algebra of derivations of (A, \cdot) is then the whole general linear algebra:

$$\mathfrak{der}(A, \cdot) = \mathfrak{gl}(A),$$

while

$$\begin{aligned}\mathfrak{der}(A, \cdot, -) &= \{f \in \mathfrak{gl}(A) : f(\bar{x}) = \overline{f(x)} \forall x\} \\ &= \{f \in \mathfrak{gl}(A) : f\mu = \mu f\} \\ &= \{f \in \mathfrak{gl}(A) : f(\mathfrak{g}_{(\bar{0}, \bar{1})}) \subseteq \mathfrak{g}_{(\bar{0}, \bar{1})}, f(\mathfrak{g}_{(\bar{1}, \bar{1})}) \subseteq \mathfrak{g}_{(\bar{1}, \bar{1})}\},\end{aligned}$$

and

$$\begin{aligned}\mathfrak{sder}(A, \cdot, -) &= \{f \in \mathfrak{gl}(A) : f\mu = -\mu f\} \\ &= \{f \in \mathfrak{gl}(A) : f(\mathfrak{g}_{(\bar{0}, \bar{1})}) \subseteq \mathfrak{g}_{(\bar{1}, \bar{1})}, f(\mathfrak{g}_{(\bar{1}, \bar{1})}) \subseteq \mathfrak{g}_{(\bar{0}, \bar{1})}\}. \quad \square\end{aligned}$$

In [EO07, Theorem 2.4] it is proved that the unital normal LRTA's are precisely the structurable algebras. In this case, the previous Proposition becomes simpler.

Corollary 1.16. *Let $(A, \cdot, -)$ be a unital algebra with involution, then $\mathfrak{sder}(A, \cdot, -) = 0$.*

In particular, the alternating module does not appear in the decomposition of $\mathfrak{ltt}(A, \cdot, -)$ as a direct sum of irreducible S_3 -modules.

Proof. Any $d \in \mathfrak{sder}(A, \cdot, -)$ satisfies $\bar{d} = -d$ and

$$d(x \cdot y) = -d(x) \cdot y - x \cdot d(y)$$

for any $x, y \in A$. With $x = y = 1$ it follows that $d(1) = 0$, and then with $y = 1$ that $d(x) = -d(x)$ for any x , so $d = 0$. \square

Remark 1.17. Given an algebra with involution $(A, \cdot, -)$, the Lie algebra $\mathfrak{l} = \mathfrak{ltt}(A, \cdot, -)$ decomposes as

$$\mathfrak{l} = \mathfrak{l}_+ \oplus \mathfrak{l}_- \oplus \mathfrak{l}_2,$$

where \mathfrak{l}_+ is the subspace of fixed elements by the action of any $\sigma \in S_3$ (that is, the sum of the trivial S_3 -submodules of \mathfrak{l}), \mathfrak{l}_- is the direct sum of the alternating S_3 -submodules of \mathfrak{l} and \mathfrak{l}_2 is the sum of the S_3 -submodules of \mathfrak{l} which are isomorphic to the unique two-dimensional irreducible module for S_3 . Then any $(d_0, d_1, d_2) \in \mathfrak{l}$ decomposes as

$$(d_0, d_1, d_2) = (d, d, d) + (f, f, f) + (f_0, f_1, f_2),$$

where $(d, d, d) \in \mathfrak{l}_+$, $(f, f, f) \in \mathfrak{l}_-$ and $(f_0, f_1, f_2) \in \mathfrak{l}_2$ are given by:

$$\begin{aligned} d &= \frac{1}{6}((d_0 + d_1 + d_2) + (\bar{d}_0 + \bar{d}_1 + \bar{d}_2)), \\ f &= \frac{1}{6}((d_0 + d_1 + d_2) - (\bar{d}_0 + \bar{d}_1 + \bar{d}_2)), \\ f_i &= \frac{1}{3}(2d_i - d_{i+1} - d_{i+2}) \quad (\text{indices modulo } 3). \end{aligned}$$

Note that indeed $(1 + \varphi + \varphi^2)(f_0, f_1, f_2) = 0$, so $(f_0, f_1, f_2) \in \mathfrak{l}_2$ and, similarly, $(d, d, d) \in \mathfrak{l}_+$ and $(f, f, f) \in \mathfrak{l}_-$. \square

2. THE TETRAHEDRON ALGEBRA

The results in the previous section, together with [Eld07], give an immediate answer to [IT07, Problem 2.10], which asks for the decomposition of the Tetrahedron algebra into a direct sum of irreducible S_4 -modules.

Recall that the Tetrahedron algebra \mathfrak{g}_{\boxtimes} has been defined in [HT07] in connection with the so called Onsager algebra introduced in [Ons44], in which the free energy of the two dimensional Ising model was computed. The Tetrahedron algebra \mathfrak{g}_{\boxtimes} is the Lie algebra over k with generators

$$\{X_{ij} : i, j \in \{0, 1, 2, 3\}, i \neq j\}$$

and relations

$$\begin{aligned} X_{ij} + X_{ji} &= 0 \text{ for } i \neq j, \\ [X_{ij}, X_{jk}] &= 2(X_{ij} + X_{jk}) \text{ for mutually distinct } i, j, k, \\ [X_{hi}, [X_{hi}, [X_{hi}, X_{jk}]]] &= 4[X_{hi}, X_{jk}] \text{ for mutually distinct } h, i, j, k. \end{aligned}$$

Consider the basis $\{x = \begin{pmatrix} -1 & 2 \\ 0 & 1 \end{pmatrix}, y = \begin{pmatrix} -1 & 0 \\ -2 & 1 \end{pmatrix}, z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\}$ of the three dimensional simple Lie algebra \mathfrak{sl}_2 . Then [HT07, Proposition 6.5 and Theorem 11.5] the Tetrahedron algebra is isomorphic to the three-point loop algebra $\mathfrak{sl}_2 \otimes \mathcal{A}$, $\mathcal{A} = k[t, t^{-1}, (t-1)^{-1}]$, by means of the isomorphism

$$\Psi : \mathfrak{g}_{\boxtimes} \rightarrow \mathfrak{sl}_2 \otimes \mathcal{A}, \tag{2.1}$$

determined by:

$$\begin{aligned}\Psi(X_{12}) &= x \otimes 1, & \Psi(X_{23}) &= y \otimes 1, & \Psi(X_{31}) &= z \otimes 1, \\ \Psi(X_{03}) &= y \otimes t + z \otimes (t - 1), \\ \Psi(X_{01}) &= z \otimes t' + x \otimes (t' - 1), \\ \Psi(X_{02}) &= x \otimes t'' + y \otimes (t'' - 1),\end{aligned}$$

where

$$t' = 1 - t^{-1} \quad \text{and} \quad t'' = (1 - t)^{-1}. \quad (2.2)$$

The symmetric group S_4 embeds naturally in the group of automorphisms $\text{Aut}(\mathfrak{g} \boxtimes \mathcal{A})$ by means of

$$\sigma(X_{ij}) = X_{\sigma(i)\sigma(j)},$$

for any $\sigma \in S_4$ and $0 \leq i < j \leq 3$. Here we identify 0 and 4. On the other hand, S_4 embeds as a group of automorphisms of $\mathfrak{g} = \mathfrak{sl}_2 \otimes \mathcal{A}$ as follows ([Eld07, Theorem 1.4]):

(i) $\varphi = \varphi_{\mathfrak{s}} \otimes \varphi_{\mathcal{A}}$, where $\varphi_{\mathfrak{s}}$ is the order 3 automorphism of \mathfrak{sl}_2 given by

$$\varphi_{\mathfrak{s}}(x) = y, \quad \varphi_{\mathfrak{s}}(y) = z, \quad \varphi_{\mathfrak{s}}(z) = x,$$

and $\varphi_{\mathcal{A}}$ is the order 3 automorphism of the k -algebra \mathcal{A} determined by

$$\varphi_{\mathcal{A}}(t) = 1 - t^{-1} = t'.$$

(ii) $\tau = \tau_{\mathfrak{s}} \otimes \tau_{\mathcal{A}}$, where $\tau_{\mathfrak{s}}$ is the order 2 automorphism of \mathfrak{sl}_2 given by

$$\tau_{\mathfrak{s}}(x) = -x, \quad \tau_{\mathfrak{s}}(y) = -z, \quad \tau_{\mathfrak{s}}(z) = -y,$$

and $\tau_{\mathcal{A}}$ is the order 2 automorphism of \mathcal{A} determined by $\tau_{\mathcal{A}}(t) = 1 - t$.

(iii) τ_1 is the automorphism of \mathfrak{g} , as a Lie algebra over \mathcal{A} , given by

$$\begin{aligned}\tau_1(x \otimes 1) &= -x \otimes 1, \\ \tau_1(y \otimes 1) &= -(z \otimes t' + x \otimes (t' - 1)), \\ \tau_1(z \otimes 1) &= x \otimes t'' + y \otimes (t'' - 1).\end{aligned} \quad (2.3)$$

(iv) τ_2 is the automorphism of \mathfrak{g} , as a Lie algebra over \mathcal{A} , given by

$$\begin{aligned}\tau_2(x \otimes 1) &= y \otimes t + z \otimes (t - 1), \\ \tau_2(y \otimes 1) &= -y \otimes 1, \\ \tau_2(z \otimes 1) &= -(x \otimes t'' + y \otimes (t'' - 1)).\end{aligned} \quad (2.4)$$

With this action of S_4 , the isomorphism Ψ in (2.1) becomes an isomorphism of Lie algebras with S_4 -action.

A particular basis $\{u_0, u_1, u_2\}$ of $\mathfrak{g} = \mathfrak{sl}_2 \otimes \mathcal{A}$, as a module for \mathcal{A} , plays a key role in [Eld07]. It is defined by:

$$\begin{aligned}u_0 &= \frac{1}{4}\Psi(X_{02} + X_{31}), \\ u_1 &= \frac{1}{4}\Psi(X_{03} + X_{12}), \\ u_2 &= \frac{1}{4}\Psi(X_{01} + X_{23}),\end{aligned} \quad (2.5)$$

and satisfies [Eld07, Theorem 1.9]:

$$[u_0, u_1] = -u_2 t, \quad [u_1, u_2] = -u_0 t', \quad [u_2, u_0] = -u_1 t''. \quad (2.6)$$

Moreover, these elements also generate \mathfrak{g} as a Lie algebra over k . The action of S_4 on this \mathcal{A} -basis is given by:

$$\begin{aligned}\tau_1(u_0) &= u_0 = -\tau_2(u_0), \\ \tau_1(u_1) &= -u_1 = -\tau_2(u_1), \\ \tau_1(u_2) &= -u_2 = \tau_2(u_2), \quad (\text{see [Eld07, Theorem 2.2]}) \\ \varphi(u_i) &= u_{i+1}, \quad (\text{indices modulo 3, see [Eld07, (1.8)]}) \\ \tau(u_1) &= \frac{1}{4}\Psi(X_{03} - X_{12}) = -[u_0, u_1] = u_2t, \quad (\text{see [Eld07, Theorem 1.9]}) \\ \tau(u_2) &= \tau\varphi(u_1) = \varphi^2\tau(u_1) = u_1t'', \\ \tau(u_0) &= \tau\varphi^2(u_1) = \varphi\tau(u_1) = u_0t'.\end{aligned}$$

Besides, the subspaces \mathfrak{t} , \mathfrak{g}_0 , \mathfrak{g}_1 and \mathfrak{g}_2 in (1.2) become:

$$\mathfrak{t} = 0, \quad \mathfrak{g}_0 = u_0\mathcal{A}, \quad \mathfrak{g}_1 = u_1\mathcal{A}, \quad \mathfrak{g}_2 = u_2\mathcal{A},$$

and the associated normal LRTA can be identified to $(\mathcal{A}, \cdot, -, \delta)$ with (see [Eld07, Proposition 3.1]):

$$\begin{aligned}a \cdot b &= (\tau_{\mathcal{A}}\varphi_{\mathcal{A}}(a))(\tau_{\mathcal{A}}\varphi_{\mathcal{A}}^2(b)), \\ \bar{a} &= -t'\tau_{\mathcal{A}}(a).\end{aligned}$$

(Here, the usual multiplication of $\mathcal{A} = k[t, t^{-1}, (1-t)^{-1}]$ is denoted by juxtaposition.)

Then Corollary 1.13, together with $\mathfrak{t} = 0$, gives the answer to [IT07, Problem 2.10]:

Proposition 2.7. *The Tetrahedron algebra is a direct sum of copies of the two irreducible three-dimensional modules for S_4 .*

But a more precise statement can be given, by providing a concrete decomposition into a direct sum of irreducible S_4 -modules. To do so, and because of Theorem 1.12, let us first compute the elements $a \in \mathcal{A}$ with $\bar{a} = \pm a$. Note that for $a \in \mathcal{A}$,

$$\begin{aligned}\bar{a} = a &\Leftrightarrow a = -t'\tau_{\mathcal{A}}(a) \\ &\Leftrightarrow ta = \tau_{\mathcal{A}}(ta), \quad \text{as } tt't'' = -1 \text{ and } \tau_{\mathcal{A}}(t) = (t'')^{-1}.\end{aligned}$$

But, as $\mathcal{A} = k[t, t^{-1}, (1-t)^{-1}] \subseteq k(t)$, any $b \in \mathcal{A}$ is written uniquely as $b = \frac{p(t)}{(t(1-t))^s}$, with $p(t)$ a polynomial in $k[t]$ with either $p(0) \neq 0$ or $p(1) \neq 0$. Since $\tau_{\mathcal{A}}(t) = 1-t$, $\tau_{\mathcal{A}}(b) = b$ if and only if $\tau_{\mathcal{A}}(p(t)) = p(t)$ or, equivalently, $p(t) = p(1-t)$. This is equivalent to the condition $p(t) \in k[t(1-t)] = k[(t - \frac{1}{2})^2]$. Note that the set $\{(t(1-t))^r : r \in \mathbb{N} \cup \{0\}\}$ is a k -basis of the polynomial ring $k[t(1-t)]$.

In a similar vein, $\tau_{\mathcal{A}}(b) = -b$ if and only if $p(t) \in (2t-1)k[t(1-t)]$. Hence, for any $a \in \mathcal{A}$:

$$\begin{aligned}\bar{a} = a &\Leftrightarrow a \in t^{-1}k[t(1-t), (t(1-t))^{-1}], \\ \bar{a} = -a &\Leftrightarrow a \in t^{-1}(2t-1)k[t(1-t), (t(1-t))^{-1}].\end{aligned}$$

Therefore, Theorem 1.12 gives:

Theorem 2.8. *For any integer s ,*

- (i) $V'_s = \text{span} \{u_0 t^{-1}(t(1-t))^s, u_1 (t')^{-1}(t'(1-t'))^s, u_2 (t'')^{-1}(t''(1-t''))^s\}$ is isomorphic, as a module for S_4 , to $V' = U' \otimes V$.
- (ii) $V_s = \text{span} \{u_0 t^{-1}(2t-1)(t(1-t))^s, u_1 (t')^{-1}(2t'-1)(t'(1-t'))^s, u_2 (t'')^{-1}(2t''-1)(t''(1-t''))^s\}$ is isomorphic, as a module for S_4 , to the standard module V .
- (iii) The decomposition

$$\mathfrak{g} = \left(\bigoplus_{s \in \mathbb{Z}} V'_s \right) \oplus \left(\bigoplus_{s \in \mathbb{Z}} V_s \right)$$

is a decomposition of $\mathfrak{g} = \mathfrak{sl}_2 \otimes \mathcal{A}$ into a direct sum of irreducible S_4 -modules.

Proof. (i) and (ii) are direct consequences of Theorem 1.12. Also, as u_0, u_1 and u_2 form a basis of \mathfrak{g} over \mathcal{A} , and

$$\mathcal{A} = k[t, t^{-1}, (1-t)^{-1}] = k[t', (t')^{-1}, (1-t')^{-1}] = k[t'', (t'')^{-1}, (1-t'')^{-1}],$$

the result in (iii) follows. \square

Remark 2.9. The symmetric group S_3 acts on \mathcal{A} with the action of φ and τ given by $\varphi_{\mathcal{A}}$ and $\tau_{\mathcal{A}}$ above. As noticed in [HT07, Lemma 6.3], $\{1\} \cup \{t^n, (t')^n, (t'')^n : n \in \mathbb{N}\}$ is a k -basis of \mathcal{A} . Now, recall that $\varphi_{\mathcal{A}}$ permutes cyclically t, t' and t'' , while $\tau_{\mathcal{A}}(t) = 1-t$, so $\tau_{\mathcal{A}}(2t-1) = -(2t-1)$ and $\tau_{\mathcal{A}}(2t'-1) = \tau_{\mathcal{A}}\varphi_{\mathcal{A}}(2t-1) = \varphi_{\mathcal{A}}^2\tau_{\mathcal{A}}(2t-1) = -(2t''-1)$. Thus, for any $n \in \mathbb{N}$:

- $W_n = \{\alpha(2t-1)^n + \alpha'(2t'-1)^n + \alpha''(2t''-1)^n : \alpha, \alpha', \alpha'' \in k, \alpha + \alpha' + \alpha'' = 0\}$ is a two dimensional irreducible module for S_3 ,
- $U_n = k\left((2t-1)^{2(n-1)} + (2t'-1)^{2(n-1)} + (2t''-1)^{2(n-1)}\right)$ is a trivial module for S_3 ,
- $U'_n = k\left((2t-1)^{2n-1} + (2t'-1)^{2n-1} + (2t''-1)^{2n-1}\right)$ is an irreducible module for S_3 isomorphic to the alternating module,

and there appears the following decomposition into a direct sum of irreducible modules:

$$\mathcal{A} = \bigoplus_{n \in \mathbb{N}} (U_n \oplus U'_n \oplus W_n). \quad \square$$

The symmetric group S_4 acts on the set of generators $\{X_{ij} : i, j \in \{0, 1, 2, 3\}, i \neq j\}$ of the Tetrahedron algebra. The subspace spanned by these generators is easily shown to split into the direct sum of two non-isomorphic irreducible S_4 -modules (recall that the index 0 is identified to 4):

$$\begin{aligned} & \text{span} \{X_{\sigma(1)\sigma(2)} + X_{\sigma(2)\sigma(3)} + X_{\sigma(3)\sigma(4)} + X_{\sigma(4)\sigma(1)} : \sigma \in S_4\} \\ &= k(X_{01} + X_{12} + X_{23} + X_{30}) + k(X_{02} + X_{23} + X_{31} + X_{10}) \\ & \quad + k(X_{03} + X_{31} + X_{12} + X_{20}), \end{aligned}$$

and

$$\begin{aligned} & \text{span} \{X_{\sigma(1)\sigma(2)} - X_{\sigma(2)\sigma(3)} + X_{\sigma(3)\sigma(4)} - X_{\sigma(4)\sigma(1)} : \sigma \in S_4\} \\ &= k(X_{01} - X_{12} + X_{23} - X_{30}) + k(X_{02} - X_{23} + X_{31} - X_{10}) \\ & \quad + k(X_{03} - X_{31} + X_{12} - X_{20}). \end{aligned}$$

Note that

$$\begin{aligned} \frac{1}{4}\Psi(X_{01} + X_{12} + X_{23} + X_{30}) &= \frac{1}{4}\Psi(X_{01} + X_{23}) + \frac{1}{4}\Psi(X_{12} - X_{03}) \\ &= u_2 - u_2t = u_2(1 - t) = u_2(t'')^{-1}. \end{aligned}$$

(See the proof of [Eld07, Theorem 1.9].) By applying φ and the fact that Ψ is an isomorphism that commutes with the actions of S_4 , we obtain:

$$\begin{aligned} \frac{1}{4}\Psi(X_{01} + X_{12} + X_{23} + X_{30}) &= u_2(t'')^{-1} =: v_2, \\ \frac{1}{4}\Psi(X_{02} + X_{23} + X_{31} + X_{10}) &= u_0t^{-1} =: v_0, \\ \frac{1}{4}\Psi(X_{03} + X_{31} + X_{12} + X_{20}) &= u_1(t')^{-1} =: v_1, \end{aligned}$$

and, similarly,

$$\begin{aligned} \frac{1}{4}\Psi(X_{01} - X_{12} + X_{23} - X_{30}) &= u_2(1 + t) =: w_2, \\ \frac{1}{4}\Psi(X_{02} - X_{23} + X_{31} - X_{10}) &= u_0(1 + t') =: w_0, \\ \frac{1}{4}\Psi(X_{03} - X_{31} + X_{12} - X_{20}) &= u_1(1 + t'') =: w_1. \end{aligned}$$

Either by looking at Theorem 2.8 or by direct computation, it follows that $kv_0 + kv_1 + kv_2$ is isomorphic to the S_4 -module V' , and that $kw_0 + kw_1 + kw_2$ is isomorphic to the standard module V . Since $1 + t$, $1 + t'$ and $1 + t''$ are not invertible in \mathcal{A} , $\{w_0, w_1, w_2\}$ is not a basis of \mathfrak{g} over \mathcal{A} . On the other hand, $\{v_0, v_1, v_2\}$ is indeed a basis of $\mathfrak{g} = \mathfrak{sl}_2 \otimes \mathcal{A}$ over \mathcal{A} . However, v_0 , v_1 and v_2 fail to generate the whole Lie algebra \mathfrak{g} , as the following two results show.

Proposition 2.10. *The k -subalgebra generated by v_0 , v_1 and v_2 is $v_0\mathcal{S} \oplus v_1\mathcal{S} \oplus v_2\mathcal{S}$, where $\mathcal{S} = k[t(1 - t), t'(1 - t'), t''(1 - t'')]$.*

Proof. Since equation (2.6) shows that:

$$\begin{aligned} [v_1, v_2] &= [u_1(t')^{-1}, u_2(t'')^{-1}] = -u_0t'(t')^{-1}(t'')^{-1} \\ &= -u_0t^{-1}t(t'')^{-1} = -v_0t(1 - t) \in v_0\mathcal{S}, \end{aligned} \tag{2.11}$$

and by applying φ one obtains too:

$$[v_2, v_0] = -v_1t'(1 - t') \in v_1\mathcal{S}, \tag{2.12}$$

$$[v_0, v_1] = -v_2t''(1 - t'') \in v_2\mathcal{S}, \tag{2.13}$$

it follows that $v_0\mathcal{S} \oplus v_1\mathcal{S} \oplus v_2\mathcal{S}$ is a subalgebra of \mathfrak{g} , and therefore, the subalgebra generated by v_0 , v_1 and v_2 is contained in $v_0\mathcal{S} \oplus v_1\mathcal{S} \oplus v_2\mathcal{S}$.

Conversely, as

$$\begin{aligned} t(1 - t) &\xrightarrow{\varphi_{\mathcal{A}}} t'(1 - t') \xrightarrow{\varphi_{\mathcal{A}}} t''(1 - t''), \\ t(1 - t) &\xrightarrow{\tau_{\mathcal{A}}} t(1 - t), \end{aligned}$$

it follows that \mathcal{S} is invariant under the action of S_3 . Also, $kv_0 + kv_1 + kv_2$ is invariant under the action of S_4 , so the algebra generated by the v_i 's: $\text{alg}\langle v_0, v_1, v_2 \rangle$, is invariant too under the action of S_4 . In particular, the action of Klein's 4-group shows that $\text{alg}\langle v_0, v_1, v_2 \rangle = v_0\mathcal{S}_0 \oplus v_1\mathcal{S}_1 \oplus v_2\mathcal{S}_2$,

where \mathcal{S}_0 , \mathcal{S}_1 and \mathcal{S}_2 are subspaces of \mathcal{A} containing the unity 1. Also, the invariance under the action of φ shows that (since $\varphi(v_i) = v_{i+1}$) $\varphi_{\mathcal{A}}(\mathcal{S}_i) \subseteq \mathcal{S}_{i+1}$ for $i = 0, 1, 2$.

Equations (2.11), (2.12) and (2.13) give

$$t(1-t)\mathcal{S}_1\mathcal{S}_2 \subseteq \mathcal{S}_0, \quad t'(1-t')\mathcal{S}_2\mathcal{S}_0 \subseteq \mathcal{S}_1, \quad t''(1-t'')\mathcal{S}_0\mathcal{S}_1 \subseteq \mathcal{S}_2. \quad (2.14)$$

But

$$\begin{aligned} [v_1, [v_1, v_2]] &= [v_0, v_1]t(1-t) = -v_2t(1-t)t''(1-t'') = -v_2t(1-t'') \\ &= -v_2t(t')^{-1} = -v_2\frac{1}{t'(1-t')}, \end{aligned}$$

as $t'' = (1-t)^{-1}$ and cyclically. Therefore $\frac{1}{t'(1-t')}\mathcal{S}_1^2\mathcal{S}_2$ is contained in \mathcal{S}_2 . In particular, since 1 is in contained in the \mathcal{S}_i 's, one has:

$$\frac{1}{t'(1-t')}\mathcal{S}_1 \subseteq \mathcal{S}_2, \quad \frac{1}{t'(1-t')}\mathcal{S}_2 \subseteq \mathcal{S}_2. \quad (2.15)$$

Equations (2.14) and (2.15) give $t'(1-t')\mathcal{S}_2 \subseteq \mathcal{S}_1 \subseteq t'(1-t')\mathcal{S}_2$, so $\mathcal{S}_1 = t'(1-t')\mathcal{S}_2$ and

$$\mathcal{S}_2 = t'(1-t')\frac{1}{t'(1-t')}\mathcal{S}_2 \subseteq t'(1-t')\mathcal{S}_2 = \mathcal{S}_1.$$

Thus, $\mathcal{S}_2 \subseteq \mathcal{S}_1$ and, by applying $\varphi_{\mathcal{A}}$ one obtains $\mathcal{S}_2 \subseteq \mathcal{S}_1 \subseteq \mathcal{S}_0 \subseteq \mathcal{S}_2$. Therefore, $\mathcal{S}_0 = \mathcal{S}_1 = \mathcal{S}_2$ and also $t'(1-t')\mathcal{S}_2 = \mathcal{S}_2$. Now (2.14) shows that \mathcal{S}_2 is a subalgebra of \mathcal{A} , and \mathcal{S}_2 contains the element $t'(1-t')$, so it contains too, by invariance under the action of S_3 , the elements $t(1-t)$ and $t''(1-t'')$, which shows that $\mathcal{S}_0 = \mathcal{S}_1 = \mathcal{S}_2 \supseteq \mathcal{S}$, as required. \square

Proposition 2.16. \mathcal{S} is a maximal subalgebra of \mathcal{A} .

Proof. Recall that $\mathcal{S} = k[t(1-t), t'(1-t'), t''(1-t'')]$. Since

$$t(1-t)t'(1-t')t''(1-t'') = t(t'')^{-1}t't^{-1}t''(t')^{-1} = 1,$$

it follows that $(t(1-t))^{-1}$, $(t'(1-t'))^{-1}$ and $(t''(1-t''))^{-1}$ are all contained in \mathcal{S} and that the subspace of \mathcal{A} fixed by $\tau_{\mathcal{A}}$, which is $\mathcal{B} = k[t(1-t), (t(1-t))^{-1}]$ is contained in \mathcal{S} . Hence

$$\mathcal{S} = \mathcal{B} \oplus \{s \in \mathcal{S} : \tau_{\mathcal{A}}(s) = -s\}.$$

Since \mathcal{A} is an integral domain, the eigenspace of eigenvalue -1 for $\tau_{\mathcal{A}}$ is the free \mathcal{B} -module $\mathcal{B}(2t-1)$, so that $\{s \in \mathcal{S} : \tau_{\mathcal{A}}(s) = -s\}$ is a \mathcal{B} -submodule of $\mathcal{B}(2t-1)$. Let us check that $\{s \in \mathcal{S} : \tau_{\mathcal{A}}(s) = -s\}$ coincides with $\mathcal{B}(2t-1)(1-t(1-t))$. Since $\mathcal{B}(2t-1) = \mathcal{B}(2t-1)(1-t(1-t)) \oplus k(2t-1)$, this will show that \mathcal{S} is a codimension one subalgebra of \mathcal{A} , and hence maximal.

Note that

$$\begin{aligned}
(t-1)^3 &= (1-2t+t^2)(t-1) = (t-1) + (2-t)t(1-t) \\
&= \frac{1}{2} \left((2t-1) - 1 + (3-(2t-1))t(1-t) \right) \\
&= \frac{1}{2} \left((2t-1)(1-t(1-t)) - 1 + 3t(1-t) \right), \\
t^3 &= (t-1)^3 + 3t^2 - 3t + 1 = (t-1)^3 - 3t(1-t) + 1 \\
&= \frac{1}{2} \left((2t-1)(1-t(1-t)) + 1 - 3t(1-t) \right),
\end{aligned}$$

and hence

$$\begin{aligned}
t'(1-t') &= t't^{-1} = (1-t^{-1})t^{-1} = \frac{t-1}{t^2} = \frac{(t-1)^3}{(t(1-t))^2} \\
&= \frac{1}{2(t(1-t))^2} \left((2t-1)(1-t(1-t)) - 1 + 3t(1-t) \right), \\
t''(1-t'') &= t''(t')^{-1} = -\frac{t}{(1-t)^2} = -\frac{t^3}{(t(1-t))^2} \\
&= \frac{-1}{2(t(1-t))^2} \left((2t-1)(1-t(1-t)) + 1 - 3t(1-t) \right),
\end{aligned}$$

so $\mathcal{S} = k[t(1-t), t'(1-t'), t''(1-t'')] \subseteq \mathcal{B} \oplus \mathcal{B}(2t-1)(1-t(1-t))$ holds.

Reciprocally, \mathcal{B} is contained in \mathcal{S} , and $(2t-1)(1-t(1-t)) = 2t'(1-t')(t(1-t))^2 + 1 - 3t(1-t)$ belongs to \mathcal{S} . Hence we have $\mathcal{S} = \mathcal{B} \oplus \mathcal{B}(2t-1)(1-t(1-t))$, as required. \square

3. LIE ALGEBRAS WITH S_3 -ACTION

Given any normal LRTA, the Lie algebra $\tilde{\mathfrak{g}}(A, \cdot, -, \delta)$ in (1.10) is endowed with an action of the symmetric group S_4 with its Lie algebra of related triples being fixed by Klein's 4-group. Thus $\mathfrak{ltt}(A, \cdot, -)$ is endowed naturally with an action of S_3 , considered in Proposition 1.14.

This section is devoted to general Lie algebras endowed with an action of S_3 by automorphisms.

First of all, consider the irreducible S_3 -modules:

- $U = ku$, the trivial module: $\sigma(u) = u$ for any $\sigma \in S_3$,
- $U' = ku'$, the alternating module: $\sigma(u') = (-1)^\sigma u'$ for any $\sigma \in S_3$,
- $W = \{(\alpha, \beta, \gamma) \in k^3 : \alpha + \beta + \gamma = 0\}$, the two dimensional irreducible module. Here $W = kw_+ + kw_-$, with $w_+ = (-1, -1, 2)$ and $w_- = (3, -3, 0)$.

Note that there are the following obvious isomorphisms of S_3 -modules

$$U \otimes M \simeq M : u \otimes m \leftrightarrow m$$

for any S_3 -module M .

Also, as S_3 -modules, U can be identified to $k(1, 1, 1) \subseteq k^3$, so $U \oplus W = k^3$ (with the natural action $\sigma((\alpha_1, \alpha_2, \alpha_3)) = (\alpha_{\sigma^{-1}(1)}, \alpha_{\sigma^{-1}(2)}, \alpha_{\sigma^{-1}(3)})$) for any

$\alpha_1, \alpha_2, \alpha_3 \in k$). The componentwise multiplication in k^3 gives the following homomorphisms of S_3 -modules (with $u = (1, 1, 1)$):

$$\begin{aligned} W \otimes W &\rightarrow W & W \otimes W &\rightarrow U = ku \\ x \otimes y &\mapsto x \bullet y = \pi_W(xy), & x \otimes y &\mapsto (x | y)u = \pi_U(xy), \end{aligned} \quad (3.1)$$

where π_W and π_U denote the projections of k^3 onto W and U respectively. Thus, \bullet is a commutative product with

$$w_+ \bullet w_+ = w_+, \quad w_+ \bullet w_- = -w_-, \quad w_- \bullet w_- = -3w_+. \quad (3.2)$$

Note that (W, \bullet) is the para-Hurwitz algebra attached to the two dimensional composition algebra $k[z] = k1 + kz$ with $z^2 = -3$. That is, $x \bullet y = \bar{x}y$ for any $x, y \in k[z]$, where \bar{x} denotes the natural involution of $k[z]$, given by $\bar{z} = -z$. This will be exploited in Section 5.

Also, $(\cdot | \cdot)$ is a nondegenerate symmetric bilinear form with:

$$(w_+ | w_+) = 2, \quad (w_- | w_-) = 6, \quad (w_+ | w_-) = 0. \quad (3.3)$$

Moreover, the S_3 -modules $\wedge^2 W$ and U' are isomorphic, so with $U' = ku'$, there is the homomorphism of S_3 -modules

$$\begin{aligned} W \otimes W &\rightarrow U' \\ x \otimes y &\mapsto \langle x | y \rangle u', \end{aligned} \quad (3.4)$$

with $\langle \cdot | \cdot \rangle$ the alternating bilinear form with $\langle w_- | w_+ \rangle = 1$. Since $W \otimes W$ is isomorphic, as a S_3 -module, to $U \oplus U' \oplus W$, the homomorphisms in (3.1) and (3.4) are, up to scalars, the only possible such homomorphisms. Besides, there are unique, up to scalars, Clebsch-Gordan homomorphisms of S_3 -modules $\diamond : U' \otimes U' \rightarrow U$ and $\diamond : U' \otimes W \rightarrow W$ given by:

$$\begin{aligned} u' \otimes u' &\mapsto u' \diamond u' = -12u, \\ u' \otimes w_+ &\mapsto u' \diamond w_+ = 2w_-, \\ u' \otimes w_- &\mapsto u' \diamond w_- = -6w_+. \end{aligned} \quad (3.5)$$

Let \mathfrak{g} be a Lie algebra endowed with an action of S_3 by automorphisms. Then by complete reducibility, \mathfrak{g} is a direct sum of copies of U , U' and W . By joining together isomorphic summands, we may write:

$$\mathfrak{g} = (U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-) \oplus (W \otimes M), \quad (3.6)$$

for vector spaces \mathfrak{d}^+ , \mathfrak{d}^- and M .

Note that $(U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-)$ is the subalgebra of \mathfrak{g} consisting of the elements fixed by the action of $\varphi = (123)$, and it is graded over \mathbb{Z}_2 by means of the action of $\tau = (12)$, which is the identity on $U \otimes \mathfrak{d}^+$ and minus the identity on $U' \otimes \mathfrak{d}^-$. The remaining brackets in \mathfrak{g} are given, by S_3 -invariance,

by:

$$\begin{aligned}
[u \otimes d_1^+, u \otimes d_2^+] &= u \otimes [d_1^+, d_2^+], \\
[u \otimes d^+, u' \otimes d^-] &= u' \otimes [d^+, d^-], \\
[u' \otimes d_1^-, u' \otimes d_2^-] &= (u' \diamond u') \otimes [d_1^-, d_2^-] = -12u \otimes [d_1^-, d_2^-], \\
[u \otimes d^+, w \otimes x] &= w \otimes d^+(x), \\
[u' \otimes d^-, w \otimes x] &= (u' \diamond w) \otimes d^-(x), \\
[w_1 \otimes x, w_2 \otimes y] &= (w_1 | w_2)u \otimes d_{x,y}^+ + \langle w_1 | w_2 \rangle u' \otimes d_{x,y}^- + w_1 \bullet w_2 \otimes xy,
\end{aligned} \tag{3.7}$$

for any $d^\pm, d_1^\pm, d_2^\pm \in \mathfrak{d}^\pm$, $x, y \in M$ and $w, w_1, w_2 \in W$, where

$$\begin{aligned}
\mathfrak{d}^\pm \times \mathfrak{d}^\pm &\rightarrow \mathfrak{d}^\pm, & (d_1^\pm, d_2^\pm) &\mapsto [d_1^\pm, d_2^\pm], \\
\mathfrak{d}^+ \times \mathfrak{d}^- &\rightarrow \mathfrak{d}^-, & (d^+, d^-) &\mapsto [d^+, d^-], \\
\mathfrak{d}^\pm \times M &\rightarrow M, & (d^\pm, x) &\mapsto d^\pm(x), \\
M \times M &\rightarrow \mathfrak{d}^\pm, & (x, y) &\mapsto d_{x,y}^\pm, \\
M \times M &\rightarrow M, & (x, y) &\mapsto xy,
\end{aligned}$$

are bilinear maps. The skew-symmetry of the bracket in \mathfrak{g} forces $[\cdot, \cdot]$ to be skew-symmetric on its arguments, $d_{x,y}^+$ and xy to be skew-symmetric too, and $d_{x,y}^-$ to be symmetric on x, y .

In this situation consider the trilinear map:

$$\begin{aligned}
\{., ., .\} : M \times M \times M &\rightarrow M \\
(x, y, z) &\mapsto \{x, y, z\} = d_{x,y}^+(z) + d_{x,y}^-(z).
\end{aligned} \tag{3.8}$$

The next result shows that, essentially, the Lie algebra \mathfrak{g} is determined by the triple $(M, xy, \{x, y, z\})$.

Theorem 3.9. *Let \mathfrak{g} be a Lie algebra endowed with an action of S_3 by automorphisms, so that φ acts nontrivially. With the notations above, for any $a, b, x, y, z, t \in M$:*

$$(xy)z = \{x, z, y\} - \{y, z, x\}. \tag{3.10a}$$

$$\{a, b, xy\} = -\{b, a, x\}y - x\{b, a, y\}, \tag{3.10b}$$

$$\{a, b, \{x, y, z\}\} - \{x, y, \{a, b, z\}\} = \{\{a, b, x\}, y, z\} - \{x, \{b, a, y\}, z\}, \tag{3.10c}$$

(that is, $(M, \{., ., .\})$ is a generalized Jordan triple system [Kan73])

$$\{xy, z, t\} + \{yz, x, t\} + \{zx, y, t\} = 0 = \{x, yz, t\} + \{y, zx, t\} + \{z, xy, t\}, \tag{3.10d}$$

Conversely, let $(M, xy, \{x, y, z\})$ be a nonzero vector space endowed with an anticommutative bilinear product xy and a trilinear product $\{x, y, z\}$ satisfying the equations (3.10). Consider the vector space

$$\mathfrak{g}(M) = (U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-) \oplus (W \otimes M), \tag{3.11}$$

with

$$\mathfrak{d}^+ = \text{span} \{ \{x, y, \cdot\} - \{y, x, \cdot\} : x, y \in M \}$$

and

$$\mathfrak{d}^- = \text{span} \{ \{x, y, \cdot\} + \{y, x, \cdot\} : x, y \in M \}$$

(thus, \mathfrak{d}^+ and \mathfrak{d}^- are subspaces of the general Lie algebra $\mathfrak{gl}(M)$). Then the subspace $\mathfrak{d}^+ + \mathfrak{d}^- = \text{span}\{\{x, y, \cdot\} : x, y \in M\}$ is a Lie subalgebra of $\mathfrak{gl}(M)$, and the vector space $\mathfrak{g}(M)$, with the bracket defined by (3.7), where $d_{x,y}^+ = \frac{1}{2}(\{x, y, \cdot\} - \{y, x, \cdot\})$ and $d_{x,y}^- = \frac{1}{2}(\{x, y, \cdot\} + \{y, x, \cdot\})$, is a Lie algebra with an action of S_3 by automorphisms defined by the action of S_3 on U , U' and W .

Proof. Without loss of generality we may extend scalars if necessary and assume that $\omega \in k$, where $\omega^3 = 1 \neq \omega$. Then W contains the basis $\{w_\omega = (\omega^2, \omega, 1), w_{\omega^2} = (\omega, \omega^2, 1)\}$, whose elements satisfy

$$\varphi(w_\omega) = \omega w_\omega, \quad \varphi(w_{\omega^2}) = \omega^2 w_{\omega^2}, \quad \tau(w_\omega) = w_{\omega^2}.$$

Also,

$$w_\omega + w_{\omega^2} = w_+ \quad \text{and} \quad w_\omega - w_{\omega^2} = \frac{\omega^2 - \omega}{3} w_-,$$

so

$$w_\omega = \frac{1}{2}\left(w_+ + \frac{\omega^2 - \omega}{3} w_-\right), \quad w_{\omega^2} = \frac{1}{2}\left(w_+ - \frac{\omega^2 - \omega}{3} w_-\right),$$

and by equations (3.2) and (3.3):

$$\begin{aligned} w_\omega \bullet w_\omega &= w_{\omega^2}, & w_{\omega^2} \bullet w_{\omega^2} &= w_\omega, & w_\omega \bullet w_{\omega^2} &= 0, \\ (w_\omega | w_\omega) &= 0 = (w_{\omega^2} | w_{\omega^2}), & (w_\omega | w_{\omega^2}) &= 1, \end{aligned}$$

and

$$\langle w_\omega | w_{\omega^2} \rangle = \frac{1}{4} \langle w_+ + \frac{\omega^2 - \omega}{3} w_- | w_+ - \frac{\omega^2 - \omega}{3} w_- \rangle = \frac{\omega^2 - \omega}{6}.$$

Also,

$$\begin{aligned} u' \diamond w_\omega &= \frac{1}{2} u' \diamond \left(w_+ + \frac{\omega^2 - \omega}{3} w_- \right) \\ &= w_- + (\omega - \omega^2) w_+ = 2(\omega - \omega^2) w_\omega \\ u' \diamond w_{\omega^2} &= \frac{1}{2} u' \diamond \left(w_+ - \frac{\omega^2 - \omega}{3} w_- \right) \\ &= w_- + (\omega^2 - \omega) w_+ = -2(\omega - \omega^2) w_{\omega^2}. \end{aligned} \tag{3.12}$$

Take $\hat{u}' = \frac{1}{2(\omega - \omega^2)} u' = \frac{\omega^2 - \omega}{6} u'$. Then

$$\begin{aligned} \hat{u}' \diamond w_\omega &= w_\omega, & \hat{u}' \diamond w_{\omega^2} &= -w_{\omega^2}, \\ \hat{u}' \diamond \hat{u}' &= -12 \frac{1}{(2(\omega - \omega^2))^2} u = u. \end{aligned} \tag{3.13}$$

Consider also the new alternating bilinear form $\langle \cdot | \cdot \rangle : W \times W \rightarrow k$ given by $\langle w_\omega | w_{\omega^2} \rangle = 1$, that is, $\langle \cdot | \cdot \rangle = \frac{6}{\omega^2 - \omega} \langle \cdot | \cdot \rangle$.

Then the brackets in (3.7) become simpler:

$$\begin{aligned}
[u \otimes d_1^+, u \otimes d_2^+] &= u \otimes [d_1^+, d_2^+], \\
[u \otimes d^+, \hat{u}' \otimes d^-] &= \hat{u}' \otimes [d^+, d^-], \\
[\hat{u}' \otimes d_1^-, \hat{u}' \otimes d_2^-] &= u \otimes [d_1^-, d_2^-], \\
[u \otimes d^+, w \otimes x] &= w \otimes d^+(x), \\
[\hat{u}', w \otimes x] &= (\hat{u}' \diamond w) \otimes d^-(x), \\
[w_1 \otimes x, w_2 \otimes y] &= (w_1 | w_2)u \otimes d_{x,y}^+ + \langle w_1 | w_2 \rangle \hat{u}' \otimes d_{x,y}^- + w_1 \bullet w_2 \otimes xy,
\end{aligned} \tag{3.14}$$

for any $d^\pm, d_1^\pm, d_2^\pm \in \mathfrak{d}^\pm$, $x, y \in M$ and $w_1, w_2 \in W$. This shows in particular that the vector space $\mathfrak{d} = \mathfrak{d}^+ \oplus \mathfrak{d}^-$ is a \mathbb{Z}_2 -graded Lie algebra (with even part \mathfrak{d}^+ and odd part \mathfrak{d}^-) with the bracket given by the brackets $[d_1^\pm, d_2^\pm]$.

So assume that \mathfrak{g} is a Lie algebra with an action of S_3 by automorphisms, decomposition as in (3.6) and Lie bracket as in (3.14). Then for any $x, y, z \in M$,

$$\begin{aligned}
[[w_{\omega^2} \otimes x, w_{\omega^2} \otimes y], w_{\omega^2} \otimes z] &= [w_\omega \otimes xy, w_{\omega^2} \otimes z] \\
&= u \otimes d_{xy,z}^+ + \hat{u}' \otimes d_{xy,z}^-.
\end{aligned}$$

The Jacobi identity then shows that

$$d_{xy,z}^\pm + d_{yz,x}^\pm + d_{zx,y}^\pm = 0,$$

and this proves (3.10d). Now,

$$\begin{aligned}
[[w_{\omega^2} \otimes x, w_{\omega^2} \otimes y], w_\omega \otimes z] &= [w_\omega \otimes xy, w_\omega \otimes z] = w_{\omega^2} \otimes (xy)z, \\
[[w_{\omega^2} \otimes y, w_\omega \otimes z], w_{\omega^2} \otimes x] &= [u \otimes d_{y,z}^+ - \hat{u}' \otimes d_{y,z}^-, w_{\omega^2} \otimes x] \\
&= w_{\omega^2} \otimes d_{y,z}^+(x) - (\hat{u}' \diamond w_{\omega^2}) \otimes d_{y,z}^-(x) \\
&= w_{\omega^2} \otimes (d_{y,z}^+(x) + d_{y,z}^-(x)) \\
&= w_{\omega^2} \otimes \{y, z, x\}, \\
[[w_\omega \otimes z, w_{\omega^2} \otimes x], w_{\omega^2} \otimes y] &= [u \otimes d_{z,x}^+ + \hat{u}' \otimes d_{z,x}^-, w_{\omega^2} \otimes y] \\
&= w_{\omega^2} \otimes d_{z,x}^+(y) + (\hat{u}' \diamond w_{\omega^2}) \otimes d_{z,x}^-(y) \\
&= w_{\omega^2} \otimes (d_{z,x}^+(y) - d_{z,x}^-(y)) \\
&= -w_{\omega^2} \otimes (d_{x,z}^+(y) + d_{x,z}^-(y)) \\
&= -w_{\omega^2} \otimes \{x, z, y\},
\end{aligned}$$

since $d_{x,y}^+$ is skew-symmetric on x, y and $d_{x,y}^-$ is symmetric. Hence the Jacobi identity shows that $(xy)z + \{y, z, x\} - \{x, z, y\} = 0$, which gives (3.10a).

Now, for any $d_1^+, d_2^+ \in \mathfrak{d}^+$ and $z \in M$:

$$\begin{aligned}
[[u \otimes d_1^+, u \otimes d_2^+], w_\omega \otimes z] &= [u \otimes [d_1^+, d_2^+], w_\omega \otimes z] \\
&= w_\omega \otimes [d_1^+, d_2^+](z), \\
[u \otimes d_1^+, [u \otimes d_2^+, w_\omega \otimes z]] &= w_\omega \otimes d_1^+(d_2^+(z)), \\
[u \otimes d_2^+, [u \otimes d_1^+, w_\omega \otimes z]] &= w_\omega \otimes d_2^+(d_1^+(z)).
\end{aligned}$$

Hence, the bilinear map $\mathfrak{d}^+ \times M \rightarrow M$, $(d^+, x) \mapsto d^+(x)$ is a representation of the Lie algebra \mathfrak{d}^+ . In the same vein, one proves:

$$\begin{aligned} [d^+, d^-](z) &= d^+(d^-(z)) - d^-(d^+(z)), \\ [d_1^-, d_2^-](z) &= d_1^-(d_2^-(z)) - d_2^-(d_1^-(z)), \end{aligned}$$

for any $d^\pm, d_1^\pm, d_2^\pm \in \mathfrak{d}^\pm$, so the linear map:

$$\begin{aligned} \rho : \mathfrak{d}^+ \oplus \mathfrak{d}^- &\rightarrow \mathfrak{gl}(M) \\ d^+ + d^- &\mapsto \rho(d^+ + d^-) : z \mapsto d^+(z) + d^-(z), \end{aligned}$$

is a representation of the Lie algebra $\mathfrak{d}^+ \oplus \mathfrak{d}^-$.

Finally, for any $d^+ \in \mathfrak{d}^+$, $w_1, w_2 \in W$ and $x, y \in M$:

$$\begin{aligned} &[u \otimes d^+, [w_1 \otimes x, w_2 \otimes y]] \\ &= [u \otimes d^+, (w_1 | w_2)u \otimes d_{x,y}^+ + \langle w_1 | w_2 \rangle \hat{u}' \otimes d_{x,y}^- + (w_1 \bullet w_2) \otimes xy] \\ &= (w_1 | w_2)u \otimes [d^+, d_{x,y}^+] + \langle w_1 | w_2 \rangle \hat{u}' \otimes [d^+, d_{x,y}^-] + (w_1 \bullet w_2) \otimes d^+(xy), \\ &[[u \otimes d^+, w_1 \otimes x], w_2 \otimes y] \\ &= [w_1 \otimes d^+(x), w_2 \otimes y] \\ &= (w_1 | w_2)u \otimes d_{d^+(x),y}^+ + \langle w_1 | w_2 \rangle \hat{u}' \otimes d_{d^+(x),y}^- + (w_1 \bullet w_2) \otimes d^+(x)y \\ &[w_1 \otimes x, [u \otimes d^+, w_2 \otimes y]] \\ &= [w_1 \otimes x, w_2 \otimes d^+(y)] \\ &= (w_1 | w_2)u \otimes d_{x,d^+(y)}^+ + \langle w_1 | w_2 \rangle \hat{u}' \otimes d_{x,d^+(y)}^- + (w_1 \bullet w_2) \otimes xd^+(y). \end{aligned}$$

Therefore,

$$d^+(xy) = d^+(x)y + xd^+(y) \quad \text{and} \quad [d^+, d_{x,y}^\pm] = d_{d^+(x),y}^\pm + d_{x,d^+(y)}^\pm, \quad (3.15)$$

for any $x, y \in M$ and $d^+ \in \mathfrak{d}^+$. In a similar vein, for any $d^- \in \mathfrak{d}^-$, $w_1, w_2 \in W$ and $x, y \in M$:

$$\begin{aligned} &[\hat{u}' \otimes d^-, [w_1 \otimes x, w_2 \otimes y]] \\ &= [\hat{u}' \otimes d^-, (w_1 | w_2)u \otimes d_{x,y}^+ + \langle w_1 | w_2 \rangle \hat{u}' \otimes d_{x,y}^- + (w_1 \bullet w_2) \otimes xy] \\ &= (w_1 | w_2)\hat{u}' \otimes [d^-, d_{x,y}^+] + \langle w_1 | w_2 \rangle \hat{u}' \otimes [d^-, d_{x,y}^-] \\ &\quad + (\hat{u}' \diamond (w_1 \bullet w_2)) \otimes d^-(xy), \\ &[[\hat{u}' \otimes d^-, w_1 \otimes x], w_2 \otimes y] \\ &= [(\hat{u}' \diamond w_1) \otimes d^-(x), w_2 \otimes y] \\ &= (\hat{u}' \diamond w_1 | w_2)u \otimes d_{d^-(x),y}^+ + \langle \hat{u}' \diamond w_1 | w_2 \rangle \hat{u}' \otimes d_{d^-(x),y}^- \\ &\quad + ((\hat{u}' \diamond w_1) \bullet w_2) \otimes d^-(x)y \\ &[w_1 \otimes x, [\hat{u}' \otimes d^-, w_2 \otimes y]] \\ &= [w_1 \otimes x, (\hat{u}' \diamond w_2) \otimes d^-(y)] \\ &= (w_1 | \hat{u}' \diamond w_2)u \otimes d_{x,d^-(y)}^+ + \langle w_1 | \hat{u}' \diamond w_2 \rangle \hat{u}' \otimes d_{x,d^-(y)}^- \\ &\quad + (w_1 \bullet (\hat{u}' \diamond w_2)) \otimes xd^-(y). \end{aligned}$$

But notice that $\langle w_1 | w_2 \rangle = (\hat{u}' \diamond w_1 | w_2) = -(w_1 | \hat{u}' \diamond w_2)$ and $(w_1 | w_2) = \langle \hat{u}' \diamond w_1 | w_2 \rangle = -\langle w_1 | \hat{u}' \diamond w_2 \rangle$. Also $\hat{u}' \diamond (w_1 \bullet w_2) = -(\hat{u}' \diamond w_1) \bullet w_2 = -w_1 \bullet (\hat{u}' \diamond w_2)$, so we conclude:

$$d^-(xy) = -d^-(x)y - xd^-(y) \quad \text{and} \quad [d^-, d_{x,y}^\pm] = d_{d^-(x),y}^\mp - d_{x,d^-(y)}^\mp, \quad (3.16)$$

for any $x, y \in M$ and $d^- \in \mathfrak{d}^+$.

Since $d_{x,y}^\pm = \mp d_{y,x}^\pm$ for any $x, y \in M$, equations (3.15) and (3.16) show that, for any $a, b, x, y \in M$:

$$\begin{aligned} \{a, b, xy\} &= d_{a,b}^+(xy) + d_{a,b}^-(xy) \\ &= d_{a,b}^+(x)y + xd_{a,b}^+(y) - d_{a,b}^-(x)y - xd_{a,b}^-(y) \\ &= -(d_{b,a}^+(x) + d_{b,a}^-(x))y - x(d_{b,a}^+(y) + d_{b,a}^-(y)) \\ &= -\{b, a, x\}y - x\{b, a, y\}, \end{aligned}$$

thus obtaining (3.10b), and

$$\begin{aligned} [d_{a,b}^+ + d_{a,b}^-, d_{x,y}^+ + d_{x,y}^-] &= d_{d_{a,b}^+(x),y}^+ + d_{x,d_{a,b}^+(y)}^+ + d_{d_{a,b}^-(x),y}^- + d_{x,d_{a,b}^-(y)}^- \\ &\quad + d_{d_{a,b}^-(x),y}^- - d_{x,d_{a,b}^-(y)}^- + d_{d_{a,b}^+(x),y}^+ - d_{x,d_{a,b}^+(y)}^+ \\ &= d_{\{a,b,x\},y}^+ + d_{\{a,b,x\},y}^- - d_{x,\{b,a,y\}}^+ - d_{x,\{b,a,y\}}^-, \end{aligned}$$

and this gives (3.10c).

The converse follows by straightforward computations of the Jacobi identity. \square

The reason behind the next definition will become clear in Theorem 4.1.

Definition 3.17. A *generalized Malcev algebra* is a triple $(M, xy, \{x, y, z\})$, where M is a vector space, $(x, y) \mapsto xy$ a bilinear anticommutative product, and $(x, y, z) \mapsto \{x, y, z\}$ a trilinear product on M , satisfying the conditions in (3.10). The Lie algebra $\mathfrak{g}(M)$ in (3.11), with the action of S_3 in Theorem 3.9, is called the *associated Lie algebra*.

Therefore, Theorem 3.9 asserts that there is a generalized Malcev algebra attached to any Lie algebra with S_3 -action, assuming that the cycle φ acts nontrivially.

Remark 3.18. Let \mathfrak{g} be a Lie algebra endowed with an action of S_3 by automorphisms, so that φ acts nontrivially, as in Theorem 3.9, and assume that $\omega \in k$, where $\omega^3 = 1 \neq \omega$. Then \mathfrak{g} decomposes as $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_\omega \oplus \mathfrak{g}_{\omega^2}$, where $\mathfrak{g}_\mu = \{x \in \mathfrak{g} : \varphi(x) = \mu x\}$ for $\mu = 1, \omega, \omega^2$. Then in the decomposition in (3.6) we have:

$$\mathfrak{g}_1 = (U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-), \quad \mathfrak{g}_\omega \oplus \mathfrak{g}_{\omega^2} = W \otimes M.$$

Besides, $\varphi(w_\omega) = \omega w_\omega$ and $\varphi(w_{\omega^2}) = \omega^2 w_{\omega^2}$, so $\mathfrak{g}_\omega = w_\omega \otimes M$ and $\mathfrak{g}_{\omega^2} = w_{\omega^2} \otimes M$. Also, for any $x, y, z \in M$:

$$\begin{aligned} [\tau(w_\omega \otimes x), \tau(w_\omega \otimes y)] &= [w_{\omega^2} \otimes x, w_{\omega^2} \otimes y] = w_\omega \otimes xy, \\ [[w_\omega \otimes x, \tau(w_\omega \otimes y)], w_\omega \otimes z] &= [[w_\omega \otimes x, w_{\omega^2} \otimes y], w_\omega \otimes z] \\ &= [u \otimes d_{x,y}^+ + \hat{u}' \otimes d_{x,y}^-, w_\omega \otimes z] \\ &= w_\omega \otimes \{x, y, z\}, \end{aligned}$$

because of (3.12), (3.13) and (3.14). Therefore M can be identified to \mathfrak{g}_ω ($x \leftrightarrow w_\omega \otimes x$) with:

$$\begin{cases} xy = [\tau(x), \tau(y)], \\ \{x, y, z\} = [[x, \tau(y)], z], \end{cases}$$

for any $x, y, z \in \mathfrak{g}_\omega$.

Conversely, given a generalized Malcev algebra $(M, xy, \{x, y, z\})$, the associated Lie algebra $\mathfrak{g}(M)$ can be identified to

$$\mathfrak{d}^+ \oplus \mathfrak{d}^- \oplus \nu_0(M) \oplus \nu_1(M),$$

where $\nu_i(M)$ denotes a copy of M (think of $\nu_0(x)$ as $w_\omega \otimes x$ and $\nu_1(x)$ as $w_{\omega^2} \otimes x$, and identify $u \otimes d^+$ to d^+ and $\hat{u}' \otimes d^-$ to d^-) with Lie bracket given by:

- $[d_1^\pm, d_2^\pm]$ is given by the usual Lie bracket in $\mathfrak{gl}(M)$,
- $[d^+, \nu_i(x)] = \nu_i(d^+(x))$,
- $[d^-, \nu_i(x)] = (-1)^i \nu_i(d^-(x))$,
- $[\nu_i(x), \nu_i(y)] = \nu_{1-i}(xy)$,
- $[\nu_0(x), \nu_1(y)] = d_{x,y}^+ + d_{x,y}^-$,

for any $d_1^\pm, d_2^\pm, d^\pm \in \mathfrak{d}^\pm$, $x \in M$, and $i = 0, 1$. □

It can be shown that some of the conditions (3.10d) in Definition 3.17 are redundant:

Proposition 3.19. *Let $(M, xy, \{x, y, z\})$ be a triple where M is a vector space, $(x, y) \mapsto xy$ a bilinear anticommutative product, and $(x, y, z) \mapsto \{x, y, z\}$ a trilinear product on M , satisfying the conditions (3.10a), (3.10b) and (3.10c). Then:*

- (i) *For any $x, y, z, t \in M$, the equation*

$$\{xy, z, t\} + \{yz, x, t\} + \{zx, y, t\} = 0$$

holds. (This is the first part of (3.10d).)

- (ii) *If either $M = M^2$ or $\{x \in M : xM^2 = 0\} = 0$ hold, then for any $x, y, z, t \in M$,*

$$\{x, yz, t\} + \{y, zx, t\} + \{z, xy, t\} = 0$$

holds too. (This is the second part of (3.10d).)

Proof. For any $x, y, z, t \in M$, equations (3.10a) and (3.10b), together with the anticommutativity of the bilinear product give:

$$\begin{aligned} ((xy)t)z &= \{xy, z, t\} - \{t, z, xy\} \\ &= \{xy, z, t\} + \{z, t, x\}y - \{z, t, y\}x. \end{aligned}$$

Therefore,

$$\begin{aligned}
& ((xy)t)z + ((yz)t)x + ((zx)t)y \\
&= \{xy, z, t\} + \{yz, x, t\} + \{zx, y, t\} \\
&\quad + \{z, t, x\}y - \{z, t, y\}x + \{x, t, y\}z \\
&\quad - \{x, t, z\}y + \{y, t, z\}x - \{y, t, x\}z \\
&= \{xy, z, t\} + \{yz, x, t\} + \{zx, y, t\} \\
&\quad + (\{y, t, z\} - \{z, t, y\})x + (\{z, t, x\} - \{x, t, z\})y \\
&\quad + (\{x, t, y\} - \{y, t, x\})z \\
&= \{xy, z, t\} + \{yz, x, t\} + \{zx, y, t\} \\
&\quad + ((xy)t)z + ((yz)t)x + ((zx)t)y,
\end{aligned}$$

by (3.10a), and this proves item (i).

Now, from equation (3.10c) we have

$$\{xy, z, \{a, t, b\}\} - \{a, t, \{xy, z, b\}\} = \{\{xy, z, a\}, t, b\} - \{a, \{z, xy, t\}, b\},$$

which, together with (i) gives

$$\{a, \{x, yz, t\} + \{y, zx, t\} + \{z, xy, t\}, b\} = 0,$$

for any $a, b, x, y, z, t \in M$, which by (3.10a) yields

$$M^2(\{x, yz, t\} + \{y, zx, t\} + \{z, xy, t\}) = 0.$$

Also, equation (3.10b), together with (i), gives

$$\{x, yz, u\} + \{y, zx, u\} + \{z, xy, u\} = 0$$

for any $x, y, z, t \in M$ and $u \in M^2$. Hence item (ii) follows. \square

Proposition 3.20. *Let \mathfrak{g} be a Lie algebra endowed with an action of S_3 by automorphisms and assume that φ acts nontrivially. Let $(M, xy, \{x, y, z\})$ be the corresponding generalized Malcev algebra, and let $\mathfrak{g}(M)$ be the associated Lie algebra. If \mathfrak{g} does not contain proper ideals invariant under the action of S_3 (in particular, if \mathfrak{g} is simple), then \mathfrak{g} and $\mathfrak{g}(M)$ are isomorphic as Lie algebras with S_3 -action.*

Proof. Write $\mathfrak{g} = (U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-) \oplus (W \otimes M)$ as in (3.6). Then

$$(U \otimes \{d \in \mathfrak{d}^+ : d(M) = 0\}) \oplus (U' \otimes \{d \in \mathfrak{d}^- : d(M) = 0\})$$

is an ideal of \mathfrak{g} invariant under the action of S_3 , and hence $\{d \in \mathfrak{d}^\pm : d(M) = 0\} = 0$, so it can be assumed that $\mathfrak{d}^+, \mathfrak{d}^-$ are contained in $\mathfrak{gl}(M)$. Besides,

$$(U \otimes d_{M,M}^+) \oplus (U' \otimes d_{M,M}^-) \oplus (W \otimes M)$$

is the ideal of \mathfrak{g} generated by M , which is necessarily the whole \mathfrak{g} . Thus, $\mathfrak{d}^\pm = d_{M,M}^\pm$ and the result follows. \square

Example 3.21. (Simple Lie algebra of type G_2)

Assume $\omega \in k$, and let E be a three dimensional vector space over k endowed with a nondegenerate symmetric bilinear form b with trivial determinant (that is $\det(b(e_i, e_j))$ is a square for any basis $\{e_1, e_2, e_3\}$ of E). Fix then an orthogonal basis $\{e_1, e_2, e_3\}$ of E with $\det(b(e_i, e_j)) = 1$ and consider the trilinear alternating map $T : E \times E \times E \rightarrow k$ determined by $T(e_1, e_2, e_3) = 1$.

Then for any $u_1, u_2, u_3 \in E$, $T(u_1, u_2, u_3)^2 = \det(b(u_i, u_j))$. There appears the attached cross product $v \times w$ determined by $b(u, v \times w) = T(u, v, w)$ for any $u, v, w \in E$, which satisfies the equation

$$(u \times v) \times w = b(u, w)v - b(v, w)v$$

for any $u, v, w \in E$ or, equivalently,

$$b(x \times y, z \times t) = \begin{vmatrix} b(x, z) & b(x, t) \\ b(y, z) & b(y, t) \end{vmatrix}$$

for any $x, y, z, t \in E$.

The split Lie algebra of type G_2 can be described (see [FH91, page 348]) as

$$\mathfrak{g} = E^* \oplus \mathfrak{sl}(E) \oplus E$$

(E^* denotes the dual vector space), with bracket given by the usual bracket in the Lie subalgebra of zero trace endomorphisms $\mathfrak{sl}(E)$ and by:

$$\left\{ \begin{array}{l} [A, e] = A(e), \quad [A, f] = -fA \text{ (composition of maps),} \\ [e_1, e_2] = -2e_1 \wedge e_2 \in E^*, \text{ where } e_1 \wedge e_2 : e \mapsto T(e_1, e_2, e), \\ [f_1, f_2] = 2f_1 \wedge f_2 \in E, \\ \quad \text{such that } T(f_1 \wedge f_2, e_1, e_2) = f_1(e_1)f_2(e_2) - f_1(e_2)f_2(e_1), \\ [e, f] = 3f(\cdot)e - f(e)I, \text{ } I \text{ denotes the identity map on } E, \end{array} \right.$$

for any $e, e_1, e_2 \in E$, $f, f_1, f_2 \in E^*$ and $A \in \mathfrak{sl}(E)$.

The symmetric group S_3 acts on \mathfrak{g} by automorphisms as follows:

$$\begin{aligned} \varphi(e) &= \omega e, \quad \varphi(f) = \omega^2 f, \quad \varphi(A) = A, \\ \tau(e) &= -b(e, \cdot), \quad \tau(A) = -A^t, \end{aligned}$$

for any $e \in E$, $f \in F$ and $A \in \mathfrak{sl}(E)$, where A^t denotes the adjoint relative to b , that is, $b(Ae, e') = b(e, A^t e')$ for any $e, e' \in E$.

Then $\mathfrak{g}_\omega = E$ is a generalized Malcev algebra (see Remark 3.18) with

$$\begin{aligned} xy &= [\tau(x), \tau(y)] = 2b(x, \cdot) \wedge b(y, \cdot) = 2x \times y, \\ \{x, y, z\} &= [[x, \tau(y)], z] = -[3b(y, \cdot)x - b(x, y)I, z] = b(x, y)z - 3b(y, z)x, \end{aligned}$$

for any $x, y, z \in E$.

By Proposition 3.20, \mathfrak{g} is isomorphic to the associated Lie algebra $\mathfrak{g}(E)$. Thus, the central simple split Lie algebra of type G_2 is completely determined by the three dimensional generalized Malcev algebra $(E, xy, \{x, y, z\})$. \square

4. LIE ALGEBRAS WITH S_3 -ACTION AND MALCEV ALGEBRAS

Some noteworthy examples of generalized Malcev algebras are provided by Malcev algebras (see [Sag61]).

Theorem 4.1. *Let $(M, xy, \{x, y, z\})$ be a generalized Malcev algebra such that $\{x, y, z\}$ is skew-symmetric on x and y . Then (M, xy) is a Malcev algebra and*

$$2\{x, y, z\} = (xy)z + x(yz) - y(xz) \quad (4.2)$$

for any $x, y, z \in M$.

Conversely, if (M, xy) is a Malcev algebra and a triple product $\{., ., .\}$ is defined on M by the formula in (4.2), then $(M, xy, \{x, y, z\})$ is a generalized Malcev algebra.

Proof. This result appears essentially, with a very different formulation, in [Mik92, Theorem 1] and in [Gri03]. We include a proof for completeness.

Assume first that $(M, xy, \{x, y, z\})$ is a generalized Malcev algebra such that $\{x, y, z\}$ is skew-symmetric on x and y . Then (3.10a) gives

$$\begin{aligned} (xy)z + x(yz) - y(xz) &= (xy)z - (yz)x + (xz)y \\ &= \{x, z, y\} - \{y, z, x\} - \{y, x, z\} + \{z, x, y\} + \{x, y, z\} - \{z, y, x\} \\ &= 2\{x, y, z\}, \end{aligned}$$

so (4.2) holds.

Now, with $J(x, y, z) = (xy)z + (yz)x + (zx)y$, this is equivalent to

$$2(xy)z - J(x, y, z) = 2\{x, y, z\}, \quad (4.3)$$

for any $x, y, z \in M$, and then

$$\begin{aligned} J(x, y, xy) &= 2(xy)(xy) - 2\{x, y, xy\} = -2\{x, y, xy\} \quad (\text{as } z^2 = 0 \text{ for any } z) \\ &= 2\{y, x, x\}y + 2x\{y, x, y\} \quad (\text{by (3.10b)}) \\ &= 2((yx)x)y + 2x((yx)y) \quad (\text{by (4.2)}) \\ &= -2J(x, y, xy). \end{aligned}$$

Hence $J(x, y, xy) = 0$ for any $x, y \in M$, so

$$J(x, y, xz) + J(x, z, xy) = 0 \quad (4.4)$$

for any $x, y, z \in M$. Now equation (3.10d) gives

$$\{xy, z, x\} + \{yz, x, x\} + \{zx, y, x\} = 0,$$

which, by (4.3) becomes

$$\begin{aligned} 0 &= J(x, y, z)x - \frac{1}{2}(J(xy, z, x) + J(yz, x, x) + J(zx, y, x)) \\ &= J(x, y, z)x - J(x, y, xz), \end{aligned}$$

because of (4.4). Hence $J(x, y, xz) = J(x, y, z)x$ for any $x, y, z \in M$, and this is equivalent to M being a Malcev algebra (see [Sag61]).

Conversely, if (M, xy) is a Malcev algebra and the triple product is defined by (4.2), then for any x, y :

$$\{x, y, .\} = \frac{1}{2}(\text{ad}_{xy} + [\text{ad}_x, \text{ad}_y]) = D(x, y),$$

$(\text{ad}_x : y \mapsto xy)$, which is known to be a derivation of (M, xy) satisfying $D(xy, z) + D(yz, x) + D(zx, y) = 0$ (see [Sag61]). Hence the conditions (3.10b), (3.10c) and (3.10d) are satisfied. Also, $2(\{x, z, y\} - \{y, z, x\}) = (xz)y + x(zy) - z(xy) - (yz)x - y(zx) + z(yx) = 2(xy)z$ for any $x, y, z \in M$ because of the anticommutativity of the product, so condition (3.10a) holds too. \square

Note that in Example 3.21 there appears a three-dimensional generalized Malcev algebra $(E, xy, \{x, y, z\})$, where (E, xy) is a Lie (and hence Malcev) algebra, but where the triple product $\{x, y, z\}$ is not given by formula (4.2), as it is not even skew symmetric in x and y .

Remark 4.5. Let (M, xy) be a Malcev algebra, and define the triple product $\{., ., .\}$ by (4.2). Then $\{x, y, .\} + \{y, x, .\}$ is identically 0, so the associated Lie algebra $\mathfrak{g}(M)$ is just

$$\mathfrak{g}(M) = (U \otimes \mathfrak{d}^+) \oplus (W \otimes M),$$

which is a *Lie algebra with triality* in the notation of [Gri03]. This is a Lie algebra with an action of S_3 by automorphisms such that the alternating module does not appear in the decomposition into a direct sum of irreducible S_3 -modules.

Besides, if (M, xy) is a Lie algebra, then $\{x, y, .\} = \frac{1}{2}(\text{ad}_{xy} + [\text{ad}_x, \text{ad}_y]) = \text{ad}_{xy}$, so $\mathfrak{g}(M) = (U \otimes \text{ad}_M) \oplus (W \otimes M)$. If the center of M is trivial, then ad_M can be identified with M itself, and hence $\mathfrak{g}(M) \simeq (U \oplus W) \otimes M = k^3 \otimes M \simeq M^3$. In this way, $\mathfrak{g}(M)$ is seen to be isomorphic to the direct sum of three copies of M , with componentwise multiplication. \square

Remark 4.6. With the same conventions as in Remark 4.5, the transposition τ provides a \mathbb{Z}_2 -grading of $\mathfrak{g}(M)$ whose even part is $(u \otimes \mathfrak{d}^+) \oplus (w_+ \otimes M)$. Since $w_+ \bullet w_+ = w_+$, $(w_+ | w_+) = 2$ and $d_{x,y}^+ = \{x, y, .\} = \frac{1}{2}D(x, y)$, it follows that this even part is isomorphic to

$$D(M, M) \oplus M,$$

with product given by the Lie bracket of derivations in $D(M, M)$, and by $[d, x] = d(x)$ and $[x, y] = D(x, y) + xy$ for any $d \in D(M, M)$ and $x, y \in M$. If the center of M is trivial, M can be identified to ad_M . Then the Lie multiplication algebra of M is $L(M) = D(M, M) \oplus \text{ad}_M$ (see [Sag61]), and the map $D(M, M) \oplus M \rightarrow L(M)$, $D(x, y) \mapsto D(x, y)$, $x \mapsto -\text{ad}_x$ is an isomorphism. That is, the fixed subalgebra of $\mathfrak{g}(M)$ by the action of τ is isomorphic to the Lie multiplication algebra of M . \square

Remark 4.7. Given a Malcev algebra M , in [PIS04] a Lie algebra $\mathcal{L}(M)$ is defined by generators $\{\lambda_a, \rho_a : a \in M\}$ and relations (see [PIS04, Eq. (1)]):

$$\begin{aligned} \lambda_{\alpha a + \beta b} &= \alpha \lambda_a + \beta \lambda_b, & \rho_{\alpha a + \beta b} &= \alpha \rho_a + \beta \rho_b, \\ [\lambda_a, \lambda_b] &= \lambda_{ab} - 2[\lambda_a, \rho_b], & [\rho_a, \rho_b] &= -\rho_{ab} - 2[\lambda_a, \rho_b], \\ [\lambda_a, \rho_b] &= [\rho_a, \lambda_b], \end{aligned}$$

for any $a, b \in M$ and $\alpha, \beta \in k$. Then S_3 acts on this Lie algebra by automorphisms as follows:

$$\begin{aligned} \tau(\lambda_a) &= -\rho_a \\ \varphi(\lambda_a) &= -\lambda_a - \rho_a, \quad \varphi(\rho_a) = \lambda_a. \end{aligned}$$

It is straightforward to check that the relations are preserved. Actually, τ coincides with the automorphism $\zeta\eta\zeta$ while φ coincides with the automorphism $\zeta\eta$ in [PIS04, p. 386]. The arguments in the proof of Proposition 3.2

in [PIS04] can be used to prove that the envelope $\mathcal{L}(M)$ is isomorphic to our $\mathfrak{g}(M)$, by means of

$$\begin{aligned}\rho_a - \lambda_a &\mapsto w_+ \otimes a, \\ \rho_a + \lambda_a &\mapsto \frac{1}{3}w_- \otimes a. \quad \square\end{aligned}$$

Recall that given any nonassociative algebra (A, \cdot) , the generalized alternative nucleus is the subspace

$$N_{alt}(A, \cdot) = \{a \in A : (a, x, y) = -(x, a, y) = (x, y, a) \ \forall x, y \in A\},$$

where $(x, y, z) = (x \cdot y) \cdot z - x \cdot (y \cdot z)$ is the associator of the elements x, y, z . The generalized alternative nucleus is a Malcev algebra under the commutator $[x, y] = x \cdot y - y \cdot x$ (see [PIS04]).

Given a unital algebra with involution $(A, \cdot, -)$, it has been shown in Corollary 1.16 that $\mathfrak{ltr}(A, \cdot, -)$ is a Lie algebra with S_3 -action ($S_3 \simeq S_4/V_4$), with no irreducible component isomorphic to the alternating module.

Proposition 4.8. *Let $(A, \cdot, -)$ be a unital algebra with involution. Then the generalized Malcev algebra associated to the Lie algebra $\mathfrak{ltr}(A, \cdot, -)$ is isomorphic to $(M, [x, y], \{x, y, z\})$, where*

$$M = \{a \in N_{alt}(A, \cdot) : \bar{a} = -a\},$$

and where $2\{x, y, z\} = [[x, y], z] + [x, [y, z]] - [y, [x, z]]$ for any $x, y, z \in M$.

Proof. The result in [AF93, Corollary 3.5] shows that

$$\begin{aligned}\mathfrak{ltr}(A, \cdot, -) &= \{(d, d, d) : d \in \mathfrak{der}(A, \cdot, -)\} \\ &\quad \oplus \{(L_{s_2} - R_{s_3}, L_{s_3} - R_{s_1}, L_{s_1} - R_{s_2}) : s_1, s_2, s_3 \in M, s_1 + s_2 + s_3 = 0\},\end{aligned}$$

with M as above. Here L and R denote the left and right multiplications in (A, \cdot) . This shows that, as vector spaces, there is an isomorphism:

$$\begin{aligned}\Psi : (U \otimes \mathfrak{der}(A, \cdot, -)) \oplus (W \otimes M) &\rightarrow \mathfrak{ltr}(A, \cdot, -) \\ u \otimes d &\mapsto (d, d, d) \\ (\alpha_1, \alpha_2, \alpha_3) \otimes s &\mapsto (L_{\alpha_2 s} - R_{\alpha_3 s}, L_{\alpha_3 s} - R_{\alpha_1 s}, L_{\alpha_1 s} - R_{\alpha_2 s}) \\ (\alpha_1 + \alpha_2 + \alpha_3 = 0)\end{aligned}$$

for $d \in \mathfrak{der}(A, \cdot, -)$, $\alpha_i \in k$, $i = 1, 2, 3$, and $s \in M$.

Now, given $w_1 = (\alpha_1, \alpha_2, \alpha_3)$, $w_2 = (\beta_1, \beta_2, \beta_3) \in W$ (so $\alpha_1 + \alpha_2 + \alpha_3 = 0 = \beta_1 + \beta_2 + \beta_3$), and $s, t \in M$, the arguments in the proof of [AF93, Corollary 3.5] give:

$$\begin{aligned}[(L_{\alpha_2 s} - R_{\alpha_3 s}, L_{\alpha_3 s} - R_{\alpha_1 s}, L_{\alpha_1 s} - R_{\alpha_2 s}), (L_{\beta_2 t} - R_{\beta_3 t}, L_{\beta_3 t} - R_{\beta_1 t}, L_{\beta_1 t} - R_{\beta_2 t})] \\ = \frac{1}{3}(D, D, D) + (L_{a_2} - R_{a_3}, L_{a_3} - R_{a_1}, L_{a_1} - R_{a_2})\end{aligned}$$

where:

$$\begin{aligned}D &= [L_{\alpha_2 s} - R_{\alpha_3 s}, L_{\beta_2 s} - R_{\beta_3 t}] \\ &\quad + [L_{\alpha_3 s} - R_{\alpha_1 s}, L_{\beta_3 s} - R_{\beta_1 t}] + [L_{\alpha_1 s} - R_{\alpha_2 s}, L_{\beta_1 s} - R_{\beta_2 t}] \\ &= (\alpha_1 \beta_1 + \alpha_2 \beta_2 + \alpha_3 \beta_3)([L_s, L_t] + [R_s, R_t]) \\ &\quad - (\alpha_2 \beta_3 + \alpha_3 \beta_1 + \alpha_1 \beta_2)[L_s, R_t] - (\alpha_3 \beta_2 + \alpha_1 \beta_3 + \alpha_2 \beta_1)[R_s, L_t].\end{aligned}$$

But

$$[L_s, R_t](x) = -(s, x, t) = (t, x, s) = -[L_t, R_s](x),$$

so $[L_s, R_t] = [R_s, L_t]$. Besides,

$$\sum_{1 \leq i \neq j \leq 3} \alpha_i \beta_j = (\alpha_1 + \alpha_2 + \alpha_3)(\beta_1 + \beta_2 + \beta_3) - \sum_{i=1}^3 \alpha_i \beta_i.$$

Hence:

$$\begin{aligned} D &= (\alpha_1 \beta_1 + \alpha_2 \beta_2 + \alpha_3 \beta_3)([L_s, L_t] + [R_s, R_t] + [L_s, R_t]) \\ &= 3(w_1 | w_2)([L_s, L_t] + [R_s, R_t] + [L_s, R_t]). \end{aligned}$$

Also, $a_1 = \frac{1}{3}(b_3 - b_2)$ (and cyclically), with $b_1 = [L_{\alpha_2 s} - R_{\alpha_3 s}, L_{\beta_2 t} - R_{\beta_3 t}](1)$ (and cyclically). Hence

$$\begin{aligned} b_1 &= (\beta_2 - \beta_3)(\alpha_2 s \cdot t - \alpha_3 t \cdot s) - (\alpha_2 - \alpha_3)(\beta_2 t \cdot s - \beta_3 s \cdot t) \\ &= (\alpha_2 \beta_2 - \alpha_3 \beta_3)[s, t], \end{aligned}$$

and

$$a_1 = \frac{1}{3}(2\alpha_1 \beta_1 - \alpha_2 \beta_2 - \alpha_3 \beta_3)[s, t].$$

Since

$$w_1 \bullet w_2 = \frac{1}{3}(2\alpha_1 \beta_1 - \alpha_2 \beta_2 - \alpha_3 \beta_3, 2\alpha_2 \beta_2 - \alpha_3 \beta_3 - \alpha_1 \beta_1, 2\alpha_3 \beta_3 - \alpha_1 \beta_1 - \alpha_2 \beta_2),$$

it follows that

$$\begin{aligned} &[\Psi(w_1 \bullet s), \Psi(w_2 \bullet t)] \\ &= \Psi((w_1 | w_2)u \otimes ([L_s, L_t] + [R_s, R_t] + [L_s, R_t])) + \Psi((w_1 \bullet w_2) \otimes [s, t]) \end{aligned}$$

and this shows that the attached generalized Malcev algebra is the triple $(M, [x, y], \{x, y, z\})$, where $\{x, y, z\} = ([L_x, L_y] + [R_x, R_y] + [L_x, R_y])(z)$. Since $(M, [., .])$ is a Malcev algebra (see [PIS04]), Theorem 4.1 finishes the proof. \square

A distinguished example is given by any octonion algebra $(\mathbb{O}, \cdot, -)$ over k with its standard involution. Since \mathbb{O} is alternative, $N_{alt}(\mathbb{O}, \cdot) = \mathbb{O}$, so $M = \mathbb{O}_0 = \{x \in \mathbb{O} : t(x) = x + \bar{x} = 0\}$. Then, it is well-known by the Principle of Triality (see [KMRT98, Chapter VIII] and references therein) that $\mathfrak{ltt}(\mathbb{O}, \cdot, -)$ is isomorphic to the orthogonal Lie algebra $\mathfrak{so}(\mathbb{O}, n)$, where n denotes the norm of \mathbb{O} ($n(x) = x \cdot \bar{x}$). The action of S_3 decomposes this Lie algebra as

$$\mathfrak{so}(\mathbb{O}, n) = (U \otimes \mathfrak{der}(\mathbb{O}, \cdot)) \oplus (W \otimes \mathbb{O}_0).$$

The associated Malcev algebra is the central simple non-Lie Malcev algebra \mathbb{O}_0 . Recall that any central simple non-Lie Malcev algebra appears in this way (see [Fil76] and [Kuz68]).

5. EXAMPLES

Other noteworthy examples of generalized Malcev algebras consist of those whose bilinear multiplication is trivial. Equations (3.10) show that these are just the Jordan triple systems (see [Mey72]).

Given a Jordan triple system T with triple product denoted by $\{x, y, z\}$, consider the Lie subalgebra of $\mathfrak{gl}(T) \oplus \mathfrak{gl}(T)$ defined by:

$$\{(d_1, d_2) : d_i(\{x, y, z\}) = \{d_i(x), y, z\} + \{x, d_{3-i}(y), z\} + \{x, y, d_i(z)\} \\ \forall x, y, z \in T, \forall i = 1, 2\}.$$

Then $\mathfrak{s}(T) = \text{span} \{(\{x, y, \cdot\}, -\{y, x, \cdot\}) : x, y \in T\}$ is a subalgebra of $\mathfrak{gl}(T) \oplus \mathfrak{gl}(T)$ and the Tits-Kantor-Koecher Lie algebra of T is the Lie algebra

$$\mathcal{K}(T) = TKK(T) = T \oplus \mathfrak{s}(T) \oplus \hat{T},$$

where \hat{T} is just a copy of T (and given an element $x \in T$, its copy in \hat{T} will be denoted by \hat{x}), with bracket given by:

$$[T, T] = [\hat{T}, \hat{T}] = 0, \quad \mathfrak{s}(T) \text{ is a subalgebra of } \mathcal{K}(T), \\ [x, \hat{y}] = (\{x, y, \cdot\}, -\{y, x, \cdot\}), \\ [(d_1, d_2), x] = d_1(x), \quad [(d_1, d_2), \hat{y}] = \widehat{d_2(y)},$$

for any $(d_1, d_2) \in \mathfrak{s}(T)$ and $x, y \in T$.

Proposition 5.1. *Assume $\omega \in k$ ($\omega^3 = 1 \neq \omega$), and let $(T, \{x, y, z\})$ be a Jordan triple system. Then $(T, xy, \{x, y, z\})$ is a generalized Malcev algebra with $xy = 0$ for any $x, y \in T$, and the linear map*

$$\begin{aligned} \mathfrak{gl}(T) &\rightarrow \mathcal{K}(T) \\ w_\omega \otimes x &\mapsto x \\ w_{\omega^2} \otimes x &\mapsto \hat{x} \\ u \otimes d^+ &\mapsto (d^+, d^+) \\ \hat{u}' \otimes d^- &\mapsto (d^-, -d^-) \end{aligned}$$

for $x \in T$, $d^+ \in \text{span} \{d_{x,y}^+ = \frac{1}{2}(\{x, y, \cdot\} - \{y, x, \cdot\}) : x, y \in T\}$ and $d^- \in \text{span} \{d_{x,y}^- = \frac{1}{2}(\{x, y, \cdot\} + \{y, x, \cdot\}) : x, y \in T\}$, is an isomorphism of Lie algebras.

Proof. The proof is straightforward from the different definitions involved. \square

Example 5.2. (Orthogonal Lie algebras)

Let us assume, for simplicity, that the ground field k is algebraically closed, and let \mathcal{V} be a vector space of dimension ≥ 3 , endowed with a nondegenerate symmetric bilinear form b . Let $\{e_1, \dots, e_n\}$ be a basis with $b(e_i, e_j) = \delta_{i,j}$. The symmetric group S_3 acts naturally by isometries by permuting the first three vectors in this basis and leaving the remaining ones fixed. Then \mathcal{V} decomposes, as a module for S_3 , into the orthogonal sum of a submodule isomorphic to the two dimensional irreducible module W and a direct sum of $n - 2$ trivial modules. Thus, we may think of \mathcal{V} as the orthogonal sum $W \perp E$ with the restriction of b to W given by the bilinear form $(\cdot | \cdot)$ in

(3.1) (by invariance under the action of S_3), and where S_3 acts trivially on E .

The orthogonal Lie algebra $\mathfrak{g} = \mathfrak{so}(\mathcal{V}, b)$ is spanned by the maps:

$$\sigma_{u,v} : w \mapsto b(u, w)v - b(v, w)u$$

for any $u, v, w \in \mathcal{V}$. Note that these maps satisfy

$$[\sigma_{u,v}, \sigma_{x,y}] = \sigma_{\sigma_{u,v}(x), y} + \sigma_{x, \sigma_{u,v}(y)}$$

for any u, v, x, y .

The action of the symmetric group S_3 on \mathcal{V} induces an action on $\mathfrak{so}(\mathcal{V}, b)$, where for any $\gamma \in S_3$ and $f \in \mathfrak{so}(\mathcal{V}, b)$, $\gamma \cdot f = \gamma f \gamma^{-1}$. In this way, $\gamma \cdot \sigma_{u,v} = \sigma_{\gamma(u), \gamma(v)}$ for any $u, v \in \mathcal{V}$. Then there is the natural decomposition $\mathfrak{g} = \mathfrak{so}(W) \oplus \mathfrak{so}(E) \oplus \sigma_{W,E} \simeq \mathfrak{so}(W) \oplus \mathfrak{so}(E) \oplus W \otimes E$, where $\mathfrak{so}(W)$ acts trivially on E and $\mathfrak{so}(E)$ acts trivially on W . Besides, the one dimensional Lie algebra $\mathfrak{so}(W)$ is spanned by $\sigma_{w_\omega, w_{\omega^2}}$.

The action of the cycle φ decomposes \mathfrak{g} into the direct sum of its eigenspaces, where $\mathfrak{g}_1 = \mathfrak{so}(W) \oplus \mathfrak{so}(E)$, $\mathfrak{g}_\omega = \sigma_{w_\omega, E}$ and $\mathfrak{g}_{\omega^2} = \sigma_{w_{\omega^2}, E}$. (Recall that $\varphi(w_\omega) = \omega w_\omega$ and $\varphi(w_{\omega^2}) = \omega^2 w_{\omega^2}$) The attached generalized Malcev algebra is defined then on \mathfrak{g}_ω by Remark 3.18. Note that for any $x, y, z \in E$:

$$\begin{aligned} [\tau(\sigma_{w_\omega, x}), \tau(\sigma_{w_\omega, y})] &= [\sigma_{w_{\omega^2}, x}, \sigma_{w_{\omega^2}, y}] \\ &= \sigma_{\sigma_{w_{\omega^2}, x}(w_{\omega^2}), y} + \sigma_{w_{\omega^2}, \sigma_{w_{\omega^2}, x}(y)} = 0, \\ [[\sigma_{w_\omega, x}, \tau(\sigma_{w_\omega, y})], \sigma_{w_\omega, z}] &= [[\sigma_{w_\omega, x}, \sigma_{w_{\omega^2}, y}], \sigma_{w_\omega, z}] \\ &= [\sigma_{x, y} + b(x, y)\sigma_{w_\omega, w_{\omega^2}}, \sigma_{w_\omega, z}] \\ &= \sigma_{w_\omega, \sigma_{x, y}(z) - b(x, y)z} \\ &= \sigma_{w_\omega, b(x, z)y - b(y, z)x - b(x, y)z}, \end{aligned}$$

which shows that the attached generalized Malcev algebra can be identified with $(E, xy, \{x, y, z\})$ with $xy = 0$ and $\{x, y, z\} = b(x, z)y - b(y, z)x - b(x, y)z$ for any $x, y, z \in E$. This is the Jordan triple system associated to the nondegenerate symmetric bilinear form $-b|_E$ (see [Mey72]). \square

Another easy example, which is neither a Malcev algebra nor a Jordan triple system is obtained with similar arguments, but with $\mathfrak{so}(\mathcal{V})$ substituted by $\mathfrak{sl}(\mathcal{V})$:

Example 5.3. (Special Lie algebras)

Let \mathcal{V} be a vector space of dimension ≥ 3 over an algebraically closed field k , and take a basis $\{e_1, \dots, e_n\}$. Then S_3 acts by permuting the first three elements of this basis. As in Example 5.2, this decomposes \mathcal{V} into a direct sum $\mathcal{V} = W \oplus E$, where W is the two dimensional irreducible module for S_3 and E is a vector space on which S_3 acts trivially. Identify the general linear Lie algebra $\mathfrak{gl}(\mathcal{V})$ with $\mathcal{V} \otimes \mathcal{V}^*$ ($v \otimes f \in \mathcal{V} \otimes \mathcal{V}^* \leftrightarrow f(\cdot)v \in \mathfrak{gl}(\mathcal{V})$, where $f(\cdot)v$ is the linear map $w \mapsto f(w)v$ for any $v, w \in \mathcal{V}$ and $f \in \mathcal{V}^*$). Recall that W is endowed with a unique (up to scalars) nondegenerate symmetric bilinear form $(\cdot | \cdot)$ invariant under the action of S_3 (see equation (3.1)) that allows us to identify W and W^* . Thus:

$$\begin{aligned} \mathfrak{gl}(W \oplus E) &\simeq (W \otimes W^*) \oplus (E \otimes E^*) \oplus (W \otimes E^*) \oplus (E \otimes W^*) \\ &\simeq \mathfrak{gl}(W) \oplus \mathfrak{gl}(E) \oplus (W \otimes E^*) \oplus (E \otimes W^*). \end{aligned}$$

The action of S_3 on \mathcal{V} induces an action of S_3 by conjugation on $\mathfrak{gl}(\mathcal{V})$, which leaves fixed $\mathfrak{gl}(E)$. Under this action, $\mathfrak{g} = \mathfrak{gl}(W \oplus E)$ decomposes as the direct sum of the eigenspaces for φ , and the ω eigenspace is, with the identifications above,

$$\mathfrak{g}_\omega = \text{span} \{w_{\omega^2} \otimes (w_{\omega^2} | \cdot)\} \oplus (w_\omega \otimes E^*) \oplus (E \otimes (w_\omega | \cdot)).$$

Besides:

$$\begin{aligned} [\tau(w_{\omega^2} \otimes (w_{\omega^2} | \cdot)), \tau(w_\omega \otimes f)] &= [w_\omega \otimes (w_\omega | \cdot), w_{\omega^2} \otimes f] \\ &= w_\omega \otimes f, \end{aligned}$$

$$\begin{aligned} [\tau(w_{\omega^2} \otimes (w_{\omega^2} | \cdot)), \tau(e \otimes (w_\omega | \cdot))] &= [w_\omega \otimes (w_\omega | \cdot), e \otimes (w_{\omega^2} | \cdot)] \\ &= -e \otimes (w_\omega | \cdot) \end{aligned}$$

$$\begin{aligned} [\tau(e \otimes (w_\omega | \cdot)), \tau(w_\omega \otimes f)] &= [e \otimes (w_{\omega^2} | \cdot), w_{\omega^2} \otimes f] \\ &= -f(e)w_{\omega^2} \otimes (w_{\omega^2} | \cdot). \end{aligned}$$

This means that the attached generalized Malcev algebra can be identified, as a vector space, to $ka \oplus E \oplus E^*$, by means of $a \leftrightarrow w_{\omega^2} \otimes (w_{\omega^2} | \cdot)$, $e \leftrightarrow e \otimes (w_\omega | \cdot)$ and $f \leftrightarrow w_\omega \otimes f$, for any $e \in E$ and $f \in E^*$. Then the anticommutative bilinear product becomes:

$$ae = -e, \quad af = f, \quad fe = f(e)a$$

for any $e \in E$ and $f \in E^*$. If $\dim E \geq 2$, this is not a Malcev algebra.

Also,

$$\begin{aligned} [w_{\omega^2} \otimes (w_{\omega^2} | \cdot), \tau(w_{\omega^2} \otimes (w_{\omega^2} | \cdot))] &= [w_{\omega^2} \otimes (w_{\omega^2} | \cdot), w_\omega \otimes (w_\omega | \cdot)] \\ &= w_{\omega^2} \otimes (w_\omega | \cdot) - w_\omega \otimes (w_{\omega^2} | \cdot), \end{aligned}$$

$$[w_{\omega^2} \otimes (w_{\omega^2} | \cdot), \tau(w_\omega \otimes f)] = 0,$$

$$[w_{\omega^2} \otimes (w_{\omega^2} | \cdot), \tau(e \otimes (w_\omega | \cdot))] = 0,$$

$$[w_\omega \otimes f_1, w_\omega \otimes f_2] = 0,$$

$$[e_1 \otimes (w_\omega | \cdot), \tau(e_2 \otimes (w_\omega | \cdot))] = 0,$$

$$\begin{aligned} [w_\omega \otimes f, \tau(e \otimes (w_\omega | \cdot))] &= [w_\omega \otimes f, e \otimes (w_{\omega^2} | \cdot)] \\ &= f(e)w_\omega \otimes (w_{\omega^2} | \cdot) - e \otimes f, \end{aligned}$$

$$\begin{aligned} [e \otimes (w_\omega | \cdot), \tau(w_\omega \otimes f)] &= [e \otimes (w_\omega | \cdot), w_{\omega^2} | \cdot] \\ &= e \otimes f - f(e)w_{\omega^2} \otimes (w_\omega | \cdot), \end{aligned}$$

for any $e, e_1, e_2 \in E$ and $f, f_1, f_2 \in E^*$. With our identification of \mathfrak{g}_ω with $ka \oplus E \oplus E^*$, it follows easily that the nonzero triple products here are given by:

$$\{a, a, a\} = 2a, \quad \{a, a, e\} = -e, \quad \{a, a, f\} = -f,$$

$$\{e, f, a\} = -f(e)a = \{f, e, a\},$$

$$\{e, f, e'\} = f(e')e + f(e)e', \quad \{f, e, e'\} = -f(e')e,$$

$$\{e, f, f'\} = -f'(e)f, \quad \{f, e, f'\} = f'(e)f + f(e)f',$$

for any $e, e' \in E$ and $f, f' \in E^*$. □

Before proceeding with more examples, let us show a final connection of the generalized Malcev algebras to other kind of algebraic structures.

Remark 5.4. Let $(M, xy, \{x, y, z\})$ be a generalized Malcev algebra and let

$$\mathfrak{g}(M) = (U \otimes \mathfrak{d}^+) \oplus (U' \otimes \mathfrak{d}^-) \oplus (W \otimes M)$$

be the associated Lie algebra with S_3 -symmetry. Then τ becomes an order two automorphism which grades $\mathfrak{g}(M)$ over \mathbb{Z}_2 . The fixed subalgebra by τ is

$$\mathfrak{g}_0(M) = (U \otimes \mathfrak{d}^+) \oplus (w_+ \otimes M),$$

whose Lie bracket works as follows:

$$\begin{aligned} [u \otimes d_1, u \otimes d_2] &= u \otimes [d_1, d_2], \\ [u \otimes d_1, w_+ \otimes x] &= w_+ \otimes d(x), \\ [w_+ \otimes x, w_+ \otimes y] &= u \otimes [x, y, \cdot] + w_+ \otimes xy, \end{aligned}$$

for any $d, d_1, d_2 \in \mathfrak{d}^+$ and $x, y \in M$, where $[x, y, \cdot] = \{x, y, \cdot\} - \{y, x, \cdot\}$. This algebra is isomorphic to the Lie algebra

$$\tilde{\mathfrak{g}}_0(M) = [M, M, \cdot] \oplus M$$

with $[M, M, \cdot] = \text{span}\{[x, y, \cdot] : x, y \in M\} \subseteq \mathfrak{g}(M)$, and bracket given by imposing that $[M, M, \cdot]$ is a subalgebra, M its natural module, and $[x, y] = [x, y, \cdot] + xy$ for any $x, y \in M$. Thus $(M, xy, [x, y, z])$ is a Lie-Yamaguti algebra (or general Lie triple system in Yamaguti's notation [Yam58]) and $\tilde{\mathfrak{g}}_0(M)$ is its standard enveloping Lie algebra (see [KW01]). \square

Another family of examples of generalized Malcev algebras comes from the second line of Freudenthal's Magic Square.

Let $K = ke + kz$ be the quadratic commutative associative algebra with $ex = x$ for any x and $z^2 = -3e$. If $\sqrt{-3} \notin k$, then K is the field extension $k[\sqrt{-3}]$, otherwise K is isomorphic to $k \times k$. This is a composition algebra relative to the norm q , that is $q(xy) = q(x)q(y)$ for any $x, y \in K$, where $q(e) = 1$, $q(z) = 3$ and $q(e, z) = (q(e+z) - q(e) - q(z)) = 0$. Let (K, \bullet, q) be the associated para-Hurwitz algebra, where $x \bullet y = \overline{xy}$ ($\bar{e} = e$, $\bar{z} = -z$), that is,

$$e \bullet e = e, \quad e \bullet z = -z = z \bullet e, \quad z \bullet z = -3e.$$

The set of nonzero idempotents of (K, \bullet) is $\{e, -\frac{1}{2}e + \frac{1}{2}z, -\frac{1}{2}e - \frac{1}{2}z\}$, and its group of automorphisms $\text{Aut}(K, \bullet)$ permutes these idempotents. From here it follows that $\text{Aut}(K, \bullet) \simeq S_3$, where the action of φ and τ are given by:

$$\begin{cases} \tau(e) = e, \quad \tau(z) = -z, \\ \varphi(e) = -\frac{1}{2}e + \frac{1}{2}z, \quad \varphi(z) = -\frac{3}{2}e - \frac{1}{2}z. \end{cases}$$

Actually, as a module for S_3 , K is isomorphic to the two dimensional module W : $e \leftrightarrow w_+$, $z \leftrightarrow w_-$. Under this linear isomorphism, the multiplication in (3.2) corresponds to the multiplication \bullet on K , while the symmetric bilinear form $(\cdot | \cdot)$ in (3.3) corresponds to the polar form $q(\cdot, \cdot)$ of the norm q of K .

The triality Lie algebra of (K, \bullet, q) is

$$\text{tri}(K, \bullet, q)$$

$$= \{(d_0, d_1, d_2) \in \mathfrak{so}(K, q) : d_0(x \bullet y) = d_1(x) \bullet y + x \bullet d_2(y) \quad \forall x, y \in K\}.$$

But the orthogonal Lie algebra $\mathfrak{so}(K, q)$ has dimension 1, being spanned by $\sigma = \sigma_{e,z} = q(e, \cdot)z - q(z, \cdot)e$. That is,

$$\sigma : \begin{cases} e \mapsto q(e, e)z - q(e, z)e = 2z, \\ z \mapsto q(e, z)z - q(z, z)e = -6e. \end{cases}$$

It follows that

$$\mathfrak{tri}(K, \bullet, q) = \{(\alpha_0\sigma, \alpha_1\sigma, \alpha_2\sigma) : \alpha_0, \alpha_1, \alpha_2 \in k, \alpha_0 + \alpha_1 + \alpha_2 = 0\}.$$

Under the identification above of W and K , the action of σ on K corresponds exactly to the action of u' on W ($u' \diamond w_+ = 2w_+$, $u' \diamond w_- = -6w_-$ because of (3.5)).

The action of S_3 on K induces an action on $\mathfrak{tri}(K, \bullet, q)$ by

$$\gamma((d_0, d_1, d_2)) = (\gamma d_0 \gamma^{-1}, \gamma d_1 \gamma^{-1}, \gamma d_2 \gamma^{-1}),$$

for any $\gamma \in S_3 = \text{Aut}(K, \bullet)$ and $(d_0, d_1, d_2) \in \mathfrak{tri}(K, \bullet, q)$. But $\tau\sigma\tau = -\sigma$ and $\varphi\sigma\varphi^2 = \sigma$, so $\mathfrak{tri}(K, \bullet, q)$ decomposes into the direct sum of two copies of the alternating module. Thus, as S_3 -modules, consider the identification:

$$\begin{aligned} \mathfrak{tri}(K, \bullet, q) &\simeq U' \otimes (k^3)_0 \\ (\alpha_0\sigma, \alpha_1\sigma, \alpha_2\sigma) &\leftrightarrow u' \otimes (\alpha_0, \alpha_1, \alpha_2), \end{aligned} \tag{5.5}$$

where $(k^3)_0 = \{(\alpha_0, \alpha_1, \alpha_2) \in k^3 : \alpha_0 + \alpha_1 + \alpha_2 = 0\}$.

Given any other symmetric composition algebra $(S, *, \hat{q})$ (see [KMRT98, Chapter VIII] and the references there in), a Lie algebra is constructed in [Eld04] on the vector space

$$\mathfrak{g}(K, S) = \mathfrak{tri}(K, \bullet, q) \oplus \mathfrak{tri}(S, *, \hat{q}) \oplus \left(\bigoplus_{i=0}^2 \iota_i(K \otimes S) \right),$$

where $\iota_i(K \otimes S)$ denotes a copy of $K \otimes S$, $i = 0, 1, 2$. The Lie bracket is defined by:

- $\mathfrak{tri}(K, \bullet, q)$ and $\mathfrak{tri}(S, *, \hat{q})$ are two commuting Lie subalgebras of $\mathfrak{g}(K, S)$ (so $[\mathfrak{tri}(K, \bullet, q), \mathfrak{tri}(S, *, \hat{q})] = 0$),
- $[(d_0, d_1, d_2), \iota_i(a \otimes x)] = \iota_i(d_i(a) \otimes x)$, $[(\hat{d}_0, \hat{d}_1, \hat{d}_2), \iota_i(a \otimes x)] = \iota_i(a \otimes \hat{d}_i(x))$, for $i = 0, 1, 2$, $a \in K$, $x \in S$, $(d_0, d_1, d_2) \in \mathfrak{tri}(K, \bullet, q)$ and $(\hat{d}_0, \hat{d}_1, \hat{d}_2) \in \mathfrak{tri}(S, *, \hat{q})$,
- $[\iota_i(a \otimes x), \iota_{i+1}(b \otimes y)] = \iota_{i+2}((a \bullet b) \otimes (x * y))$, for $i = 0, 1, 2$ (indices modulo 3), $a, b \in K$ and $x, y \in S$,
- $[\iota_i(a \otimes x), \iota_i(b \otimes y)] = \hat{q}(x, y)\theta^i(t_{a,b}) + q(a, b)\theta^i(t_{x,y})$ for $i = 0, 1, 2$, $a, b \in K$ and $x, y \in S$, where $\theta : (d_0, d_1, d_2) \mapsto (d_2, d_0, d_1)$, $t_{a,b} = (\sigma_{a,b}, \frac{1}{2}q(a, b)I - R_a L_b, \frac{1}{2}q(a, b)I - L_a R_b)$ and similarly for $t_{x,y}$. As usual R and L denote right and left multiplications, I is the identity mapping, and $\sigma_{a,b} = q(a, \cdot)b - q(\cdot, a)$.

Depending on the dimension of S being 1, 2, 4 or 8, the Lie algebra $\mathfrak{g}(K, S)$ is a form respectively of $\mathfrak{sl}_3(k)$, $\mathfrak{sl}_3(k) \oplus \mathfrak{sl}_3(k)$, $\mathfrak{sl}_6(k)$ or the simple Lie algebra of type E_6 . That is, we get the second line of Freudenthal's Magic Square [Eld04].

Now, the action of S_3 by automorphisms of K extends to an action of S_3 by automorphisms of $\mathfrak{g}(K, S)$, with the action of any $\gamma \in S_3$ is given by:

- $\gamma(\iota_i(a \otimes x)) = \iota_i(\gamma(a) \otimes x)$, for any $a \in K$, $x \in S$, $i = 0, 1, 2$,

- $\gamma((d_0, d_1, d_2)) = (\gamma d_0 \gamma^{-1}, \gamma d_1 \gamma^{-1}, \gamma d_2 \gamma^{-1})$ for any (d_0, d_1, d_2) in $\mathbf{tri}(K, \bullet, q)$,
- $\gamma((\hat{d}_0, \hat{d}_1, \hat{d}_2)) = (\hat{d}_0, \hat{d}_1, \hat{d}_2)$ for any $(\hat{d}_0, \hat{d}_1, \hat{d}_2) \in \mathbf{tri}(S, *, \hat{q})$.

Thus, $\mathbf{tri}(S, *, q)$ is a trivial module for S_3 , $\mathbf{tri}(K, \bullet, q)$ is already known to be isomorphic to two copies of the alternating module by (5.5): $\mathbf{tri}(K, \bullet, q) \simeq U' \otimes (k^3)_0$, while $\bigoplus_{i=0}^2 \iota_i(K \otimes S)$ is a sum of $3 \times \dim S$ copies of the two dimensional irreducible module $W \simeq K$.

Consider then the natural identification:

$$\begin{aligned} \mathfrak{g}(K, S) &\simeq (U \otimes \mathbf{tri}(S, *, q)) \oplus (U' \otimes (k^3)_0) \oplus (W \otimes (\bigoplus_{i=0}^2 \iota_i(S))), \\ (\hat{d}_0, \hat{d}_1, \hat{d}_2) &\leftrightarrow u \otimes (\hat{d}_0, \hat{d}_1, \hat{d}_2), \\ (\alpha_0 \sigma, \alpha_1 \sigma, \alpha_2 \sigma) &\leftrightarrow u' \otimes (\alpha_0, \alpha_1, \alpha_2), \\ \iota_i(a \otimes x) &\leftrightarrow a \otimes \iota_i(x), \end{aligned}$$

for $(\hat{d}_0, \hat{d}_1, \hat{d}_2) \in \mathbf{tri}(S, *, \hat{q})$, $(\alpha_0, \alpha_1, \alpha_2) \in (k^3)_0$, $\sigma = \sigma_{e,z}$ as above, $i = 0, 1, 2$, $a \in K \simeq W$, and $x \in S$.

The Lie bracket on $\mathfrak{g}(K, S)$ translates into a bracket on the right which proves the validity of the following example:

Example 5.6. (Second line of Freudenthal's Magic Square)

Let $(S, *, \hat{q})$ be a symmetric composition algebra and let $M = \bigoplus_{i=0}^2 \iota_i(S)$, be the direct sum of three copies of S . Then $(M, XY, \{X, Y, Z\})$ is a generalized Malcev algebra, where for $i, j, k \in \{0, 1, 2\}$ and $x, y, z \in S$:

- $\iota_i(x) \iota_{i+1}(y) = \iota_{i+2}(x * y)$,
- $\iota_i(x) \iota_i(y) = 0$,
- $\{\iota_i(x), \iota_j(y), \iota_k(z)\} = 0$, if $i \neq j$,
- $\{\iota_i(x), \iota_i(y), \iota_{i+1}(z)\} = -\iota_{i+1}((y * z) * x)$,
- $\{\iota_i(x), \iota_i(y), \iota_{i+2}(z)\} = -\iota_{i+2}(x * (z * y))$,
- $\{\iota_i(x), \iota_i(y), \iota_i(z)\} = \iota_i(\hat{q}(x, z)y - \hat{q}(y, z)x + \hat{q}(x, y)z)$. □

Notice that, for example, for $x, y \in S$,

$$[\iota_i(e \otimes x), \iota_i(e \otimes y)] = q(e, e) \theta^i(t_{x,y}) = 2\theta^i(t_{x,y}),$$

and this translates to

$$[w_+ \otimes x, w_+ \otimes y] = 2u \otimes \theta^i(t_{x,y}) = (w_+ | w_+) u \otimes \theta^i(t_{x,y}).$$

This gives $\iota_i(x) \iota_i(y) = 0$ and

$$d_{\iota_i(x), \iota_i(y)}^+(\iota_j(z)) = \begin{cases} \iota_i(\sigma_{x,y}(z)) & \text{if } j = i, \\ \iota_{i+1}(\frac{1}{2}\hat{q}(x, y)z - (y * z) * x) & \text{if } j = i + 1 \text{ (modulo 3)}, \\ \iota_{i+2}(\frac{1}{2}\hat{q}(x, y)z - x * (z * y)) & \text{if } j = i + 2 \text{ (modulo 3)}. \end{cases}$$

Also $[\iota_i(e \otimes x), \iota_i(z \otimes y)] = \hat{q}(x, y) \theta^i(t_{e,z})$, and since $t_{e,z} = (\sigma, -\frac{1}{2}\sigma, -\frac{1}{2}\sigma)$, this gives $[w_+ \otimes x, w_- \otimes y] = \hat{q}(x, y) u' \otimes \theta^i(1, -\frac{1}{2}, -\frac{1}{2})$ (where again θ denotes the cyclic permutation of the components). Thus,

$$d_{\iota_i(x), \iota_i(y)}^-(\iota_j(z)) = \begin{cases} \iota_i(\hat{q}(x, y)z) & \text{if } j = i, \\ \iota_{i+1}(-\frac{1}{2}\hat{q}(x, y)z) & \text{if } j = i + 1 \text{ (modulo 3)}, \\ \iota_{i+2}(-\frac{1}{2}\hat{q}(x, y)z) & \text{if } j = i + 2 \text{ (modulo 3)}. \end{cases}$$

The different items in the example above follow at once from these computations.

REFERENCES

- [All78] Bruce N. Allison, *A class of nonassociative algebras with involution containing the class of Jordan algebras*, Math. Ann. **237** (1978), no. 2, 133–156.
- [AF93] Bruce N. Allison and John R. Faulkner, *Nonassociative coefficient algebras for Steinberg unitary Lie algebras*, J. Algebra **161** (1993), no. 1, 1–19.
- [Eld04] Alberto Elduque, *The magic square and symmetric compositions*, Rev. Mat. Iberoamericana **20** (2004), no. 2, 475–491.
- [Eld07] ———, *The S_4 -action on the Tetrahedron algebra*, Proc. Royal Soc. Edinburgh **137A** (2007), 1227–1248. arXiv:math.RA/0604218.
- [EO07] Alberto Elduque and Susumu Okubo, *Lie algebras with S_4 -action and structurable algebras*, J. Algebra **307** (2007), no. 2, 864–890. arXiv:math.RA/0508558.
- [Fil76] Valerii T. Filippov, *Central simple Malcev algebras*, Algebra and Logic **15** (1976), no. 2, 147–151.
- [FH91] William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics.
- [Gri03] Alexander Grishkov, *Lie algebras with triality*, J. Algebra **266** (2003), no. 2, 698–722.
- [HT07] Brian Hartwig and Paul Terwilliger, *The tetrahedron algebra, the Onsager algebra, and the \mathfrak{sl}_2 loop algebra*, J. Algebra **308** (2007), no. 2, 840–863. arXiv:math-ph/0511004.
- [IT07] Tatsuro Ito and Paul Terwilliger, *Finite-dimensional irreducible modules for the three-point \mathfrak{sl}_2 loop algebra*, arXiv:0707.2313.
- [Kan73] Issai L. Kantor, *Models of the exceptional Lie algebras*, Dokl. Akad. Nauk SSSR **208** (1973), 1276–1279.
- [KW01] Michael K. Kinyon and Alan Weinstein, *Leibniz algebras, Courant algebroids, and multiplications on reductive homogeneous spaces*, Amer. J. Math. **123** (2001), no. 3, 525–550.
- [KMRT98] Max-Albert Knus, Alexander Merkurjev, Markus Rost, and Jean-Pierre Tignol, *The book of involutions*, American Mathematical Society Colloquium Publications, vol. 44, American Mathematical Society, Providence, RI, 1998, With a preface in French by J. Tits.
- [Kuz68] Eugene N. Kuzmin, *Malcev algebras and their representations*, Algebra and Logic **7** (1968), no. 4, 48–69.
- [Mey72] Kurt Meyberg, *Lectures on algebras and triple systems*, The University of Virginia, Charlottesville, Va., 1972, Notes on a course of lectures given during the academic year 1971–1972.
- [Mik92] Pavel O. Mikheev, *On the embedding of Mal'tsev algebras into Lie algebras*, Algebra i Logika **31** (1992), no. 2, 167–173, 221.
- [Oku05] Susumu Okubo, *Symmetric triality relations and structurable algebras*, Linear Algebra Appl. **396** (2005), 189–222.
- [Ons44] Lars Onsager, *Crystal statistics. I. A two-dimensional model with an order-disorder transition*, Phys. Rev. (2) **65** (1944), 117–149.
- [PIS04] José M. Pérez-Izquierdo and Ivan P. Shestakov, *An envelope for Malcev algebras*, J. Algebra **272** (2004), no. 1, 379–393.
- [Sag61] Arthur A. Sagle, *Malcev algebras*, Trans. Amer. Math. Soc. **101** (1961), 426–458.
- [Yam58] Kiyosi Yamaguti, *On the Lie triple system and its generalization*, J. Sci. Hiroshima Univ. Ser. A **21** (1957/1958), 155–160.

DEPARTAMENTO DE MATEMÁTICAS E INSTITUTO UNIVERSITARIO DE MATEMÁTICAS Y
APLICACIONES, UNIVERSIDAD DE ZARAGOZA, 50009 ZARAGOZA, SPAIN
E-mail address: `elduque@unizar.es`

DEPARTMENT OF PHYSICS AND ASTRONOMY, UNIVERSITY OF ROCHESTER, ROCHESTER,
NY 14627, USA
E-mail address: `okubo@pas.rochester.edu`