

Multi-Cycle HST Treasury Program for STIS: Mapping the Galactic Environment of the Sun

Priscilla C. Frisch¹

1 Treasury Program Goal: High S/N and Resolution Survey of Interstellar Clouds within 20 pc of Sun

We do not understand the galactic environment of the Sun. Interstellar clouds form the cosmic "ecosystem" through which the Sun moves. The only way that the physical properties of nearby interstellar material (ISM) can be *surveyed* is through observations of interstellar absorption lines in the ultraviolet and far ultraviolet regions. Over 500 B, A, F, G, and K stars within 20 pc are brighter than 9th mag, and 225 brighter than 6th mag (Fig. 1). In addition, at least two white dwarf stars are in the upwind direction and within 20 pc of the Sun, with $V \sim 11$ mag. Many of these stars could be used to survey the nearby interstellar medium. The key cloud parameters that need to be measured are the cloud density, ionization, temperature and velocity. Each of these variables has a significant effect on the heliosphere boundary conditions.

A survey-mode study of nearby ISM is required to understand the solar environment because most available target stars for this program are cool and faint, and high S/N is needed. An optimal survey would use both STIS E140H and E230H data, at $S/N \geq 50$, for ~ 100 stars, yielding data on Mg II, Mg I, H I or D I, C I, and S II, Zn II, and where possible C II, and C II* and others. Interesting regions consist of those where there are no current data, where the velocities of different clouds degenerate, where there currently is some evidence but no hard data on an ionization gradient, or where an expected velocity component is not seen. In addition, some stars that have already been observed by HST should be reobserved, in search of time-variation in the hydrogen wall that would show that the solar magnetic activity cycle impacts the hydrogen wall, an important signature of the ISM-heliosphere interaction.

The minimum data set per star would include the 2800 Å Mg II and Mg I lines, from which the interstellar electron density can be derived for both warm and cool clouds because of the high dielectronic recombination rate of Mg II in warm $T > 6000$ K clouds. Additional information on either H I, D I, and an element with first ionization potential (FIP) less than 13.6 eV such as sulfur or zinc, then allows reconstruction of the ionization of the sightline. Both the *Copernicus* and HST satellites have successfully observed these species towards both warm and cool stars, however to acquire a data set of consistent quality at high resolution and high-S/N requires a large number of orbits to observe faint stars. STIS MAMA count rates limit possible observations for A and B stars, so that the grating settings will vary between stars. An example of an A-type target star observed at high resolution and near- and far-UV, would require 2–8 orbits per star, for $S/N > 100$. Another example of a star for this program is an 11th magnitude DA white dwarf star, which would require medium resolution data, including both long and short wavelengths, for 5-6 orbits per star.

A survey program that consists of ~ 100 stars will require a large number of orbits such

¹University of Chicago, Department of Astronomy and Astrophysics, 5640 South Ellis Avenue, Chicago, IL 60637 USA; Phone: 773-702-0181; Email: frisch@oddjob.uchicago.edu




Figure 1: Left: Spatial distribution of B, A, F, G, and K stars with 20 pc of the Sun. The spectral types are color coded. Right: Distribution of same set of stars over distance. All stars are in the Hipparcos catalog.

as would be possible under a Multi-Cycle Treasury program. The second reason for advancing a survey of local ISM as a MCTP program is that understanding our galactic environment is a high-impact program, with goals that are difficult to achieve piecemeal through small proposals.

2 Interstellar Clouds and the Solar Galactic Environment

The Sun left the deepest void of the Local Bubble sometime within the past 50,000–120,000 years and entered an interstellar wind with an upwind direction towards the galactic center region. The ISM now surrounding the Sun is warm, partially ionized, and very low density, $n_{\text{H}} \sim 0.24$, $\text{H}^+/\text{H}_{\text{total}} \sim 0.2$, however the velocity of the ISM measured *inside* of the solar system is not seen towards nearby stars in the upwind direction. This and other indicators suggest the Sun crosses into different clouds on the order of $\sim 1/40,000$ years locally. Over much shorter timescales the Sun travels through strong ionization gradients in the ISM, because of both the low cloud opacities and the distribution of ionization sources. The ISM surrounding the Sun shapes the heliosphere, which in turn modulates the galactic cosmic ray flux at Earth, see Fig. 2. The cosmic ray record is traced by the radioisotope record on Earth, which now provides coverage for the past ~ 0.5 Myrs, during which time the Sun has left the Local Bubble interior and moved $\sim 8 - 10$ pc through space, and possibly encountered different types of ISM. An accurate map of the distribution of nearby ISM, along with the physical properties of that ISM, potentially will allow the history of the Earth’s recent exposure to galactic cosmic rays to be calculated from physical properties of clouds that have crossed our cosmic trail in space.

The present day interaction between the Sun and surrounding ISM is traced by: (1) Direct observations of ISM inside of the heliosphere, including H, He, N, O, Ne, Ar, and interstellar dust grains. Observations of the florescence of solar $\text{Ly}\alpha$ radiation off of interstellar H° inside of the heliosphere by *Copernicus* during the mid-1970s, and HST during the mid-1990’s, show that the Sun has been in the same interstellar cloud for the past ~ 20 years, during which time





Figure 2: Left: The heliosphere today, with a weak Mach 1 bow shock (purple), heliopause, termination shock (greenish semispherical surface), and heliotail (figure from <http://antwarp.gsfc.nasa.gov/apod/ap020624.html>). Right: The importance of the configuration of the outer heliosphere for galactic cosmic ray modulation, can be seen, since today most of the GCR modulation is in the inner heliosheath whose properties are strongly affected by the surrounding ISM (figure from Wimmer-Schweingruber, McNutt, et al. 2007, submitted to *Experimental Astronomy*).

the heliosphere has sampled a pathlength of $\sim 100AU$ of this cloud. (2) Lyman α observations of the "hydrogen wall", formed where interstellar protons and hydrogen-atoms couple by charge exchange as the ions are compressed and diverted around the heliosphere. (3) Energetic neutral atoms (ENAs) formed by charge-exchange between interstellar H° atoms and the solar wind. The Interstellar Boundary EXplorer (IBEX) mission to be launched in 2008 will directly measure the properties of the ISM-heliosphere interaction through measurements of ENAs formed in the outer heliosphere.

3 The Heliosphere and Terrestrial Climate

Interstellar clouds form the boundary conditions of the heliosphere. There are major differences in the way two hypothetical interstellar clouds with the same heliocentric velocities, V_{HC} , same temperature and total density, but different ionizations, will interact with the heliosphere. This is because interstellar neutrals enter the heliosphere, become ionized, and mass-load the solar wind, thereby enhancing the solar wind barrier to galactic cosmic rays, whereas in contrast interstellar ions increase the external ram pressure of the ISM on the heliosphere. The net result is that the flux of galactic cosmic rays at Earth varies with the solar galactic environment. The cosmic ray attenuation by the heliosphere is illustrated in Fig. 2. There is evidence that galactic cosmic rays seed the formation of terrestrial clouds, thereby cooling Earth.² There is strong circumstantial evidence that increased fluxes of galactic cosmic rays at Earth (as measured by ^{10}Be isotopes in ice cores) are correlated with cold climate spells, such as the Maunder Minimum $\sim 1640 - 1700$.

²A concise reference describing the physical processes that link the galactic environment of the Sun with the terrestrial climate is the book *Solar Journey: The Significance of Our Galactic Environment for the Heliosphere and Earth*, Ed. P. C. Frisch, Springer 2006.

4 Appendix: End Notes

The above summary was prepared in response to a request for 3-page white papers discussing Multi-Cycle Treasury Programs for the Hubble Space Telescope. A more complete list of references for the items listed above are given in this end note. Fig. 1: The stars that are plotted are from the Hipparcos star catalog [4]. Credits for the magnetohydrodynamic heliosphere model visualization in Fig. 2, left, are A. Hanson, P. Fu, P. Frisch, and [2]. An additional reference for Fig. 2, right, is [5]. Discussions of the changing galactic environment of the Sun over short time scales of $\sim 10^5$ years can be found in [1]. Additional information about the IBEX mission can be found in [3] and at the URL <http://www.ibex.swri.edu>. The ionization gradient in the cloud surrounding the Sun, and constancy of the cloud's velocity over ~ 20 years, are discussed in [6]; this reference also cites references for measurements of interstellar H, He, O, N, Ne, and Ar inside of the heliosphere. Hydrogen-wall data are discussed in [7]. The processes that relate our galactic environment to the terrestrial climate are discussed in chapters of the book *Solar Journey*, listed in footnote 2.

References

- [1] P. C. Frisch and J. D. Slavin. The Sun's Journey through the Local Interstellar Medium: The PaleoLISM and Paleoheliosphere. *Astrophysics and Space Sciences Transactions*, 2:53–61, 2006.
- [2] T. J. Linde. *A Three-Dimensional Adaptive Multifluid MHD Model of the Heliosphere*. PhD thesis, Univ. of Michigan, Ann Arbor, 1998. <http://hpcc.engin.umich.edu/CFD/publications>.
- [3] D. McComas, F. Allegrini, P. Bochsler, M. Bzowski, M. Collier, H. Fahr, H. Fichtner, P. Frisch, H. Funsten, S. Fuselier, G. Gloeckler, M. Gruntman, V. Izmodenov, P. Knappenberger, M. Lee, S. Livi, D. Mitchell, E. Möbius, T. Moore, D. Reisenfeld, E. Roelof, N. Schwadron, M. Wieser, M. Witte, P. Wurz, and G. Zank. The Interstellar Boundary Explorer (IBEX). In *AIP Conf. Proc. 719: Physics of the Outer Heliosphere*, pages 162–181, 2004.
- [4] M. A. C. Perryman. The HIPPARCOS Catalogue. *A&A* , 323:L49–L52, 1997.
- [5] N. A. Schwadron, C. Goodrich, H. Spence, L. Townsend, F. Cucinotta, M. Weyland, M. Gollightly, M. Desai, A. Posner, D. Hassler, D. Krauss-Varban, J. Luhmann, J. Miller, B. Heber, and T. Onsager. "Earth-Moon-Mars radiation environment module (EMMREM), Aerospace Conference, 2007". *IEEE*, pages 1–10, 2007.
- [6] J. D. Slavin and P. C. Frisch. The Boundary Conditions of the Heliosphere: Photoionization Models Constrained by Interstellar and In Situ Data. *A&A* , *submitted*, 000:0000–0000, 2007.
- [7] B. E. Wood, S. Redfield, J. L. Linsky, H.-R. Müller, and G. P. Zank. Stellar Ly α Emission Lines in the Hubble Space Telescope Archive: Intrinsic Line Fluxes and Absorption from the Heliosphere and Astrospheres. *ApJS* , 159:118–140, 2005.