

REMARKS ON LEMPERT FUNCTIONS OF BALANCED DOMAINS

NIKOLAI NIKOLOV AND PETER PFLUG

ABSTRACT. This note should clarify how the behavior of certain invariant objects reflects the geometric convexity of balanced domains.

1. INTRODUCTION AND RESULTS

By \mathbb{D} we denote the unit disc in \mathbb{C} . Let D be a domain in \mathbb{C}^n . Recall first the definitions of the Carathéodory pseudodistance and the Lempert function of D :

$$c_D(z, w) = \sup\{\tanh^{-1} |f(w)| : f \in \mathcal{O}(D, \mathbb{D}) : f(z) = 0\},$$

$$\tilde{k}_D(z, w) = \inf\{\tanh^{-1} |\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) : \varphi(0) = z, \varphi(\alpha) = w\}.$$

The Kobayashi pseudodistance k_D can be defined as the largest pseudodistance below of \tilde{k}_D . Note that if $k_D^{(m)}$ denotes the m -th Lempert function of D , $m \in \mathbb{N}$, that is,

$$k_D^{(m)}(z, w) = \inf\left\{\sum_{j=1}^m \tilde{k}_D(z_{j-1}, z_j) : z_0, \dots, z_m \in D, z_0 = z, z_m = w\right\},$$

then

$$k_D(z, w) = \inf_m k_D^{(m)}(z, w).$$

If l_D is any one of the introduced functions from above, we set $l_D^* = \tanh l_D$.

Recall that D is said to be balanced if $\lambda z \in D$ for any $\lambda \in \overline{\mathbb{D}}$ and any $z \in D$. Denoting by h_D the Minkowski function of D , i.e.,

$$h_D(z) = \inf\{t > 0 : z/t \in D\}, \quad z \in \mathbb{C}^n,$$

2000 *Mathematics Subject Classification.* 32F45.

Key words and phrases. Balanced domain, Minkowski function, Lempert functions, Kobayashi pseudodistance, Carathéodory pseudodistance.

This note was written during the stay of the first named author at the Universität Oldenburg supported by a grant from the DFG, Az. PF 227/9-1 (October 2007). He thanks both institutions for their support.

then

$$D = D_h = \{z \in \mathbb{C}^n : h_D(z) < 1\}.$$

We point out that D is pseudoconvex if and only if $\log h_D$ is a plurisubharmonic function. Set $\widehat{h}_D = h_{\widehat{D}}$, where \widehat{D} is the convex hull of D .

Let us summarize some well-known facts about relations between h_D , \widehat{h}_D , and the functions from above, where one of their arguments is the origin.

Proposition 1. (cf. [1]) *Let $D \subset \mathbb{C}^n$ be a balanced domain and $a \in \mathbb{D}$. Then*

- (i) $\widehat{h}_D \leq c_D^*(0, \cdot) \leq \widetilde{k}_D^*(0, \cdot) \leq h_D$;
 - (ii) $k_D^*(0, a) = h_D(a) \iff h_D(a) = \widehat{h}_D(a)$.
- If, in addition, D is pseudoconvex, then

$$\widetilde{k}_D^*(0, \cdot) = h_D.$$

Proposition 1 shows that at a point $a \in D$ the value of $k_D(0, \cdot)$ is maximal if and only if D is "convex in the direction of a ", i.e., $h_D(a) = \widehat{h}_D(a)$. In fact, more is true as the following result shows.*

Proposition 2. *Let $D \subset \mathbb{C}^n$ be a balanced domain and $a \in D$. Then*

$$(k_D^{(3)}(0, a))^* = h_D(a) \iff h_D(a) = \widehat{h}_D(a).$$

Remark. We do not know if the number 3 can be replaced by 2.

Conversely, one may ask whether the fact that $l_D(0, a)$ (l_D as above) is "minimal" (i.e., $l_D(0, a) = h_G(a)$ for some domain G containing D) implies also some convexity property. Here we start with the following result.

Proposition 3. *Let $D \subset \mathbb{C}^n$ be a bounded balanced domain and $G \subset \mathbb{C}^n$ be a pseudoconvex balanced domain with $G \supset D$. Assume that h_D is continuous at some $a \in D$ and \overline{G} contains no nontrivial analytic discs through $a/h_G(a)$, $h_G(a) \neq 0$. Then*

$$\widetilde{k}_D^*(0, a) = h_G(a) \iff h_D(a) = h_G(a).$$

Remarks. (a) Since the envelope of holomorphy $\mathcal{E}(D)$ of a balanced domain D is balanced (see [2], Remark 3.1.2(b)), one may apply the above result for D and $\mathcal{E}(D)$.

(b) If h_G is continuous near a and ∂G contains no nontrivial discs through $a/h_G(a)$, then the maximum principle implies that \overline{G} contains no nontrivial analytic discs through $a/h_G(a)$, too.

*The proofs of the following propositions and examples will be given in Section 2.

(c) In light of Proposition 3, it is natural to ask whether there is a non-pseudoconvex balanced domain D such that $h_D = \tilde{k}_D^*(0, \cdot)$ on D . The authors do not know the answer.

The following examples show that the assumptions about continuity of h_D at a and discs in Proposition 3 are essential.

Example 4. *If $D = \mathbb{D}^2 \setminus \{(t, t) : |t| \geq 1/2\}$, $d = (t, t)$, $|t| < 1/2$, then*

$$h_D(d) = 2|t| \quad \text{but} \quad \tilde{k}_D^*(0, d) = |t|.$$

On the other hand, $\overline{\mathbb{D}^2}$ contains no nontrivial analytic discs through any point of $\partial\mathbb{D} \times \partial\mathbb{D}$.

Moreover, the following example gives a balanced Reinhardt domain D such that $G = \mathcal{E}(D) = \widehat{D}$ has nontrivial analytic discs in its boundary for which Proposition 3 fails to hold. Note that in this case h_D and h_G are continuous functions.

Example 5. *Let $0 < a < 1$ and*

$$D = \{z \in \mathbb{D}^2 : |z_2|^2 - a^2 < 2(1 - a^2)|z_1|\}.$$

Then D is a balanced Reinhardt domain and $\mathcal{E}(D) = \mathbb{D}^2$ (cf. [2]). On the other hand, if $c = (0, d)$, $|d| < a$, then

$$h_D(c) = \frac{|d|}{a} > \tilde{k}_D^*(0, c) = |d| = \widehat{h}_D(c).$$

For a balanced domain D and $a \in D$ set

$$\mathbb{D}_{D,a} := \{\lambda a : |\lambda| h_D(a) < 1\}.$$

Assuming minimality along the whole slice $\mathbb{C}a \cap D = \mathbb{D}_{D,a}$ for some $a \in D$ we have

Proposition 6. *Let $D \subset \mathbb{C}^n$ be a taut balanced domain, $a \in D$ and $m \in \mathbb{N}$. Then*

$$(k_D^{(m)}(0, \tilde{a}))^* = \widehat{h}_D(\tilde{a}), \quad \tilde{a} \in \mathbb{D}_{D,a} \iff h_D(a) = \widehat{h}_D(a).$$

Recall that a domain $D \subset \mathbb{C}^n$ is said to be taut if $\mathcal{O}(\mathbb{D}, D)$ is a normal family. Note that a balanced domain D is taut if and only if h_D is a continuous plurisubharmonic function and $(h_D)^{-1}(0) = 0$. Weakening the continuity assumption for h_D the following statement remains true.

Proposition 7. *Let $D \subset \mathbb{C}^n$ be a bounded pseudoconvex balanced domain, $a \in D$ and $m \in \mathbb{N}$. Assume that h_D is continuous at a and ∂D contains no nontrivial analytic discs through $a/h_D(a)$. Then*

$$(k_D^{(m)}(0, \tilde{a}))^* = \widehat{h}_D(\tilde{a}), \quad \tilde{a} \in \mathbb{D}_{D,a} \iff h_D(a) = \widehat{h}_D(a).$$

Remark. We do not know if the condition about discs is superfluous or not. On the other hand, continuity and pseudoconvexity are essential as Examples 4 and 5 have shown (see also Example 9 below).

Recall that a boundary point b of a domain D in \mathbb{C}^n is said to be a *local weak barrier point* if there are a neighborhood U of b and a negative plurisubharmonic function u on $D \cap U$ such that $\lim_{D \ni z \rightarrow b} u(z) = 0$.

Proposition 7 is a consequence of the following

Proposition 8. *Let b be a local weak barrier point of a bounded domain $D \subset \mathbb{C}^n$. If ∂D contains no nontrivial analytic discs through b , then*

$$\lim_{w \rightarrow b} k_D^{(m)}(z, w) = \infty, \quad z \in D, \quad m \in \mathbb{N}.$$

We point out that the assumption about tautness is essential in Proposition 6 as the following examples show.

Example 9. *The unit ball \mathbb{B} in \mathbb{C}^2 contains a proper non-taut pseudoconvex balanced domain D with $\widehat{D} = \mathbb{B}$ such that*

$$(k_D^{(2)}(0, \cdot))^* = \|\cdot\|.$$

Example 10. *There is an unbounded pseudoconvex balanced Reinhardt domain D in \mathbb{C}^2 with continuous Minkowski function and a point $a \in D$ such that $\widehat{h}_D(a) > 0$, ∂D contains a nontrivial analytic disc through $a/h_D(a)$,*

$$(k_D^{(2)}(0, \lambda a))^* = |\lambda| \widehat{h}_D(a), \quad |\lambda| \leq 1,$$

but even

$$c_D^*(0, \lambda a) > |\lambda| \widehat{h}_D(a), \quad 1 < |\lambda| < 1/h(a).$$

Remark. Despite of these examples, we do not know any example of a taut balanced domain D such that $(k_D^{(m)}(0, a))^* = \widehat{h}_D(a)$ for some $m \in \mathbb{N}$ and some $a \in D$, but $h_D(a) > \widehat{h}_D(a)$.

2. PROOFS

Proof of Proposition 2. We have only to prove that

$$(k_D^{(3)}(0, a))^* = h_D(a) \Rightarrow h_D(a) = \widehat{h}_D(a).$$

First, we shall show that

$$(1) \quad (k_D^{(2)}(0, \lambda a))^* = |\lambda| h_D(a), \quad \lambda \in \mathbb{D}.$$

We may assume that $h_D(a) \neq 0$. Taking the disc $\mathbb{D} \ni t \rightarrow ta/h_D(a)$ as a competitor for $\tilde{k}_D(\lambda a, a)$ gives

$$\tilde{k}_D(\lambda a, a) \leq p(h_D(\lambda a), h_D(a)),$$

where p denotes the Poincaré distance. This and the inequality

$$p(0, h_D(a)) = k_D^{(3)}(0, a) \leq k_D^{(2)}(0, \lambda a) + \tilde{k}_D(\lambda a, a)$$

imply that

$$p(0, |\lambda|h_D(a)) = p(0, h_D(a)) - p(|\lambda|h_D(a), h_D(a)) \leq k_D^{(2)}(0, \lambda a).$$

So

$$(k_D^{(2)}(0, \lambda a))^* \geq |\lambda|h_D(a)$$

and the opposite inequality always holds.

It follows from (1) that

$$\lim_{\lambda \rightarrow 0} \frac{k_D^{(2)}(0, \lambda a)}{|\lambda|} = h_D(a).$$

On the other hand, by Proposition 2 in [4], this limit does not exceed

$$\kappa_D^{(2)}(0; a) := \inf\{\kappa_D(0; a_1) + \kappa_D(0; a_2) : a_1 + a_2 = a\},$$

where κ_D denotes the Kobayashi–Royden pseudometric of D . Since $\kappa_D(0; \cdot) \leq h_D$ (cf. [1]), we conclude that

$$h_D(a) \leq h_D(a_1) + h_D(a_2) \text{ if } a_1 + a_2 = a,$$

which means that $h_D(a) = \widehat{h}_D(a)$. □

Proof of Proposition 3. We have only to prove that

$$\tilde{k}_D^*(0, a) = h_G(a) \Rightarrow h_D(a) \leq h_G(a).$$

Let $(\varphi_j) \subset \mathcal{O}(\mathbb{D}, D)$ and $\alpha_j \rightarrow h_G(a)$ be such that $\varphi_j(0) = 0$ and $\varphi_j(\alpha_j) = a$. Writing φ_j in the form $\varphi_j(\lambda) = \lambda\psi_j(\lambda)$, $\psi_j \in \mathcal{O}(\mathbb{D}, \mathbb{C}^n)$, it follows by the maximum principle that $h_G \circ \psi_j \leq 1$ and hence $\psi_j \in \mathcal{O}(\mathbb{D}, \overline{G})$. Since D is bounded, we may assume that $\varphi_j \rightarrow \varphi \in \mathcal{O}(\mathbb{D}, \overline{D})$ and then $\psi_j \rightarrow \psi \in \mathcal{O}(\mathbb{D}, \overline{G})$. On the other hand, since \overline{G} contains no nontrivial analytic discs through $\psi(h_G(a)) = b$, where $b = a/h_G(a)$, it follows that $\psi(\mathbb{D}) = b$. Using that h_D is continuous at b , we get that

$$1 > h_D(\varphi_j(\lambda)) \rightarrow |\lambda|h_D(b), \quad \lambda \in \mathbb{D}.$$

Letting $\lambda \rightarrow 1$ leads to $h_D(b) \leq 1$ which is the desired inequality. □

Proof of Example 4. We have only to prove that

$$\tilde{k}_D^*(0, d) \leq |t|.$$

For any $r \in (|t|, 1)$ we may choose $\alpha \in \mathbb{D}$ such that $t = \varphi(t/r)$, where $\varphi(\lambda) = \lambda \frac{\lambda - \alpha}{1 - \bar{\alpha}\lambda}$. Then $\psi = (\text{rid}, \varphi) \in \mathcal{O}(\mathbb{D}, D)$ is a competitor for $\tilde{k}_D^*(0, d)$ which shows that $\tilde{k}_D^*(0, d) \leq |t|/r$. It remains to let $r \rightarrow 1$. □

Remarks. (a) Let D be the domain from Example 4. Note that even

$$\tilde{k}_D(0, \cdot) = \tilde{k}_{\mathbb{D}^2}(0, \cdot).$$

It is enough to prove that $\tilde{k}_D(0, a) \leq |a_1|$ for $a = (a_1, a_2) \in D$, $a_1 \neq a_2$, $|a_1| \geq |a_2|$. For this, take the discs $\psi(\lambda) = (\lambda, \lambda a_2/a_1)$ as a competitor for $\tilde{k}_D(0, a)$.

On the other hand, if $a_1 = (0, b)$ and $a_2 = (b, 0)$, $b \in \mathbb{D}$, then

$$\tilde{k}_D(a_1, a_2) = \tilde{k}_{\mathbb{D}^2}(a_1, a_2) \iff |b| \leq 4/5.$$

Indeed, using the Möbius transformation $\psi_b(\lambda) = \frac{\lambda-b}{1-b\lambda}$, we get that

$$\tilde{k}_D(a_1, a_2) = \tilde{k}_{D_b}(0, a),$$

where $a = (b, -b)$ and $D_b = \mathbb{D}^2 \setminus \{(\psi_b(\lambda), \lambda) : 1/2 \leq |\lambda| < 1\}$.

For $|b| < 4/5$, it is easy to check that $\varphi = (\text{id}, -\text{id}) \in \mathcal{O}(\mathbb{D}, D_b)$. This implies that $\tilde{k}_{D_b}^*(0, a) \leq |b|$ and hence $\tilde{k}_D(a_1, a_2) = \tilde{k}_{\mathbb{D}^2}(a_1, a_2)$.

To get the same for $|b| = 4/5$, it is enough to take $r\varphi$, $r \in (0, 1)$, as a competitor for $\tilde{k}_{D_b}^*(0, a)$ and then to let $r \rightarrow 1$.

Assume now that $|b| > 4/5$ and $\tilde{k}_D(a_1, a_2) = \tilde{k}_{\mathbb{D}^2}(a_1, a_2)$. Then we may find discs $\varphi_j \in \mathcal{O}(\mathbb{D}, D_b)$ such that $\varphi_j(0) = 0$ and $\varphi_j(\alpha_j) = a$, where $\alpha_j \rightarrow b$. It follows by the Schwarz-Pick lemma that $\varphi_j \rightarrow \varphi$. On the other hand, $\varphi(\mathbb{D}) \cap \{(\psi_b(\lambda), \lambda) : 1/2 < |\lambda| < 1\}$ is a singleton which contradicts to Hurwitz's theorem.

(b) Note that (a) shows that Theorem 3.4.2 in [1] is in some sense sharp. On the other hand, this theorem implies that, if $D_n = \mathbb{D}^n \setminus \{(t, \dots, t) : |t| \geq 1/2\}$, $n \geq 3$, then

$$\tilde{k}_{D_n} = \tilde{k}_{\mathbb{D}^n}.$$

Proof of Example 5. We have only to prove that $\tilde{k}_D^*(0, c) \leq d$, $d \in (0, a)$. For this, it is enough to show that $\varphi = (\psi, \text{id}) \in \mathcal{O}(\mathbb{D}, D)$, where $\psi(\lambda) = \lambda \frac{\lambda-d}{1-d\lambda}$. Since $|\psi(\lambda)| \geq x \frac{x-d}{1-dx}$ for $x = |\lambda|$, we have to check that

$$(x^2 - a^2)(1 - dx) < 2(1 - a^2)x(x - d), \text{ i.e.,}$$

$$dx^3 + (1 - 2a^2)x^2 - d(2 - a^2)x + a^2 > 0.$$

Note that this inequality is true for $x = 0$. Using that $x \in (0, 1)$ and $d \in (0, a)$, it suffices to prove that

$$ax^3 + (1 - 2a^2)x^2 - a(2 - a^2)x + a^2 \geq 0$$

which is equivalent to the obvious inequality $(x - a)^2(ax + 1) \geq 0$. \square

Proof of Proposition 6. It is enough to show that if $\mathbb{D}_{D,a} \ni a_k \rightarrow a/h_D(a)$, $a \neq 0$, and $(k_D^{(m)}(0, \tilde{a}_k))^* = \widehat{h}_D(\tilde{a}_k)$, then $h_D(a) = \widehat{h}_D(a)$. For this, recall that if D is a taut domain, then (cf. [1], Proposition 3.2.1)

$$\lim_{w \rightarrow \partial D} k_D^{(m)}(z, w) = \infty, \quad w \in D.$$

In our case this implies that $\widehat{h}_D(\tilde{a}_k) \rightarrow 1$ and hence $\widehat{h}_D(a/h(a)) = 1$. \square

Proof of Proposition 8. We shall proceed by induction on m . For this, we shall need the following.

Lemma 11. *Under the assumptions of Proposition 8, for any $(\varphi_k) \subset \mathcal{O}(\mathbb{D}, D)$ with $\varphi_k(0) \rightarrow b$ one has that $\varphi_k \rightarrow b$ locally uniformly on \mathbb{D} .*

Assuming Lemma 11 easily implies that Proposition 8 is true for $m = 1$. Suppose that this statement is true for some $m - 1 \in \mathbb{N}$ but false for m . Then we may find $z \in D$ and sequences $(z_{j,k})_k \subset D$, $0 \leq j \leq m$, such that $z_{0,k} = z$, $z_{m,k} \rightarrow b$ and

$$\sup_k \sum_{j=1}^m \tilde{k}_D(z_{j-1,k}, z_{j,k}) < \infty.$$

In virtue of our induction hypothesis one has that $z_{m-1,k} \not\rightarrow b$. Passing to a subsequence, we may assume that $z_{m-1,k} \rightarrow a \in \overline{D}$, $a \neq b$ and $(\tilde{k}_D(z_{m-1,k}, z_{m,k}))^* \rightarrow r < 1$. Then there are $\varphi_k \in \mathcal{O}(\mathbb{D}, D)$ with $\varphi_k(0) = z_{m,k}$, $\varphi_k(r_k) = z_{m-1,k}$ and $r_k \rightarrow r$ which contradicts Lemma 11. \square

Proof of Lemma 11. Since D is bounded, it is easily seen that for any neighborhood U of b there is another neighborhood $V \subset U$ of b and a number $s \in (0, 1]$ such that, if $\varphi \in \mathcal{O}(\mathbb{D}, D)$ with $\varphi(0) \in V$, then $\varphi(s\mathbb{D}) \subset D \cap U$.

Now we choose U such that there is a negative plurisubharmonic function u on $D \cap U$ with $\lim_{D \ni z \rightarrow b} u(z) = 0$. Applying Proposition 3.4 in [3] implies that b is a t -point for $D \cap D$ which means that

(*) $(\varphi_j|_{s\mathbb{D}})$ is compactly divergent w.r.t. $D \cap U$.

Assume now that (φ_j) does not converge to b . Passing to a subsequence, we may assume that $\varphi_j \rightarrow \varphi \in \mathcal{O}(\mathbb{D}, \overline{D})$ and $\varphi(\mathbb{D}) \neq \{b\}$. It follows by (*) that $\varphi(s\mathbb{D}) \subset \partial(D \cap U)$. Since $\varphi(0) = b \in \partial D \cap U$, we may find $s' \in (0, s]$ such that $\varphi(s'\mathbb{D}) \subset \partial D$. Using that ∂D contains no nontrivial analytic discs through b , we get that $\varphi(s'\mathbb{D}) = \{b\}$ which contradicts the identity principle. \square

Proof of Example 9. J. Siciak (cf. [1], Example 3.1.12) constructed a plurisubharmonic function $\psi : \mathbb{C}^2 \rightarrow [0, \infty)$ such that $\psi(\lambda z) = |\lambda|\psi(z)$ ($\lambda \in \mathbb{C}, z \in \mathbb{C}^n$), $\psi \not\equiv 0$, but $\psi = 0$ on a dense subset S of \mathbb{C}^n . Set

$$D = \{z \in \mathbb{C}^2 : \|z\| + \psi(z) < 1\}.$$

For any $a \in D$ we may choose a sequence $S \cap D \supset (z_k) \rightarrow a$. Then

$$\tanh^{-1} \|a\| = k_{\mathbb{B}_2}^{(2)}(0, a) \leq k_D^{(2)}(0, a) \leq \tilde{k}_D(0, z_k) + \tilde{k}_D(z_k, a).$$

Letting $k \rightarrow \infty$ gives $k_D^{(2)}(0, a) = \tanh^{-1} \|a\|$. \square

Proof of Example 10. Setting

$$D = \{z \in \mathbb{C}^2 : |z_1| < 1, |z_1^{34} z_2^{55}| < 1\},$$

then

$$c_D^*(0, z) = \max\{|z_1|, |z_1 z_2|, |z_1^2 z_2^3|, |z_1^5 z_2^8|, |z_1^{13} z_2^{21}|, |z_1^{34} z_2^{55}|\}$$

(see [1], Example 2.7.12). Since $\widehat{D} = \mathbb{D} \times \mathbb{C}$, we have that

$$c_D^*(0, z) > \widehat{h}_D(z) = |z_1| \iff |z_2| > 1.$$

On the other hand, note that $\{0\} \times \mathbb{C} \subset D$ and $\mathbb{D} \times \{z_2\} \subset D, |z_2| \leq 1$. So if $|z_1| < 1, |z_2| \leq 1$, then

$$k_D^{(2)}(0, (z_1, z_2)) \leq k_D(0, (0, z_2)) + k_D((0, z_2), (z_1, z_2)) \leq k_{\mathbb{D}}(0, z_1)$$

and hence

$$(k_D^{(2)}(0, (z_1, z_2)))^* \leq |z_1| = \widehat{h}_D((z_1, z_2)).$$

Since the opposite inequality always holds and ∂D contains a nontrivial analytic disc through any point $b = (b_1, b_2) \in \partial D$ with $|b_1| \neq |b_2|$, it follows that any point $a = (a_1, a_2)$ with $|a_1| < |a_2| = 1$ has the desired properties. \square

REFERENCES

- [1] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter Exp. Math. 9, de Gruyter, Berlin, New York, 1993.
- [2] M. Jarnicki, P. Pflug, *Extension of holomorphic functions*, de Gruyter Exp. Math. 34, de Gruyter, Berlin, New York, 2000.
- [3] N. Nikolov, P. Pflug, *Local vs. global hyperconvexity, tautness or k -completeness for unbounded open sets in \mathbb{C}^n* , Ann. Scuola Norm. Sup. Pisa Cl. Sci (5), 4, 601 – 618 (2005).
- [4] N. Nikolov, P. Pflug, *On the derivatives of the Lempert functions*, Ann. Mat. Pura Appl., DOI 10.1007/s10231-007-0056-z.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA

E-mail address: `nik@math.bas.bg`

CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, INSTITUT FÜR MATHEMATIK, FAKULTÄT V, POSTFACH 2503, D-26111 OLDENBURG, GERMANY

E-mail address: `pflug@mathematik.uni-oldenburg.de`