

Subregular characters of the unitriangular group over a finite field

Mikhail V. Ignatyev*

0. Introduction

Let k be a field and n be a natural number. By $G_n(k) = \text{UT}(n, k)$ we denote the group of all unipotent lower-triangular $n \times n$ -matrices with coefficients from k ; this group is called a *unitriangular* group. By $\mathfrak{g}_n(k) = \mathfrak{ut}(n, k)$ we denote its Lie algebra over k ; this Lie algebra consists of all nilpotent lower-triangular matrices with coefficients in k .

If $k = \mathbb{F}_q$ is a finite field, then $G_n(q) = G_n(k)$ is a finite group; so, there are finitely many classes of equivalency of irreducible complex representations of this group. A description of all irreducible characters (or of some series of them) is a classical problem of representation theory. The orbit method of A.A. Kirillov [K1], [K2] allows to reduce the similar problem of description of unitary irreducible representations of Lie groups to the problem of classification of coadjoint orbits; this method is also valid for $G_n(q)$ (see. [Ka]), but complete classification of coadjoint orbits for an arbitrary n is unknown.

A description of orbits of the principal series (i.e., orbits of maximum dimension) of Lie groups was presented in the pioneering work on the orbit method [K2]; it's also valid over a finite field [K3]. In C. Andre's works (see [A1], [A2]) so-called *basic characters* are described (in particular, an exact formula for characters of the principal series is found).

The problem of description of orbits, representations and characters of sub-maximum dimension is a natural generalisation of these results; such orbits, representations and characters are called *subregular*. They play an important role in algebraic geometry and K -theory (see, for example, [Lu]). Subregular orbits are described in [IP]. The main goal of this paper is to give an exact formula for the corresponding characters. This formula shows that subregular characters (as characters of the principal series) can be described in terms of coefficients of minors of the characteristic matrix.

The paper is organized as follows. In section 1 we collect some basic facts about the group G_n and discuss Andre's formulae for characters of the principal series (Theorem 1.12). In section 2 we formulate the Main Theorem including exact formulas for subregular characters (Theorem 2.9). More precisely, for an arbitrary subregular character we find the elements such that the value of this character on the conjugacy classes of these elements is non-zero and compute this value. Section 3 is devoted to the description of these conjugacy classes (Theorem 3.5). In section 4 we recall some facts about semi-direct decomposition of the group G_n , which are needed for the proof of the main Theorem, which is given in section 5. Finally, section 6 includes some remarks about discussed problems.

*This research was supported by Samara regional grant for students and young scientists no. 23E2.1D

1. Characters of the principal series

Let $k = \mathbb{F}_q$ and $\text{char } k = p$, i.e., $q = p^r$, where p is a prime number. Throughout the paper, we suppose $p \geq n$. According to [Ka], under these assumptions the orbit method is valid: there is a one-to one correspondence between the set of all irreducible complex characters of our group and the set of all coadjoint orbits, i.e, $G_n(q)$ -orbits in the dual space $\mathfrak{g}_n^*(q)$. Moreover, to each orbit $\Omega \subset \mathfrak{g}_n(q)$ the character

$$\chi_\Omega(\exp a) = q^{-\frac{1}{2} \dim \Omega} \cdot \sum_{f \in \Omega} \theta(f(a)), \quad a \in \mathfrak{g}_n(q), \quad (1)$$

is assigned (here $\theta: \mathbb{F}_q \rightarrow \mathbb{C}$ is a non-trivial character of the additive group of the field \mathbb{F}_q). On the other hand, there are no exact formulas that allow to find the value of the character of a given orbit on a given element of the unitriangular group.

Remark 1.1. There exists a form $\langle A, B \rangle = \text{tr}(AB)$ which is non-degenerate on $\mathfrak{gl}_n(k)$. This allows to identify $\mathfrak{ut}^*(n, q)$ with the space of all nilpotent upper-triangular matrices by the formula $f(x) = \langle f, x \rangle = \sum_{i,j} \xi_{ji} x_{ij}$, where $x = (x_{ij}) \in \mathfrak{g}_n(q)$, $f = (\xi_{ij}) \in \mathfrak{g}_n^*(q)$. Then the coadjoint action

$$K(g): \mathfrak{g}_n^* \rightarrow \mathfrak{g}_n^*: (K(g)f)(x) = f(\text{Ad}_{g^{-1}}x), \quad g \in G_n, \quad f \in \mathfrak{g}_n^*, \quad x \in \mathfrak{g}_n$$

is given by

$$K(g): x \mapsto (g x g^{-1})_{\text{high}}, \quad x \in \mathfrak{g}_n^*, \quad g \in G_n,$$

where $(a)_{\text{high}}$ denotes the following matrix: its elements below the main diagonal coincide with the corresponding elements of the matrix a and its elements on the main diagonal and below are equal to zero.

Representations of maximum dimension play an important role in this theory. These representations (and their orbits and characters) form the so-called *principal series*; orbits of maximum dimension are called *regular*. We'll use the following

Notation 1.2. Let $n \in \mathbb{N}$. Set $n_0 = [n/2]$, $n_1 = [(n-1)/2]$. Note that $n = n_0 + n_1 + 1$.

Notation 1.3. Let $g = (y_{ij}) \in \text{Mat}(n, k)$. By $\Delta_{i_1, \dots, i_k}^{j_1, \dots, j_k}(g)$ we denote the minor of the matrix g with the rows i_1, \dots, i_k and the columns j_1, \dots, j_k in the given order (for a given $1 \leq k \leq n, 1 \leq i_1, \dots, i_k, j_1, \dots, j_k \leq n$). In particular, for an arbitrary $1 \leq d \leq n_0$ we set

$$\Delta_d(g) = \Delta_{n-d+1, n-d+2, \dots, n}^{1, 2, \dots, d}(g).$$

We also denote $\Delta^X(g) = \Delta_{\sigma(i_1, \dots, i_k)}^{\tau(j_1, \dots, j_k)}(g)$, where X is the set of the pairs $X = \{(i_1, j_1), \dots, (i_k, j_k)\}$ and σ, τ are permutations such that $\sigma(i_1) < \dots < \sigma(i_k)$ and $\tau(j_1) < \dots < \tau(j_k)$.

A complete description of orbits of the principal series is well-known [K3]:

Theorem 1.4. Let $k = \mathbb{F}_q$ be a finite field. An arbitrary regular orbit has the following defining equations:

$$\Delta_d({}^t a) = \beta_d, \quad a \in \mathfrak{g}_n^*(q), \quad 1 \leq d \leq n_0$$

(here ${}^t a$ is the transpose of the matrix a , β_d are arbitrary scalars from \mathbb{F}_q , $\beta_1, \dots, \beta_{n_0-1} \in \mathbb{F}_q^*$ and $\beta_{n_0} \neq 0$ for odd n). \square

Corollary 1.5. For an arbitrary orbits of the principal series there exists the unique *canonical form of a regular orbit*, i.e., the matrix of the form

$$f = \begin{pmatrix} 0 & \dots & 0 & \xi_{1,n} \\ 0 & \dots & \xi_{2,n-1} & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & 0 & 0 \end{pmatrix},$$

where $\xi_{1,n} = \beta_1, \xi_{d,n-d+1} = \frac{\beta_d}{\beta_{d-1}}, d = 2, \dots, n_0$, that consists in this orbit. \square

Moreover, dimension of a representation (and an orbit) of the principal series is known [Le]:

Theorem 1.6. Let $\mu(n) = (n - 2) + (n - 4) + \dots$ and T_Ω be the representation that corresponds to an orbit Ω of the principal series. Then $\dim \Omega = 2\mu(n)$, $\dim T_\Omega = q^{\mu(n)}$. \square

Note that there are exactly $q^{2\mu(n)}$ points on an orbit Ω of dimension $2\mu(n)$ [K4].

In this section we recall the formula for characters of the principal series, i.e., characters of the form $\chi = \chi_f$, where $f \in \Omega_f \subset \mathfrak{g}_n^*$ is the canonical form on a regular orbit. More precisely, let χ be an irreducible character of the group G_n . By $\text{Supp } \chi$ denote its *support* (i.e., the set $\{g \in G_n \mid \chi(g) \neq 0\}$); obviously, the support is a union of certain conjugacy classes and the value of the character on an arbitrary conjugacy class is constant. So, it's enough to describe the support explicitly and compute the value of the character on an arbitrary conjugacy class containing in the support.

We'll use the following notations from Andre's paper [A2]:

Notation 1.7. We denote by $\Phi(n)$ the set of all pairs $\{(i, j) \mid 1 \leq j < i \leq n\}$ (we call them *roots*, because to each $(i, j) \in \Phi(n)$ the root vector e_{ji} , i.e., the matrix with 1 in the (j, i) -th entry and zeroes elsewhere, is assigned). Let $D \subset \Phi(n)$ be a subset that contains at most one element from each row and at most one element from each column; then this subset is called *basic*. If a basic subset consists from the roots of the form $(n - j + 1, j)$ then it's called *regular*.

Example 1.8. Here we draw one of regular subsets: $D = \{(6, 1), ((4, 3))\} \subset \Phi(6)$. The entries $(i, j) \in D$ are marked by the symbol \otimes :

	1	2	3	4	5	6
1						
2						
3						
4			\otimes			
5						
6	\otimes					

Notation 1.9. For an arbitrary basis subset $D \subset \Phi(n)$ and arbitrary map $\varphi: D \rightarrow \mathbb{F}_q^*$ consider the element of the group $G_n(q)$ of the form

$$x_D(\varphi) = 1_n + \sum_{(n-j+1, j) \in D} \varphi(n-j+1, j) e_{n-j+1, j},$$

where 1_n is the identity $n \times n$ -matrix (if $D = \emptyset$ then $x_D(\varphi) = 1_n$). By $\mathcal{K}_D(\varphi)$ denote the conjugacy class of this element and by \mathcal{K}_{reg} denote the (disjoint) union of $\mathcal{K}_D(\varphi)$ such that D is a regular subset and $\varphi: D \rightarrow \mathbb{F}_q^*$ is a map.

Definition 1.10. Let D be a subset of $\Phi(n)$. A root $(i, j) \in \Phi(n)$ is called *D-regular*, if $(i, k) \notin D$ and $(k, j) \notin D$ for any $i > k > j$. By $R(D)$ denote the set of all *D-regular* roots.

Example 1.11. Let $n = 6$; $D = \{(3, 2), (6, 4)\} \subset \Phi(6)$ is a basic subset. Here we mark the roots $(i, j) \notin R(D)$:

	1	2	3	4	5	6
1						
2						
3		⊗				
4						
5				⊗		
6						

Now, set $\Phi_{\text{reg}} = \{(i, j) \in \Phi(n) \mid i > n - j + 1\}$, $m_D = |R(D) \cap \Phi_{\text{reg}}|$ for an arbitrary regular subset $D \subset \Phi(n)$. Fix also any non-trivial character $\theta: \mathbb{F}_q \rightarrow \mathbb{C}$ of the additive group of the ground field.

Theorem 1.12. Let $\Omega = \Omega_f \subset \mathfrak{g}_n^*(q)$ be a regular orbit, let $f = (\xi_{ij})$ be the canonical form of this orbit and $\chi = \chi_f$ be the corresponding character. Then

1. $\text{Supp } \chi = \mathcal{K}_{\text{reg}}$.
2. $\chi(g) = q^{m_D} \cdot \theta_f(e_D(\varphi))$ for any $g \in \mathcal{K}_D(\varphi) \subset \mathcal{K}_{\text{reg}}$, where $e_D(\varphi) = x_D(\varphi) - 1_n \in \mathfrak{g}_n(q)$ and $\theta_f: \mathfrak{g}_n(q) \rightarrow \mathbb{C}$ is given by the formula

$$\theta_f(x) = \theta(f(x)) = \prod_{(i,j) \in \Phi(n)} \theta(\xi_{ji} x_{ij}), \quad x = (x_{ij}) \in \mathfrak{g}_n(q).$$

Proof. This is a special case of [A2, Theorem 5.1]. \square

In Andre's work [A2] one can find explicit description of $\mathcal{K}_D(\varphi)$ for any basic subset $D \subset \Phi(n)$. More precisely, $g \in \mathcal{K}_D(\varphi) \subset G_n$ if and only if

$$\Delta^{R_D(i,j)}(g) = \Delta^{R_D(i,j)}(x_D(\varphi)) \quad (2)$$

for all $(i, j) \in R(D)$. Here $R_D(i, j) = \{(i, j)\} \cup \{(k, l) \in D \mid l > j \text{ and } k < i\}$.

In particular, if $m = \max_{(i,j) \in D} j$ then

$$y_{ij} = 0 \text{ if } j > m \text{ or } i < n - m + 1. \quad (3)$$

One can see that characters of the principal series and regular orbits can be described in terms of minors of matrices from G_n ; by definition, this is also true for all Andre's basic characters. In the next section we'll see that this is *not* true for subregular characters in general (see [A2]).

2. Subregular characters (statements)

It follows from Theorem 1.4 that a coadjoint orbit "in general position" has maximum dimension $2\mu(n)$. More precisely, the set of points such that their orbits are regular is a dense open subset of $\mathfrak{g}_n^*(q)$ in Zariski topology. Denote this subset by \mathcal{O}_{reg} , it's given by the set of inequalities $\Delta_d \neq 0$, $1 \leq d \leq n_1$ (that's why we use the term "regular orbits"). Recall that $n = n_0 + n_1 + 1$, where $n_0 = \lfloor n/2 \rfloor$, $n_1 = \lfloor (n-1)/2 \rfloor$.

On the other hand, for an arbitrary $1 \leq d \leq n_1$ one can consider the hyper-surface $\mathcal{O}_d \subset \mathfrak{g}_n^*$ given by $\Delta_d = 0$. This hyper-surface splits into the union of certain orbits.


Definition 2.1. An orbit $\Omega \in \mathfrak{g}_n^*$ (and the corresponding representation T_Ω and character χ_Ω) is called *subregular*, if $\dim \Omega = 2\mu(n) - 2$ (resp., $\dim T_\Omega = q^{\mu(n)-1}$), i.e., it has dimension = dimension of a regular orbit $- 2$ (in the other words, this orbit has sub-maximum dimension, because any orbit is even-dimensional [K4]).

Each subregular orbit contains in the unique \mathcal{O}_d and has maximum dimension among all orbits containing in \mathcal{O}_d . Hence, we say that a subregular orbit is d -subregular if it contains in $\Omega \subset \mathcal{O}_d$. In [IP] defining equations of subregular orbits are found, but we will not use them in the sequel; we only list here elements of \mathfrak{g}_n^* such that their orbits are subregular.

Definition 2.2. An element $f = (\xi_{ij}) \in \mathfrak{g}_n^*$ is called a canonical form of a subregular orbit (of the first, second or third type resp.) if

1. There are a number $1 \leq d < n_1$ and $\beta_1, \dots, \beta_{d-1}, \beta', \beta'', \beta_{d+1}, \dots, \beta_{n_0-1} \in k^*$, $\beta_{n_0}, \beta \in k$, where $\beta_{n_0} \neq 0$ for odd n , such that $\xi_{j, n-j+1} = \beta_j$ for any $1 \leq j \leq d-1$ and $d+2 \leq j \leq n_0$, $\xi_{d, n-d} = \beta'$, $\xi_{d+1, n-d+1} = \beta''$, $\xi_{n-d, n-d+1} = \beta$, and $\xi_{ij} = 0$ for all other roots.
2. n is odd and there are $\beta_1, \dots, \beta_{n_1-1} \in k^*$, $\beta', \beta'' \in k$ such that $\xi_{j, n-j+1} = \beta_j$ for any $1 \leq j \leq n_1 - 1$, $\xi_{n_1, n_0+1} = \beta'$, $\xi_{n_1+1, n_0+2} = \beta''$, and $\xi_{ij} = 0$ for all other roots.
3. n is even and there are $\beta_1, \dots, \beta_{n_1-1}, \beta \in k^*$, $\beta', \beta'' \in k$, such that $\xi_{j, n-j+1} = \beta_j$ for any $1 \leq j \leq n_1 - 1$ and either $\xi_{n_1, n_0+1} = \beta$, $\xi_{n_1+1, n_0+2} = \beta'$, $\xi_{n_1+2, n_0+2} = \beta''$ (and all other $\xi_{ij} = 0$), or $\xi_{n_1, n_0} = \beta'$, $\xi_{n_1+1, n_0+2} = \beta$ (and all other $\xi_{ij} = 0$).

Example 2.3. Here we draw some canonical forms of subregular orbits for $n = 8$, $d = 1$ and $d = 2$. The symbol \otimes marks the roots (i, j) such that $\xi_{ji} \neq 0$:



Theorem 2.4. The orbit of a canonical form is subregular. More over, for each subregular orbit there exists the unique canonical form that contains in this orbit.

Proof. See [IP] (there the case $\text{char } k = 0$ is considered, but the proof is still valid in our case $k = \mathbb{F}_q$ if $\text{char } k = p \geq n$). \square

This allows to define d -subregular forms and subregular orbits of the first, second or third type by the obvious way. Clearly, if $1 \leq d < n_1$, then d -subregular orbit is of the first type, and n_1 -subregular orbits are of the second type, if n is odd, and the third one, if n is even.

So, the main goal is to find exact formulas for all characters of the form χ_f , where f is a canonical form of a subregular orbit. As for characters of the principal series, we'll describe the support of a given subregular character and compute the value of this characters on an arbitrary conjugacy class contained in the support (we'll firstly consider the case $1 \leq d < n_1$ and then (in section 6) the case of n_1 -subregular orbits).

Notation 2.5. For an arbitrary $1 \leq d < n_1$ by $D_0(d)$ and $D_1(d)$ denote one of the following sets resp.:

$$\begin{aligned}
 D_0(d) : & \quad \emptyset, \{(n-d+1, d)\}, \{(n-d, d)\}, \{(n-d+1, d+1)\}, \\
 & \quad \{(n-d, d), (n-d+1, d+1)\}, \\
 D_1(d) : & \quad \{(d+1, d), (n-d+1, n-d)\}, \\
 & \quad \{(d+1, d), (n-d+1, n-d), (n-d, d), (n-d+1, d+1)\}.
 \end{aligned}$$

Definition 2.6. A subset $D \subset \Phi(n)$ is called d -subregular ($1 \leq d < n_1$) if it has the form $D = D' \cup D_i(d)$, where either $i = 0$ or $i = 1$ and D' is a regular subset that doesn't consist the roots $(n - d + 1, d)$ and $(n - d + 1, d + 1)$. Note that subregular subsets consisting $D_1(d)$ are not basic in general.

Remark 2.7. Denote $\text{Supp } f = \{(i, j \in \Phi(n)) \mid f(e_{ij}) \neq 0\}$. One can see that $D \subset \text{Supp } f \cup (n - d + 1, n - d) \cup (n - d + 1, d)$, if f is a canonical form of a subregular orbit.

Example 2.8. Here we draw one of 1-subregular subsets for $n = 6$:

$$D = \{(2, 1), (5, 1), (6, 2), (4, 3), (6, 5)\} \subset \Phi(6).$$

	1	2	3	4	5	6
1						
2	⊗					
3						
4			⊗			
5	⊗					
6		⊗			⊗	

Define $R(D)$, $x_D(\varphi)$, $e_D(\varphi)$, $\mathcal{K}_D(\varphi)$, θ and θ_f , as for regular orbits. For a subregular orbit $\Omega_f \subset \mathcal{O}_d$ denote

$$\mathcal{K}_f = \bigcup_{(D, \varphi)} \mathcal{K}_D(\varphi),$$

where the union is over all d -subregular subsets D and over maps $\varphi: D \rightarrow \mathbb{F}_q^*$ such that

$$\xi_{d, n-d} \cdot \varphi(d + 1, d) = \xi_{d+1, n-d+1} \cdot \varphi(n - d + 1, n - d), \text{ where } f = (\xi_{ij}). \quad (4)$$

For a d -subregular subset D containing $D_1(d)$ we denote

$$\Phi_d = \{(i, j) \in \Phi(n) \mid i > n - j + 1, j \notin \{d, n - d\}, i \notin \{n - d + 1, n - d\}\}.$$

Finally, for a d -subregular subset D let

$$m_D = \begin{cases} |R(D) \cap \Phi_{\text{reg}}| - 1, & \text{if } D \supset D_0(d), \\ |R(D) \cap \Phi_d| + n - 2d - 1, & \text{if } D \supset D_1(d). \end{cases}$$

Theorem 2.9. Let $1 \leq d < n_1$, let $\Omega_f \subset \mathcal{O}_d \subset \mathfrak{g}_n^*(q)$ be a subregular orbit, and $\chi = \chi_f$ be the corresponding character. Then

1. $\text{Supp } \chi = \mathcal{K}_f$.
2. $\chi(g) = q^{m_D} \cdot \theta_f(e_D(\varphi))$ for an arbitrary $g \in \mathcal{K}_D(\varphi) \subset \mathcal{K}_f$.

Sketch of the proof. In Lemma 3.2 we present certain equations the element $x_D(\varphi)$ satisfies, and show that the ideal J of $k[G_n]$ generated by these equations is invariant under the adjoint action. In Lemma 3.3 we prove that J is a prime ideal. In Lemma 3.4 we find the stabilizer $\mathcal{C} = \text{Stab}_{G_n}(x_D(\varphi))$ and prove that

$$\dim \mathcal{K}_D(\varphi) = \text{codim } \mathcal{C} = \dim V(J)$$

(here $V(J) = \{g \in G_n \mid F(g) = 0 \text{ for all } F \in J\}$, as usual). This shows that $\mathcal{K}_D(\varphi) = V(J)$, i.e., our equations are exactly the defining equations for the conjugacy class of the element $x_D(\varphi)$ (Theorem 3.5).

On the other hand, in Lemmas 5.1 — 5.3 we prove that if $\chi(g) \neq 0$, then g satisfies these equations (and the additional equation (4)), and compute the value $\chi(g)$ in this case. This concludes the proof. \square

Remark 2.10. If $D \supset D_1(d)$ is d -subregular, $\varphi, \tilde{\varphi}$ are maps from D to \mathbb{F}_q^* , and

$$\begin{aligned} & \varphi(d+1, d) \cdot \varphi(n-d+1, d+1) + \varphi(n-d, d) \cdot \varphi(n-d+1, n-d) = \\ & = \tilde{\varphi}(d+1, d) \cdot \tilde{\varphi}(n-d+1, d+1) + \tilde{\varphi}(n-d, d) \cdot \tilde{\varphi}(n-d+1, n-d), \end{aligned}$$

then $\mathcal{K}_D(\varphi) = \mathcal{K}_D(\tilde{\varphi})$ (see section 3). But one can see (using the results of section 3) that in this case $\theta_f(e_D(\varphi)) = \theta_f(e_D(\tilde{\varphi}))$. Since m_D is independent of φ , the character value in the formulation of the Theorem is well-defined.

It's easy to prove a variant of this Theorem for the case of n_1 -subregular orbits by some modifications of definitions and formulations (see section 6).

3. Conjugacy classes $\mathcal{K}_D(\varphi)$

Here we present an explicit description of the conjugacy class $\mathcal{K}_D(\varphi)$ of an element $x_D(\varphi)$. Fix an arbitrary $1 \leq d < n_1$ and a subregular subset $D \supset D_1(d)$ (in this section and in the two next sections we consider the case $D(d) = D_1(d)$; the case $D(d) = D_0(d)$ is similar, see section 6).

Notation 3.1. It's convenient to split a d -subregular subset into the union

$$D = D^- \sqcup D_1(d) \sqcup D^+,$$

where $D^- = \{(i, j) \in D \mid j < d\}$, $D^+ = \{(i, j) \in D \mid d < j < n-d\}$.

Recall that $D' = D \setminus D_1(d) = D^- \sqcup D^+$ is a regular subset (see Definition 2.6). Let $D'' = D \setminus \{(n-d, d), (n-d+1, d+1)\}$ (it's a basic subset of $\Phi(n)$, and $D' \subset D'' \subset D$).

It's impossible to describe $\mathcal{K}_D(\varphi)$ in terms of minors of matrices $g = (y_{ij}) \in G_n$. Let $(i, j) \in D^+$ and $m = \max_{(i,j) \in D^+} j$. Consider the following polynomials:

$$\alpha_{ij} = \sum_{l=n-m+1}^{n-d} y_{n-d+1,l} y_{l,j}, \quad \beta_{ij} = \sum_{l=d+1}^m y_{i,l} y_{l,d}, \quad \gamma = \sum_{l=d+1}^{n-d} y_{n-d+1,l} y_{l,d}$$

(more conceptual description of these polynomials is given in section 6). For simplicity, we'll write Δ_{ij} instead of $\Delta^{R_{D''}(i,j)}(g)$, and y_β, y_α instead of $y_{d+1,d}, y_{n-d+1,n-d}$ resp. ($R_{D''}(i, j)$ are defined similar to $R_D(i, j)$ in (2)).

Let us now start with proving Theorem 2.9. Let $c_0 \in k$, $c_\alpha, c_\beta, c_{ij} \in k^*$, where $(i, j) \in D'$, are arbitrary scalars. Consider the ideal J of $k[G_n]$ generated by the elements

$$\begin{aligned} & y_\alpha - c_\alpha, y_\beta - c_\beta, \gamma - c_0, \\ & \alpha_{ij}, \beta_{ij}, \quad (i, j) \in D^+, \\ & \Delta_{ij} - c_{ij}, \quad (i, j) \in D', \\ & \Delta_{ij}, \quad (i, j) \in R(D'') \setminus D'. \end{aligned} \tag{5}$$

Lemma 3.2. J is G_n -invariant, i.e., if $g \in V(J)$, then $xgx^{-1} \in V(J)$ for all $x \in G_n$.

Proof. Since the set D'' is basic, all Δ_{ij} invariant [A2, Lemma 2.1].

Let $g = (y_{ij}) \in V(J)$. Denote $x_{rs}(\lambda) = 1 + \lambda e_{rs}$. Since every element $x \in G_n$ can be written as a product $x = x_{r_1 s_1}(\lambda_1) \dots x_{r_m s_m}(\lambda_m)$ for a certain m and $(r_i, s_i) \in \Phi(n)$, $\lambda_i \in k$, it's enough to prove that if $x = x_{rs}(\lambda)$, then $xgx^{-1} \in V(J)$. But if x has this form, then

$$(xgx^{-1})_{ij} = \begin{cases} y_{ij}, & \text{if } i \neq r \text{ and } j \neq s, \\ y_{ij}, & \text{if } i = r \text{ and } j \geq s, \text{ or } j = s \text{ and } i \leq r, \\ y_{rj} + \lambda y_{sj}, & \text{if } i = r, j < s, \\ y_{is} - \lambda y_{ir}, & \text{if } j = s, i > r. \end{cases}$$

Hence, y_α, y_β are invariant obviously, and the proof of invariance of other elements is by direct enumeration of possible values of r and s .

For example, consider the polynomial α_{ij} . If $s \leq j$ or $r > n - d + 1$, then all coordinate functions involved in this polynomial are invariant themselves; this is also true, if $r = n - d + 1$, $s \leq n - m + 1$. If $r = n - d + 1, s > n - m + 1$, then

$$\begin{aligned} \alpha_{ij}(xgx^{-1}) &= \sum_{l=n-m+1}^{s-1} (y_{n-d+1,l} + \lambda y_{s,l}) y_{l,j} + \sum_{l=s}^{n-d} y_{n-d+1,l} y_{l,j} = \\ &= \alpha_{ij}(g) + \lambda \cdot \sum_{l=n-m+1}^{s-1} y_{s,l} y_{l,j} = \alpha_{ij}(g), \end{aligned}$$

because $y_{s,l} = 0$ for all $l \geq n - m + 1 \geq n - n_0 + 1 > m$ (it's a particular case of (3)).

If $r < n - d + 1, s = j$, then α_{ij} is invariant for the same reasons. Finally, if $r < n - d + 1, s > j$, then

$$\begin{aligned} \alpha_{ij}(xgx^{-1}) &= \sum_{l \neq r, s} y_{n-d+1,l} y_{l,j} + (y_{r,j} + \lambda y_{s,j}) y_{n-d+1,r} + y_{s,j} (y_{n-d+1,s} - \lambda y_{n-d+1,r}) = \\ &= \alpha_{ij}(g) + \lambda \cdot (y_{s,j} y_{n-d+1,r} - y_{s,j} y_{n-d+1,r}) = \alpha_{ij}(g). \end{aligned}$$

Invariance of β_{ij} and γ can be proved similarly. \square

This means that $V(J)$ is a union of cojugacy classes. It's easy to see that $x_D(\varphi) \in V(J)$, if

$$\begin{aligned} c_{ij} &= \Delta_{ij}(x_D(\varphi)), \\ c_0 &= \gamma(x_D(\varphi)), \\ c_\alpha &= \varphi(n - d + 1, n - d), \\ c_\beta &= \varphi(d + 1, d). \end{aligned} \tag{6}$$

Thus, $\mathcal{K}_D(\varphi) \subset V(J)$ for this values of constants.

For any root $\xi = (i, j) \in \Phi(n)$, we define its *level* as the number $u(\xi) = i - j$. Then the formula

$$\xi = (i_1, j_1) < \eta = (i_2, j_2) \Leftrightarrow \text{either } u(\xi) < u(\eta) \text{ or } u(\xi) = u(\eta), j_1 < j_2, \tag{7}$$

defines a complete order on the set of all roots. For each root $\xi = (i, j) \in \Phi(n)$ let I_ξ be the ideal in $k[G_n]$ generated by all $y_\eta, \eta < \xi$, and $\xi_0 = (i_0, j_0)$ be the maximal root in D'' , which is less than ξ (if exists).

Lemma 3.3. The ideal J is a prime ideal of $\bar{k}[y_{ij}]$ (here \bar{k} is the algebraic closure of k).

Proof. Consider the following transformation of coordinates:

$$\begin{aligned}\tilde{y}_{n-d+1, n-d} &= y_\alpha - c_\alpha, & \tilde{y}_{d+1, d} &= \tilde{y}_\beta - c_\beta, & \tilde{y}_{n-d+1, d+1} &= \gamma - c_0, \\ \tilde{y}_{n-d+1, i} &= \alpha_{ij}, & \tilde{y}_{i, d+1} &= \beta_{ij}, & (i, j) &\in D^+, \\ \tilde{y}_{ij} &= \Delta_{ij} - c_{ij}, & (i, j) &\in D', \\ \tilde{y}_{ij} &= \Delta_{ij}, & (i, j) &\in R(D'') \setminus D'.\end{aligned}\tag{8}$$

(it's easy to see that $J = \langle \tilde{y}_\xi \rangle_{\xi \in B}$, where $B \subset \Phi(n)$ denotes the set of all roots (i, j) from the left-side hand of (8)). Note that for any $\xi \in B$ we have

$$\tilde{y}_\xi \equiv y_\xi^0 \cdot y_\xi + a_\xi \pmod{I_\xi},\tag{9}$$

where y_ξ^0 is an invertible element of $\bar{k}[y_{ij}]/J$, and $a_\xi \in k$ is a certain scalar. Indeed, this is evident for $\tilde{y}_{n-d+1, n-d}$, $\tilde{y}_{d+1, d}$. For other roots we have:

$$\begin{aligned}\tilde{y}_{n-d+1, d+1} &= \gamma - c_0 = y_{n-d+1, d+1} y_{d+1, d} + \dots, \\ \tilde{y}_{n-d+1, i} &= \alpha_{ij} = y_{n-d+1, i} y_{i, j} + \dots, & (i, j) &\in D^+, \\ \tilde{y}_{i, d+1} &= \beta_{ij} = y_{i, d+1} y_{d+1, d} + \dots, & (i, j) &\in D^+, \\ \tilde{y}_{ij} &= \Delta_{ij} - c_{ij} = y_{ij} \cdot \Delta_{i_0, j_0} + \dots, & (i, j) &\in D', \\ \tilde{y}_{ij} &= \Delta_{ij} = y_{ij} \cdot \Delta_{i_0, j_0} + \dots, & (i, j) &\in R(D'') \setminus D',\end{aligned}$$

where for any \tilde{y}_ξ , $\xi \in B$, dots denote elements, equal to zero modulo I_ξ , and scalars (we assume that $\Delta_{i_0, j_0} = 1$, if the root ξ_0 does not exist for a given $\xi \in \Phi(n)$). But one can easily obtain the following equatities modulo J :

$$\begin{aligned}y_{d+1, d} &\equiv c_\beta \neq 0, & y_{n-m+1, m} &\equiv c_{n-m+1, m} \neq 0, \\ y_{i, j} &\equiv c_{ij}/c_{i_0, j_0} \neq 0, & (i, j) &\in D^+, j < m, \\ \Delta_{i_0, j_0} &\equiv c_{i_0, j_0} \neq 0, & (i_0, j_0) &\in D''\end{aligned}$$

(recall that $m = \max_{(i, j) \in D^+} j$). This concludes the proof of (9).

Hence, in $\bar{k}[y_{ij}]/J$ all y_ξ are polynomial in \tilde{y}_ξ (for each $\xi \in B$). Consequently,

$$\bar{k}[y_{ij}]/J = \bar{k}[y_\xi]_{\xi \in \Phi(n)} / \langle \tilde{y}_\xi \rangle_{\xi \in B} \cong \bar{k}[\{y_\xi\}_{\xi \notin B} \cup \{\tilde{y}_\xi\}_{\xi \in B}] / \langle \tilde{y}_\xi \rangle_{\xi \in B} \cong \bar{k}[\tilde{y}_\xi]_{\xi \in B}.$$

In particular, $\bar{k}[y_{ij}]/J$ is a domain, so J is a prime ideal. \square

This Lemma shows that $V(J)$ is an irreducible subvariety in $G_n(\bar{k})$, because $\bar{k}[G_n] \cong \bar{k}[y_{ij}]$.

Lemma 3.4. Let $\mathcal{C} = \text{Stab}_{G_n}(x_D(\varphi)) = \{g \in G_n \mid gx_D(\varphi) = x_D(\varphi)g\}$ be the stabilizer (the centralizer) of $x_D(\varphi)$. Then $\dim \mathcal{C} = \text{codim } V(J)$, where J is generated by the elements (5) with the scalars given by (6).

Proof. Firstly, we'll present the defining equations of the centralizer \mathcal{C} . For any $\xi = (i, j) \in \Phi(n)$, let Φ_ξ be the union of all roots from the i -th column and the j -th row. We put

$$\begin{aligned}\Phi_\alpha &= \{(n-d+1, i) \mid (i, j) \in D^+\}, & \tilde{\Phi}_\alpha &= \{(n-d+1, j) \mid j < d\}, \\ \Phi_\beta &= \{(i, d+1) \mid (i, j) \in D^+\}, & \tilde{\Phi}_\beta &= \{(i, d) \mid i > n-d+1\}, \\ \Phi_\gamma &= \{(n-d, d)\}, & \Phi_\delta &= (\cup_{\xi \in D} \Phi_\xi) \cap (R(D) \setminus D), \\ A &= \Phi_\alpha \sqcup \tilde{\Phi}_\alpha \sqcup \Phi_\beta \sqcup \tilde{\Phi}_\beta \sqcup \Phi_\gamma \sqcup \Phi_\delta \subset \Phi(n)\end{aligned}$$

(these sets are really disjoint). It's easy to check that \mathcal{C} is given by the following equations:

$$\begin{aligned}
(\alpha) \quad & y_{n-d+1,i} a_{ij} = a_{n-d+1,n-d} y_{n-d,j}, \quad (i,j) \in D^+, \\
(\beta) \quad & y_{i,d+1} a_{d+1,d} = a_{ij} y_{jd}, \quad (i,j) \in D^+, \\
(\tilde{\alpha}) \quad & a_{n-d+1,n-d} y_{n-d,j} + a_{n-d+1,d+1} y_{d+1,j} = 0, \quad j < d, \\
(\tilde{\beta}) \quad & y_{i,d+1} a_{d+1,d} + y_{i,n-d} a_{n-d,d} = 0, \quad i > n-d+1, \\
(\gamma) \quad & y_{n-d+1,n-d} a_{n-d,d} + y_{n-d+1,d+1} a_{d+1,d} = a_{n-d+1,n-d} y_{n-d,d} + a_{n-d+1,d+1} y_{d+1,d}, \\
(\delta) \quad & y_{ij} = 0, \quad (i,j) \in \Phi_\delta,
\end{aligned} \tag{10}$$

(here we write $a_{ij} = \varphi(i,j) \in k^*$ for simplicity).

So, \mathcal{C} is defined by the equations (10), which are labeled by roots from A , and $V(J)$ is defined by the equations (5), which are labeled by roots from B . Consider the maps $\sigma_A: A \rightarrow \Phi(n)$ and $\sigma_B: B \rightarrow \Phi(n)$, given by the formulas

$$\sigma_A(\xi) = \begin{cases} (n-i+1, j), & \xi = (i, j), (n-i+1, i) \in D'', j \neq d+1, i < n-j+1, \\ (i, n-j+1), & \xi = (i, j), (n-j+1, j) \in D'', i \neq n-d, i > n-j+1, \\ (i, d), & \xi = (i, d+1) \in \Phi_\delta, i \neq n-d, \\ (n-d+1, j), & \xi = (n-d, j) \in \Phi_\delta, j \neq d+1, \\ (n-d+1, d), & \xi = (n-d, d+1) \in \Phi_\delta, \\ \xi, & \text{elsewhere,} \end{cases}$$

$$\sigma_B(\xi) = \begin{cases} (n-d, j), & \xi = (n-d+1, i), (i, j) \in D^+, \\ \xi, & \text{elsewhere.} \end{cases}$$

They define the map $\sigma: A \sqcup B \rightarrow \Phi(n)$ (here $A \sqcup B$ is the disjoint union of A and B) by the rule $\sigma|_A = \sigma_A$, $\sigma|_B = \sigma_B$; it's easy to see that this map is a bijection. Since $\dim \mathcal{K}_D(\varphi) = \text{codim } \mathcal{C} = |A|$, $\text{codim } V(J) = |B|$, this concludes the proof. \square

Theorem 3.5. The defining ideal J of the conjugacy class $\mathcal{K}_D(\varphi) \subset G_n(k)$ of an element $x_D(\varphi)$ is generated by the elements (5) with the scalars given by (6).

Proof. Since conjugacy classes of $G_n(\bar{k})$ are Zariski-closed [S, Proposition 2. 5], previous Lemmas follow that $\mathcal{K}_D(\varphi) = V(J)$ over \bar{k} . Hence, the sets of their k -points also coincide. \square

4. A semi-direct decomposition of G_n

The next goal is to describe the support of a subregular character $\chi = \chi_f$ explicitly (we'll see that the support coincides with \mathcal{K}_f) and to compute the value of this character on a conjugacy class contained in the support. We'll use induction on dimension of the group and the Mackey's method of semi-direct decomposition (see, f.e., [Le]). In this section we collect some basic facts which are needed for the sequel.

Let G be a finite group, A, B be its subgroups and $G = A \rtimes B$ be their semi-direct product (i.e., $G = AB$ and $A \triangleleft G$).

Definition 4.1. Let $G = A \rtimes B$ be a finite group and A be abelian. For a given irreducible character ψ of the group A , the subset $B^\psi = \{b \in B \mid \psi \circ \tau_b = \psi\}$ of the group B is said to be the *centralizer* of this character (here $\tau_b: A \rightarrow A: a \mapsto bab^{-1}$).

The following Theorem satisfies [Le]:

Theorem 4.2. Let $G = A \rtimes B$ be a finite group and A be abelian. Then every irreducible representation τ of the group G has the form $\tau = \text{Ind}_{A \rtimes B}^G \psi \otimes \tilde{\tau}$, where ψ is a certain irreducible character of the group A , and $\tilde{\tau}$ is a certain irreducible representation of the centralizer B^ψ . Hence, every irreducible character χ of the group G has the form $\chi = \text{Ind}_{A \rtimes B}^G \psi \tilde{\chi}$, where $\tilde{\chi}$, ψ are certain irreducible characters of the groups A , B_ψ resp. On the other hand, any character $\text{Ind}_{A \rtimes B}^G \psi \tilde{\chi}$ is an irreducible character of G . \square

We denote

$$\begin{aligned} P_n &= \{g = (y_{ij}) \in G_n \mid y_{ij} = 0 \text{ for } j \neq 1\}, \\ G_{n-1} &\cong \{g = (y_{ij}) \in G_n \mid y_{ij} = 0 \text{ for } j = 1\} \hookrightarrow G_n. \end{aligned}$$

Then $G_n = P_n \rtimes G_{n-1}$, moreover, the group $P_n \cong \mathbb{F}_q^{n-1}$ is abelian; hence, all conditions of Theorem 4.2 are satisfied.

We fix a non-trivial additive character $\theta: \mathbb{F}_q \rightarrow \mathbb{C}$. Any irreducible character of the group P_n has the form

$$p = (p_{ij}) \in P_n \mapsto \theta(s_2 p_{21}) \cdot \dots \cdot \theta(s_n p_{n1}),$$

where $s_i \in \mathbb{F}_q$, $2 \leq i \leq n$, are arbitrary scalars. We'll consider the case $\theta_n(p) = \theta(s_n p_{n1})$ and the case $\theta_{n-1}(p) = \theta(s_{n-1} p_{n-1,1})$. One can see that their centralizers in the subgroup G_{n-1} are

$$\begin{aligned} G_{n-1}^{\theta_n} &= \{g = (y_{ij}) \in G_{n-1} \subset G_n \mid y_{nj} = 0 \text{ for } 1 \leq j \leq n-1\}, \\ G_{n-1}^{\theta_{n-1}} &= \{g = (y_{ij}) \in G_{n-1} \subset G_n \mid y_{n-1,j} = 0 \text{ for } 1 \leq j \leq n-2\}. \end{aligned}$$

Note that $G_{n-1}^{\theta_n} \cong G_{n-2}$, and $G_{n-1}^{\theta_{n-1}} \cong \tilde{G}_{n-2} = G_{n-2} \times \mathbb{F}_q$ (we'll consider only these embeddings of these subgroups into G_n).

For a linear function $f \in \mathfrak{g}_n^*$, let $\pi(f) \in \mathfrak{g}_{n-2}^*$ and $\tilde{\pi}(f)$ denote its restrictions to \mathfrak{g}_{n-2} and $\tilde{\mathfrak{g}}_{n-2}$ respectively (here $\tilde{\mathfrak{g}}_{n-2} = \text{Lie } \tilde{G}_{n-2}$ and embeddings of subalgebras into \mathfrak{g}_n correspond to embeddings of subgroups into G_n). Due to the Mackey's method, if $1 \leq d < n_1$ and $f = (\xi_{ij}) \in \mathcal{O}_d$ is the canonical form of a d -subregular orbit, then

$$\begin{aligned} T_f &= \text{Ind}_{P_n \rtimes G_{n-2}}^{G_n} \theta_n \otimes T_{\pi(f)}, \text{ where } s_n = \xi_{1n}, \quad \text{if } d > 1, \\ T_f &= \text{Ind}_{P_n \rtimes \tilde{G}_{n-2}}^{G_n} \theta_{n-1} \otimes T_{\tilde{\pi}(f)}, \text{ where } s_{n-1} = \xi_{1,n-1}, \quad \text{if } d = 1, \end{aligned} \tag{11}$$

where T_f (resp. $T_{\pi(f)}$ and $T_{\tilde{\pi}(f)}$) denotes the representation of the group G_n (resp. G_{n-2} and \tilde{G}_{n-2}), corresponding to the orbit $\Omega_f \subset \mathfrak{g}_n^*$ (resp. $\Omega_{\pi(f)} \subset \mathfrak{g}_{n-2}^*$ and $\Omega_{\tilde{\pi}(f)} \subset \tilde{\mathfrak{g}}_{n-2}^*$). All terms in the last equality are well-defined, because the orbit method is valid for the group \tilde{G}_{n-2} . So, the problem can be reduced to the study of representations of dimension less than the first one has, and we can use an inductive argument.

Finally, certain coset decompositions of the group G_n are needed for construction of induced representations. It's easy to check that

$$\begin{aligned} H_n &= \{h = (t_{ij}) \in G_{n-1} \mid t_{ij} = 0 \text{ for } i \neq n\}, \\ \tilde{H}_n &= \{h = (t_{ij}) \in G_{n-1} \mid t_{ij} = 0 \text{ for } i \neq n-1\}. \end{aligned}$$

are complete systems of representatives of $G_n/(P_n \rtimes G_{n-2})$ and $G_n/(P_n \rtimes \tilde{G}_{n-2})$ resp.

Remark 4.3. Since $\tilde{G}_{n-2} \cong G_{n-2} \times \mathbb{F}_q$, we have a complete description of the representation $T_{\tilde{\pi}(f)}$. Precisely, we note that G_{n-2} is isomorphic to the subgroup $G_{n-2} \times 0$ of \tilde{G}_{n-2} (this induces

the embedding of Lie algebras $\mathfrak{g}_{n-2} \hookrightarrow \tilde{\mathfrak{g}}_{n-2}$; let ψ be the projection $\tilde{G}_{n-2} \rightarrow G_{n-2} = G_{n-2} \times 0$ and $g = (y_{ij}) \in \tilde{G}_{n-2}$. Then the character of the representation $T_{\tilde{\pi}(f)}$ has the form $g \mapsto \chi(\psi(g)) \cdot \theta(\xi_{n-1,n} y_{n,n-1})$, where χ is the character of the principal series of G_{n-2} , corresponding to the orbit $(\tilde{\pi}(f))|_{\mathfrak{g}_{n-2}}$.

Remark 4.4. There is another ("symmetric") decomposition $G_n = P'_n \rtimes G'_{n-1}$, where

$$\begin{aligned} P'_n &= \{g = (y_{ij}) \in G_n \mid y_{ij} = 0 \text{ for } j \neq 1\}, \\ G'_{n-1} &= \{g = (y_{ij}) \in G_n \mid y_{ij} = 0 \text{ for } j = 1\}. \end{aligned}$$

Reflecting other subgroups and subsets in the anti-diagonal, we get \tilde{G}'_{n-2} , H'_n and \tilde{H}'_n (G_{n-2} , embedded into G_n as above, is invariant under this reflection, i.e. $G'_{n-2} = G_{n-2}$). Then the irreducible character $\theta'_{n-1}: P'_n \rightarrow \mathbb{C}$ has the form $P'_n \ni p = (p_{ij}) \mapsto \theta(\xi_{2,n} p_{n,2} + \xi_{n-1,n} p_{n,n-1})$.

Now we are able to conclude the proof of Theorem 2.9.

5. The proof of the Main Theorem

Let $1 \leq d < n_1$, $f \in \mathcal{O}_d$ be a canonical form of a d -subregular orbit and $\chi = \chi_f$ be the corresponding character (as above). We'll prove in this section that its support coincides with \mathcal{K}_f and compute its value on an arbitrary conjugacy class $\mathcal{K}_D(\varphi) \subset \mathcal{K}_f$. We've proved in section 3 that the subvariety $\mathcal{K}_D(\varphi)$ of G_n is defined by the equations (5) with the scalars given by (6); in the other words, $V(J)$ is the defining ideal of $\mathcal{K}_D(\varphi)$ in $k[G_n]$.

The following proof is by induction on n ; the base can be checked directly (f.e., using (1)). We'll assume that if $g = (y_{ij}) \in G_n$, then $y_{n-d+1,n-d} \neq 0$ (this means that $D \supset D_1(d)$). Firstly, we have to consider the case $d = 1$. For convenience, we denote

$$\begin{aligned} \Phi_{i_0} &= \{(i, j) \in \Phi(n) \mid i = i_0\}, & \tilde{\Phi}_{i_0} &= \cup_{i \geq i_0} \Phi_i, \\ \Phi^{j_0} &= \{(i, j) \in \Phi(n) \mid j = j_0\}, & \tilde{\Phi}^{j_0} &= \cup_{j \leq j_0} \Phi^j. \end{aligned}$$

Lemma 5.1. Let $f \in \mathcal{O}_1$ be a canonical form of a 1-subregular orbit, and $g \in G_n$. If $\chi(g) \neq 0$, then $g \in \mathcal{K}_D(\varphi)$ for a certain $x_D(\varphi)$ satisfying (4).

Proof. Recall that

$$T_f = \text{Ind}_{P_n \rtimes \tilde{G}_{n-2}}^{G_n} \theta_{n-1} \otimes T_{\tilde{\pi}(f)}$$

(see (11)). Since $G_n = P_n \rtimes G_{n-1}$, an arbitrary element $g \in G_n$ can be uniquely represented as $g = pg'$, $p \in P_n$, $g' \in G_{n-1}$ (in fact, p and g' are given by replacing the corresponding entries of g by zeroes). Hence, for a given $g \in G_n$, the element $p(g) \in P_n$ is well-defined. By $\tilde{\pi}(g)$ we denote the element of \tilde{G}_{n-2} , which is given by replacing the corresponding entries of g by zeroes. We have

$$\chi(g) = \chi_f(g) = \text{Ind}_{P_n \rtimes \tilde{G}_{n-2}}^{G_n} \theta_{n-1}(p(g)) \cdot \chi_{\tilde{\pi}(f)}(\tilde{\pi}(g)) = \sum_{h \in \tilde{H}_n} \theta_{n-1}(p(h^{-1}gh)) \cdot \chi_{\tilde{\pi}(f)}(\tilde{\pi}(h^{-1}gh)) \quad (12)$$

(here the summation is over all $h \in \tilde{H}_n$ such that $h^{-1}gh \in P_n \rtimes \tilde{G}_{n-2}$).

Note that for any $h = (t_{ij}) \in \tilde{H}_n$,

$$\begin{aligned} (h^{-1}gh)_{n,j} &= y_{n,j} + y_{n,n-1}t_{n-1,j}, & \text{if } 2 \leq j \leq n-2, \\ (h^{-1}gh)_{n-1,j} &= y_{n-1,j} - \sum_{i=j+1}^{n-2} t_{n-1,i}y_{i,1}, & \text{if } 1 \leq j \leq n-3, \\ (h^{-1}gh)_{ij} &= y_{ij} & \text{for all other } 1 \leq j < i \leq n. \end{aligned} \quad (13)$$

If $\chi_{\tilde{\pi}(f)}(\tilde{\pi}(g)) \neq 0$, then it follows from the equations of regular orbits (Theorem 1.4), formulas for characters of the principal series (Theorem 1.12) and remark 4.3 that

$$\chi_{\tilde{\pi}(f)}(\tilde{\pi}(g)) = q^s \cdot \chi_{\pi^2(f)}(\pi^2(g)) \cdot \theta(\xi_{2,n} \cdot (-1)^{|X_1|-1} \cdot \frac{\Delta^{X_1}(g)}{\Delta^{X_2}(g)}) \cdot \theta(\xi_{n-1,n} y_{n,n-1}). \quad (14)$$

Here $X_1 = D \setminus (\Phi^1 \cup \Phi^{n-1})$, $X_2 = X_1 \setminus \Phi_n$, $s = |(R(D^+) \cap \Phi_n) \setminus (\tilde{\Phi}^2 \cup \Phi^{n-1})|$ and π^2 denotes the maps $G_n \rightarrow G_{n-4}$ and $\mathfrak{g}_n^* \rightarrow \mathfrak{g}_{n-4}^*$ (we assume that G_{n-4} is embedded into G_n like $G_{n-4} \subset G_{n-2} \subset G_n$). In particular, $\chi_{\pi^2(f)}$ is a character of the principal series of the group G_{n-4} .

On the other hand, (13) shows that $\chi_{\pi^2(f)}(\pi^2(g)) = \chi_{\pi^2(f)}(\pi^2(h^{-1}gh))$ for all $h \in \tilde{H}_n$. Substituting this to (12) and using (14), we obtain

$$\begin{aligned} \chi(g) &= q^s \cdot \chi_{\pi^2(f)}(\pi^2(g)) \times \\ &\times \sum_{h \in \tilde{H}_n} \theta_{n-1}(p(h^{-1}gh)) \cdot \theta(\xi_{2,n} \cdot (-1)^{|X_1|-1} \cdot \frac{\Delta^{X_1}(h^{-1}gh)}{\Delta^{X_2}(h^{-1}gh)} + \xi_{n-1,n} y_{n,n-1}). \end{aligned} \quad (15)$$

Furthermore,

$$\begin{aligned} \theta_{n-1}(p(h^{-1}gh)) &= \theta(\xi_{1,n-1} \cdot (y_{n-1,1} - \sum_{i=2}^{n-2} t_{n-1,i} y_{i,1})), \\ \frac{\Delta^{X_1}(h^{-1}gh)}{\Delta^{X_2}(h^{-1}gh)} &= \frac{\Delta^{X_1}(g) + y_{n,n-1} \cdot \Delta^{X_1}(g_t)}{\Delta^{X_2}(g)}, \end{aligned} \quad (16)$$

where g_t denotes the matrix given by replacing y_{nj} by t_{nj} for $2 \leq j \leq n-2$ in the matrix g (the last equality can be checked directly).

Suppose now that $\chi(g) \neq 0$. Thus $\chi_{\pi^2(f)}(\pi^2(g)) \neq 0$ (see (15)), some conditions from the equations (5), defining $\mathcal{K}_D(\varphi)$, are satisfied automatically. Precisely, we have to check only the following equalities:

$$\begin{aligned} \xi_{1,n-1} y_{2,1} &= \xi_{2,n} y_{n,n-1}, \quad (\text{condition (4)}) \\ \alpha_{ij} &= \beta_{ij} = 0, \quad (i, j) \in D^+, \\ \Delta_{ij} &= 0, \quad (i, j) \in (R(D'') \setminus D') \cap (\tilde{\Phi}^2 \cup \tilde{\Phi}_{n-1}). \end{aligned}$$

But $(n, n-1), (2, 1) \in D''$, so $R(D'') \cap (\Phi^1 \cup \Phi_n) = \emptyset$, hence, the last set of equalities in fact has the form $\Delta_{n-1,j} = 0, \Delta_{i,2} = 0$.

Consider the equalities $\Delta_{n-1,j} = 0$ firstly. If $j > m = \max_{(i,j) \in D} j$, then they have the form $\Delta_{n-1,j} = y_{n-1,j} = 0$, and other equalities (with $j \leq m$) are exactly Kronecker-Capelli conditions of compatibility of the system of linear in $t_{n-1,j}$ equations

$$y_{n-1,j} - \sum_{i=j+1}^{n-2} t_{n-1,i} y_{i,1} = 0, \quad 2 \leq j \leq m. \quad (17)$$

In both cases equalities under consideration are satisfied, because they express the condition $h^{-1}gh \in P_n \times \tilde{G}_{n-2}$ (see (13)).

On the other hand, it follows from this condition and (13) that $t_{n-1,j} = -\frac{y_{n,j}}{y_{n,n-1}}$ for all $m < j \leq n-2$. Substituting this to (17), we obtain that $\alpha_{ij} = 0$ for any $(i, j) \in D^+$ (all other

equalities of this form are satisfied because of compatibility of the system (17)). The equalities $\beta_{ij} = 0$, $(i, j) \in D^+$, and the other equalities $\Delta_{i,2} = 0$ can be obtained by using symmetric semi-direct decomposition of G_n (see remark 4.4).

Finally, we'll prove that (4) is satisfied. Note that $\sum_{c \in \mathbb{F}_q} \theta(ct) = 0$ for all $t \in \mathbb{F}_q^*$. Hence, the coefficients of $t_{n-1,j}$ in the character formula, that vary independently, have to be zero. For example, the coefficient of $t_{n-1,2}$ equals zero. Since $\Delta^{X_1}(g^t) = (-1)^{|X_1|-2} \cdot t_{n-1,2} \cdot \Delta^{X_2}(g) + \dots$ (members without $t_{n-1,2}$), the coefficient of $t_{n-1,2}$ in the character formula is equal to

$$-\xi_{1,n-1}y_{21} + \xi_{2,n} \cdot (-1)^{|X_1|} \cdot y_{n,n-1} \cdot \frac{(-1)^{|X_1|-2} \cdot \Delta^{X_2}(g)}{\Delta^{X_2}(g)} = -\xi_{1,n-1}y_{21} + \xi_{2,n}y_{n,n-1},$$

so, $-\xi_{1,n-1}y_{21} + \xi_{2,n}y_{n,n-1} = 0$. \square

Lemma 5.2. Let $f \in \mathcal{O}_1$ be a canonical form of a 1-subregular orbit, $g \in \mathcal{K}_D(\varphi) \subset \mathcal{K}_f$. Then $\chi(g) = q^{m_D} \cdot \theta_f(e_D(\varphi))$.

Proof. We'll compute the value of the character under the assumption that this value isn't equal to zero. Substituting (16) to (15), we have:

$$\begin{aligned} \chi(g) &= q^s \cdot \chi_{\pi^2(f)}(\pi^2(g)) \cdot \sum_{h \in \tilde{H}_n} \theta(\xi_{1,n-1} \cdot (y_{n-1,1} - \sum_{i=2}^{n-2} t_{n-1,i}y_{i,1}) + \\ &+ \xi_{2,n} \cdot (-1)^{|X_1|} \cdot \frac{\Delta^{X_1}(g) + y_{n,n-1} \cdot \Delta^{X_1}(g_t)}{\Delta^{X_2}(g)} + \xi_{n-1,n}y_{n,n-1}), \end{aligned}$$

if $\chi_{\tilde{\pi}(f)}(\tilde{\pi}(g)) \neq 0$. If $\chi(g) \neq 0$, then the following four groups of conditions are satisfied.

1. The first condition $\chi_{\pi^2(f)}(\pi^2(g)) \neq 0$ is satisfied automatically, because $g \in \mathcal{K}_D(\varphi)$. Indeed, it's evident, that $\pi^2(D) \in \Phi(n-4)$ is a regular subset and $\pi^2(g) \in \mathcal{K}_{\pi^2(D)}(\varphi|_{\pi^2(D)})$. But $\chi_{\pi^2(f)}(\pi^2(g))$ is a character of the principal series of G_{n-4} , so, Theorem 1.12 shows that

$$\chi_{\pi^2(f)}(\pi^2(g)) = q^{m_{\pi^2(D)}} \cdot \theta_{\pi^2(f)}(\pi^2(e_D(\phi)))$$

(the map $\pi^2: \mathfrak{g}_n \rightarrow \mathfrak{g}_{n-4}$ is defined by the obvious way).

2. The second group of conditions provides the inequality $\chi_{\tilde{\pi}(f)}(\tilde{\pi}(g)) \neq 0$. Since $\chi_{\pi^2(f)}(\pi^2(g)) \neq 0$, it's enough to prove that $\Delta_{ij}(h^{-1}gh) = 0$ for $(i, j) \in (R(D'') \setminus D) \cap (\Phi^2 \cup \Phi_n)$. More precisely, these conditions have the form

$$\begin{aligned} \Delta_{i,2}(g) &= 0, \quad (i, 2) \in R(D)'' \setminus D \\ \Delta_{nj}(h^{-1}gh) &= \Delta_{nj}(g) + y_{n,n-1} \cdot \Delta_{nj}(g_t) = 0, \quad (n, j) \in R(D)'' \setminus D. \end{aligned}$$

The first set of equalities is contained in the Kronecker-Capelli conditions of compatibility of the system (17). Since $t_{n-1,j}$, $3 \leq j \leq m$, contain only in the second set of equalities, it follows that $t_{n-1,j}$, $3 \leq j \leq m$, $(n-1, j) \notin R(D'')$, may be arbitrary, and other $t_{n-1,j}$ are uniquely determined by the formulas

$$t_{n-1,j} = (-1)^{|X_1|-3} \cdot \frac{-\Delta_{n,j}(g) - \sum_{\substack{r>j \\ (n-1,r) \notin R(D'')}} \pm t_{n-1,r} \cdot \Delta^{Y_r(j)}(g)}{y_{n,n-1} \cdot \Delta^{Y(j)}}, \quad (18)$$

where $Y(j) = D^+ \setminus \tilde{\Phi}^j$, $Y_r(j) = (D^+ \setminus (\Phi^r \cup \tilde{\Phi}^{j-1})) \cup \{(n-r+1, j)\}$, and signs of $t_{n-1,r}$ alternates.

3. The third condition says that we must consider $h \in \tilde{H}_n$, such that $h^{-1}gh \in P_n \rtimes \tilde{G}_{n-2}$. This condition is always satisfied, if $g \in \mathcal{K}_D(\varphi)$ (see the proof of Lemma 5.1).

4. Finally, the fourth condition says that the coefficients of $t_{n-1,j}$ in the character formula, that vary independently, are equal to zero. These are $t_{n-1,2}$ and $t_{n-1,j}$, $3 \leq j \leq m$, $(n-1, j) \notin R(D'')$. The end of the proof of Lemma 5.1 shows that the condition (4) is satisfied if and only if the coefficient of $t_{n-1,2}$ equals zero; hence, we must prove only that the coefficients of all $t_{n-1,j}$, $3 \leq j \leq m$, $(n-1, j) \notin R(D'')$, are equal to zero. Simplifying (18) under all our assumptions, one can check that this is equivalent to the set of equalities $\beta_{ij} = 0$, $(i, j) \in D^+$, which are satisfied, because $g \in \mathcal{K}_D(\varphi)$.

So, the character formula contains only the following expression:

$$\begin{aligned} & \xi_{n-1,n} y_{n,n-1} + \xi_{1,n-1} \cdot (y_{n-1,1} - \sum_{i=m+1}^{n-2} y_{i,1} \cdot (-\frac{y_{n-1,i}}{y_{n,n-1}})) + \xi_{2,n} \cdot \dots = \\ & = \xi_{n-1,n} y_{n,n-1} + \xi_{1,n-1} \cdot \frac{\sum_{i=n-m+1}^{n-1} y_{n,i} y_{i,1}}{y_{n,n-1}} + \xi_{1,n-1} \cdot \frac{\sum_{i=m+1}^{n-m} y_{n,i} y_{i,1}}{y_{n,n-1}} + \xi_{2,n} \cdot \dots, \end{aligned} \quad (19)$$

where the numerator of every summand of the group marked by dots contains the only one of the elements $y_{i,1}$, $2 \leq i \leq n-m$, and these elements aren't contained in other summands. On the other hand, the symmetric decomposition of the group G_n (see remark 4.4) give the following expression in the character formula

$$\begin{aligned} & \xi_{n-1,n} y_{n,n-1} + \xi_{2,n} \cdot \frac{\sum_{j=2}^{n-m} y_{n,j} y_{j,1}}{y_{21}} + \xi_{1,n-1} \cdot \dots = \\ & = \xi_{n-1,n} y_{n,n-1} + \xi_{2,n} \cdot \frac{\sum_{j=2}^m y_{n,j} y_{j,1}}{y_{21}} + \xi_{2,n} \cdot \frac{\sum_{j=m+1}^{n-m} y_{n,j} y_{j,1}}{y_{21}} + \xi_{1,n-1} \cdot \dots, \end{aligned} \quad (20)$$

which has to coincide with (19) (here the numerator of every summand of the group marked by dots contains the only one of the elements $y_{n,j}$, $n-m+1 \leq j \leq n-1$, and these elements aren't contained in other summands). Since $\xi_{1,n-1} y_{21} = \xi_{2,n} y_{n,n-1}$, we have that, in fact, (19) and (20) have the common form

$$\xi_{1,n-1} \cdot \frac{\sum_{i=n-m+1}^{n-1} y_{n,i} y_{i,1}}{y_{n,n-1}} + \xi_{2,n} \cdot \frac{\sum_{j=2}^{n-m} y_{n,j} y_{j,1}}{y_{21}} + \xi_{n-1,n} y_{n,n-1} = \xi_{1,n-1} \cdot \frac{\gamma(g)}{y_{n,n-1}} + \xi_{n-1,n} y_{n,n-1}.$$

We also have

$$\sum_{h \in \tilde{H}_n} \theta(\xi_{1,n-1} \cdot \frac{\gamma(g)}{y_{n,n-1}} + \xi_{n-1,n} y_{n,n-1}) = q^{s_1} \cdot \theta(\xi_{1,n-1} \cdot \frac{\gamma(g)}{y_{n,n-1}} + \xi_{n-1,n} y_{n,n-1}), \quad (21)$$

where $s_1 = |R(D) \cap \Phi_{n-1} \cap \tilde{\Phi}^m|$. Indeed, $t_{n-1,j}$, $2 \leq j \leq m$, $(n-1, j) \in R(D)$, vary independently, and if we fix them, then all other $t_{n-1,j}$ are uniquely determined.

Note that

$$\begin{aligned} & \xi_{1,n-1} \cdot \frac{\gamma(g)}{y_{n,n-1}} = \xi_{1,n-1} \cdot \frac{\gamma(x_D(\varphi))}{y_{n,n-1}} = \xi_{1,n-1} \cdot \frac{a_{n-d+1,n-d} a_{n-d,1} + a_{n-d+1,2} a_{2,1}}{a_{n,n-1}} = \\ & = \xi_{1,n-1} a_{n-1,1} + \frac{\xi_{1,n-1}}{a_{n,n-1}} \cdot a_{n-d+1,2} a_{2,1} = \xi_{1,n-1} a_{n-1,1} + \frac{\xi_{2,n-d+1}}{a_{2,1}} \cdot a_{n-d+1,2} a_{2,1} = \\ & = \xi_{1,n-1} a_{n-1,1} + \xi_{2,n-d+1} a_{n-d+1,2} \end{aligned} \quad (22)$$

(here $a_{ij} = \varphi(i, j)$, as above).

Finally, substitute (21) and (22) to (15):

$$\begin{aligned}\chi_{n,f}(g) &= q^s \cdot \chi_{\pi^2(f)}(\pi^2(g)) \cdot \sum_{h \in \tilde{H}_n} \theta(\xi_{1,n-1} \cdot \frac{\gamma(g)}{y_{n,n-1}} + \xi_{n-1,n} y_{n,n-1}) = \\ &= q^{s+s_1} \cdot \chi_{\pi^2(f)}(\pi^2(g)) \cdot \theta(\xi_{n-1,1} a_{n,n-1} + \xi_{1,n-1} a_{n-1,1} + \xi_{2,n-d+1} a_{n-d+1,2}) = q^{m_D} \cdot \theta_f(e_D(\varphi)),\end{aligned}$$

because $m_{\pi^2(D)} + s + s_1 = m_D$. The proof is complete. \square

Lemma 5.3. Let $1 < d < n_1$, $f \in \mathcal{O}_d$ be a canonical form of a d -subregular, $\chi = \chi_f$ be the corresponding character, and $g \in G_n$. Then $\chi(g) \neq 0$ if and only if $g \in \mathcal{K}_D(\varphi) \subset \mathcal{K}_f$; in this case $\chi(g) = q^{m_D} \cdot \theta_f(e_D(\varphi))$.

Proof. Suppose that the conditions of the Lemma are satisfied. According to (11),

$$T_f = \text{Ind}_{P_n \rtimes G_{n-2}}^{G_n} \theta_n \otimes T_{\pi(f)},$$

hence,

$$\chi(g) = \sum_{h \in H_n} \theta_n(p(h^{-1}gh)) \cdot \chi_{\pi(f)}(\pi(h^{-1}gh))$$

(here the summation is over $h \in H_n$ such that $h^{-1}gh \in P_n \rtimes G_{n-2}$). Since $\pi(f)$ is a canonical form of a subregular orbit of G_{n-2} , we may assume that Theorem 2.9 is valid for $\chi_{\pi(f)}$ (inductive assumption).

Using the same arguments, as in two previous Lemmas (Kronecker-Capelli criterion of compatibility of systems of linear equations, conditions of the form $h^{-1}gh \in P_n \rtimes G_{n-2}$, and vanishing the coefficients of independent variables in the character formula), one can obtain the required result. \square

6. Remarks and generalizations

Above we consider d -subregular subsets D with $D(d) = D_1(d)$ and assume that $d < n_1$. Here we'll consider all other cases.

First, if $D \supset D_0(d)$, then the subset D is basic and the conjugacy class $\mathcal{K}_D(\varphi)$ is given by the equations (2) (see [A2]).

Second, we should consider subregular orbits of the second and the third types (these are n_1 -subregular orbits for even and odd n respectively). Note that if $f = (\xi_{ij}) \in \mathcal{O}_{n_1}$ is a canonical form of a subregular orbit of the second type, then the set $D = \{(i, j) \in \Phi(n) \mid \xi_{ji} \neq 0\}$ is basic itself, so in this case the subregular character is a basic character (in the sense of Andre); a complete description of such characters is given in [A2]. This is also true for subregular orbits of the third type, if $\xi_{n_1, n_0+1} = 0$ (see definition 2.2).

If \mathcal{O}_f is a subregular orbit of the third type and $\xi_{n_1, n_0+1} \neq 0$, then all statements are the same as in the case of orbits of the first type (except that the ideal J doesn't contain the polynomial $\gamma - c_0$). All proofs are quite similar to proofs in case of orbits of the first type, so we don't reply them.

Finally, we should explain the genesis of polynomials α_{ij} , β_{ij} and γ from our description of subregular characters of the first type with $D \supset D_1(d)$: (these polynomials are *not* minors!). Consider the following *characteristic matrix*

$$M(t) = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ t \cdot y_{21} & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ t \cdot y_{n-1,1} & t \cdot y_{n-1,2} & \dots & 1 & 0 \\ t \cdot y_{n,1} & t \cdot y_{n,2} & \dots & t \cdot y_{n,n-1} & 1 \end{pmatrix}.$$

A minor of the matrix $M(t)$ is polynomial on t , and its coefficients are polynomial on y_{ij} (hence, we can consider them as function on G_n or as elements of $\mathbb{F}_q[\mathfrak{g}_n]$).

Notation 6.1. By $M_{j_1, \dots, j_k}^{i_1, \dots, i_k}(t^d, g)$ we denote the value of the coefficient of t^d in the minor of the matrix $M(t)$ with rows $1 \leq i_1, \dots, i_k \leq n$ and columns $1 \leq j_1, \dots, j_k \leq n$ on an element $g = (y_{ij}) \in G_n$. We also denote $M^X(d, g) = M_{\tau(j_1, \dots, j_k)}^{\sigma(i_1, \dots, i_k)}(t^d, g)$, where X is the set of pairs of the form $X = \{(i_1, j_1), \dots, (i_k, j_k)\}$, and σ, τ are permutations such that $\sigma(i_1) < \dots < \sigma(i_k)$ and $\tau(j_1) < \dots < \tau(j_k)$.

For example, for any $1 \leq d \leq n_0$ we have $M^X(d, g) = \Delta_d(g)$, where $X = \{(n, 1), (n-1, 2), \dots, (n_1+2, n_0)\}$; hence, all minors considered above are also coefficients of minors of characteristic matrix. On the other hand, if $(i, j) \in D^+$ (where $D \supset D_1(d)$ is a d -subregular subset and $1 \leq d < n_1$), $m = \max_{(i,j) \in D} j$, then

$$\begin{aligned} \alpha_{ij}(g) &= \pm M^{X_{ij}^\alpha}(2, g), & X_{ij}^\alpha &= \{n-d+1, j\} \cup \bigcup_{d < i \leq m} \{(i, i)\}, \\ \beta_{ij}(g) &= \pm M^{X_{ij}^\beta}(2, g), & X_{ij}^\beta &= \{i, d\} \cup \bigcup_{d < j \leq m} \{(j, j)\}, \\ \gamma(g) &= \pm M^{X^\gamma}(2, g), & X^\gamma &= \{n-d+1, d\} \cup \bigcup_{d < i < n-d+1} \{(i, i)\} \end{aligned}$$

(the choice of signs depends on the cardinality of D^+).

We see that subregular characters can be described in terms of coefficients of minors of the characteristic matrix. In fact, subregular orbits, all orbits for $n \leq 7$ [IP] and all irreducible characters for $n \leq 5$ [I] can be described in these terms too. So, we may conjecture that all orbits and characters of the unitriangular group of an arbitrary dimension can be describes in terms of coefficients of minors of the characteristic matrix.

The author expresses his sincere gratitude to professor A.N. Panov for suggesting the ideas that gave rise to this work.

References

- [A1] Andre C.A.M. Basic characters of the unitriangular group. J. Algebra, v. **175**, 1995, p. 287-319.
- [A2] Andre C.A.M. The basic character table of the unitriangular group. J. Algebra, v. **241**, 2001, p. 437-471.
- [I] Ignatev M.V. Characters of the unitriangular group over a finite field. Proceedings of the XIII International Conference for Undergraguate and Graduate Students and Young Scientists "Lomonosov". Vol. IV. Moscow, Moscow University Press, 2006, p. 65-66.
- [IP] Ignatev M.V., Panov A.N. Coadjoint orbits of the group $UT(7, K)$. arXiv: math.RT/0603649.
- [Ka] Kazhdan D. Proof of Springer's hypothesis. Israel J. Math., v. **28**, 1977, p. 272-286.

- [K1] Kirillov A.A. Lectures on the orbit method. Novosibirsk, Nauchnaya kniga IDMI, 2002.
- [K2] Kirillov A.A. Unitary representations of nilpotent Lie groups. Uspekhi Mat. Nauk, v. **17**, 1962, p. 57-110.
- [K3] Kirillov A.A. The orbit method and finite groups. Moscow, MCCME, MC IUM, 1998.
- [K4] Kirillov A.A. Variations on the triangular theme. Amer. Math. Soc. Transl, v. **169**, 1995, p. 43-73.
- [Le] Lehrer G.I. Discrete series and the unipotent subgroup. Composito Math., v. **28**, fasc. 1, 1974, p. 9-19.
- [Lu] Lusztig G. Subregular nilpotent elements and bases in K -theory. Canad, J. Math., v. **51(6)**, 1999, p.1194-1225.
- [S] Steinberg R. Conjugasy classes in algebraic groups. Lecture Notes in Mathematics, v. **366**. New York and Berlin, Springer, 1974.

Department of Algebra and Geometry, Samara State University, ul. Akad. Pavlova, 1, Samara 443011, Russia

E-mail address: mihail.ignatev@gmail.com