

# WHEN EVERY MULTILINEAR MAPPING IS MULTIPLE SUMMING

Geraldo Botelho\* and Daniel Pellegrino†

## Abstract

In this paper we give a systematized treatment to some coincidence situations for multiple summing multilinear mappings which extend, generalize and simplify the methods and results obtained thus far. The application of our general results to the pertinent particular cases gives several new coincidences as well as easier proofs of some known results.

## Introduction

Multiple summing multilinear mappings between Banach spaces have been proved to be a very important and very useful nonlinear generalization of the ideal of absolutely summing linear operators (see [1, 3, 5, 8, 9, 10, 11, 12, 13, 14, 16]). This class was introduced, independently, by Matos [8] (under the terminology *fully summing multilinear mappings*) and Bombal, Pérez-García and Villanueva [3]. The original methods and deep results due to Pérez-García [11], which were a source of inspiration to us in this paper, have played a crucial role in the development of the theory.

A coincidence situation for multiple summing mappings is a situation in which every  $n$ -linear mapping from  $E_1 \times \cdots \times E_n$  to  $F$ , where  $E_1, \dots, E_n$  and  $F$  are fixed Banach spaces, is multiple  $(q; p_1, \dots, p_n)$ -summing for some numbers  $q, p_1, \dots, p_n$ . It happens that the condition enjoyed by multiple summing mappings by definition (see Definition 1.1) is a very restrictive one, so coincidence situations are supposed to be very rare. Nevertheless, some situations like that are known (along the paper we will come through some of them) and in this paper we will prove some more. Such multilinear coincidence theorems are usually proved with the help of linear coincidence situations. In this paper we give a unified treatment to this approach, in the sense that we identify general linear conditions from which multilinear coincidences will follow. In this fashion we obtain new multilinear coincidence situations as well as generalizations and simplifications of some known ones.

---

\*Supported by CNPq Project 202162/2006-0.

†Supported by CNPq Projects 471054/2006-2 and 308084/2006-3.

2000 Mathematics Subject Classification. Primary 46G25; Secondary 47B10.

# 1 Background and notation

Throughout this paper  $n$  is a positive integer,  $E_1, \dots, E_n, E$  and  $F$  will stand for Banach spaces over  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ , and  $E'$  is the dual of  $E$ . By  $\mathcal{L}(E_1, \dots, E_n; F)$  we denote the Banach space of all continuous  $n$ -linear mappings from  $E_1 \times \dots \times E_n$  to  $F$  with the usual sup norm. If  $E_1 = \dots = E_n = E$ , we write  $\mathcal{L}(^n E; F)$  and if  $F = \mathbb{K}$  we simply write  $\mathcal{L}(E_1, \dots, E_n)$  and  $\mathcal{L}(^n E)$ . For the general theory of multilinear mappings we refer to Dineen [7].

Let  $p \geq 1$ . By  $\ell_p(E)$  we mean the Banach space of all absolutely  $p$ -summable sequences  $(x_j)_{j=1}^\infty$ ,  $x_j \in E$  for all  $j$ , with the norm  $\|(x_j)_{j=1}^\infty\|_p = \left(\sum_{j=1}^\infty \|x_j\|^p\right)^{1/p}$ .  $\ell_p^w(E)$  denotes the Banach space of all sequences  $(x_j)_{j=1}^\infty$ ,  $x_j \in E$  for all  $j$ , such that  $(\varphi(x_j))_{j=1}^\infty \in \ell_p$  for every  $\varphi \in E'$  with the norm

$$\|(x_j)_{j=1}^\infty\|_{w,p} = \sup\{\|(\varphi(x_j))_{j=1}^\infty\|_p : \varphi \in E', \|\varphi\| \leq 1\}.$$

**Definition 1.1.** Let  $1 \leq p_j \leq q$ ,  $j = 1, \dots, n$ . An  $n$ -linear mapping  $A \in \mathcal{L}(E_1, \dots, E_n; F)$  is *multiple  $(q; p_1, \dots, p_n)$ -summing* if there is a constant  $C \geq 0$  such that

$$\left(\sum_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \|A(x_{j_1}^{(1)}, \dots, x_{j_n}^{(n)})\|^q\right)^{\frac{1}{q}} \leq C \prod_{k=1}^n \|(x_j^{(k)})_{j=1}^{m_k}\|_{w,p_j}$$

for every  $m_1, \dots, m_n \in \mathbb{N}$  and any  $x_{j_k}^{(k)} \in E_k$ ,  $j_k = 1, \dots, m_k$ ,  $k = 1, \dots, n$ . It is clear that we may assume  $m_1 = \dots = m_n$ . The infimum of the constants  $C$  working in the inequality is denoted by  $\pi_{q;p_1, \dots, p_n}(A)$ .

The subspace  $\Pi_{q;p_1, \dots, p_n}^n(E_1, \dots, E_n; F)$  of  $\mathcal{L}(E_1, \dots, E_n; F)$  of all multiple  $(q; p_1, \dots, p_n)$ -summing becomes a Banach space with the norm  $\pi_{q;p_1, \dots, p_n}(\cdot)$ . If  $p_1 = \dots = p_n = p$  we say that  $A$  is multiple  $(q; p)$ -summing and write  $A \in \Pi_{q;p}^n(E_1, \dots, E_n; F)$ . The symbols  $\Pi_{q;p_1, \dots, p_n}^n(^n E; F)$ ,  $\Pi_{q;p}^n(^n E; F)$ ,  $\Pi_{q;p_1, \dots, p_n}^n(E_1, \dots, E_n)$ ,  $\Pi_{q;p}^n(E_1, \dots, E_n)$ ,  $\Pi_{q;p_1, \dots, p_n}^n(^n E)$  and  $\Pi_{q;p}^n(^n E)$  are defined in the obvious way.

Making  $n = 1$  we recover the classical ideal of absolutely  $(q; p)$ -summing linear operators, for which the reader is referred to Diestel, Jarchow and Tonge [6]. For the space of absolutely  $(q; p)$ -summing linear operators from  $E$  to  $F$  we shall write  $\Pi_{q;p}(E; F)$  rather than  $\Pi_{q;p}^1(E; F)$ .

**Remark 1.2.** Throughout the paper we will obtain multilinear coincidences from linear ones. On the other hand, it must be clear that multilinear coincidences always imply linear ones. More precisely, it is not difficult to prove (see the proof of [10, Theorem 4.3]) that if  $\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;p_1, \dots, p_n}^n(E_1, \dots, E_n; F)$ , then  $\mathcal{L}(E_j; F) = \Pi_{q;p_j}(E_j; F)$ ,  $j = 1, \dots, n$ .

## 2 General results

Our first result establishes the conditions from which several (known and new) coincidence theorems will follow.

**Theorem 2.1.** Let  $p, r \in [1, q]$  and let  $F$  be a Banach space. By  $B(p, q, r, F)$  we mean the collection of all Banach spaces  $E$  such that

$$\mathcal{L}(E; F) = \Pi_{q;p}(E; F) \text{ and } \mathcal{L}(E; \ell_q(F)) = \Pi_{q;r}(E; \ell_q(F)).$$

Then, for every  $n \geq 2$ ,

$$\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;r, \dots, r, p}^n(E_1, \dots, E_n; F)$$

whenever  $E_1, \dots, E_n \in B(p, q, r, F)$ .

*Proof.* We proceed by induction on  $n$ . Case  $n = 2$ : let  $E_1, E_2 \in B(p, q, r, F)$ . By the open mapping theorem there are constants  $C_1$  and  $C_2$  such that

$$\pi_{q;p}(u) \leq C_1 \|u\| \text{ for every } u \in \mathcal{L}(E_2; F) \text{ and}$$

$$\pi_{q;r}(v) \leq C_2 \|v\| \text{ for every } v \in \mathcal{L}(E_1; \ell_q(F)).$$

Let  $A \in \mathcal{L}(E_1, E_2; F)$ . Given two sequences  $(x_j^{(1)})_{j=1}^\infty \in \ell_r^w(E_1)$  and  $(x_j^{(2)})_{j=1}^\infty \in \ell_p^w(E_2)$ , fix  $m \in \mathbb{N}$  and consider the continuous linear operator

$$A_1^{(m)} : E_1 \longrightarrow \ell_q(F) : A_1^{(m)}(x) = (A(x, x_1^{(2)}), \dots, A(x, x_m^{(2)}), 0, 0, \dots).$$

So,  $A_1^{(m)}$  is  $(q; r)$ -summing and  $\pi_{q;r}(A_1^{(m)}) \leq C_2 \|A_1^{(m)}\|$ . For each  $x \in B_{E_1}$ , consider the continuous linear operator

$$A_x : E_2 \longrightarrow F : A_x(y) = A(x, y).$$

So,  $A_x$  is  $(q; p)$ -summing and  $\pi_{q;p}(A_x) \leq C_1 \|A_x\| \leq C_1 \|A\| \|x\| \leq C_1 \|A\|$ . Therefore,

$$\begin{aligned} \left( \sum_{j=1}^m \sum_{k=1}^m \|A(x_j^{(1)}, x_k^{(2)})\|^q \right)^{\frac{1}{q}} &= \left( \sum_{j=1}^m \|A_1^{(m)}(x_j^{(1)})\|^q \right)^{\frac{1}{q}} \\ &\leq \pi_{q;r}(A_1^{(m)}) \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ &\leq C_2 \|A_1^{(m)}\| \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ &= C_2 \sup_{x \in B_{E_1}} \left( \sum_{k=1}^m \|A(x, x_k^{(2)})\|^q \right)^{\frac{1}{q}} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ &= C_2 \sup_{x \in B_{E_1}} \left( \sum_{k=1}^m \|A_x(x_k^{(2)})\|^q \right)^{\frac{1}{q}} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ &\leq C_2 \sup_{x \in B_{E_1}} \pi_{q;p}(A_x) \left\| (x_k^{(2)})_{k=1}^m \right\|_{w,p} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ &\leq C_1 C_2 \|A\| \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \left\| (x_k^{(2)})_{k=1}^m \right\|_{w,p}, \end{aligned}$$

which shows that  $A$  is multiple  $(q; r, p)$ -summing and  $\pi_{q;r,p}(A) \leq C_1 C_2 \|A\|$ .

Suppose now that the result holds for  $n$ , that is: for every  $E_1, \dots, E_n \in B(p, q, r, F)$ ,  $\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;r,\dots,r,p}^n(E_1, \dots, E_n; F)$ . To prove the case  $n + 1$ , let  $E_1, \dots, E_{n+1} \in B(p, q, r, F)$ . As  $E_2, \dots, E_{n+1}$  are  $n$  Banach spaces in  $B(p, q, r, F)$ ,  $\mathcal{L}(E_2, \dots, E_{n+1}; F) = \Pi_{q;r,\dots,r,p}^n(E_2, \dots, E_{n+1}; F)$  by the induction hypotheses, so we can select a constant  $C_1$  such that

$$\pi_{q;r,\dots,r,p}(B) \leq C_1 \|B\| \text{ for every } B \in \mathcal{L}(E_2, \dots, E_{n+1}; F).$$

Since  $E_1 \in B(p, q, r, F)$ , there is a constant  $C_2$  such that

$$\pi_{q;r}(v) \leq C_2 \|v\| \text{ for every } v \in \mathcal{L}(E_1; \ell_q(F)).$$

Let  $A \in \mathcal{L}(E_1, \dots, E_{n+1}; F)$ . Given sequences  $(x_j^{(1)})_{j=1}^\infty \in \ell_r^w(E_1), \dots, (x_j^{(n)})_{j=1}^\infty \in \ell_r^w(E_n)$  and  $(x_j^{(n+1)})_{j=1}^\infty \in \ell_p^w(E_{n+1})$ , fix  $m \in \mathbb{N}$  and consider the continuous linear operator

$$A_1^{(m)} : E_1 \longrightarrow \ell_q(F) : A_1^{(m)}(x) = \left( (A(x, x_{j_2}^{(2)}, \dots, x_{j_{n+1}}^{(n+1)})_{j_2, \dots, j_{n+1}=1}^m, 0, 0, \dots) \right).$$

So,  $A_1^{(m)}$  is  $(q; r)$ -summing and  $\pi_{q;r}(A_1^{(m)}) \leq C_2 \|A_1^{(m)}\|$ . For each  $x \in B_{E_1}$ , consider the continuous  $n$ -linear mapping

$$A_x^n : E_2 \times \dots \times E_{n+1} \longrightarrow F : A_x^n(x_2, \dots, x_{n+1}) = A(x, x_2, \dots, x_{n+1}).$$

So,  $\pi_{q;r,\dots,r,p}(A_x^n) \leq C_1 \|A_x^n\| \leq C_1 \|A\| \|x\| \leq C_1 \|A\|$ . Hence,

$$\begin{aligned} & \left( \sum_{j_1=1}^m \dots \sum_{j_{n+1}=1}^m \left\| A(x_{j_1}^{(1)}, \dots, x_{j_{n+1}}^{(n+1)}) \right\|^q \right)^{\frac{1}{q}} = \left( \sum_{j=1}^m \left\| A_1^{(m)}(x_j^{(1)}) \right\|^q \right)^{\frac{1}{q}} \\ & \leq \pi_{q;r}(A_1^{(m)}) \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \leq C_2 \|A_1^{(m)}\| \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ & = C_2 \sup_{x \in B_{E_1}} \left( \sum_{j_2=1}^m \dots \sum_{j_{n+1}=1}^m \left\| A(x, x_{j_2}^{(2)}, \dots, x_{j_{n+1}}^{(n+1)}) \right\|^q \right)^{\frac{1}{q}} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ & = C_2 \sup_{x \in B_{E_1}} \left( \sum_{j_2=1}^m \dots \sum_{j_{n+1}=1}^m \left\| A_x^n(x_{j_2}^{(2)}, \dots, x_{j_{n+1}}^{(n+1)}) \right\|^q \right)^{\frac{1}{q}} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ & \leq C_2 \sup_{x \in B_{E_1}} \pi_{q;r,\dots,r,p}(A_x^n) \left( \prod_{k=2}^n \left\| (x_j^{(k)})_{j=1}^m \right\|_{w,r} \right) \left\| (x_j^{(n+1)})_{j=1}^m \right\|_{w,p} \left\| (x_j^{(1)})_{j=1}^m \right\|_{w,r} \\ & \leq C_1 C_2 \|A\| \left( \prod_{k=1}^n \left\| (x_j^{(k)})_{j=1}^m \right\|_{w,r} \right) \left\| (x_j^{(n+1)})_{j=1}^m \right\|_{w,p}, \end{aligned}$$

which shows that  $A$  is multiple  $(q; r, \dots, r, p)$ -summing and completes the proof.  $\square$

Rewriting the proof above for  $E_1 = \dots = E_n = E$ , we obtain

**Theorem 2.2.** *Let  $p, r \in [1, q]$  and let  $E$  and  $F$  be Banach spaces such that  $\mathcal{L}(E; F) = \Pi_{q;p}(E; F)$  and  $\mathcal{L}(E; \ell_q(F)) = \Pi_{q;r}(E; \ell_q(F))$  with*

$$\pi_{q;p}(u) \leq C_1 \|u\| \text{ for every } u \in \mathcal{L}(E; F) \text{ and}$$

$$\pi_{q;r}(v) \leq C_2 \|v\| \text{ for every } v \in \mathcal{L}(E; \ell_q(F)).$$

Then, for every  $n \geq 2$ ,

$$\mathcal{L}({}^n E; F) = \Pi_{q;r,\dots,r,p}^n({}^n E; F) \text{ and}$$

$$\pi_{q;r,\dots,r,p}(A) \leq C_1 C_2^{n-1} \|A\| \text{ for every } A \in \mathcal{L}({}^n E; F).$$

For scalar-valued mappings we get the following particular cases:

**Corollary 2.3.** *Given  $1 \leq r \leq q$ , by  $B(r, q)$  we mean the collection of all Banach spaces  $E$  such that  $\mathcal{L}(E; \ell_q) = \Pi_{q;r}(E; \ell_q)$ . Then, for every  $n \geq 2$ ,*

$$\mathcal{L}(E_1, \dots, E_n) = \Pi_{q;r,\dots,r,q}^n(E_1, \dots, E_n)$$

whenever  $E_1, \dots, E_n \in B(r, q)$ .

**Corollary 2.4.** *Let  $1 \leq r \leq q$  and let  $E$  be a Banach space such that  $\mathcal{L}(E; \ell_q) = \Pi_{q;r}(E; \ell_q)$  with  $\pi_{q;r}(v) \leq C \|v\|$  for every  $v \in \mathcal{L}(E; \ell_q)$ . Then, for every  $n \geq 2$ ,*

$$\mathcal{L}({}^n E) = \Pi_{q;r,\dots,r,q}^n({}^n E) \text{ and } \pi_{q;r,\dots,r,q}(A) \leq C^{n-1} \|A\| \text{ for every } A \in \mathcal{L}({}^n E).$$

### 3 Applications

We start by showing that some known coincidence theorems are easy combinations of our general results with linear ones. From now on,  $n$  will always be an integer not smaller than 2.

**Proposition 3.1.** [3, Theorem 3.2] *If  $F$  has cotype  $q$  and  $E_1, \dots, E_n$  are arbitrary Banach spaces, then*

$$\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;1}^n(E_1, \dots, E_n; F) \text{ and}$$

$$\pi_{q;1}(A) \leq C_q(F)^n \|A\| \text{ for every } A \in \mathcal{L}(E_1, \dots, E_n; F),$$

where  $C_q(F)$  is the cotype  $q$  constant of  $F$ .

*Proof.* Both  $F$  and  $\ell_q(F)$  have cotype  $q$  (see [6, Theorem 11.12]), so  $\mathcal{L}(E; F) = \Pi_{q;1}(E; F)$  and  $\mathcal{L}(E; \ell_q(F)) = \Pi_{q;1}(E; \ell_q(F))$  for every Banach space  $E$  by [6, Corollary 11.17]. The desired coincidence follows from Theorem 2.1 and the estimate for the norms from its proof.  $\square$

**Proposition 3.2.** [3, Theorem 5.1] *If  $E_1, \dots, E_n$  are  $\mathcal{L}_1$ -spaces and  $H$  is a Hilbert space, then*

$$\begin{aligned} \mathcal{L}(E_1, \dots, E_n; H) &= \Pi_{2;2}^n(E_1, \dots, E_n; H) \text{ and} \\ \pi_{2;2}(A) &\leq K_G^n \|A\| \text{ for every } A \in \mathcal{L}(E_1, \dots, E_n; H), \end{aligned}$$

where  $K_G$  stands for the Grothendieck constant.

*Proof.* Both  $H$  and  $\ell_2(H)$  are  $\mathcal{L}_2$ -spaces (see [4, Ex. 23.17(a)]), so  $\mathcal{L}(E; H) = \Pi_{2;2}(E; H)$  and  $\mathcal{L}(E; \ell_2(H)) = \Pi_{2;2}(E; \ell_2(H))$  for every  $\mathcal{L}_1$ -space  $E$  by [6, Theorems 3.1 and 2.8]. As before, the result follows from Theorem 2.1 and its proof.  $\square$

**Proposition 3.3.** [3, Theorem 3.1] *If  $F$  has cotype 2 and  $E_1, \dots, E_n$  are  $\mathcal{L}_\infty$ -spaces, then  $\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{2;2}^n(E_1, \dots, E_n; F)$ .*

*Proof.* Both  $F$  and  $\ell_2(F)$  have cotype 2 [6, Theorem 11.12], so  $\mathcal{L}(E; F) = \Pi_{2;2}(E; F)$  and  $\mathcal{L}(E; \ell_2(F)) = \Pi_{2;2}(E; \ell_2(F))$  for every  $\mathcal{L}_\infty$ -space  $E$  by [6, Theorem 11.14(a)]. Call on Theorem 2.1 once more.  $\square$

Using [6, Theorem 11.14(b)] instead of [6, Theorem 11.14(a)] in the proof above we obtain

**Proposition 3.4.** *If  $F$  has cotype  $q > 2$ ,  $E_1, \dots, E_n$  are  $\mathcal{L}_\infty$ -spaces and  $r < q$ , then  $\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;r}^n(E_1, \dots, E_n; F)$ .*

Now we derive some coincidence situations which, as far as we know, are new. The first one complements nice information given in [14, Corollary 3.20].

**Proposition 3.5.** *If  $E_1, \dots, E_n$  are arbitrary Banach spaces and  $q \geq 2$ , then*

$$\begin{aligned} \mathcal{L}(E_1, \dots, E_n) &= \Pi_{q;1, \dots, 1, q}^n(E_1, \dots, E_n) \text{ and} \\ \pi_{q;1, \dots, 1, q}(A) &\leq C_q (\ell_q)^{n-1} \|A\| \text{ for every } A \in \mathcal{L}(E_1, \dots, E_n). \end{aligned}$$

*Proof.* Since  $\ell_q$  has cotype  $q$ ,  $\mathcal{L}(E; \ell_q) = \Pi_{q;1}(E; \ell_q)$  for every Banach space  $E$ , so the desired coincidence follows from Corollary 2.3 and the estimate for the norms follows from the proof of Theorem 2.1.  $\square$

In order to prove that there is no general inclusion theorem for multiple summing multilinear mappings (that is,  $q \leq p \not\Rightarrow \Pi_{q;q}^n \subseteq \Pi_{p;p}^n$ ), in [13, Theorem 3.6] the authors show that  $\Pi_{q;q}^2(^2\ell_1) \neq \mathcal{L}(^2\ell_1)$  for every  $q > 2$ , whereas  $\Pi_{2;2}^2(^2\ell_1) = \mathcal{L}(^2\ell_1)$  (cf. Proposition 3.2). Next proposition shows that the non-coincidence  $\Pi_{q;q}^2(^2\ell_1) \neq \mathcal{L}(^2\ell_1)$ ,  $q > 2$ , is quite sharp.

**Proposition 3.6.** *If  $E_1, \dots, E_n$  are  $\mathcal{L}_1$ -spaces and  $2 \leq r < q$ , then  $\mathcal{L}(E_1, \dots, E_n) = \Pi_{q;r, \dots, r, q}^n(E_1, \dots, E_n)$ .*

*Proof.* For every  $\mathcal{L}_1$ -space  $E$  and  $2 \leq r < q$ ,  $\mathcal{L}(E; \ell_q) = \Pi_{q;r}(E; \ell_q)$  by a result due to Bennet [2, Proposition 5.2(iv)]. The result follows from Corollary 2.3.  $\square$

Using [2, Proposition 5.1(ii)] instead of [2, Proposition 5.2(iv)] we get:

**Proposition 3.7.** *If  $E_1, \dots, E_n$  are  $\mathcal{L}_\infty$ -spaces and  $q > r, q > 2$ , then  $\mathcal{L}(E_1, \dots, E_n) = \Pi_{q;r, \dots, r, q}^n(E_1, \dots, E_n)$ .*

## 4 Multiple summing mappings on $\mathcal{L}_1$ -spaces

We have already obtained some applications of our results to multiple summing mappings on  $\mathcal{L}_1$ -spaces. In this section we go a little further in this direction. Before using a new approach, we apply our general results a couple of times more.

**Proposition 4.1.** *Let  $E_1, \dots, E_n$  be  $\mathcal{L}_1$ -spaces.*

(a)  $\mathcal{L}(E_1, \dots, E_n; H) = \Pi_{q;1,\dots,1,q}^n(E_1, \dots, E_n; H)$  for every Hilbert space  $H$  and any  $q \geq 2$ .

(b)  $\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{q;1,\dots,1,r}^n(E_1, \dots, E_n; F)$  for every  $\mathcal{L}_q$ -space  $F$  and any  $2 \leq r < q$ .

*Proof.* (a) Let  $E$  be an  $\mathcal{L}_1$ -space and  $H$  be a Hilbert space. [6, Theorems 3.1 and 2.8] yield that  $\mathcal{L}(E; H) = \Pi_{q;q}(E; H)$ , and  $\mathcal{L}(E; \ell_q(H)) = \Pi_{q;1}(E; \ell_q(H))$  because  $\ell_q(H)$  has cotype  $q$ . Theorem 2.1 gives the result.

(b) Let  $E$  be an  $\mathcal{L}_1$ -space and  $F$  be an  $\mathcal{L}_q$ -space. Using [2, Proposition 5.2(iv)] once more we know that  $\mathcal{L}(E; F) = \Pi_{q;r}(E; F)$ .  $\mathcal{L}(E; F) = \Pi_{q;1}(E; \ell_q(F))$  as  $\ell_q(F)$  has cotype  $q$ , so Theorem 2.1 completes the proof.  $\square$

Next result allows us to go a little bit further.

**Theorem 4.2.** *Let  $r \geq s$ . If  $\mathcal{L}(\ell_1; F) = \Pi_{r;s}(\ell_1; F)$ , then*

$$\mathcal{L}(E_1, \dots, E_n; F) = \Pi_{r; \min\{s, 2\}}^n(E_1, \dots, E_n; F)$$

for every  $n \in \mathbb{N}$  and any  $\mathcal{L}_1$ -spaces  $E_1, \dots, E_n$ .

*Proof.* By a standard localization argument we may assume  $E_1 = \dots = E_n = \ell_1$ . Let  $(x_j^{(1)})_{j=1}^{m_1}, \dots, (x_j^{(n)})_{j=1}^{m_n}$  be  $n$  finite sequences in  $\ell_1$ .

Claim: For every  $1 \leq p \leq 2$ ,

$$\left\| (x_{j_1}^{(1)} \otimes \dots \otimes x_{j_n}^{(n)})_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \right\|_{w,p} \leq K_G^{2n-2} \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \cdots \left\| (x_j^{(n)})_{j=1}^{m_n} \right\|_{w,p}$$

on the completed  $n$ -fold projective tensor product  $\widehat{\otimes}_\pi^n \ell_1$ .

Proof of the claim: we proceed by induction on  $n$ . Given  $A \in \mathcal{L}(\ell_1)$ , by [13, Theorem 3.4] we know that  $A \in \Pi_{p;p}^2(\ell_1)$  and  $\pi_{p;p}(A) \leq K_G^2 \|A\|$ . Denoting by  $\sigma$  the canonical bilinear mapping from  $\ell_1 \times \ell_1$  to  $\ell_1 \widehat{\otimes}_\pi \ell_1$ ,  $\sigma(x, y) = x \otimes y$ , and taking the supremum over all  $\varphi \in B_{(\ell_1 \widehat{\otimes}_\pi \ell_1)'}$ , it follows that

$$\begin{aligned} \sup_{\varphi} \left( \sum_{j_1, j_2=1}^{m_1, m_2} \left| \varphi(x_{j_1}^{(1)} \otimes x_{j_2}^{(2)}) \right|^p \right)^{\frac{1}{p}} &= \sup_{\varphi} \left( \sum_{j_1, j_2=1}^{m_1, m_2} \left| \varphi \circ \sigma(x_{j_1}^{(1)}, x_{j_2}^{(2)}) \right|^p \right)^{\frac{1}{p}} \\ &\leq \sup_{\varphi} \pi_{p;p}(\varphi \circ \sigma) \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \left\| (x_j^{(2)})_{j=1}^{m_2} \right\|_{w,p} \\ &\leq K_G^2 \sup_{\varphi} \|\varphi \circ \sigma\| \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \left\| (x_j^{(2)})_{j=1}^{m_2} \right\|_{w,p} \\ &= K_G^2 \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \left\| (x_j^{(2)})_{j=1}^{m_2} \right\|_{w,p}. \end{aligned} \quad (1)$$

Now suppose that the desired inequality holds for  $k$  and let us prove that it holds for  $k + 1$ . We are assuming that

$$\left\| (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_k}^{(k)})_{j_1, \dots, j_k=1}^{m_1, \dots, m_k} \right\|_{w,p} \leq K_G^{2k-2} \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \cdots \left\| (x_j^{(k)})_{j=1}^{m_k} \right\|_{w,p}.$$

Using that  $\widehat{\otimes}_\pi^k \ell_1$  is isometrically isomorphic to  $\ell_1$ , applying first (1) then the induction hypotheses, we get

$$\begin{aligned} & \left\| (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_k}^{(k)} \otimes x_{j_{k+1}}^{(k+1)})_{j_1, \dots, j_{k+1}=1}^{m_1, \dots, m_{k+1}} \right\|_{w,p} \\ &= \left\| \left( (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_k}^{(k)}) \otimes x_{j_{k+1}}^{(k+1)} \right)_{j_1, \dots, j_{k+1}=1}^{m_1, \dots, m_{k+1}} \right\|_{w,p} \\ &\leq K_G^2 \left\| (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_k}^{(k)})_{j_1, \dots, j_k=1}^{m_1, \dots, m_k} \right\|_{w,p} \cdot \left\| (x_{j_{k+1}}^{(k+1)})_{j_{k+1}=1}^{m_{k+1}} \right\|_{w,p} \\ &\leq K_G^2 K_G^{2k-2} \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \cdots \left\| (x_j^{(k)})_{j=1}^{m_k} \right\|_{w,p} \left\| (x_j^{(k+1)})_{j=1}^{m_{k+1}} \right\|_{w,p} \\ &= K_G^{2(k+1)-2} \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w,p} \cdots \left\| (x_j^{(k)})_{j=1}^{m_k} \right\|_{w,p} \left\| (x_j^{(k+1)})_{j=1}^{m_{k+1}} \right\|_{w,p}, \end{aligned}$$

completing the proof of the claim.

Let  $A \in \mathcal{L}(^n \ell_1; F)$ . By  $A_L$  we mean the linearization of  $A$  on  $\widehat{\otimes}_\pi^n \ell_1$ , that is  $A_L \in \mathcal{L}(\widehat{\otimes}_\pi^n \ell_1; F)$  and  $A_L(x_1 \otimes \cdots \otimes x_n) = A(x_1, \dots, x_n)$  for every  $x_j \in \ell_1$ . Since  $\widehat{\otimes}_\pi^n \ell_1$  is isometrically isomorphic to  $\ell_1$ , by assumption we have that  $A_L$  is  $(r; s)$ -summing and  $\pi_{r;s}(A_L) \leq M \|A_L\| = M \|A\|$ , where  $M$  is a constant independent of  $A$ . Using the claim with  $p = \min\{s, 2\}$  we get

$$\begin{aligned} & \left( \sum_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \left\| A(x_{j_1}^{(1)}, \dots, x_{j_n}^{(n)}) \right\|^r \right)^{\frac{1}{r}} \\ &= \left( \sum_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \left\| A_L(x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_n}^{(n)}) \right\|^r \right)^{\frac{1}{r}} \\ &\leq \pi_{r;s}(A_L) \left\| (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_n}^{(n)})_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \right\|_{w,s} \\ &\leq \pi_{r;s}(A_L) \left\| (x_{j_1}^{(1)} \otimes \cdots \otimes x_{j_n}^{(n)})_{j_1, \dots, j_n=1}^{m_1, \dots, m_n} \right\|_{w, \min\{s, 2\}} \\ &\leq M \|A\| K_G^{2n-2} \left\| (x_j^{(1)})_{j=1}^{m_1} \right\|_{w, \min\{s, 2\}} \cdots \left\| (x_j^{(n)})_{j=1}^{m_n} \right\|_{w, \min\{s, 2\}}, \end{aligned}$$

which shows that  $A$  is multiple  $(r; \min\{s, 2\})$ -summing.  $\square$



**Corollary 4.3.** *Let  $1 \leq p \leq 2$ ,  $r \geq p$  and let  $F$  be a Banach space. The following assertions are equivalent:*

- (a)  $\mathcal{L}(\ell_1; F) = \Pi_{r;p}(\ell_1; F)$ .
- (b)  $\mathcal{L}({}^n\ell_1; F) = \Pi_{r;p}^n({}^n\ell_1; F)$  for every  $n \in \mathbb{N}$ .
- (c)  $\mathcal{L}({}^n\ell_1; F) = \Pi_{r;p}^n({}^n\ell_1; F)$  for some  $n \in \mathbb{N}$ .

*Proof.* (a)  $\implies$  (b) follows from Theorem 4.2, (b)  $\implies$  (c) is obvious and (c)  $\implies$  (a) follows from Remark 1.2.  $\square$

The last result of the paper makes clear how our methods systematize the subject and generalize and simplify the known results. The coincidence

$$\mathcal{L}({}^n\ell_1; \ell_2) = \Pi_{p;p}^n({}^n\ell_1; \ell_2) \text{ for } n \in \mathbb{N} \text{ and } 1 \leq p \leq 2 \quad (2)$$

was proved in [11] in the following fashion: the author proves first that  $\mathcal{L}({}^n\ell_1; \ell_2) = \Pi_{2;2}^n({}^n\ell_1; \ell_2)$  (a result we reobtained in Proposition 3.2), then uses this to prove that  $\mathcal{L}({}^n\ell_1; \ell_2) = \Pi_{1;1}^n({}^n\ell_1; \ell_2)$  (see also [3, Theorem 5.2]), and finally uses this last coincidence to obtain (2) (cf. [11, Corolario 5.24]). Several auxiliary results are used along the way. On the other hand, (2) is nothing but a particular case of the next corollary, which is a straightforward combination of Theorem 4.2 with [2, Theorem 5.2].

**Corollary 4.4.** *Let  $E_1, \dots, E_n$  be  $\mathcal{L}_1$ -spaces and let  $F$  be an  $\mathcal{L}_q$ -space,  $1 \leq q < +\infty$ . Then*

$$\mathcal{L}(E_1 \dots, E_n; F) = \Pi_{r;p}^n(E_1 \dots, E_n; F)$$

*if either*

- (a)  $q < 2$ ,  $r \geq q^*$  and  $p = 2$ , where  $\frac{1}{q} + \frac{1}{q^*} = 1$ , or
- (b)  $q > 2$ ,  $r \geq q$  and  $p = 2$ , or
- (c)  $q = 2$  and  $1 \leq p \leq r \leq 2$ .

ACKNOWLEDGEMENT. Part of this paper was written while G.B. was a CNPq Postdoctoral Fellow in the Departamento de Análisis Matemático at Universidad de Valencia. He thanks Pilar Rueda and the members of the department for their kind hospitality.

## References

- [1] M. Acosta, D. García and M. Maestre. *A multilinear Lindenstrauss theorem*, J. Funct. Anal. **235** (2006), 122-136.
- [2] G. Bennett. *Schur multipliers*, Duke Math. J. **44** (1977), 603-639.
- [3] F. Bombal, D. Pérez-García and I. Villanueva. *Multilinear extensions of Grothendieck's theorem*, Quart. J. Math. **55** (2004), 441-450.
- [4] A. Defant and K. Floret. *Tensor norms and operator ideals*, North-Holland Mathematical Studies 174, North-Holland, 1993.

- [5] A. Defant and D. Pérez-García. *A tensor norm preserving unconditionality for  $\mathcal{L}_p$ -spaces*, Trans. Amer. Math. Soc., to appear.
- [6] J. Diestel, H. Jarchow and A. Tonge. *Absolutely summing operators*, Cambridge University Press, Cambridge, 1995.
- [7] S. Dineen. *Complex Analysis on Infinite Dimensional Spaces*, Springer-Verlag, London, 1999.
- [8] M. C. Matos. *Fully absolutely summing mappings and Hilbert-Schmidt operators*, Collect. Math. **54** (2003), 111-136.
- [9] M. C. Matos and D. Pellegrino. *Fully summing mappings between Banach spaces*, Studia Math. **178** (2007), 47-61.
- [10] D. Pellegrino and M. Souza. *Fully summing multilinear and holomorphic mappings into Hilbert spaces*, Math. Nachr. **278** (2005), 877-887.
- [11] D. Pérez-García. *Operadores multilineales absolutamente sumantes*, Thesis, Univ. Complutense de Madrid, 2003.
- [12] D. Pérez-García. *The inclusion theorem for multiple summing operators*, Studia Math. **165** (2004), 275-290.
- [13] D. Pérez-García and I. Villanueva. *Multiple summing operators on Banach spaces*, J. Math. Anal. Appl. **285** (2003), 86-96.
- [14] D. Pérez-García and I. Villanueva. *Multiple summing operators on  $C(K)$  spaces*, Ark. Mat. **42** (2004), 153-171.
- [15] A. Pietsch. *Ideals of multilinear functionals*, Proceedings of the Second International Conference on Operator Algebras, Ideals and Their Applications in Theoretical Physics, 185-199, Teubner-Texte, Leipzig, 1983.
- [16] M. Souza. *Aplicações multilineares completamente absolutamente somantes*, Thesis, Univ. Estadual de Campinas, 2003.

[Geraldo Botelho] Faculdade de Matemática, Universidade Federal de Uberlândia, 38.400-902 - Uberlândia, Brazil, e-mail: botelho@ufu.br.

[Daniel Pellegrino] Departamento de Matemática, Universidade Federal da Paraíba, 58.051-900 - João Pessoa, Brazil, e-mail: dmpellegrino@gmail.com.