# EXTENSIONS OF LIE-RINEHART ALGEBRAS AND COTANGENT BUNDLE REDUCTION

J. HUEBSCHMANN, M. PERLMUTTER, AND T. S. RATIU

ABSTRACT. Let Q denote a smooth manifold acted upon smoothly by a Lie group G. The G-action lifts to an action on the total space  $T^*Q$  of the cotangent bundle of Q and hence on the standard symplectic Poisson algebra of smooth functions on  $T^*Q$ . The Poisson algebra of G-invariant functions on  $T^*Q$  yields a Poisson structure on the space  $(T^*Q)/G$  of G-orbits. We relate this Poisson algebra with extensions of Lie-Rinehart algebras and derive an explicit formula for this Poisson structure in terms of differentials. We then show, for the particular case where the G-action on Q is principal, how an explicit description of the Poisson algebra derived in the literature by an ad hoc construction is essentially a special case of the formula for the corresponding extension of Lie-Rinehart algebras. By means of various examples, we also show that this kind of description breaks down when the G-action does not define a principal bundle.

MSC 2010: Primary: 53D20; Secondary: 17B63 17B65 17B66 17B81 53D17

#### 1. Introduction

Let Q denote a smooth manifold acted upon smoothly by a Lie group G. The G-action lifts to a symplectic action on the total space  $T^*Q$  of the cotangent bundle of Q, and hence G acts on the ordinary symplectic Poisson algebra of smooth functions on  $T^*Q$ . Collective symplectic reduction then leads to the Poisson algebra of G-invariant functions on  $T^*Q$ , and the question arises of determining this Poisson algebra in terms of the space of G-orbits in Q and the requisite additional data.

The simplest non-trivial case one could perhaps think of arises when we take Q to be the group G itself, with G-action via left translation. In this case, the Poisson algebra of G-invariant functions on  $T^*Q$  simply comes down to the familiar Lie-Poisson algebra of smooth functions on the dual  $\mathfrak{g}^*$  of the Lie algebra  $\mathfrak{g}$  of G. A non-degenerate G-invariant scalar product on  $\mathfrak{g}$  then defines a G-invariant Riemannian metric or, equivalently, a kinetic energy on Q = G, and the associated equations of motion, that is, the equations defining the geodesics, are determined by the Euler-Poincaré equations on  $\mathfrak{g}$ . For  $G = SO(3, \mathbb{R})$ , viewed as the configuration space of a rigid body moving freely about its center of mass, the inertia tensor of the body is precisely such a quadratic form and the Euler-Poincaré equations are the familiar Euler equations for the motion of a free rigid body [1], [3], [30].

More generally, when  $Q \to B$  is a principal G-bundle, the manifold  $(T^*Q)/G$  carries a natural Poisson structure. Decomposing it in terms of the canonical Poisson structures on the total space  $T^*(Q/G)$  of the cotangent bundle of the shape space Q/G and the Lie-Poisson structure on the dual  $\mathfrak{g}^*$  of the Lie algebra  $\mathfrak{g}$  of G requires as an additional ingredient a principal connection. Under that hypothesis, the Poisson bracket has been determined in [31],[32] and used to study the motion of a particle in a Yang-Mills field (the Wong equations) and various continuum mechanical problems. A coordinate free proof of the resulting formula was given in [44]. A characterization of the symplectic leaves of this Poisson structure was found in [28], based on previous results in [41], [43].

For the case where the G-action is not free (and in particular not principal), the problem has been studied, is the subject of ongoing research, and partial results are known. The singular symplectic reduced space at zero, see [40], has additional structure induced by the cotangent bundle character of  $T^*Q$ ; this structure has been completely described in [37]. For the particular case where Q has a single G-orbit type, see [34], the same program has been carried out both in the symplectic and Poisson categories for arbitrary momentum values in [11], [36]. In the completely general case of a proper non-free action of G

1

Key words and phrases. Lie-Poisson algebra, Lie-Rinehart algebra, extension of Lie-Rinehart algebra, singular reduction, singular cotangent bundle reduction, semi-algebraic set, Poisson structure, tautological Poisson structure, symplectic leaf, fiber bundle, connection.

on Q, the description of the Poisson geometry of the stratified space  $(T^*Q)/G$  is built on the previously quoted results and elaborated on in work in progress.

In the present paper we will show that the constructions in the quoted references admit a simple explanation in terms of what we refer to as the  $tautological\ Poisson\ structure\ of\ a\ Lie-Rinehart\ algebra\ and, furthermore, in terms of extensions of Lie-Rinehart algebras. One of us has already explored this tautological Poisson structure in [12] (without having introduced that terminology). At the present stage, suffice it to say the following: Take <math>(A,L)$  to be the Lie-Rinehart algebra  $(C^{\infty}(M),\mathfrak{X}(M))$  which consists of the smooth functions  $C^{\infty}(M)$  and the smooth vector fields  $\mathfrak{X}(M)$  on a smooth manifold M; now the tautological Poisson algebra is the algebra of smooth functions on the total space  $T^*M$  of the cotangent bundle of M that are polynomial on the fibers, endowed with the (negative of the) ordinary cotangent bundle Poisson structure. For clarity we note that the tautological Poisson structure associated to a general Lie-Rinehart algebra is a purely algebraic construction; the algebraicity of the construction is here reflected in the fact that in the particular case under discussion we take functions that are polynomial on the fibers. We hope that the terminology 'tautological Poisson structure' does not conceal from the browsing reader that this concept is a substantial generalization of the Poisson structure on the dual of a Lie algebroid; we are indebted to the referee for having observed this risk.

We mention in passing that Lie-Rinehart algebras yield substantial new insight elsewhere, see [12]–[19] and, in particular, are a crucial tool in a systematic approach to singular Kähler reduction [20]–[22]. In particular Lie-Rinehart algebras provide the appropriate algebraic tool for a satisfactory description of the behavior of the theory at singularities. The corresponding singular Kähler quantization was developed in [23] and a particular gauge model for quantum mechanics on a stratified space was developed in [24]; an unforeseen physical phenomenon (a tunneling) arose out of that model.

At various stages in the paper, to make the paper somewhat more easily accessible to a wider audience, we make explicit the necessary transition from the algebra to the geometry that underlies our initial problem. In this vein, in Section 2, we reproduce the relationship between formal differentials, derivations, and vector fields. A good understanding of this relationship is indispensable for the rest of the paper. In Section 3 we explain the tautological Poisson algebra of a Lie-Rinehart algebra. In Section 4 we develop the behavior of the tautological Poisson algebra relative to an extension of Lie-Rinehart algebras. Here the general algebraic result is Theorem 4.2; it spells out the corresponding Poisson algebra in terms of data that characterize the underlying extension of Lie-Rinehart algebras. In Section 5 we then proceed to describe the Poisson structure arising from an extension of Lie-Rinehart algebras in terms of the formal differentials resulting from the kernel, of those resulting from the quotient and, furthermore, in terms of the requisite additional data; see Theorem 5.6 for details for the case where the quotient is a general Lie-Rinehart algebra and Theorem 5.9 for the particular case where the quotient is the Lie-Rinehart algebra of derivations of a commutative algebra A. These are structure results in the purely algebraic theory of Lie-Rinehart algebras that are interesting in their own right.

As for our initial problem, that of describing the Poisson structure of G-invariant functions on  $T^*Q$  in terms of the space of G-orbits in Q and the requisite additional data, when the action of G on Q is principal, the Poisson algebra of G-invariant functions on  $T^*Q$  can be completely characterized in terms of the tautological Poisson structure of a suitable Lie-Rinehart algebra and therefore in terms of extensions of Lie-Rinehart algebras, cf. Proposition 5.13. This observation recovers the familiar expressions for the gauged Lie-Poisson bracket. This kind of description generalizes to Lie algebroids with constant rank structure map, see Theorem 5.17, in particular, to transitive Lie algebroids, as explained in Corollary 5.18.

However, in the situation when the G-action on Q is not free, this kind of description and hence the characterizations given in the quoted references can only partially recover the Poisson algebra of G-invariant functions on  $T^*Q$ . In Section 6 below, we justify this observation with a number of examples. The description of the Poisson algebra in terms of the tautological Poisson algebra associated to a Lie-Rinehart algebra that we elaborate upon in the present paper opens the door to a systematic treatment of singular cotangent bundle reduction by means of techniques from real algebraic geometry; the examples in Section 6 only hint at a huge unexplored territory here, similar to that arising in singular Kähler reduction and singular Kähler quantization.

Our approach is essentially algebraic: rather than working with the Poisson 2-tensor we exploit the corresponding Poisson 2-form, and the requisite algebra provides a road map through the jungle of covariant derivatives, see, e.g., Theorems 5.6 and 5.9. Our approach immediately translates to other categories, for example, it applies to the orbit space of the total space of the cotangent bundle on a non-singular algebraic variety Q, endowed with the associated Poisson structure on the structure sheaf, when an algebraic group acts principally on Q. We spare the reader and ourselves these added troubles here. We plan to explore, at another occasion, the significance of our approach for the equations of motion.

Part of the present apparatus was first developed for the smooth case, phrased in the language of Lie algebroids, in papers of T. Courant, A. Weinstein, and others; see e. g. [5], [8], [9], [25], and the references there.

## Conventions.

We will write the de Rham operator on a smooth manifold as **d**. In the standard formalism, given a smooth symplectic manifold  $(M, \omega)$ , the Hamiltonian vector field  $X_h$  associated to the function  $h \in C^{\infty}(M)$  is defined by the identity

$$\mathbf{i}_{X_h}\omega := \omega(X_h, \cdot) = \mathbf{d}h;$$

given another smooth function h on M, the associated Poisson bracket of f and h is defined by

$$\{f, h\} := \omega(X_f, X_h) = X_h(f) = -X_f(h).$$

In particular, when M is the total space  $T^*Q$  of the cotangent bundle an a smooth manifold Q (the configuration space of a mechanical system) and  $\vartheta$  the tautological 1-form on  $T^*Q$ , the standard cotangent bundle symplectic form is defined by  $\omega := -\mathbf{d}\vartheta$  (see, e.g., [1, 3, 26, 27, 28, 29, 34]), so that for  $h \in C^{\infty}(T^*Q)$ , the Hamiltonian vector field  $X_h \in \mathfrak{X}(T^*Q)$  is given by the identity  $-\mathbf{i}_{X_h}\mathbf{d}\vartheta = \mathbf{d}h$ . This convention yields the standard Poisson brackets  $\{q, p\} = 1$  etc. and Hamilton's equation in the form  $\dot{f} = \{f, h\}$ ; in particular, with  $h = \frac{p^2}{2}$ , we get the standard relationship  $\dot{q} = \{q, h\} = p$ , and the Poisson bracket of the two functions f and h on  $T^*Q$  is given by  $\{f, h\} = -\mathbf{d}\vartheta(X_f, X_h)$ .

However the Poisson bracket  $\{\cdot,\cdot\}$  on  $C^{\infty}(T^*Q)$  we shall use in Section 3 assigns to  $f, h \in C^{\infty}(T^*Q)$  the function  $\{f,h\} = \mathbf{d}\vartheta(X_f,X_h)$ . In standard cotangent bundle coordinates  $(q^i,p_i)$  on  $T^*Q$  the value  $\{f,h\}$  of the Poisson bracket of f and h is then given by the expression

(1.1) 
$$\{f, h\} = \sum_{i=1}^{\dim M} \left( \frac{\partial f}{\partial p_i} \frac{\partial h}{\partial q^i} - \frac{\partial h}{\partial p_i} \frac{\partial f}{\partial q^i} \right)$$

and Hamilton's equations defined by  $h \in C^{\infty}(T^*Q)$  then take the form  $\dot{f} = \{h, f\}$ .

Throughout, we use the symbol ♦ to indicate the end of a remark.

# 2. Derivations and formal differentials

Let R be a commutative ring with identity, A an R-algebra with identity, at first not necessarily commutative, and M an A-bimodule. We recall that an M-valued R-derivation or M-valued derivation over R (relative Hochschild 1-cocycle)  $d: A \to M$  on A is a morphism of R-modules satisfying the additional requirement

(2.1) 
$$d(ab) = (da)b + a(db), \ a, b \in A.$$

When A is commutative and M an ordinary A-module, the requisite left and right A-module structures on M being identified, the identity (2.1) simplifies to

$$(2.2) d(ab) = b(da) + a(db), \ a, b \in A.$$

The alerted reader will notice that this is a purely algebraic identity and not just beginning calculus.

Let now A be a commutative R-algebra with unit and  $\mathcal{A}$  a commutative A-algebra with unit. We denote the R-module of derivations of A over R by  $\operatorname{Der}(A)$ , that of derivations of  $\mathcal{A}$  over R by  $\operatorname{Der}(\mathcal{A})$ , and the A-module of derivations of  $\mathcal{A}$  over A by  $\operatorname{Der}(\mathcal{A}|A)$ ; actually  $\operatorname{Der}(A)$  acquires an A-module and an R-Lie algebra structure and  $\operatorname{Der}(\mathcal{A})$  an  $\mathcal{A}$ -module and an A-Lie algebra structure. More generally given an A-module M, we denote the A-module of M-valued R-derivations on A by  $\operatorname{Der}(A, M)$  and, likewise, given an A-module M, we denote the A-module of M-valued R-derivations on A by  $\operatorname{Der}(A, M)$  and the A-module of M-valued A-derivations on A by  $\operatorname{Der}(A, M)$ .

Let  $D_A$  denote the A-module of formal differentials of A over R, likewise  $D_A$  the A-module of formal differentials of A over R, and  $D_{A|A}$  the A-module of formal differentials of A over A. We will write the

elements of  $D_A$  as da ( $a \in A$ ) etc. By definition, the A-module  $D_A$  represents the derivation functor  $Der(A, \cdot)$  on the category of A-modules: The universal derivation  $d: A \to D_A$  that assigns  $da \in D_A$  to  $a \in A$  has the property that, given an A-module M, the A-module morphism

(2.3) 
$$\operatorname{Hom}_A(D_A, M) \longrightarrow \operatorname{Der}(A, M), \ \varphi \mapsto \varphi \circ d, \ \varphi \in \operatorname{Hom}_A(D_A, M),$$

is an isomorphism of A-modules, plainly natural in M.

We will now recall a number of facts that we will subsequently use. To this end, let V be a finitely generated projective A-module and let  $S = S_A[V]$ , the symmetric A-algebra on V. Without a hypothesis of the kind that V be finitely generated and projective, a number of sequences that we exploit below need no longer be exact; moreover the theory to be developed below involves the A-dual of V and, without that hypothesis it would be hard to handle that A-dual effectively. In fact, without a hypothesis of that kind, many of the basic constructions in the present paper break down; this is, perhaps, the reason why the presence of singularities spoils the description of the Poisson algebra we are looking for. Indeed, in geometry, the projectivity hypothesis (of the space of derivations of the algebra of functions, viewed as a module over that algebra of functions) corresponds to the smoothness of the underlying space.

2.1. The sequence of inclusions  $R \subseteq A \subseteq \mathcal{A}$  determines a canonical exact sequence

$$(2.4) \mathcal{A} \otimes_A \mathcal{D}_A \longrightarrow \mathcal{D}_A \longrightarrow \mathcal{D}_{A|A} \longrightarrow 0$$

of A-modules.

2.2. The exact sequence (2.4) dualizes to a canonical exact sequence

$$(2.5) 0 \longrightarrow \operatorname{Der}(A|A) \longrightarrow \operatorname{Der}(A) \longrightarrow \operatorname{Der}(A,A)$$

of A-modules.

2.3. The canonical S-module morphism

(2.6) 
$$\operatorname{Hom}_{A}(V, S) \longrightarrow \operatorname{Der}(S|A), \quad \phi \mapsto d_{\phi}, \quad d_{\phi}(\alpha) := \phi(\alpha), \quad \alpha \in V,$$

is an isomorphism. Further, the canonical morphism  $S \otimes_A \operatorname{Hom}_A(V, A) \to \operatorname{Hom}_A(V, S)$  is an isomorphism of S-modules, and therefore the canonical S-module morphism

$$(2.7) S \otimes_A \operatorname{Hom}_A(V, A) \longrightarrow \operatorname{Der}(S|A)$$

is an isomorphism of S-modules as well. Consequently the assignment to  $\alpha \in V$ , viewed as a member of  $S = S_A[V]$ , of the differential  $d\alpha \in D_{S|A}$  induces an isomorphism

$$(2.8) S \otimes_A V \longrightarrow D_{S|A}$$

of S-modules. The isomorphism (2.6) is inverse to the dual of (2.8).

2.4. With S substituted for A, the exact sequence (2.4) of S-modules refines to

$$(2.9) 0 \longrightarrow S \otimes_A D_A \longrightarrow D_S \longrightarrow S \otimes_A V \longrightarrow 0.$$

Here the second unlabeled arrow sends a member of  $S \otimes_A D_A$  of the kind  $s \otimes_A da$  ( $s \in S$  and  $a \in A$ ) to the formal differential sda of S over R; the injectivity of this arrow is, perhaps, not entirely obvious, and we justify it in the appendix. Further, given  $\alpha \in V$ , the third arrow sends the formal differential  $d\alpha$  over R to the differential of S over A induced by  $\alpha$  relative to the A-module structure on V, the S-module  $D_{S|A}$  being identified with  $S \otimes_A V$  via (2.8).

2.5. With S substituted for A, the exact sequence (2.5) of S-modules refines to

$$(2.10) 0 \longrightarrow \operatorname{Hom}_{A}(V, S) \longrightarrow \operatorname{Der}(S) \longrightarrow S \otimes_{A} \operatorname{Der}(A) \longrightarrow 0;$$

in terms of the canonical isomorphisms  $\operatorname{Hom}_A(V, \mathbb{S}) \to \operatorname{Der}(\mathbb{S}|A)$  and  $\mathbb{S} \otimes_A \operatorname{Der}(A) \to \operatorname{Der}(A, \mathbb{S})$ , the two unlabeled arrows in (2.10) are given by the obvious associations:  $\operatorname{Der}(\mathbb{S}|A) \longrightarrow \operatorname{Der}(\mathbb{S})$  is the obvious inclusion and  $\operatorname{Der}(\mathbb{S}) \longrightarrow \operatorname{Der}(A, \mathbb{S})$  the obvious restriction. We note that the sequence (2.10) arises from (2.9) by dualization. However, when we dualize the sequence (2.10), we do not in general get back the sequence (2.9) unless the A-module  $D_A$  of formal differentials of A is reflexive.

- 2.6. Let B be a smooth manifold, and let  $A = C^{\infty}(B)$ . A smooth version  $D_A^{\text{smooth}}$  of the A-module  $D_A$  of formal differentials is the A-module  $D_A^{\text{smooth}} := \Gamma(\tau_B^*)$  of differential forms on B or, equivalently, smooth sections of the cotangent bundle  $\tau_B^* : T^*B \to B$  of B. However the canonical morphism  $D_A \to D_A^{\text{smooth}}$  of A-modules is surjective but not an isomorphism. For example, over the real line, the formal differential  $d\sin(x) \cos(x)dx$  is non-zero but is zero as a differential form.
- 2.7. To spell out a geometric interpretation of the exact sequences (2.9) and (2.10), let B be a smooth manifold and  $\lambda \colon \mathcal{L} \to B$  a smooth vector bundle on B. Let  $\rho \colon \mathcal{L}^* \to B$  denote the vector bundle on B that is dual to  $\lambda$  and, for  $b \in B$ , let  $\mathcal{L}_b^* := \rho^{-1}(b)$  be the fiber over  $b \in B$ . When we view the total space  $\mathcal{L}^*$  of the vector bundle  $\rho$  merely as a smooth manifold we denote that total space by N.

The injective smooth vector bundle map Ver:  $N \times_B \mathcal{L}^* \to TN$  over N given by

$$\operatorname{Ver}(\alpha_b, \beta_b) := \frac{d}{dt}\Big|_{t=0} (\alpha_b + t\beta_b) =: \operatorname{Ver}_{\alpha_b}(\beta_b)$$

has range the total space V of the vertical bundle  $V \to N$  and hence induces an isomorphism of vector bundles over N from  $N \times_B \mathcal{L}^* \to N$  to  $V \to N$ . Notice that, for every  $b \in B$ , the map  $\operatorname{Ver}_{\alpha_b}$  has the form  $\operatorname{Ver}_{\alpha_b} \colon \mathcal{L}_b^* \to T_{\alpha_b} N$ . Thus we obtain the following exact sequence of vector bundles with base N, spelled out here for the total spaces:

$$(2.11) 0 \longrightarrow N \times_B \mathcal{L}^* \longrightarrow TN \longrightarrow N \times_B TB \longrightarrow 0 (\alpha_b, \beta_b) \longmapsto \operatorname{Ver}_{\alpha_b}(\beta_b) \longrightarrow (n, T_n \rho(v_n))$$

where  $n \in N$  and  $\rho(n) = b$ . This is the geometric analog of the sequence (2.10).

Dualizing the sequence (2.11) we obtain an exact sequence of vector bundles over N; written out in terms of the total spaces, this sequence has the form

where  $n \in N$  and  $\rho(n) = b$ . This is the geometric analog of the sequence (2.9).

Indeed, over the reals  $\mathbb{R}$  as ground ring, let  $A = C^{\infty}(B)$  and take the spaces of sections in the smooth category; relative to the resulting sequence  $\mathbb{R} \subseteq A \subseteq C^{\infty}(\mathcal{L}^*)$  of real algebras, (2.12) thus yields precisely an extension of the kind (2.9) and (2.11) one of the kind (2.10) except that the algebra  $C^{\infty}(\mathcal{L}^*)$  of smooth functions on  $\mathcal{L}^*$  has now been substituted for S. We note that the two sequences (2.11) and (2.12) are dual to each other.

- 2.8. Abstracting from the previous example, as before, let A be a unital commutative R-algebra and suppose that, as an A-module, Der(A) is finitely generated and projective. Over a field as ground ring, when A is an affine algebra, this property can be taken as that defining the notion of regularity of a commutative algebra. In the present general case, the double A-dual  $D_A^{**}$  of  $D_A$ , that is, the A-dual  $Der(A)^* = Hom_A(Der(A), A)$  of Der(A), is a finitely generated projective A-module as well. This module represents the derivation functor  $Der(A, \cdot)$  on the category of finitely generated projective A-modules. In particular, under heading (2.6), the A-module  $D_A^{**}$  is precisely the double A-dual  $D_A^{**}$  of  $D_A$ .
  - 3. The Tautological Poisson algebra of a Lie-Rinehart algebra
- 3.1. **Motivation.** To motivate the abstract notion of tautological Poisson algebra associated to a Lie-Rinehart algebra, we will first explain a special case.

Let Q denote a smooth finite dimensional manifold and  $\tau_Q \colon \mathrm{T}Q \to Q$  and  $\tau_Q^* \colon \mathrm{T}^*Q \to Q$  its tangent and cotangent bundles, respectively. The 1-form  $\vartheta \in \Omega^1(\mathrm{T}^*Q)$  defined as the composite

$$\begin{array}{ll} \mathrm{T}\mathrm{T}^{*}Q & \longrightarrow \mathrm{T}Q \times_{Q} \mathrm{T}^{*}Q & \longrightarrow \mathbb{R} \\ V_{\alpha_{q}} & \longmapsto \left(\mathrm{T}_{\alpha_{q}}\tau_{Q}^{*}\left(V_{\alpha_{q}}\right),\alpha_{q}\right) & \longmapsto \left\langle\alpha_{q},\mathrm{T}_{\alpha_{q}}\tau_{Q}^{*}\left(V_{\alpha_{q}}\right)\right\rangle, \ q \in Q, \ \alpha_{q} \in \mathrm{T}_{q}^{*}Q, \end{array}$$

is the tautological 1-form on  $T^*Q$ . In standard cotangent bundle coordinates  $(q^j, p_j)$ , the 1-form  $\vartheta$  takes the form  $p_j \mathbf{d} q^j$ . The standard symplectic form on  $T^*Q$  is the 2-form  $-\mathbf{d}\vartheta$ . For formal reasons, we will instead work with the symplectic form  $\mathbf{d}\vartheta$  on  $T^*Q$ , and we will denote the associated Poisson bracket by  $\{\cdot,\cdot\}$ . This is the Poisson bracket which, in Darboux coordinates, is given by (1.1).

It is well known that, relative to the obvious filtration of the algebra of globally defined differential operators on Q, the associated graded algebra is canonically isomorphic to the algebra of smooth functions on  $T^*Q$  that are polynomial on the fibers. In terms of the notation  $(A, L) = (C^{\infty}(Q), \mathfrak{X}(Q))$ , this graded algebra is the symmetric A-algebra  $\mathcal{S}_A[L]$  on L. It is a standard fact that the commutator of differential operators induces a Poisson bracket on  $\mathcal{S}_A[L]$ , in fact, the restriction of the Poisson bracket  $\{\cdot, \cdot\}$  on  $C^{\infty}(T^*Q)$  to the subalgebra  $\mathcal{S}_A[L]$ .

We will now recall some of the details: The algebra  $C^{\infty}(T^*Q)$  being endowed with the Poisson bracket  $\{\cdot,\cdot\}$ , the vector space  $\operatorname{Lin}(T^*Q)$  of smooth functions on  $T^*Q$  that are linear on the fibers of the cotangent bundle  $\tau_Q^* \colon T^*Q \to Q$  of Q is a Lie (not Poisson) subalgebra of  $C^{\infty}(T^*Q)$ . Each smooth vector field X on Q defines a smooth momentum function  $\delta_X \colon T^*Q \ni \alpha_q \mapsto \langle \alpha_q, X(q) \rangle \in \mathbb{R}$   $(q \in Q)$  and the assignment to the vector field X on Q of the function  $\delta_X$  is a Lie algebra homomorphism  $\delta \colon \mathfrak{X}(Q) \to \operatorname{Lin}(T^*Q)$ . Indeed, write the action of the Lie algebra  $\mathfrak{X}(Q)$  of vector fields on Q on the algebra  $C^{\infty}(Q)$  of smooth functions on Q as

$$\mathfrak{X}(Q) \times C^{\infty}(Q) \longrightarrow C^{\infty}(Q), \quad (X, f) \longmapsto X(f);$$

now, for any  $f, h \in C^{\infty}(Q)$  and  $X, Y \in \mathfrak{X}(Q)$ , the Poisson bracket on  $\mathcal{S}_A[L]$  is determined by the following relations:

$$(3.1) \{f \circ \tau_Q^*, h \circ \tau_Q^*\} = 0, \{\delta_X, f \circ \tau_Q^*\} = X(f), \{\delta_X, \delta_Y\} = \delta_{[X,Y]}.$$

We will refer to these relations later when we study the general situation.

Via the ring operations in  $C^{\infty}(T^*Q)$  and the action of  $\mathfrak{X}(Q)$  on  $C^{\infty}(Q)$ , the functions on  $T^*Q$  of the kind  $f \circ \tau_Q^*$ , where f ranges over functions in  $C^{\infty}(Q)$ , together with the functions on  $T^*Q$  of the kind  $\delta_X$ , where X ranges over smooth vector fields on Q, completely determine the Poisson bracket  $\{\cdot, \cdot\}$  on  $C^{\infty}(T^*Q)$ .

**Proposition 3.1.** The Poisson bracket on  $C^{\infty}(T^*Q)$  determined by (3.1) is the symplectic Poisson bracket associated to the 1-form  $d\vartheta$ .

*Proof.* This is immediate: By construction, in Darboux coordinates  $q_1, \ldots, q_m, p_1, \ldots, p_m$ , the Poisson bracket (3.1) is given by  $\{p_j, q_k\} = \delta_{j,k}, 1 \leq j, k \leq m$ .

We note that here and henceforth we do not distinguish in notation between vector fields on Q and derivations of  $C^{\infty}(Q)$ .

Remark 3.2. For a general smooth manifold Q, with our conventions, given a smooth function f, the assignment to f of the smooth vector field  $\{f, \cdot\}$  on  $T^*Q$  is a Lie algebra map from the Lie algebra underlying the Poisson algebra to the Lie algebra of smooth vector fields whereas, in the standard formalism, the ordinary Hamiltonian vector field map is an anti Lie morphism; at the technical level, the present convention has the advantage that a structure map to be introduced later (the morphism  $\pi^{\sharp}$  in Proposition 3.4 below) is compatible with the Lie-Rinehart structures, and this will greatly simplify the exposition.

**Remark 3.3.** Let G be a connected Lie group, and let  $\mathfrak{g}$  denote its Lie algebra, more precisely, the Lie algebra of left-invariant vector fields on G. These are the fundamental vector fields for the right translation action of G on itself, and left translation

$$G \times \mathfrak{g} \longrightarrow \mathrm{T} G$$

trivializes the tangent bundle of G. Let  $X,Y \in \mathfrak{g}$ ; the Poisson bracket  $\{\delta_X,\delta_Y\}$  of the associated momentum functions on  $T^*G$  is given by  $\{\delta_X,\delta_Y\}=\delta_{[X,Y]}$ ; this is a mere tautology since the Lie bracket  $[X,Y] \in \mathfrak{g}$  is defined in terms of the commutator bracket of vector fields on G.

Let G act on Q from the right. Then the fundamental vector field map  $\mathfrak{g} \to \mathfrak{X}(Q)$  is a morphism of Lie algebras, and the composite

$$\delta \colon \mathfrak{g} \longrightarrow C^{\infty}(\mathrm{T}^*Q)$$

of the fundamental vector field map with the momentum function  $\delta$  yields the *comomentum* associated to the corresponding momentum mapping  $\mu \colon \mathrm{T}^*Q \to \mathfrak{g}^*$ . The G-action being on the right of Q, the G-equivariance of the momentum mapping is equivalent to the momentum mapping being a Poisson map relative to the Poisson bracket  $\{\cdot,\cdot\}$  on  $\mathrm{T}^*Q$  chosen above (associated to  $\mathrm{d}\vartheta$ , that is, the negative of

the standard bracket) and the ordinary Lie-Poisson bracket on  $\mathfrak{g}^*$ . Hence the comomentum (3.2) is a morphism of Lie algebras. We will write  $\mathfrak{g}_+^*$  when we take  $\mathfrak{g}^*$  to be endowed with the standard Lie-Poisson structure and  $\mathfrak{g}_-^*$  when we think of  $\mathfrak{g}^*$  as being endowed with the negative of the standard Lie-Poisson structure.

We will now take Q to be the group G itself. Let  $\{\cdot, \cdot\}_+$  denote the standard cotangent bundle Poisson bracket on  $T^*G$  so that  $\{\cdot, \cdot\}$  is the negative of the standard cotangent bundle Poisson bracket. It is well known that the composite

$$(3.3) T^*G \longrightarrow \mathfrak{g}^* \times G \longrightarrow \mathfrak{g}^*$$

of right translation

$$T^*G \longrightarrow \mathfrak{g}^* \times G, \ \alpha_x \longmapsto (\alpha_x \circ \rho_x, x), \ x \in G,$$

with the projection to the first component yields the momentum mapping

(3.4) 
$$\mu_{\ell} \colon \mathrm{T}^* G \longrightarrow \mathfrak{g}^*, \ \alpha_x \longmapsto \alpha_x \circ \rho_x$$

for the G-action on  $T^*G$  by left translation. The G-equivariance of the momentum mapping  $\mu_\ell$  is well known to be equivalent to  $\mu_\ell$  being a Poisson map relative to  $\{\cdot,\cdot\}_+$ ; hence  $\mu_\ell$  induces isomorphisms

$$((\mathbf{T}^*G)/G, \{\cdot, \cdot\}_+) \longrightarrow \mathfrak{g}_+^*, \quad ((\mathbf{T}^*G)/G, \{\cdot, \cdot\}) \longrightarrow \mathfrak{g}_-^*$$

of Poisson manifolds.

3.2. The formal notion. The pair  $(A, L) := (C^{\infty}(Q), \mathfrak{X}(Q))$  is among the standard examples of a *Lie-Rinehart* algebra over the real numbers as ground ring [12], [13], [14], [38] and, abstracting from the cotangent bundle situation, we are led to the *tautological Poisson algebra* associated to a Lie-Rinehart algebra that we now describe.

Fix the commutative ring with identity R; the unadorned tensor product symbol  $\otimes$  will always refer to the tensor product over R. Further, denote by A a unital commutative R-algebra. An (R,A)-Lie algebra [38] is a Lie algebra L over R which acts on A (from the left) by derivations (the action being written as  $(\alpha \otimes a) \mapsto \alpha(a)$ ) and is also an A-module (the structure map being written as  $(a \otimes \alpha) \mapsto a\alpha$ ), in such a way that suitable compatibility conditions are satisfied which generalize standard properties of the Lie algebra of vector fields on a smooth manifold viewed as a module over the ring of functions; these conditions read

$$(3.6) (a\alpha)(b) = a(\alpha(b)), \quad \alpha \in L, a, b \in A,$$

$$[\alpha, a\beta] = a[\alpha, \beta] + \alpha(a)\beta, \quad \alpha, \beta \in L, a \in A.$$

The pair (A, L) is then referred to a Lie-Rinehart algebra. Given two Lie-Rinehart algebras (A, L) and (A', L'), a morphism  $(\phi, \psi) \colon (A, L) \longrightarrow (A', L')$  of Lie-Rinehart algebras consists of morphisms  $\phi \colon A \to A'$  and  $\psi \colon L \to L'$  in the appropriate categories that are compatible with the additional structure. With this notion of morphism, Lie-Rinehart algebras constitute a category. Apart from the example of smooth functions and smooth vector fields on a smooth manifold, a related (but more general) example is the pair consisting of a commutative algebra A and the R-module Der(A) of derivations of A with the obvious A-module structure and the commutator bracket on Der(A); here the commutativity of A is crucial.

Take a Lie-Rinehart algebra (A, L), that is, A is a commutative algebra and L an (R, A)-Lie algebra. Denote by  $\mathcal{S}_A[L]$  the symmetric A-algebra on L. The L-action on A and the R-Lie bracket operation on L induce a Poisson bracket

$$\{\cdot,\cdot\}\colon \mathbb{S}_A[L]\otimes \mathbb{S}_A[L]\longrightarrow \mathbb{S}_A[L]$$

on  $S_A[L]$  completely determined by the relations

(3.9) 
$$\{\alpha, \beta\} = [\alpha, \beta], \quad \alpha, \beta \in L,$$

$$\{\alpha, a\} = \alpha(a) \in A, \quad a \in A, \ \alpha \in L,$$

$$\{u, vw\} = \{u, v\}w + v\{u, w\}, \quad u, v, w \in \mathcal{S}_A[L].$$

In particular, taking  $u=a, v=b\in A$  and  $w=\alpha\in L, \alpha\neq 0$  in (3.11), we obtain  $\{a,b\}=0$ . We refer to the Poisson algebra  $(\mathcal{S}_A[L],\{\cdot,\cdot\})$  as the tautological Poisson algebra associated to the Lie-Rinehart algebra (A,L). We note that, for the standard example  $(C^{\infty}(Q),\mathfrak{X}(Q))$  of a Lie-Rinehart algebra, the tautological Poisson bracket just defined is the same as the Poisson bracket given in (3.1).

The commutative R-algebra A being fixed, the tautological Poisson algebra is plainly functorial in (R, A)-Lie algebras: A morphism  $\phi: L_1 \to L_2$  of (R, A)-Lie algebras induces a morphism

$$(3.12) \qquad (S_A[L_1], \{\cdot, \cdot\}) \longrightarrow (S_A[L_2], \{\cdot, \cdot\})$$

of Poisson algebras.

We will now justify the terminology "tautological Poisson algebra". To this end, we recall briefly how, for an arbitrary Poisson algebra, an appropriate Lie-Rinehart algebra serves as a replacement for the tangent bundle of a smooth symplectic manifold.

Let  $(\mathcal{P}, \{\cdot, \cdot\})$  be a Poisson algebra. The association  $(du, dv) \mapsto \pi(du, dv) := \{u, v\}$ , as u and v range over  $\mathcal{P}$ , yields a  $\mathcal{P}$ -valued  $\mathcal{P}$ -bilinear skew-symmetric 2-form

$$(3.13) \pi = \pi_{\{\cdot,\cdot,\cdot\}} \colon \mathcal{D}_{\mathcal{P}} \otimes \mathcal{D}_{\mathcal{P}} \longrightarrow \mathcal{P}$$

on  $D_{\mathcal{P}}$ ; this 2-form is said to be the *Poisson 2-form* of  $(\mathcal{P}, \{\cdot, \cdot\})$ . The adjoint

(3.14) 
$$\pi^{\sharp} : \mathcal{D}_{\mathcal{P}} \longrightarrow \mathrm{Der}(\mathcal{P}) = \mathrm{Hom}_{\mathcal{P}}(\mathcal{D}_{\mathcal{P}}, \mathcal{P}), \qquad \pi^{\sharp}(du)(dv) = \{u, v\},$$

of  $\pi$  is a morphism of  $\mathcal{P}$ -modules and the formula

$$[adu, bdv] := a\{u, b\}dv + b\{a, v\}du + abd\{u, v\}, \quad a, b, u, v \in \mathcal{P},$$

yields a Lie bracket  $[\cdot, \cdot]$  on  $D_{\mathcal{P}}$ , viewed as an R-module (see [12] for more details). For the record we recall the following result, established in [12, Theorem 3.8].

**Proposition 3.4.** The  $\mathcal{P}$ -module structure on  $D_{\mathcal{P}}$ , the R-bilinear bracket  $[\cdot, \cdot]$  on  $D_{\mathcal{P}}$ , and the morphism  $\pi^{\sharp} \colon D_{\mathcal{P}} \to \operatorname{Der}(\mathcal{P})$  of  $\mathcal{P}$ -modules endow the pair  $(\mathcal{P}, D_{\mathcal{P}})$  with a Lie-Rinehart algebra structure in such a way that  $\pi^{\sharp} \colon D_{\mathcal{P}} \to \operatorname{Der}(\mathcal{P})$  is a morphism of  $(R, \mathcal{P})$ -Lie algebras.

Given a  $\mathcal{P}$ -module D, we will use the notation  $\operatorname{Alt}_{\mathcal{P}}(D,\mathcal{P})$  for the graded  $\mathcal{P}$ -algebra of  $\mathcal{P}$ -valued  $\mathcal{P}$ -multilinear alternating forms on D. When  $(\mathcal{P},D)$  is a Lie-Rinehart algebra (over the ground ring R), the standard Cartan-Chevalley-Eilenberg operator d (equivalently: the de Rham operator evaluated formally), endows  $\operatorname{Alt}_{\mathcal{P}}(D,\mathcal{P})$  with a differential graded R-algebra structure, cf. [12]. Denote by  $D_{\{\cdot,\cdot\}}$  the  $(R,\mathcal{P})$ -Lie algebra  $(D_{\mathcal{P}},[\cdot,\cdot],\pi^{\sharp})$ . The 2-form  $\pi_{\{\cdot,\cdot\}}$ , which is defined for every Poisson algebra, is a 2-cocycle in the Rinehart algebra  $(\operatorname{Alt}_{\mathcal{P}}(D_{\{\cdot,\cdot\}},\mathcal{P}),d)$ . As opposed to other definitions that involve a defining chain complex and only refer to a Poisson algebra of smooth functions on a smooth manifold, in [12], the Poisson cohomology  $H^*_{\operatorname{Poisson}}(\mathcal{P},\mathcal{P})$  of the Poisson algebra  $(\mathcal{P},\{\cdot,\cdot\})$  was defined as the cohomology of the Lie-Rinehart algebra  $D_{\{\cdot,\cdot\}}$ , *i.e.*,

When  $\mathcal{P}$  is the algebra of smooth functions on a smooth symplectic manifold M, endowed with the symplectic Poisson structure, the inverse of the symplectic structure induces a vector bundle isomorphism from  $\tau_M^*$  onto  $\tau_M$  and hence an isomorphism from the smooth version  $\mathrm{D}^{\mathrm{smooth}}_{\{\cdot,\cdot\}}\cong\Omega^1(M)$  (the  $C^\infty$ -module of ordinary 1-forms on M) of  $\mathrm{D}_{\{\cdot,\cdot\}}$  onto the ordinary  $(\mathbb{R},\mathcal{P})$ -Lie algebra of smooth vector fields on M which identifies the 2-form  $\pi$  on  $\mathrm{D}_{\mathcal{P}}$  with the symplectic structure. Furthermore, under the canonical isomorphism between  $\mathrm{Alt}_{\mathcal{P}}(\mathrm{D}^{\mathrm{smooth}}_{\mathcal{P}},\mathcal{P})$  and the exterior  $\mathcal{P}$ -algebra  $\wedge_{\mathcal{P}}(\mathfrak{X}(M))$  on  $\mathfrak{X}(M)$ , the 2-form  $\pi$  corresponds to the familiar Poisson 2-tensor. In addition, when P is a Poisson manifold and d the de Rham operator, the bracket (3.15) coincides with the Lie bracket on 1-forms on P; see [1, Proposition 3.3.8] for the symplectic case and [12, Subsection 3.12] and [42, Theorem 4.1] for the general Poisson case. We emphasize, cf. [12], that the abstract construction given above defines the 2-form  $\pi$  for an arbitrary Poisson algebra, whether or not it arises from a smooth symplectic manifold.

We return to a general Lie-Rinehart algebra (A, L). For the algebra  $\mathcal{S} := \mathcal{S}_A[L]$ , endowed with the Poisson structure (3.8), the Poisson 2-form  $\pi \colon D_{\mathcal{S}} \otimes_{\mathcal{S}} D_{\mathcal{S}} \to \mathcal{S}$  given by (3.13) is a Poisson coboundary, that is,  $\{\cdot, \cdot\}$  admits a Poisson potential [12]. Indeed, as an algebra,  $\mathcal{S}$  is generated by the elements of A and those of L. Hence, as an  $\mathcal{S}$ -module,  $D_{\mathcal{S}}$  is generated by the formal differentials da,  $a \in A$ , and  $d\alpha$ ,  $\alpha \in L$ . Let  $D_{\mathcal{S}|A}$  denote the  $\mathcal{S}$ -module of formal differentials of  $\mathcal{S}$  over A; as an  $\mathcal{S}$ -module,  $D_{\mathcal{S}|A}$  is an induced module of the form  $\mathcal{S} \otimes_A L$  generated by the formal differentials  $d\alpha$ , where  $\alpha$  ranges over L.

A straightforward calculation (see [12, p. 92]) shows that the 1-form  $\vartheta \colon D_{\mathcal{S}} \to \mathcal{S}$  given through the projection to  $D_{\mathcal{S}|A}$  by

(3.17) 
$$\vartheta(d\alpha) := \alpha, \quad \alpha \in L,$$

is a Poisson potential for  $\{\cdot, \cdot\}$ , that is,

(3.18) 
$$\pi = d\vartheta \in \operatorname{Alt}_{S}^{2}(D_{\{\cdot,\cdot\}}, S).$$

For later reference, we note that the 2-form  $\pi$  is given by the identities

(3.19) 
$$\pi(d\alpha, d\beta) = [\alpha, \beta], \quad \alpha, \beta \in L,$$

(3.20) 
$$\pi(d\alpha, db) = \alpha(b), \quad \alpha \in L, \ b \in A.$$

This description of the 2-form  $\pi$  is consistent in the following sense: Given  $a \in A$  and  $\alpha \in L$ , when  $a\alpha$  is viewed as a member of  $S = S_A[L]$ , the differential  $d(a\alpha)$  over the ground ring R takes the form

$$d(a\alpha) = (da)\alpha + a(d\alpha)$$

and, given  $\beta \in L$ , the identities

$$\pi (d(a\alpha), d\beta) = [a\alpha, \beta] = a[\alpha, \beta] - \beta(a)\alpha$$
$$\pi ((da)\alpha + ad\alpha, \beta) = \alpha\pi (da, \beta) + a[\alpha, \beta] = -\beta(a)\alpha + a[\alpha, \beta]$$

hold. We emphasize that the 2-form  $\pi$  is  $S_A[L]$ -bilinear. In particular, the following observation is immediate; we spell it out, for later reference.

**Proposition 3.5.** The Poisson structure (3.8) on  $S_A[L]$  recovers, in fact, is equivalent to, the Lie-Rinehart structure on (A, L): Given  $\alpha, \beta \in L$ , the bracket  $[\alpha, \beta]$  is determined by the identity (3.9) and, given  $a \in A$  and  $\alpha \in L$ , the value  $\alpha(a)$  of the action of  $\alpha$  on a is determined by the identity (3.10). Thus the Lie-Rinehart structure can be reconstructed from the Poisson structure.

Remark 3.6. Suppose that  $\mathcal{P}$  is the algebra of smooth functions on the total space  $N=\mathrm{T}^*B$  of the cotangent bundle on a smooth manifold B, endowed with the standard symplectic Poisson structure. Then the smooth version of the 1-form  $\vartheta$  given in (3.17) is identical to the standard 1-form on  $N=\mathrm{T}^*B$ , and the resulting symplectic structure is therefore occasionally referred to as the tautological symplectic structure on  $N=\mathrm{T}^*B$ . Actually, in this case, with the notation  $(A,L)=(C^\infty(B),\mathfrak{X}(B))$ , in the appropriate Fréchet topology,  $\mathcal{S}_A[L]$  is dense in the algebra  $\mathcal{P}=C^\infty(\mathrm{T}^*B)$  of smooth functions on  $N=\mathrm{T}^*B$ : Indeed, using a locally affine open cover of B, in the standard manner, we endow  $C^\infty(\mathrm{T}^*B)$  with the locally convex Fréchet algebra structure which, on the total space of the cotangent bundle of any of the affine open subsets of B in the cover, is given by the standard construction involving the corresponding families of submultiplicative semi-norms. In the resulting topology,  $\mathcal{S}_A[L]$  is dense in  $C^\infty(\mathrm{T}^*B)$ . Under the canonical isomorphism between

$$Alt_{\mathcal{P}}(D_{\mathcal{P}}^{smooth}, \mathcal{P}) = Alt_{\mathcal{P}}(\Omega^{1}(N), \mathcal{P})$$

and the exterior  $\mathcal{P}$ -algebra  $\wedge_{\mathcal{A}}(\mathfrak{X}(N))$  on  $\mathfrak{X}(N)$  recalled above for a general smooth symplectic manifold written there as M, the formulas (3.19) and (3.20) then yield the Poisson 2-tensor for the cotangent bundle Poisson structure on  $T^*B$ . The Poisson structure (3.8) on the algebra  $\mathcal{S}_A[L]$  relative to a general Lie-Rinehart algebra (A, L) is formally of the same kind.

For an ordinary real Lie algebra  $\mathfrak{g}$ , viewed as an  $(\mathbb{R}, \mathbb{R})$ -Lie algebra with trivial  $\mathfrak{g}$ -action on  $\mathbb{R}$ , the tautological Poisson structure plainly comes down to the algebraic version of the ordinary Lie-Poisson structure on the dual  $\mathfrak{g}^*$  of  $\mathfrak{g}$ .

These observations are intended to justify the terminology tautological Poisson structure.

# 4. Extensions of Lie-Rinehart algebras and tautological Poisson algebra

The algebraic analog of an "Atiyah sequence" (associated to a principal bundle) or of a "transitive Lie algebroid" (see below for details) is an extension of Lie-Rinehart algebras [16]. We note that transitive Lie algebroids which do *not* arise from a principal bundle abound; see, e.g., [2]. We recall at the present stage that a Lie algebroid is said to be *transitive* when its anchor map is fiberwise surjective.

Let L', L, L'' be (R, A)-Lie algebras. An extension of (R, A)-Lie algebras is a short exact sequence

$$\mathbf{e} \colon 0 \longrightarrow L' \longrightarrow L \stackrel{p}{\longrightarrow} L'' \longrightarrow 0$$

in the category of (R, A)-Lie algebras; notice, in particular, that the Lie algebra L' necessarily acts trivially on A.

Recall that an **e**-connection associated to the extension (4.1) is an A-module section  $\omega \colon L'' \to L$  of the projection  $p \colon L \to L''$ . The **e**-curvature of  $\omega$  is the A-bilinear map  $\Omega \colon L'' \times L'' \to L'$  of A-modules defined by the identity

$$[\omega(\alpha''), \omega(\beta'')] = \omega\left([\alpha'', \beta'']_{L''}\right) + \Omega(\alpha'', \beta''), \ \alpha'', \beta'' \in L'',$$

where  $[\alpha'', \beta'']_{L''}$  refers to the Lie bracket in L''.

As discussed below, the notions of e-connection and e-curvature generalize the standard concepts of principal connection and principal curvature. For a general extension (4.1) of (R, A)-Lie algebras, a choice of connection  $\omega$  induces an A-module decomposition

$$L = L' \oplus \omega(L'') \cong L' \oplus L''$$

and, in terms of this decomposition, the Lie bracket  $[\cdot,\cdot]$  on L takes the form

$$(4.3) \qquad [(\alpha', \alpha''), (\beta', \beta'')] = ([\alpha', \beta']_{L'} + [\alpha', \beta''] + [\alpha'', \beta'] + \Omega(\alpha'', \beta''), [\alpha'', \beta'']_{L''});$$

here  $[\alpha', \beta']_{L'}$  refers to the Lie bracket in L'. When L'' is a projective A-module, one can always find an e-connection.

Since  $\omega$  is a section of p, viewed as a morphism of A-modules, it follows from (4.2) that the values of  $\Omega$  lie in L'. Thus  $\Omega$  is an L'-valued alternating A-bilinear 2-form on L''. We note that, similarly as in standard Schreier extension theory for groups, suitably interpreted,  $\Omega$  is a non-abelian 2-cocycle.

**Example 4.1.** Here we take the ground ring R to be the field  $\mathbb{R}$  of real numbers. Let B be a smooth finite-dimensional manifold and denote by A the algebra  $C^{\infty}(B)$  of smooth functions on B. Let G be a Lie group and  $\xi \colon Q \to B$  a right principal G-bundle. The vertical subbundle  $\tau_Q | V \colon V \to Q$  of the tangent bundle  $\tau_Q \colon \mathrm{T}Q \to Q$  is well known to be trivial (beware, not equivariantly trivial), having as fiber the Lie algebra  $\mathfrak{g}$  of G, that is,  $V \cong Q \times \mathfrak{g}$ . These data fit into the fundamental exact sequence

where  $\lambda_Q(q)$ :  $=\frac{d}{dt}\Big|_{t=0} q \cdot \exp(t\lambda) \in \mathrm{T}_q Q$  defines the fundamental vector field  $\lambda_Q \in \mathfrak{X}(Q)$ . We note that the range of the second arrow is the total space  $V \subset \mathrm{T}Q$  of the subbundle of the tangent bundle whose sections are vertical vector fields. All arrows are equivariant vector bundle maps relative to the diagonal action

$$(q,\lambda)\cdot x\colon = (q\cdot x, \operatorname{Ad}_{x^{-1}} x), \quad q\in Q, \ \lambda\in\mathfrak{g}, \ x\in G,$$

on  $Q \times \mathfrak{g}$  and the tangent lifted action on TQ. Dividing out the G-actions, we obtain an extension

$$(4.5) 0 \longrightarrow \operatorname{ad}(\xi) \longrightarrow \tau_O/G \longrightarrow \tau_B \longrightarrow 0$$

of vector bundles over B, where  $\tau_B$  is the tangent bundle of B, and

is a transitive Lie algebroid over B. Here the notation  $\mathrm{ad}(\xi)$  refers to the vector bundle  $Q \times_G \mathfrak{g} \to B$  associated to the principal bundle  $\xi$  by the adjoint representation of G on its Lie algebra  $\mathfrak{g}$ . The sequence (4.5) was introduced by Atiyah [4] and is now usually referred to as the *Atiyah sequence* of the principal bundle  $\xi$ .

The spaces of smooth sections  $\Gamma_B(\operatorname{ad}(\xi))$  and  $\Gamma_B(\tau_Q/G)$  over B inherit obvious Lie algebra structures, in fact  $(\mathbb{R}, A)$ -Lie algebra structures, and

$$(4.7) 0 \longrightarrow \Gamma_B(\operatorname{ad}(\xi)) \longrightarrow \Gamma_B(\tau_Q/G) \longrightarrow \mathfrak{X}(B) \longrightarrow 0$$

is an extension of  $(\mathbb{R}, A)$ -Lie algebras; here  $\mathfrak{X}(B)$  is the ordinary Lie algebra of vector fields on B and  $\Gamma_B(\mathrm{ad}(\xi))$  is in an obvious way the Lie algebra of the *group* of *gauge transformations* of  $\xi$ .

Pick a connection on the principal bundle  $\xi \colon Q \to Q/G = B$ . The associated horizontal lift operator hor:  $Q \times_B TB \to TQ$ , thought of as a G-equivariant vector bundle morphism over Q, is a section of the sequence (4.4). Taking the quotient by G we obtain a section [hor]:  $TB \to (TQ)/G$  of the Atiyah sequence (4.5). Conversely, any section  $\sigma \colon TB \to (TQ)/G$  of the Atiyah sequence (4.5) induces a G-equivariant section of (4.4). Thus, taking  $L' = \Gamma_B(\mathrm{ad}(\xi))$ ,  $L = \Gamma_B(\tau_Q/G)$ , and  $L'' = \mathfrak{X}(B)$ , we see that the chosen connection determines a unique e-connection  $\omega \colon L'' \to L$  described above. We note that we can identify L' with the G-invariant vertical vector fields on Q, and L'' with the G-invariant horizontal vector fields on Q.

Recall that the curvature of the chosen connection, thought of as a 2-form on B with values in the adjoint bundle  $\operatorname{ad}(\xi)$ , is the  $C^{\infty}(B)$ -bilinear map  $\Omega \colon \mathfrak{X}(B) \times \mathfrak{X}(B) \to \Gamma_B(\operatorname{ad}(\xi))$  of  $C^{\infty}(B)$ -modules characterized by the identity

$$[[hor]X, [hor]Y] = [hor]([X, Y]) + \Omega(X, Y).$$

**Theorem 4.2.** Given an extension  $\mathbf{e} \colon 0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$  of (R,A)-Lie algebras together with an  $\mathbf{e}$ -connection  $\omega \colon L'' \to L$  and with  $\mathbf{e}$ -curvature  $\Omega \colon L'' \otimes_A L'' \to L'$  of  $\omega$ , the tautological Poisson structure

$$\{\cdot,\cdot\}\colon S_A[L]\otimes S_A[L]\longrightarrow S_A[L]$$

on  $S_A[L]$  (defined in (3.9), (3.10), (3.11)) is determined by the following identities:

$$\{\alpha', \beta'\} = [\alpha', \beta']_{L'}, \quad \alpha', \beta' \in L',$$

(4.9) 
$$\{\alpha'', \beta''\} = [\alpha'', \beta'']_{L''} + \Omega(\alpha'', \beta''), \quad \alpha'', \beta'' \in L'',$$

$$\{\alpha', \beta''\} = [\alpha', \beta''], \quad \alpha' \in L', \beta'' \in L'',$$

$$\{\alpha'', \beta'\} = [\alpha'', \beta'], \quad \alpha'' \in L'', \beta' \in L',$$

$$\{\alpha'', a\} = \alpha''(a) \in A, \quad a \in A, \alpha'' \in L'',$$

$$\{\alpha', a\} = 0 \in A, \quad a \in A, \ \alpha' \in L',$$

$$\{u, vw\} = \{u, v\}w + v\{u, w\}, \quad u, v, w \in \mathcal{S}_A[L].$$

*Proof.* This is an immediate consequence of the identities (3.9)–(3.11) and (4.3).

We note that  $\{a, b\} = 0$  for every  $a, b \in A$ .

Corollary 4.3. Under the hypotheses of Theorem 4.2, the extension e of (R, A)-Lie algebras can be reconstructed from the Poisson structure on  $S_A[L]$  in the sense that the extension is determined by the relations (4.8)–(4.11).

*Proof.* This is an immediate consequence of Theorem 4.2 and Proposition 3.5.  $\Box$ 

In particular, when we specialize to the case where the ground ring R is the field  $\mathbb{R}$  of real numbers, under the circumstances of Example 4.1, the tautological Poisson structure is determined by the identities given in Theorem 4.2. Thus this description of the tautological Poisson bracket (4.2) in Theorem 4.2 completely recovers the Poisson structure on  $(T^*Q)/G$ .

#### 5. Description of the Poisson structure in terms of differentials

Our ultimate goal is to reconcile our description of the Poisson structure on  $(T^*Q)/G$  that results from Theorem 4.2 with that given in Theorem IV.1 of [44]. To this end, we will now develop a description, in terms of differentials, of the Poisson bracket given in Theorem 4.2. As before, A denotes a commutative unital algebra over a general commutative unital ground ring R.

5.1. Algebraic generalities. We will momentarily abstract from the particular situation spelled out in Theorem 4.2 and consider a finitely generated projective A-module M together with an A-module decomposition  $M = M' \oplus M''$  into a direct sum of two projective A-modules M' and M''. We will simplify the exposition somewhat and use the notation

$$S := S_A[M], \quad S' := S_A[M'], \quad S'' := S_A[M''].$$

The A-module decomposition of M induces a canonical decomposition  $S \cong S' \otimes_A S''$  as the tensor product of the commutative A-algebras S' and S'', and we will also identify S with the symmetric S''-algebra on  $S'' \otimes_A M'$  as well as with the symmetric S'-algebra on  $S' \otimes_A M''$ .

In view of (2.8),  $D_{\mathfrak{S}'|A} \cong \mathfrak{S}' \otimes_A M'$ ,  $D_{\mathfrak{S}''|A} \cong \mathfrak{S}' \otimes_A M''$ ,  $D_{\mathfrak{S}|A} \cong \mathfrak{S} \otimes_A M$ , and the S-module  $D_{\mathfrak{S}|A}$  of formal differentials of S over the algebra A decomposes as the direct sum

$$(5.1) D_{S|A} = (S \otimes_{S'} D_{S'|A}) \oplus (S \otimes_{S''} D_{S''|A}) \cong S \otimes_A M' \oplus S \otimes_A M''.$$


Hence the exact sequence (2.9) of S-modules takes the form

$$(5.2) 0 \longrightarrow S \otimes_A D_A \longrightarrow D_S \longrightarrow S \otimes_A M' \oplus S \otimes_A M'' \longrightarrow 0.$$

To make the basic constructions later in the paper more easily comprehensible for the reader, we will now somewhat refine the exact sequence (5.2):

In view of (2.8),  $D_{S'|A} \cong S' \otimes_A M'$  and  $D_{S''|A} \cong S'' \otimes_A M''$ , canonically. The inclusions  $R \subseteq A \subseteq S'$ ,  $R \subseteq A \subseteq S''$ ,  $R \subseteq S' \subseteq S$ , and  $R \subseteq S'' \subseteq S$  determine exact sequences of S-modules of the kind (2.9).

**Proposition 5.1.** These exact sequence fit into the following commutative diagram of S-modules having exact rows and columns:



*Proof.* These sequences are all of the kind (2.9). In view of the naturality of the constructions, commutativity of the diagram is therefore immediate. We leave the details to the reader.

5.2. The geometric situation. We will illustrate the diagram (5.3) in terms of vector bundles. Consider two vector bundles  $p_{E_1} : E_1 \to B$  and  $p_{E_2} : E_2 \to B$ . Form the Whitney sum

$$p_{E_1} \oplus p_{E_2} \colon E_1 \oplus E_2 \to B.$$

Denote by  $p_{E_1^*}: E_1^* \to B$ ,  $p_{E_2^*}: E_2^* \to B$ , and  $p_{E_1^*} \oplus p_{E_2^*}: E_1^* \oplus E_2^* \to B$  the corresponding dual bundles. When thought of as the total space of a vector bundle over  $E_1^*$  or as the total space of a vector bundle over  $E_2^*$ , instead of  $E_1^* \oplus E_2^*$  we shall write  $N := E_1^* \times_B E_2^*$  as in the following diagram:

$$N \xrightarrow{\tau_{E_1^*}^*} E_1^*$$

$$\downarrow^{\tau_{E_2^*}^*} \qquad \downarrow^{p_{E_1^*}}$$

$$E_2^* \xrightarrow{p_{E_2^*}} B$$

Below we will use all three vector bundle structures of N over B,  $E_1^*$ , and  $E_2^*$ .

**Proposition 5.2.** The following commutative diagram of vector bundles with base N has all rows and columns exact:

Proof. We note first that the diagram is symmetric in  $E_1$  and  $E_2$ ; thus the first row and the first column as well as the second row and the second column coincide when  $E_1$  and  $E_2$  are interchanged. Thus, it suffices to give the maps for the first two rows. Let  $\lambda_b^{E_1}, \mu_b^{E_1} \in E_1^*, \lambda_b^{E_2} \in E_2^*$ , so  $(\lambda_b^{E_1}, \lambda_b^{E_2}) \in N$ , and  $v_{\lambda_b^{E_1}} \in T_{\lambda_b^{E_1}} E_1^*, v_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \in T_{(\lambda_b^{E_1}, \lambda_b^{E_2})} N$ . The maps are:

$$\begin{split} N \times_B E_1{}^* &\ni \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), \mu_b^{E_1} \right) \longmapsto \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), \operatorname{Ver}_{\lambda_b^{E_1}} (\mu_b^{E_1}) \right) \in N \times_{E_1{}^*} \operatorname{T}(E_1{}^*) \\ N \times_{E_1{}^*} \operatorname{T}(E_1{}^*) &\ni \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), v_{\lambda_b^{E_1}} \right) \longmapsto \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), T_{\lambda_b^{E_1}} p_{E_1{}^*} \left( v_{\lambda_b^{E_1}} \right) \right) \in N \times_B \operatorname{T}B \\ N \times_B E_1{}^* &\ni \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), \mu_b^{E_1} \right) \longmapsto \operatorname{Ver}_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \left( \mu_b^{E_1} \right) \in \operatorname{T}N \\ \operatorname{T}N &\ni v_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \longmapsto \left( (\lambda_b^{E_1}, \lambda_b^{E_2}), \operatorname{T}_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \tau_{E_2{}^*}^* \left( v_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \right) \right) \in N \times_{E_2{}^*} \operatorname{T}(E_2{}^*), \end{split}$$

where

$$\begin{aligned} \operatorname{Ver}_{\lambda_b^{E_1}}(\mu_b^{E_1}) &= \left. \frac{d}{dt} \right|_{t=0} \left( \lambda_b^{E_1} + t \mu_b^{E_1} \right) \in \mathcal{T}_{\lambda_b^{E_1}}(E_1^*), \\ \operatorname{Ver}_{(\lambda_b^{E_1}, \lambda_b^{E_2})} \left( \mu_b^{E_1} \right) &= \left. \frac{d}{dt} \right|_{t=0} \left( \lambda_b^{E_1} + t \mu_b^{E_1}, \lambda_b^{E_2} \right) \in \mathcal{T}_{(\lambda_b^{E_1}, \lambda_b^{E_2})} N. \end{aligned}$$

The commutativity of the diagram and the exactness of the rows and columns is an easy verification.  $\Box$ 

Dualizing diagram (5.4) we obtain the commutative diagram

of vector bundles over N having exact rows and columns.

We next show that the diagram that we obtain when we take sections in (5.5) is the  $C^{\infty}$ -analog of (5.3). We will use the notation  $\Gamma$  for the smooth sections functor. Recall the familiar Serre-Swan equivalence: Given a smooth (paracompact) manifold M, the assignment to a smooth vector bundle  $\zeta$  on M of its  $C^{\infty}(M)$ -module  $\Gamma(\zeta)$  of smooth sections is an equivalence of categories between smooth vector bundles on M and finitely generated projective  $C^{\infty}(M)$ -modules.

**Lemma 5.3.** For vector bundles  $\mathscr{E}_1 \to M$  and  $\mathscr{E}_2 \to M$ , the fiber product  $\mathscr{E}_1 \times_M \mathscr{E}_2 \to \mathscr{E}_1$  being regarded as a vector bundle over  $\mathscr{E}_1$ , the Serre-Swan equivalence yields a canonical isomorphism

$$\Gamma_{\mathscr{E}_1}(\mathscr{E}_1 \times_M \mathscr{E}_2) \cong C^{\infty}(\mathscr{E}_1) \otimes_{C^{\infty}(M)} \Gamma_M(\mathscr{E}_2)$$

of  $C^{\infty}(\mathcal{E}_1)$ -modules.

For a proof, the reader may consult, e. g., [10] (2.26).

Applying the lemma to the exact sequence that we obtain by taking sections in (5.5), we get the following commutative diagram of  $C^{\infty}(N)$ -modules

with all rows and columns exact. This is the precise geometric analog (5.3), with  $(R, A) = (\mathbb{R}, C^{\infty}(B))$ ,  $M' = \Gamma_B(E_1)$ ,  $M'' = \Gamma_B(E_2)$ , the algebras  $C^{\infty}(E_1^*)$ ,  $C^{\infty}(E_2^*)$ , and  $C^{\infty}(N)$  being the  $C^{\infty}$ -completions of the respective rings S', S'' and S.

5.3. **Basic example.** We will study the diagram (5.5) in the context of Example 4.1. Let  $\xi \colon Q \to B$  be a right principal G-bundle and  $\widetilde{\xi} \colon \widetilde{Q} = Q \times_B \mathrm{T}^*B \to \mathrm{T}^*B$  its pull back relative to the cotangent bundle projection  $\tau_B^* \colon \mathrm{T}^*B \to B$ . Thus, in the commutative diagram

(5.7) 
$$\begin{split} \widetilde{Q} & \xrightarrow{\widetilde{\xi}} & \mathbf{T}^* B \\ \downarrow \xi^*(\tau_B^*) & \downarrow \tau_B^* \\ Q & \xrightarrow{\xi} & B, \end{split}$$

the vertical projection map  $\xi^*(\tau_B^*) \colon \widetilde{Q} \to Q$  is a vector bundle projection with fibers  $T_b^*B$   $(b \in B)$  and  $\widetilde{\xi} \colon \widetilde{Q} \to T^*B$  is a principal G-bundle.


Consider the tangent bundles  $\tau_Q \colon \mathrm{T}Q \to Q$  and  $\tau_B \colon \mathrm{T}B \to B$  of Q and B, respectively, as well as the induced vector bundle  $\xi^*(\tau_B) \colon Q \times_B \mathrm{T}B \to Q$  on Q, the pull back of  $\tau_B$  via  $\xi$ . Viewed as a map over Q, the G-equivariant surjective map  $\mathrm{T}Q \ni v_q \longmapsto (q, \mathrm{T}_q \xi(v_q)) \in Q \times_B \mathrm{T}B \ (q \in Q)$  is a morphism of vector bundles on Q, spelled out here for the total spaces.

5.3.1. The Sternberg space. The total space of the coadjoint bundle  $\operatorname{ad}^*(\widetilde{\xi}): N_{\operatorname{St}}: = \widetilde{Q} \times_G \mathfrak{g}^* \to \operatorname{T}^*B$  is occasionally referred to in the literature as the Sternberg space (associated to the data).

We will now apply the construction in Subsection 5.2 above: We take the vector bundles  $p_{E_1} : E_1 \to B$  and  $p_{E_2} : E_2 \to B$  to be the adjoint bundle  $\mathrm{ad}(\xi) : Q \times_G \mathfrak{g} \to B$  of  $\xi : Q \to B$  and the tangent bundle  $\tau_B : \mathrm{T}B \to B$  of B, respectively. Then  $N := (Q \times_G \mathfrak{g}^*) \oplus \mathrm{T}^*B$  is not only the total space of a vector bundle over B but the natural projections  $N \to Q \times_G \mathfrak{g}^*$  and  $N \to \mathrm{T}^*B$  also define vector bundles. In what follows we shall use all three vector bundle structures. We note that

$$(5.8) N\ni ([q,\mu],\alpha_{[q]})\longleftrightarrow [(\alpha_{[q]},q),\mu]\in N_{\mathrm{St}},\ q\in Q,\ \mu\in\mathfrak{g}^*,\ \alpha_{[q]}\in \mathrm{T}_{[q]}^*B,$$

yields a canonical isomorphism of vector bundles from  $N \to T^*B$  to  $N_{St} \to T^*B$ . In view of this isomorphism, in the situation under discussion, the diagrams (5.5) and (5.6) now specialize to (5.9)



The last diagram is the  $C^{\infty}$ -analog of diagram (5.3) in the present special case.

To be precise,  $(R, A) = (\mathbb{R}, C^{\infty}(B))$ , the algebra  $C^{\infty}(N_{\mathrm{St}})$  of ordinary smooth functions on  $N_{\mathrm{St}}$  is substituted for the algebra  $\mathcal{S}$ , as an A-module,  $M' = \Gamma_B(Q \times_G \mathfrak{g})$  is the space of sections of the adjoint bundle  $\mathrm{ad}(\xi) \colon Q \times_G \mathfrak{g} \to B$  on B, as an A-module,  $M'' = \Gamma_B(\mathrm{T}B) = \mathfrak{X}(B)$ , the smooth vector fields on B, the algebra  $\mathcal{S}''$  is the algebra of smooth functions on  $\mathrm{T}^*B$  that are polynomial on the fibers of the cotangent bundle  $\mathrm{T}^*B \to B$  of B. Since (5.7) is a pull back diagram, the obvious projection  $\widetilde{Q} \times_G \mathfrak{g} \to \mathrm{T}^*B$  is a vector bundle on  $\mathrm{T}^*B$ ; hence the  $\mathcal{S}''$ -module  $\mathcal{S}'' \otimes_A M'$  is naturally isomorphic to the space of sections of the adjoint bundle  $\mathrm{ad}(\widetilde{\xi}) \colon \widetilde{Q} \times_G \mathfrak{g} \to \mathrm{T}^*B$  of  $\widetilde{\xi}$  that are polynomial on the fibers of the cotangent bundle  $\mathrm{T}^*B \to B$  of B. The algebras  $C^{\infty}(Q \times_G \mathfrak{g}^*)$ ,  $C^{\infty}(\mathrm{T}^*B)$ , and  $C^{\infty}(N_{\mathrm{St}})$  are the  $C^{\infty}$ -completions of the respective rings  $\mathcal{S}'$ ,  $\mathcal{S}''$  and  $\mathcal{S}$ . Diagram (5.10) now arises from (5.9) by the operation of taking smooth sections of the corresponding vector bundles on  $N_{\mathrm{St}}$ .

5.3.2. The Weinstein space. The smooth manifold  $N_W := (T^*Q)/G$  is occasionally referred to in the literature as the Weinstein space; it inherits a Poisson structure from the ordinary symplectic Poisson structure of  $T^*Q$ . The projection  $(T^*Q)/G \ni [\alpha_q] \mapsto [q] \in Q/G = B \ (q \in Q)$  turns  $N_W$  into a vector bundle over B.

Viewed as a map over Q, the obvious G-equivariant surjective map  $TQ \longrightarrow Q \times_B TB$  is a morphism of vector bundles on Q, spelled out here for the total spaces. A principal connection for  $\xi$  is precisely a G-equivariant vector bundle section for this surjection. Thus, pick a principal connection for  $\xi$ . Taking duals in the fiber directions, we then obtain a G-equivariant map

$$T^*Q \longrightarrow Q \times_B T^*B = \widetilde{Q},$$

indeed a morphism of G-vector bundles on Q, spelled out here for the total spaces. This map combines with the G-momentum mapping  $T^*Q \to \mathfrak{g}^*$  to yield a G-equivariant diffeomorphism

$$(5.11) T^*Q \longrightarrow \widetilde{Q} \times \mathfrak{g}^*;$$

in particular, the G-equivariant projection  $T^*Q \to \widetilde{Q}$  to the first component is now seen to be the projection of a G-vector bundle on  $\widetilde{Q}$  which, as an ordinary vector bundle on  $\widetilde{Q}$ , is trivial (not equivariantly trivial). Thus the diffeomorphism (5.11), in turn, descends to a diffeomorphism

$$(5.12) N_W = (T^*Q)/G \longrightarrow \widetilde{Q} \times_G \mathfrak{g}^* = N_{St}.$$

This identifies  $N_{\rm W}$  with  ${\rm T}^*(Q/G)\oplus (Q\times_G\mathfrak{g}^*)$ , and the latter space acquires, as we have seen in (5.8), a vector bundle structure over each of the three base spaces  $B, Q\times_G\mathfrak{g}^*$ , and  ${\rm T}^*B$  in a natural way. One of the main goals of this paper, carried out in Subsection 5.7, is to express the natural quotient Poisson bracket on  $({\rm T}^*Q)/G$  via this vector bundle isomorphism over B on the space  ${\rm T}^*(Q/G)\oplus (Q\times_G\mathfrak{g}^*)\cong N_{\rm St}$ . In order to do this, we step back and consider the general situation presented in Subsection 5.1.

5.4. Linear connections. Consider a vector bundle  $\lambda \colon \mathcal{L} \to B$ ; let  $V = \mathcal{L} \times_B \mathcal{L} \to \mathcal{L}$  denote its vertical subbundle. Recall that a *linear* or *Ehresmann* connection on  $\lambda$  is given by a section  $\mathcal{L} \times_B TB \to T\mathcal{L}$  for the associated extension

$$(5.13) 0 \longrightarrow V \longrightarrow T\mathcal{L} \longrightarrow \mathcal{L} \times_B TB \longrightarrow 0$$

of vector bundles on  $\mathcal{L}$  or, equivalenty, by a retraction  $T^*\mathcal{L} \to \mathcal{L} \times_B T^*B$  for the extension

$$0 \longleftarrow V^* \longleftarrow T^* \mathcal{L} \longleftarrow \mathcal{L} \times_B T^* B \longleftarrow 0$$

of the dual vector bundles on  $\mathcal{L}$ .

In the algebraic setting, accordingly, let V be a projective A-module, and let  $S_A[V]$  be the symmetric A-algebra on V. We define a linear connection on the A-module V to be a retraction

$$(5.14) r: D_{\mathcal{S}_A[V]} \longrightarrow \mathcal{S}_A[V] \otimes_A D_A$$

for the extension

$$(5.15) 0 \longrightarrow S_A[V] \otimes_A D_A \longrightarrow D_{S_A[V]} \longrightarrow S_A[V] \otimes_A V \longrightarrow 0$$

in the category of  $S_A[V]$ -modules. Using the canonical isomorphism

$$\operatorname{Hom}_{\mathcal{S}_A[V]}(\mathcal{S}_A[V] \otimes_A V, \mathcal{S}_A[V]) \longrightarrow \operatorname{Der}(\mathcal{S}_A[V]|A)$$

of  $S_A[V]$ -modules, cf. (2.6), we see that a linear connection on V determines a section

$$(5.16) s: S_A[V] \otimes_A \operatorname{Der}(A) \longrightarrow \operatorname{Der}(S_A[V])$$

for the dual extension

$$(5.17) 0 \longrightarrow \operatorname{Der}(S_A[V]|A) \longrightarrow \operatorname{Der}(S_A[V]) \longrightarrow S_A[V] \otimes_A \operatorname{Der}(A) \longrightarrow 0$$

of  $S_A[V]$ -modules (but, beware, the retraction r is not determined by the section s unless  $D_A$  is reflexive, i.e., unless the canonical A-module morphism from  $D_A$  to its double A-dual  $D_A^{**}$  is an isomorphism). A linear connection, i. e. retraction r of the kind (5.14), for V induces the direct sum decomposition

in the category of  $S_A[V]$ -modules, and the projection to  $S_A[V] \otimes_A V$  restricts to an isomorphism

$$\ker(r) \longrightarrow S_A[V] \otimes_A V$$

of  $S_A[V]$ -modules. We will use the notation

$$(D_{S_A[V]})_{\text{vert}} = \ker(r), \quad (D_{S_A[V]})_{\text{hor}} = S_A[V] \otimes_A D_A,$$

and we will refer to  $(D_{S_A[V]})_{\text{vert}}$  and  $(D_{S_A[V]})_{\text{hor}}$  as the *vertical* and *horizontal* components, respectively, of the resulting decomposition (5.18) of  $D_{S_A[V]}$ , so that this decomposition takes the form

Since V is supposed to be projective as an A-module, a linear connection on V exists.

We now return to the situation of Subsection 5.1 above. Thus M' and M'' are projective A-modules,  $M = M' \oplus M''$ ,  $S = S_A[M]$ ,  $S' = S_A[M']$ ,  $S'' = S_A[M'']$ . Let r be a linear connection, i. e. retraction

 $r: D_{S'} \longrightarrow S' \otimes_A D_A$  of the kind (5.14), for M'; such a retraction exists since M' is projective as an A-module. Let

be the corresponding direct sum decomposition (5.18) in the category of S'-modules. Then

$$S \otimes_{S'} r : S \otimes_{S'} D_{S'} \longrightarrow S \otimes_A D_A$$

is a retraction for the upper horizontal exact sequence

$$(5.21) 0 \longrightarrow S \otimes_A D_A \longrightarrow S \otimes_{S'} D_{S'} \longrightarrow S \otimes_A M' \longrightarrow 0$$

in (5.3), and

$$\ker(S \otimes_{S'} r) = S \otimes_{S'} \ker(r).$$

Moreover, since the projection from  $D_{S'}$  to  $S' \otimes_A M'$  restricts to an isomorphism  $\ker(r) \longrightarrow S' \otimes_A M'$  of S'-modules, the projection from  $S \otimes_{S'} D_{S'}$  to  $S \otimes_A M'$  restricts to an isomorphism

$$(5.22) S \otimes_{S'} \ker(r) \longrightarrow S \otimes_A M'$$

of S-modules. Hence, when we view  $S \otimes_{S'} \ker(r)$  as an S-submodule of  $D_S$  via the injection  $S \otimes_{S'} D_{S'} \to D_S$  (arising as an arrow of the middle row of (5.3)), the isomorphism (5.22) is also the restriction of the projection from  $D_S$  to  $S \otimes_A M'$  (arising as an arrow of the middle column of (5.3)). Consequently the chosen linear connection r for M' induces corresponding direct sum decompositions of  $S \otimes_{S'} D_{S'}$  and  $D_S$  into the following vertical and horizontal components:

$$(S \otimes_{S'} D_{S'})_{\text{vert}} = (D_S)_{\text{vert}} = S \otimes_{S'} \ker(r),$$
  
$$(S \otimes_{S'} D_{S'})_{\text{hor}} = S \otimes_A D_A, (D_S)_{\text{hor}} = S \otimes_{S''} D_{S''},$$

and we will refer to  $(S \otimes_{S'} D_{S'})_{vert}$  as the *vertical* component of  $(S \otimes_{S'} D_{S'})$ , to  $(S \otimes_{S'} D_{S'})_{hor}$  as its *horizontal* component, to  $(D_S)_{vert}$  as the *vertical* component of  $D_S$ , and to  $(D_S)_{hor}$  as its *horizontal* component. Thus the chosen linear connection r for M' induces the direct sum decompositions

$$(5.23) S \otimes_{S'} D_{S'} = (S \otimes_{S'} D_{S'})_{hor} \oplus (S \otimes_{S'} D_{S'})_{vert}$$

Further, the projection from  $S \otimes_{S'} D_{S'}$  to  $S \otimes_A M'$  (in the upper horizontal row of (5.3)) restricts to an isomorphism  $(S \otimes_{S'} D_{S'})_{\text{vert}} \to S \otimes_A M'$  of S-modules and the projection from  $D_S$  to  $S \otimes_A M'$  (in the middle horizontal row of (5.3)) restricts to an isomorphism  $(D_S)_{\text{vert}} \to S \otimes_A M'$  of S-modules; these isomorphisms are precisely the isomorphism (5.22) of S-modules. Thus, all told, somewhat more formally, the resulting retraction

$$D_{\mathcal{S}} \longrightarrow (D_{\mathcal{S}})_{hor} = \mathcal{S} \otimes_{\mathcal{S}''} D_{\mathcal{S}''}$$

for the middle horizontal exact sequence

$$(5.25) 0 \longrightarrow S \otimes_{S''} D_{S''} \longrightarrow D_S \longrightarrow S \otimes_A M' \longrightarrow 0$$

in (5.3) is a linear connection on the induced S"-module S"  $\otimes_A M'$ , and the notation is consistent.

By construction, in view of the direct sum decomposition  $M = M' \oplus M''$ , any  $\alpha' \in M'$  is a "vertical" element of S in an obvious sense; however the differential  $d\alpha'$  refers to the differential of  $\alpha'$  viewed as a member of S, taken as an R-algebra, i. e. over the ground ring R (rather than as an A-algebra). Hence, in general, given  $\alpha' \in M'$ , the differential  $d\alpha' \in D_S$  has, beyond a vertical component  $(d\alpha')_{\text{vert}} \in (D_S)_{\text{vert}}$ , a horizontal component  $(d\alpha')_{\text{hor}} \in (D_S)_{\text{hor}}$ ; now,  $d\alpha'$  is actually the image in  $D_S$  of the differential of  $\alpha'$  in  $D_{S'}$  and, under the projection  $D_S \longrightarrow S \otimes_A M' \oplus S \otimes_A M''$  in (5.2), the horizontal component  $(d\alpha')_{\text{hor}}$  goes to zero whence  $(d\alpha')_{\text{hor}}$  is a member of  $S \otimes_A D_A$ .

**Remark 5.4.** At the present stage, the theory is symmetric in M' and M'', that is, we could have chosen a linear connection for M'' as well and developed the same kind of decomposition of  $D_{\mathcal{S}}$  as (5.24).

**Example 5.5.** Consider the trivial principal bundle  $Q \to B$  where  $Q = B \times G$ , endowed with the trivial connection so that, with the notation established earlier,  $N = (T^*B) \times \mathfrak{g}^*$ . Let f be a function on  $T^*B$ , and let  $X \in \mathfrak{g}$ , viewed as a linear function on  $\mathfrak{g}^*$ ; view X, f and fX as functions on N in the obvious way. The ordinary differential  $\mathbf{d}(fX)$  of the function fX decomposes as

$$\mathbf{d}(fX) = (\mathbf{d}f)X + f\mathbf{d}X, \quad (\mathbf{d}(fX))_{\text{vert}} = f\mathbf{d}X, \quad (\mathbf{d}(fX))_{\text{hor}} = (\mathbf{d}f)X.$$

Here the notation  $\mathbf{d}f$  refers to the ordinary differential of f, viewed as a function on  $N=(\mathrm{T}^*B)\times\mathfrak{g}^*$  that depends only on the first variable, and  $\mathbf{d}X$  refers to the ordinary differential of X, viewed as a function on  $N=(\mathrm{T}^*B)\times\mathfrak{g}^*$  that depends only on the second variable.

5.5. The Poisson structure. We return to the situation of Section 4 above. Thus

$$e: 0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$$

is an extension of (R, A)-Lie algebras. Choose an e-connection for the extension  $\mathbf{e}$ , i.e., an A-module section  $\omega \colon L'' \to L$ . The connection  $\omega$  determines (and is determined by) an A-module decomposition  $L = L' \oplus L''$ ; here we identify L'' with its image  $\omega(L'')$  in L, with a slight abuse of notation. We also choose a linear connection  $r \colon D_{S'} \to S' \otimes D_A$  on the A-module L' (see (5.14)) and hence, by (5.24),  $D_S = (D_S)_{\text{hor}} \oplus (D_S)_{\text{vert}}$ , as S-modules, where

$$(D_{\mathcal{S}})_{\text{vert}} = \mathcal{S} \otimes_{\mathcal{S}'} \ker(r) \cong \mathcal{S} \otimes_A L', \qquad (D_{\mathcal{S}})_{\text{hor}} : = \mathcal{S} \otimes_{\mathcal{S}''} D_{\mathcal{S}''}.$$

We will now apply the material developed before, with M, M', M'' the A-modules that underlie, respectively, L, L', and L'' and, to simplify the notation, we will denote these underlying A-modules by L, L', and L'' as well.

To develop the promised description, in terms of differentials, of the Poisson bracket given in Theorem 4.2, all we have to do now is to evaluate the formulas (3.19) and (3.20) in terms of the direct sum decomposition (5.24). The algebras  $S' = S_A[L']$  and  $S'' = S_A[L'']$  carry the tautological Poisson structures that correspond to the A-Lie algebra structure of L' and to the Lie-Rinehart structure of (A, L''), respectively, and the corresponding Poisson 2-form (3.13) is defined for each of these algebras. We denote these 2-forms by

$$\pi' : D_{S'|A} \otimes_{S'} D_{S'|A} \longrightarrow S', \ \pi'' : D_{S''} \otimes_{S''} D_{S''} \longrightarrow S''$$

and, relative to the decomposition  $D_{\mathcal{S}} = (D_{\mathcal{S}})_{hor} \oplus (D_{\mathcal{S}})_{vert}$  (cf. (5.24)), with a slight abuse of notation, we will likewise denote the S-bilinear extensions to the corresponding S-valued forms by

$$\pi': (D_{\mathcal{S}})_{\text{vert}} \otimes_{\mathcal{S}} (D_{\mathcal{S}})_{\text{vert}} \longrightarrow \mathcal{S}, \ \pi'': (D_{\mathcal{S}})_{\text{hor}} \otimes_{\mathcal{S}} (D_{\mathcal{S}})_{\text{hor}} \longrightarrow \mathcal{S}$$

as well. Furthermore, as before, let  $\Omega: L'' \otimes_A L'' \to L'$  denote the **e**-curvature of the **e**-connection  $\omega$  (see (4.2)). For  $\alpha'', \beta'' \in L''$ , let  $d\alpha'', d\beta'' \in D_{8''}$  be their formal differentials over the ground ring R, and denote by

$$[d\alpha''], [d\beta''] \in D_{S'' \mid A} \cong S'' \otimes_A L''$$

the associated formal differentials over A, that is, the images of  $d\alpha''$  and  $d\beta''$  under the projection  $D_{S''} \to D_{S''|A}$  in (2.9), with the notation adjusted suitably. We will write

(5.26) 
$$\Omega^{\sharp}([d\alpha''],[d\beta'']) = \Omega(\alpha'',\beta''),$$

and we extend  $\Omega^{\sharp}$  to an S-bilinear alternating 2-form

$$\Omega^{\sharp} \colon \left( \mathbb{S} \otimes_{\mathbb{S}''} \mathcal{D}_{\mathbb{S}''|A} \right) \otimes_{\mathbb{S}} \left( \mathbb{S} \otimes_{\mathbb{S}''} \mathcal{D}_{\mathbb{S}''|A} \right) \longrightarrow \mathbb{S}$$

in the obvious manner. We then write the S-bilinear alternating 2-form on  $(D_S)_{hor} = S \otimes_{S''} D_{S''}$  induced via the projection

$$(D_{\mathcal{S}})_{hor} = \mathcal{S} \otimes_{\mathcal{S}''} D_{\mathcal{S}''} \longrightarrow \mathcal{S} \otimes_{\mathcal{S}''} D_{\mathcal{S}''|A} \cong \mathcal{S} \otimes_A L''$$

as

$$\widetilde{\Omega} \colon (D_{\mathcal{S}})_{\mathrm{hor}} \otimes_{\mathcal{S}} (D_{\mathcal{S}})_{\mathrm{hor}} \longrightarrow \mathcal{S}.$$

**Theorem 5.6.** Given  $f, h \in S = S_A[L]$ , the Poisson bracket  $\{f, h\} \in S$  is given by the expression

(5.27) 
$$\{f, h\} = \pi'((df)_{\text{vert}}, (dh)_{\text{vert}}) + \pi''((df)_{\text{hor}}, (dh)_{\text{hor}}) + \widetilde{\Omega}((df)_{\text{hor}}, (dh)_{\text{hor}}) + \pi((df)_{\text{vert}}, (dh)_{\text{hor}}) + \pi((df)_{\text{hor}}, (dh)_{\text{vert}}).$$

Furthermore, the Poisson 2-form  $\pi$  on  $D_S$ , restricted to  $(D_S)_{hor} = S \otimes_{S''} D_{S''}$ , is given by the sum  $\pi'' + \Omega$  and, restricted to  $(D_S)_{vert}$ , by  $\pi'$ .

*Proof.* Let  $f, h \in S = S_A[L]$ . We will show that

(5.28) 
$$\pi((df)_{\text{vert}}, (dh)_{\text{vert}}) = \pi'((df)_{\text{vert}}, (dh)_{\text{vert}})$$

and

(5.29) 
$$\pi((df)_{\text{hor}}, (dh)_{\text{hor}}) = \pi''((df)_{\text{hor}}, (dh)_{\text{hor}}) + \widetilde{\Omega}((df)_{\text{hor}}, (dh)_{\text{hor}}).$$

Since

$$\{f, h\} = \pi(df, dh) = \pi((df)_{\text{hor}} + (df)_{\text{vert}}, (dh)_{\text{hor}} + (dh)_{\text{vert}})$$

$$= \pi((df)_{\text{vert}}, (dh)_{\text{vert}}) + \pi((df)_{\text{hor}}, (dh)_{\text{hor}}) + \pi((df)_{\text{vert}}, (dh)_{\text{hor}}) + \pi((df)_{\text{hor}}, (dh)_{\text{vert}}),$$

(5.28) and (5.29) imply (5.27).

As an R-algebra, S is generated by A, L', and L''. It suffices to establish the identities (5.28) and (5.29) (and hence the identity (5.27)) when f and h range over  $a \in A$ ,  $\alpha' \in L'$ , and  $\alpha'' \in L''$ . By construction,

$$\begin{cases}
da \in \mathbb{S} \otimes_{A} \mathcal{D}_{A} \subseteq (\mathcal{D}_{\mathbb{S}})_{\text{hor}} \\
d\alpha'' \in \mathbb{S} \otimes_{\mathbb{S}''} \mathcal{D}_{\mathbb{S}''} = (\mathcal{D}_{\mathbb{S}})_{\text{hor}} \\
d\alpha' \in \mathbb{S} \otimes_{\mathbb{S}'} \mathcal{D}_{\mathbb{S}'} \text{ and hence } d\alpha' = (d\alpha')_{\text{vert}} + (d\alpha')_{\text{hor}}, \text{ where} \\
(d\alpha')_{\text{vert}} \in (\mathbb{S} \otimes_{\mathbb{S}'} \mathcal{D}_{\mathbb{S}'})_{\text{vert}} = (\mathcal{D}_{\mathbb{S}})_{\text{vert}} \\
(d\alpha')_{\text{hor}} = r(d\alpha') \in \mathbb{S}' \otimes_{A} \mathcal{D}_{A} \subset \mathbb{S} \otimes_{A} \mathcal{D}_{A} \subset \mathbb{S} \otimes_{\mathbb{S}''} \mathcal{D}_{\mathbb{S}''} = (\mathcal{D}_{\mathbb{S}})_{\text{hor}}.
\end{cases}$$

To prove (5.28) and (5.29), we need to check them for the pairs (a,b),  $(a,\alpha')$ ,  $(a,\alpha'')$ ,  $(\alpha',\beta')$ ,  $(\alpha',\alpha'')$ ,  $(\alpha'',\beta'')$ , where  $a,b\in A,\alpha',\beta'\in L'$ ,  $\alpha'',\beta''\in L''$ .

In view of (5.30), the only non-zero pair on which either side of (5.28) does not vanish is  $(\alpha', \beta')$ . In view of (3.9), by the definition of the tautological Poisson structure on L', the right-hand side of (5.28) equals  $[\alpha', \beta']_{L'}$  whereas, by (4.8), the left-hand side has the same value.

Next, we prove (5.29) by verifying it on each pair listed above. On (a,b) both sides vanish. Indeed, in view of the definition of the tautological Poisson bracket,  $\pi(da,db)=\pi'(da,db)$  is zero. In addition,  $\widetilde{\Omega}(da,db)=0$  since, by construction, the 2-form  $\widetilde{\Omega}$  factors through the 2-form  $\Omega^{\sharp}$ , and  $\Omega^{\sharp}$  vanishes on generators of the form da.

On the pair  $(a, \alpha')$  we proceed in the following way. Since  $(d\alpha')_{\text{hor}} \in \mathcal{S} \otimes_A D_A$  we need to check equality for a pair (sda, db), where  $s \in \mathcal{S}$ , and  $a, b \in A$ . By construction,  $\pi(sda, db) = s\{a, b\} = 0$  and, similarly,  $\pi''(sda, db) = s\{a, b\} = 0$ . The term  $\widetilde{\Omega}(sda, db) = s\Omega^{\sharp}(da, db)$  is zero as before.

On the pair  $(a, \alpha'')$ , by (4.12), the left-hand side is  $\pi(da, d\alpha'') = \{a, \alpha''\} = \alpha''(a)$ . By (3.10), the right-hand side equals  $\pi''(da, d\alpha'') + \widetilde{\Omega}(da, d\alpha'') = \pi''(da, d\alpha'') = \alpha''(a)$ .

To compute both sides of (5.29) on the pair  $(\alpha', \beta')$ , in view of (5.30), we just need to evaluate them on elements of the form  $\alpha' = s_1 da_1$  and  $\beta' = s_2 da_2$  for  $s_1, s_2 \in \mathcal{S}$  and  $a_1, a_2 \in A$ . Both sides of (5.29) vanish.

To compute both sides of (5.29) on the pair  $(\alpha', \alpha'')$ , in view of (5.30), we just need to evaluate them on elements of the form sda and  $d\alpha''$  for  $s \in \mathcal{S}$ ,  $a \in \mathcal{S}$ , and  $\alpha'' \in L''$ . The left-hand side equals  $\pi(sda, d\alpha'') = s\alpha''(a)$  by (4.12). By (3.10), the right-hand side equals

$$\pi''(sda, d\alpha'') + \widetilde{\Omega}(sda, d\alpha'') = s\pi''(da, d\alpha'') = s\alpha''(a).$$

On the pair  $(\alpha'', \beta'')$ , by (4.9), the left-hand side equals

$$\pi(d\alpha'',d\beta'')=\{\alpha'',\beta''\}=[\alpha'',\beta'']_{L''}+\Omega(\alpha'',\beta'').$$

By the definition of the tautological Poisson bracket on L'' (see (3.9) and (5.26)), the right-hand side equals

$$\pi''(d\alpha'',d\beta'') + \widetilde{\Omega}(d\alpha'',d\beta'') = \pi''(d\alpha'',d\beta'') + \Omega^{\sharp}([d\alpha''],[d\beta'']) = [\alpha'',\beta'']_{L''} + \Omega(\alpha'',\beta'').$$

Since the differentials of two members  $\alpha''$  and  $\beta''$  of L'', viewed as elements of S, have zero vertical components, the Poisson 2-form  $\pi$  on  $D_S$ , restricted to  $(D_S)_{hor} = S \otimes_{S''} D_{S''}$ , is obviously given by the sum  $\pi'' + \widetilde{\Omega}$ . The remaining 'furthermore' statement is a consequence of Lemma 5.7 below.

**Lemma 5.7.** Given  $\alpha', \beta' \in L'$ , the value  $\pi((d\alpha')_{\text{vert}}, (d\beta')_{\text{hor}})$  is zero.

*Proof.* Recall from (5.30) that  $(d\beta')_{hor} \in S \otimes_A D_A$  and

$$(d\alpha')_{\text{vert}} \in (D_{\mathcal{S}})_{\text{vert}} = (\mathcal{S} \otimes_{\mathcal{S}'} D_{\mathcal{S}'})_{\text{vert}} \subset \mathcal{S} \otimes_{\mathcal{S}'} D_{\mathcal{S}'}.$$

Therefore, we can assume that  $(d\alpha')_{\text{vert}} = sd\beta''$  or  $(d\alpha')_{\text{vert}} = sda$ , for some  $a \in A$ ,  $\beta'' \in L'$ , and  $s \in S$ . Similarly, we can assume that  $(d\beta')_{\text{hor}} = \bar{s}db$  for some  $b \in A$ ,  $\bar{s} \in S$ . Therefore  $\pi(sd\beta'', \bar{s}db) = s\bar{s}\{\beta'', b\}$  vanishes, by (4.13). Likewise,  $\pi(sda, \bar{s}db) = s\bar{s}\{a, b\} = 0$ .

5.6. The special case L'' = Der(A). We will show that, in this case, the e-connection  $\omega$  determines a unique linear connection  $r_{\omega}$  for L' and that, furthermore, with this linear connection, the terms  $\pi((df)_{\text{vert}}, (dh)_{\text{hor}})$  and  $\pi((df)_{\text{hor}}, (dh)_{\text{vert}})$  in formula (5.27) vanish. This amounts to showing that, given  $\alpha' \in L'$  and  $\alpha'' \in L''$ , the Poisson bracket

$$\{\alpha'', \alpha'\} = [\alpha'', \alpha'],$$

cf. (4.11), is completely absorbed in the term  $\pi''(d\alpha'', (d\alpha')_{hor})$ , that is, the component  $\pi(d\alpha'', (d\alpha')_{vert})$  is zero. In the case at hand, this observation greatly simplifies the formula (5.27).

For illustration we note that, in the circumstances of Example 5.5, given a function f on B, a vector  $X \in \mathfrak{g}$ , and  $Y \in \mathfrak{X}(B)$ , when fX and Y are viewed as functions on N in the obvious manner as in Example 5.5,

$${Y, fX} = Y(f)X = \pi''(dY, (d(fX))_{hor}).$$

Thus the Poisson bracket  $\{Y, fX\}$  is manifestly absorbed in the term  $\pi''(dY, (d(fX))_{hor})$ . The reasoning below involving the linear connection induced from the chosen e-connection somewhat reduces the argument to what corresponds, at the infinitesimal level, algebraically to that special case.

We return to the situation of Section 4 above. The extension of (R, A)-Lie algebras under discussion now has the form

$$\mathbf{e} \colon 0 \longrightarrow L' \longrightarrow L \longrightarrow \mathrm{Der}(A) \longrightarrow 0$$

but we continue to write  $L'' = \operatorname{Der}(A)$ . The sequence of inclusions  $R \subseteq A \subseteq S'$  determines the exact sequence

$$(5.32) 0 \longrightarrow S' \otimes_A D_A \longrightarrow D_{S'} \longrightarrow S' \otimes_A L' \longrightarrow 0;$$

this is a particular case of the exact sequence (2.9). Taking S'-duals, we obtain the exact sequence

$$(5.33) 0 \longrightarrow \operatorname{Der}(\mathcal{S}'|A) \longrightarrow \operatorname{Der}(\mathcal{S}') \longrightarrow \mathcal{S}' \otimes_A L'' \longrightarrow 0$$

with  $L'' = \text{Der}(A) \cong \text{Hom}_A(D_A, A)$ ; the sequence (5.33) is a particular case of (2.10). Notice that, when we induce up the sequence (5.32) to one of S-modules (by taking the tensor product with S over S'), we obtain the upper horizontal exact sequence of S-modules in the commutative diagram (5.3). Taking S'-duals once more, we arrive at a commutative diagram

the unlabeled vertical maps being the canonical maps to the double duals, the double A-dual of  $D_A$  being written as  $D_A^{**}$  and, likewise, the double S'-dual of  $D_{S'}$  being written as  $D_{S'}^{**}$ .

**Proposition 5.8.** An **e**-connection, i.e., an A-module section  $\omega$ :  $Der(A) \to L$  for the extension (5.31) of (R, A)-Lie algebras, gives rise to a linear connection  $r_{\omega} \colon D_{S'} \to S' \otimes_A D_A$  on the A-module L' (equivalently: S'-module retraction for (5.32)) in such a way that the S'-module section

$$(5.35) s_{\omega} \colon \mathcal{S}' \otimes_{A} L'' \longrightarrow \mathrm{Der}(\mathcal{S}')$$

for (5.33) dual to  $r_{\omega}$  is uniquely determined by  $\omega$ .

*Proof.* Let  $\omega: L'' \to L$  be an **e**-connection, i.e., A-module section for the extension **e** of (R, A)-Lie algebras. Given  $\alpha'' \in L''$ , consider the association

$$L' \longrightarrow L', \ (\alpha'', \alpha') \longmapsto [\omega(\alpha''), \alpha'] \in L'$$

where the notation  $[\omega(\alpha''), \alpha']$  refers to the commutator in L. This defines an R-linear map

$$(5.36) L'' \longrightarrow \operatorname{End}(L').$$

Since L' acts trivially on A, given  $\alpha'' \in L''$ , the formulas

$$d_{\alpha''}(\alpha') = [\omega(\alpha''), \alpha'], \ d_{\alpha''}(a) = \alpha''(a), \ \alpha' \in L', \ a \in A,$$

yield an R-derivation  $d_{\alpha''}: \mathcal{S}' \to \mathcal{S}'$ , the assignment to  $\alpha'' \in L''$  of  $d_{\alpha''}$  yields an A-linear map from L'' to  $Der(\mathcal{S}')$ , and the extension

$$(5.37) s_{\omega} \colon S' \otimes_A L'' \longrightarrow \operatorname{Der}(S')$$

thereof to an S'-linear map is an S'-module section for (5.33), plainly uniquely determined by  $\omega$ . With the notation

$$(5.38) (\operatorname{Der}(S'))_{\operatorname{vert}} = \operatorname{Der}(S'|A), (\operatorname{Der}(S'))_{\operatorname{hor}} = s_{\omega}(S' \otimes_{A} L''),$$

we obtain an S'-module direct sum decomposition

$$\operatorname{Der}(S') = (\operatorname{Der}(S'))_{\operatorname{vert}} \oplus (\operatorname{Der}(S'))_{\operatorname{hor}}$$

of Der(S') into a vertical component  $(Der(S'))_{vert}$  and a horizontal component  $(Der(S'))_{hor}$ . The dual of (5.37) is a retraction

$$\widetilde{r}_{\omega} \colon \mathcal{D}_{\mathcal{S}'}^{**} \longrightarrow \mathcal{S}' \otimes_{A} \mathcal{D}_{A}^{**}$$

for the bottom row extension in diagram (5.34). This retraction  $\widetilde{r}_{\omega}$  induces the direct sum decomposition  $D_{S'}^{**} = (S' \otimes_A D_A^{**}) \oplus \ker(\widetilde{r}_{\omega})$  in the category of S'-modules, and the restriction to  $\ker(\widetilde{r}_{\omega})$  of the projection to  $S' \otimes_A L'$  is an isomorphism of S'-modules. Lifting the resulting section  $S' \otimes_A L' \to D_{S'}^{**}$  to a section  $S' \otimes_A L' \to D_{S'}$  for (5.32), we obtain a direct sum decomposition

$$D_{\mathcal{S}'} \cong (\mathcal{S}' \otimes_A D_A) \oplus \ker(\widetilde{r}_\omega) \cong (\mathcal{S}' \otimes_A D_A) \oplus \mathcal{S}' \otimes_A L',$$

and the associated retraction to  $S' \otimes_A D_A$  is a retraction

$$r_{\omega} \colon \mathcal{D}_{\mathcal{S}'} \longrightarrow \mathcal{S}' \otimes_A \mathcal{D}_A$$

for the extension (5.32) in the category of S'-modules of the kind we are looking for.

We will refer to a connection of the kind  $r_{\omega}$  in Proposition 5.8 as a linear connection on L' associated to the **e**-connection  $\omega$ .

**Theorem 5.9.** Suppose that the quotient L'' in the extension  $\mathbf{e}$  of (R, A)-Lie algebras under discussion is the (R, A)-Lie algebra  $\mathrm{Der}(A)$  and that the requisite linear connection on L' is a connection  $r_{\omega}$  associated to the  $\mathbf{e}$ -connection  $\omega \colon L'' \to L$ . Then the terms  $\pi((df)_{\mathrm{vert}}, (dh)_{\mathrm{hor}})$  and  $\pi((df)_{\mathrm{hor}}, (dh)_{\mathrm{vert}})$  in the formula (5.27) vanish. Hence, given  $f, h \in \mathcal{S} = \mathcal{S}_A[L]$ ,

$$(5.39) {f,h} = \pi'((df)_{\text{vert}}, (dh)_{\text{vert}}) + \pi''((df)_{\text{hor}}, (dh)_{\text{hor}}) + \widetilde{\Omega}((df)_{\text{hor}}, (dh)_{\text{hor}}).$$

*Proof.* We return to the proof of Theorem 5.6. When the members f and h of S range over the members of A, L', and L'', the only cases where the value  $\pi((df)_{\text{vert}}, (dh)_{\text{hor}})$  does not obviously vanish are those where  $(f,h) = (\alpha',\beta')$  and  $(f,h) = (\alpha',\alpha'')$ , with  $\alpha',\beta' \in L'$  and  $\alpha'' \in L''$ .

In view of Lemma 5.7,  $\pi((d\alpha')_{\text{vert}}, (d\beta')_{\text{hor}}) = 0$ . Thus, it remains to consider the case  $(f, h) = (\alpha', \alpha'')$ , i.e., since  $d\alpha'' \in (D_S)_{\text{hor}}$ , we need to show that  $\pi((d\alpha')_{\text{vert}}, d\alpha'') = 0$ .

As before, we will now identify L'' with  $\omega(L'')$  so that, as A-modules,  $L = L' \oplus L''$ , and  $\omega$  is the canonical injection. In view of Proposition 5.8, the decomposition (5.20) now takes the form

Accordingly, with the notation  $\tilde{r}_{\omega}$  used in the proof of Proposition 5.8,

$$D_{S'}^{**} = (S' \otimes_A D_A^{**}) \oplus \ker(\widetilde{r}_{\omega}).$$

With a slight abuse of notation, given a formal differential  $\beta \in D_{S'}$ , we will denote its image in  $D_{S'}^{**}$ , i.e., the induced S'-linear map from Der(S') to S', by  $\beta \colon Der(S') \to S'$  as well.

By the definition of  $\widetilde{r}_{\omega}$ , the kernel  $\ker(\widetilde{r}_{\omega})$  of  $\widetilde{r}_{\omega}$  consists of the S'-linear maps  $\beta \colon \operatorname{Der}(S') \to S'$  such that the composition

$$(5.42) S' \otimes_A L'' \xrightarrow{s_{\omega}} \operatorname{Der}(S') \xrightarrow{\beta} S'$$

is zero. By construction, then,  $d_{\alpha''} = s_{\omega}(\alpha'') \in (\text{Der}(S'))_{\text{hor}}$ , in view of the decomposition (5.38), and  $(d\alpha')_{\text{vert}} : \text{Der}(S') \to S'$  has the following property:

$$(d\alpha')_{\text{vert}}(d\alpha'') = ((d\alpha')_{\text{vert}} \circ s_{\omega})(\alpha'') = 0.$$

Given  $\beta \in D_{S'}$  and  $\alpha'' \in L''$ , so that  $d_{\alpha''} \in Der(S')$ , we will show that

(5.43) 
$$\beta(d_{\alpha''}) = \pi(d\alpha'', \beta).$$

Indeed, as an R-module,  $D_{S'}$  is generated by the elements  $sd\alpha'$  and tda, where  $s, t \in S'$ ,  $\alpha' \in L'$ , and  $a \in A$ . Now

$$(sd\alpha')(d_{\alpha''}) = s(d\alpha'(d_{\alpha''})) = s(d_{\alpha''}(\alpha')) = s[\alpha'', \alpha'] = s\{\alpha'', \alpha'\} = s\pi(d\alpha'', d\alpha') = \pi(d\alpha'', sd\alpha')$$
$$(tda)(d_{\alpha''}) = t(da(d_{\alpha''})) = t(d_{\alpha''}(a)) = t\alpha''(a) = t\{\alpha'', a\} = t\pi(d\alpha'', da) = \pi(d\alpha'', tda)$$

which proves (5.43).

By construction, in view of (5.43),

$$\pi(d\alpha'',(d\alpha')_{\mathrm{hor}}) = (d\alpha')_{\mathrm{hor}}(d\alpha'') = ((d\alpha')_{\mathrm{vert}} + (d\alpha')_{\mathrm{hor}})(d\alpha'') = (d\alpha')(d\alpha'') = d\alpha''(\alpha') = [\alpha'',\alpha'].$$

On the other hand, by (4.11),

$$[\alpha'', \alpha'] = {\alpha'', \alpha'} = \pi(d\alpha'', d\alpha') = \pi(d\alpha'', (d\alpha')_{\text{hor}}) + \pi(d\alpha'', (d\alpha')_{\text{vert}}).$$

This shows that  $\pi(d\alpha'', (d\alpha')_{\text{vert}}) = 0$ .

Remark 5.10. In Theorem 5.9, relative to the direct sum decomposition  $D_{\mathcal{S}} = (D_{\mathcal{S}})_{hor} \oplus (D_{\mathcal{S}})_{vert}$ , cf. (5.24), the S-valued S-bilinear alternating form  $\pi$  on  $D_{\mathcal{S}}$ —the Poisson form—has the special feature that it decomposes into the two separate components  $\pi'' + \widetilde{\Omega}$  on  $(D_{\mathcal{S}})_{hor}$  and  $\pi'$  on  $(D_{\mathcal{S}})_{vert}$ , that is, though defined on all of  $D_{\mathcal{S}}$ , the 2-form  $\pi$  has no non-zero cross term on  $(D_{\mathcal{S}})_{vert} \times (D_{\mathcal{S}})_{hor}$ .

Remark 5.11. In the situation of Theorem 5.6, suppose that the Lie-Rinehart structure map from L'' to Der(A) is injective, and let  $\mathcal{R} = A^{L''}$ :  $= \{a \in A \mid \alpha''(a) = 0, \alpha'' \in L''\}$ . By definition,  $L'' = Der(A|\mathcal{R})$ . For example, when A is the ring of smooth functions on a smooth manifold B, in view of the Frobenius theorem, an  $(\mathbb{R}, A)$ -Lie algebra L'' of the kind under discussion defines a foliation of B, and  $\mathcal{R}$  is then the algebra of functions that are constant on the leaves. In the general case, we can then build the theory with  $\mathcal{R}$  as ground ring rather than just R: Relative to the sequence of inclusions  $\mathcal{R} \subseteq A \subseteq \mathcal{S}$ , the exact sequence (2.9) has the form

$$0 \longrightarrow S \otimes_A D_{A|\mathcal{R}} \longrightarrow D_{S|\mathcal{R}} \longrightarrow S \otimes_A L \longrightarrow 0$$

where we keep in mind that the canonical S-module morphism  $S \otimes_A L \to D_{S|A}$  is an isomorphism. That sequence fits into the commutative diagram

having the two vertical arrows surjective; the top row is precisely the exact sequence (2.9) relative to the ground ring R. With  $\Re$  as ground ring, Theorem 5.9 applies, that is, the Poisson bracket is given by the formula (5.39), and there are no non-zero cross terms, even when the injection of L'' into Der(A) is not an isomorphism. This observation justifies building the theory over a ground ring more general than a field.  $\blacklozenge$ 

**Example 5.12.** Consider the situation in Example 5.5, that is,  $P = B \times G$  and  $P \to B$  is the trivial principal G-bundle endowed with the trivial principal connection. In this case  $R = \mathbb{R}$ ,  $A = C^{\infty}(B)$  and  $L'' = \mathfrak{X}(B) = \mathrm{Der}(C^{\infty}(B))$ , so Theorem 5.9 applies. The algebra  $S = S_A[L]$  comes down to the tensor product algebra  $S[\mathfrak{g}] \otimes S_A[L'']$ . Let  $a \in C^{\infty}(B)$ ,  $\lambda \in \mathfrak{g}$ , and  $Y \in \mathfrak{X}(B)$ , and view  $a, \lambda, Y$  as functions on  $N = T^*B \times \mathfrak{g}^*$ . The algebra  $S' = S_A[L']$  decomposes as the tensor product algebra  $A \otimes S[\mathfrak{g}]$  and, as an S'-module,  $D_{S'}$  decomposes canonically as the direct sum  $S[\mathfrak{g}] \otimes D_A \oplus S' \otimes \mathfrak{g}$ . The projection to  $S[\mathfrak{g}] \otimes D_A$  yields the requisite linear connection, and it is immediate that  $(d\lambda)_{hor} = 0$ . Thus, by (5.39), the Poisson bracket on  $T^*B \times \mathfrak{g}^*$  is the sum of the canonical (minus) Poisson bracket on  $T^*B$  and the

minus Lie-Poisson bracket on  $\mathfrak{g}^*$ , cf. Remark 3.3 for the Lie-Poisson bracket on  $\mathfrak{g}^*$  that comes into play here.

In the next subsection we will determine the bracket for a general principal bundle.

5.7. The gauged Lie-Poisson bracket. Return to the situation of Example 4.1: Thus  $\xi \colon Q \to B$  is a right principal G-bundle. The ground ring R is now that of the real numbers,  $\mathbb{R}$ , and the algebra A is the algebra  $C^{\infty}(B)$ .

Let L' denote the A-Lie algebra of sections of the adjoint bundle  $\operatorname{ad}(\xi)\colon Q\times_G\mathfrak{g}\to B$  of  $\xi$ ; as before, we will view L' as an  $(\mathbb{R},A)$ -Lie algebra with trivial L'-action on A. ¿From the standard (plus) Lie-Poisson structure of  $\mathfrak{g}^*$ , the total space of the coadjoint bundle  $\operatorname{ad}^*(\xi)\colon Q\times_G\mathfrak{g}^*\to B$  of  $\xi$  acquires a smooth Poisson structure, the symmetric A-algebra  $\mathfrak{S}'=\mathfrak{S}_A[L']$  on L' carries the tautological Poisson structure introduced in Section 3 above, and the Poisson algebra  $\mathfrak{S}'$  embeds into  $C^\infty(Q\times_G\mathfrak{g}^*)$  as a Poisson algebra in an obvious manner via the dual pairing between  $\mathfrak{g}$  and  $\mathfrak{g}^*$ . Indeed,  $\mathfrak{S}'=\mathfrak{S}_A[L']$  is precisely the A-algebra of functions on  $Q\times_G\mathfrak{g}^*$  that are polynomial on the fibers of the coadjoint bundle projection  $\operatorname{ad}^*(\xi)$  and  $\mathfrak{S}'$  thus embeds into  $C^\infty(Q\times_G\mathfrak{g}^*)$  as a Fréchet dense subalgebra.

As in Subsection 5.3, let  $\widetilde{\xi} \colon \widetilde{Q} = Q \times_B \mathrm{T}^*B \to \mathrm{T}^*B$  denote the pull back bundle relative to the cotangent bundle projection  $\tau_B^* \colon \mathrm{T}^*B \to B$  and  $N_{\mathrm{St}} = \widetilde{Q} \times_G \mathfrak{g}^*$  the total space of the coadjoint bundle  $\mathrm{ad}^*(\widetilde{\xi}) \colon \widetilde{Q} \times_G \mathfrak{g}^* \to \mathrm{T}^*B$ , the Sternberg space (associated to the data). Exactly as before, let  $\widetilde{L}'$  denote the  $C^{\infty}(\mathrm{T}^*B)$ -Lie algebra of sections of the adjoint bundle of  $\widetilde{\xi}$ ; the Sternberg space  $N_{\mathrm{St}}$ , being the total space of the coadjoint bundle of  $\widetilde{\xi}$ , acquires a smooth Poisson structure, the symmetric  $C^{\infty}(\mathrm{T}^*B)$ -algebra  $\widetilde{S}'$  on  $\widetilde{L}'$  carries the tautological Poisson structure, and the Poisson algebra  $\widetilde{S}'$  embeds into  $N_{\mathrm{St}}$  as a Fréchet dense Poisson subalgebra. We will refer to the present Poisson structure on  $N_{\mathrm{St}}$  as the Lie-Poisson structure on  $N_{\mathrm{St}}$ .

Pick a connection on  $\xi \colon Q \to B$ . This choice of connection induces the corresponding diffeomorphism  $N_W \to N_{\mathrm{St}}$  given above as (5.12). From the obvious Poisson structure on  $N_W = (\mathrm{T}^*Q)/G$ , the Sternberg space  $N_{\mathrm{St}}$  thus inherits a Poisson structure and, cf. Remark 5.16 below, we shall see that the new Poisson structure perturbs the Lie-Poisson structure on  $N_{\mathrm{St}}$  in a very precise way. We are about to show that the new Poisson structure depends on the Lie-Poisson structure, the cotangent bundle Poisson structure on  $\mathrm{T}^*B$ , and the choice of connection; we therefore refer to the resulting Poisson bracket on  $N_{\mathrm{St}}$  as a gauged Lie-Poisson bracket.

Introduce the following notation:

- $\widehat{z}$ : for a point of  $N_{\mathrm{St}} = \widetilde{Q} \times_G \mathfrak{g}^*$ ,
- $\vartheta_B \colon \mathrm{TT}^*B \to \mathbb{R}$  for the tautological 1-form on  $\mathrm{T}^*B$ ,
- $\pi_{\mathfrak{g}^*}: N_{\operatorname{St}} = \widetilde{Q} \times_G \mathfrak{g}^* \to \mathrm{T}^*B$  for the bundle projection,
- $\beta = \pi_{\mathfrak{g}^*}(\widehat{z}) \in \mathrm{T}^*B$ ,
- $\omega \colon TQ \to \mathfrak{g}$  for the connection form on TQ,
- $\Omega$  for the curvature thereof,
- $\widetilde{\omega} : T\widetilde{Q} \to \mathfrak{g}$  for the lifted connection form on  $T\widetilde{Q}$ ,
- $\widetilde{\Omega}$  for the lifted curvature,
- $\widehat{\mu}$  for the variable in the fiber  $\mathfrak{g}_{\widehat{z}}^* = \pi_{\mathfrak{g}^*}^{-1}(\pi_{\mathfrak{g}^*}(\widehat{z})) \subseteq N_{\operatorname{St}}$  through the point  $\widehat{z}$  of  $N_{\operatorname{St}}$ ,
- $\langle \cdot, \cdot \rangle$  for the canonical pairing between a vector space and its dual,
- $\sharp$ :  $T^*(T^*B) \longrightarrow T(T^*B)$  for the isomorphism of vector bundles on  $T^*B$  induced by the symplectic form  $d\theta_B$ , the isomorphism being spelled out here on the total spaces.

The principal connection  $\omega$  on  $\xi$  induces a linear connection on the coadjoint bundle  $\operatorname{ad}^*(\xi)$  of  $\xi$  and hence on the lifted coadjoint bundle  $\operatorname{ad}^*(\widetilde{\xi})$ :  $N_{\operatorname{St}} = \widetilde{Q} \times_G \mathfrak{g}^* \to \operatorname{T}^*B$ . Given a function f on  $N_{\operatorname{St}} = \widetilde{Q} \times_G \mathfrak{g}^*$  and a point  $\widehat{z}$  of  $N_{\operatorname{St}}$ , the differential  $\operatorname{d} f(\widehat{z})$  decomposes uniquely in the form

$$\mathbf{d}f(\widehat{z}) = \mathbf{d}_{\widetilde{\omega}}f(\widehat{z}) + \mathbf{d}_{\widehat{\mu}}f(\widehat{z})$$

into a vertical summand  $\mathbf{d}_{\widehat{\mu}}f(\widehat{z})$  and a horizontal summand  $\mathbf{d}_{\widetilde{\omega}}f(\widehat{z})$ . This decomposition corresponds precisely to the decomposition (5.24) above, with the same significance of the terms *horizontal* and *vertical*.

We will now substitute the extension (4.7) of  $(\mathbb{R}, A)$ -Lie algebras for the extension (5.31) of  $(\mathbb{R}, A)$ -Lie algebras and apply the reasoning in Subsection 5.6 above. Thus, as an  $(\mathbb{R}, A)$ -Lie algebra,  $L'' = \text{Der}(A) = \mathfrak{X}(B)$ , the algebra  $\mathfrak{S}'' = \mathfrak{S}_A[L'']$  is the algebra of smooth functions on  $T^*B$  that are polynomial on the

fibers of the cotangent bundle projection  $T^*B \to B$ , as an  $(\mathbb{R}, A)$ -Lie algebra, L is that of G-equivariant vector fields on Q, and the algebra S is the algebra of smooth functions on  $N_{St} = \widetilde{Q} \times_G \mathfrak{g}^*$  that are polynomial on the fibers of the projection  $\widetilde{Q} \times_G \mathfrak{g}^* \to T^*B$ . Formula (5.39) then yields formula (5.45) below; up to sign, this formula coincides with the corresponding formula in [44]. We therefore label the subsequent result as a proposition rather than as a theorem.

**Proposition 5.13.** The Poisson bracket  $\{f_1, f_2\}$  of two functions  $f_1$  and  $f_2$  on  $\widetilde{Q} \times_G \mathfrak{g}^*$ , evaluated at the point  $\widehat{z}$  of  $N_{St} = \widetilde{Q} \times_G \mathfrak{g}^*$ , is given by the following expression where, as before,  $\beta = \pi_{\mathfrak{g}^*}(\widehat{z}) \in T^*B$ :

$$\{f_{1}, f_{2}\}(\widehat{z}) = \mathbf{d}\vartheta_{B}(\beta) \left(\mathbf{d}_{\widetilde{\omega}} f_{1}^{\sharp}(\widehat{z}), \mathbf{d}_{\widetilde{\omega}} f_{2}^{\sharp}(\widehat{z})\right)$$

$$+ \left\langle \widehat{z}, \widetilde{\Omega}(\beta) \left(\mathbf{d}_{\widetilde{\omega}} f_{1}^{\sharp}(\widehat{z}), \mathbf{d}_{\widetilde{\omega}} f_{2}^{\sharp}(\widehat{z})\right) \right\rangle$$

$$+ \left\langle \widehat{z}, \left[\mathbf{d}_{\widehat{\mu}} f_{1}(\widehat{z}), \mathbf{d}_{\widehat{\mu}} f_{2}(\widehat{z})\right]_{\widehat{z}} \right\rangle.$$

In this formula,  $\mathbf{d}\vartheta_B$  is the resulting symplectic structure on  $\mathrm{T}^*B$ , the expression  $\mathbf{d}_{\widetilde{\omega}}f^{\sharp}(\widehat{z}) \in \mathrm{T}_{\beta}(\mathrm{T}^*B)$  refers to the vector in  $\mathrm{T}_{\beta}(\mathrm{T}^*B)$  associated to  $\mathbf{d}_{\widetilde{\omega}}f(\widehat{z})$  under  $\sharp$  and, in  $\langle \widehat{z}, \widetilde{\Omega}(\beta) \left( \mathbf{d}_{\widetilde{\omega}}f_1^{\sharp}(\widehat{z}), \mathbf{d}_{\widetilde{\omega}}f_2^{\sharp}(\widehat{z}) \right) \rangle$  and  $\langle \widehat{z}, [\mathbf{d}_{\widehat{\mu}}f_1(\widehat{z}), \mathbf{d}_{\widehat{\mu}}f_2(\widehat{z})]_{\widehat{z}} \rangle$ , the point  $\widehat{z}$  is viewed as a vector in the vector space  $\mathfrak{g}_{\widehat{z}}^*$ .

**Remark 5.14.** In [44], the tautological 1-form on B is written as  $\alpha_B$ ; furthermore, in that reference, the group acts on the total space Q from the left (written there as P) and, accordingly, the term  $\langle \hat{z}, [\mathbf{d}_{\widehat{\mu}} f_1(\widehat{z}), \mathbf{d}_{\widehat{\mu}} f_2(\widehat{z})]_{\widehat{z}} \rangle$  comes with a minus sign.  $\blacklozenge$ 

Proof. This is a special case of formula (5.39). The terms  $\pi'((df)_{\text{vert}}, (dh)_{\text{vert}})$ ,  $\pi''((df)_{\text{hor}}, (dh)_{\text{hor}})$ ,  $\widetilde{\Omega}((df)_{\text{hor}}, (dh)_{\text{hor}})$  on the right-hand side of (5.39) correspond to, respectively,  $\langle \widehat{z}, [\mathbf{d}_{\widehat{\mu}} f_1(\widehat{z}), \mathbf{d}_{\widehat{\mu}} f_2(\widehat{z})]_{\widehat{z}} \rangle$ ,  $\mathbf{d}\vartheta_B(\beta) \left(\mathbf{d}_{\widetilde{\omega}} f_1^{\sharp}(\widehat{z}), \mathbf{d}_{\widetilde{\omega}} f_2^{\sharp}(\widehat{z})\right)$ , and  $\langle \widehat{z}, \widetilde{\Omega}(\beta) \left(\mathbf{d}_{\widetilde{\omega}} f_1^{\sharp}(\widehat{z}), \mathbf{d}_{\widetilde{\omega}} f_2^{\sharp}(\widehat{z})\right) \rangle$ , in (5.45).

A straightforward though slightly tedious comparison shows that the formula (5.45) for the value of the Poisson bracket  $\{f_1, f_2\}$  at the point  $\hat{z}$  is indeed a special case of the formula (5.39).

Remark 5.15. Our approach provides a simple explanation for the formula (5.45) for the Poisson structure: The term  $d\vartheta_B(\beta)\left(\mathbf{d}_{\widetilde{\omega}}f_1^\sharp(\widehat{z}),\mathbf{d}_{\widetilde{\omega}}f_2^\sharp(\widehat{z})\right)$  comes from the symplectic Poisson structure associated to the 2-form  $d\vartheta_B$  on the base  $T^*B$  of the bundle projection  $\pi_{\mathfrak{g}^*}\colon N_{\mathrm{St}}\to T^*B$  and thus corresponds to the first term on the right-hand side of (4.9) and to the identity (4.12) above; the term  $\left\langle\widehat{z},\widetilde{\Omega}(\beta)\left(\mathbf{d}_{\widetilde{\omega}}f_1^\sharp(\widehat{z}),\mathbf{d}_{\widetilde{\omega}}f_2^\sharp(\widehat{z})\right)\right\rangle$  comes from the curvature of the connection and therefore corresponds to the second term on the right-hand side of (4.9); and the term  $\left\langle\widehat{z},\left[\mathbf{d}_{\widehat{\mu}}f_1(\widehat{z}),\mathbf{d}_{\widehat{\mu}}f_2(\widehat{z})\right]_{\widehat{z}}\right\rangle$  comes from the Lie-Poisson structure of  $\mathfrak{g}^*$  and therefore corresponds to the identity (4.8). The two terms in the Poisson tensor beyond that coming from the cotangent bundle Poisson structure on  $T^*B$  simply reconstruct the Atiyah sequence (4.5) of the principal bundle or, equivalently, the corresponding extension of  $(\mathbb{R}, C^{\infty}(B))$ -Lie algebras, in terms of the connection.

**Remark 5.16.** The summand  $\langle \widehat{z}, [\mathbf{d}_{\widehat{\mu}} f_1(\widehat{z}), \mathbf{d}_{\widehat{\mu}} f_2(\widehat{z})]_{\widehat{z}} \rangle$  in (5.45) yields the Lie-Poisson bracket on the Sternberg space  $N_{\mathrm{St}} = \widetilde{Q} \times_G \mathfrak{g}^*$ . Thus the formula (5.45) manifestly perturbs the Lie-Poisson structure of the Sternberg space  $N_{\mathrm{St}}$  in a very precise way.

5.8. General Lie algebroids. Let  $\lambda \colon \mathcal{L} \to B$  be a general Lie algebroid; then the pair  $(A, L) = (C^{\infty}(B), \Gamma_B(\lambda))$  acquires a Lie-Rinehart algebra structure. A particular example is a Lie algebroid of the kind reproduced in Example 4.1. For a general Lie algebroid  $\lambda$ , the algebra  $\mathcal{S}_A[L]$  is in an obvious manner isomorphic to the algebra of smooth functions on the total space  $\mathcal{L}^*$  of the dual bundle  $\lambda^* \colon \mathcal{L}^* \to B$  that are polynomial on the fibers of  $\lambda^*$ , the algebra  $C^{\infty}(\mathcal{L}^*)$  of ordinary smooth functions on  $\mathcal{L}^*$  acquires a Poisson structure in an obvious way, and the injection of  $\mathcal{S}_A[L]$  into  $C^{\infty}(\mathcal{L}^*)$  maps the former algebra onto a Fréchet dense subalgebra of the latter and is plainly compatible with the Poisson structures. The special case of the tangent bundle has been presented in Subsection 5.7. In the general case, the image L'' of L under the morphism  $L \to \mathfrak{X}(B)$  of  $(\mathbb{R}, A)$ -Lie algebras, which is part of the Lie algebroid structure of  $\lambda$ , is an  $(\mathbb{R}, A)$ -Lie algebra, and the morphism  $L \to \mathfrak{X}(B)$  of  $(\mathbb{R}, A)$ -Lie algebras induces a morphism  $\mathcal{S}_A[L] \to \mathcal{S}_A[\mathfrak{X}(B)]$  of Poisson algebras, the latter algebra being the ordinary Poisson

algebra of smooth functions on the total space  $T^*B$  of the cotangent bundle of B that are polynomial on the fibers. Moreover, the projection from L to L'' fits into an extension

$$\mathbf{e} \colon 0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$$

of  $(\mathbb{R}, A)$ -Lie algebras of the kind (4.1). A special case is the key example (4.7) above.

Under a suitable additional hypothesis, the construction in the previous section yields the following description of the Poisson algebra  $S_A[L]$  in terms of A, the extension  $\mathbf{e}$ , and an  $\mathbf{e}$ -connection thereof.

**Theorem 5.17.** Suppose that the anchor  $\rho: \mathcal{L} \to TB$  has constant rank, so that L'' defines a foliation  $\mathcal{F}$  on B, and denote by  $A^{\mathcal{F}}$  the algebra of smooth functions on B that are constant on the leaves of  $\mathcal{F}$ . Then the Poisson algebras  $\mathcal{S}_A[L]$  and  $\mathcal{S}_A[L'']$  are Poisson algebras even over  $A^{\mathcal{F}}$ , the functions in  $A^{\mathcal{F}}$  being Casimir functions, the extension  $\mathbf{e}$  splits in the category of A-modules and, once an  $\mathbf{e}$ -connection (i. e., A-module splitting of  $\mathbf{e}$ ) has been chosen, the identities (4.8)–(4.14) yield an explicit description of the Poisson algebra  $\mathcal{S}_A[L]$  of smooth functions on  $\mathcal{L}^*$ . Furthermore, in terms of differentials, the tautological Poisson bracket on  $\mathcal{S}_A[L]$  is then as well given by the formula (5.39).

*Proof.* Since the anchor  $\rho: \mathcal{L} \to TB$  has constant rank,  $\rho$  maps  $\mathcal{L}$  onto the total space of a subbundle of the tangent bundle of B. As an A-module, L'' is the space of sections of this vector bundle whence L'' is necessarily projective as an A-module. Working over  $A^{\mathcal{F}}$  as ground ring, cf. Remark 5.11, we conclude that the tautological Poisson bracket on  $\mathcal{S}_A[L]$  is still given by the formula (5.39). This implies the assertion.

Here is a special case.

Corollary 5.18. Suppose that  $\lambda \colon \mathcal{L} \to B$  is a transitive Lie algebroid. Then L'' coincides with the  $(\mathbb{R}, A)$ -Lie algebra  $\mathfrak{X}(B)$  of ordinary smooth vector fields on B, the extension  $\mathbf{e}$  splits in the category of A-modules and, once an  $\mathbf{e}$ -connection has been chosen, the identities (4.8)–(4.14) yield an explicit description of the Poisson algebra  $S_A[L]$  of smooth functions on  $\mathcal{L}^*$  and hence a complete description of the Poisson algebra  $(C^{\infty}(\mathcal{L}^*), \{\cdot, \cdot\})$  of ordinary smooth functions on  $\mathcal{L}^*$ . Furthermore, in terms of differentials, the tautological Poisson bracket on  $S_A[L]$  is then as well given by the formula (5.39) and hence by a suitable variant of (5.45).

The Corollary applies, in particular, to the transitive Lie algebroid (4.6), but that special case has been dealt with before.

# 6. Non-regular quotients of cotangent bundles

Let Q denote a smooth manifold, G a group acting smoothly on Q from the right (as before), and lift the action to the total space TQ of the tangent bundle. We no longer suppose that the action of G on Q is principal and form the non-regular quotient  $(T^*Q)/G$ . Endow the orbit space B = Q/G with the quotient topology, let  $\pi \colon Q \to B$  be the canonical projection, and let  $(A_Q, L_Q) := (C^\infty(Q), \mathfrak{X}(Q))$ . Moreover, let  $A = C^\infty(Q)^G$ , the algebra of smooth G-invariant functions on G0, viewed as an algebra of continuous functions on G1, and let G2, the Lie algebra of smooth G2-invariant vector fields on G3. The Lie-Rinehart structure on G4, G7 in particular, an action G7 in G8 and G9 in G9. The Lie-Rinehart structure on G9 in G9 derivations. In this section we will show that, in the present more general case, an algebra of the kind G9 in G9. The functions on the quotient G9 in an obvious way, this embedding cannot be onto a Fréchet dense subalgebra. Consequently, a statement of the kind spelled out in Theorem 4.2 no longer suffices to recover the Poisson algebra on the quotient G9.

6.1. The Poisson structure on  $S_A[L]$ . Let L'' denote the image of L in Der(A). Similarly, the pair (A, L'') acquires a Lie-Rinehart structure, and the surjection from L to L'' fits into an extension

(6.1) 
$$\mathbf{e} \colon 0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$$

of  $(\mathbb{R}, A)$ -Lie algebras of the kind (4.1). When G is a Lie group and when the G-action on Q is principal, the extension (6.1) is the sequence of the spaces of sections of the Atiyah sequence (4.5). In the general case, the orbit space  $(T^*Q)/G$  need no longer be a smooth manifold, the algebra  $\mathcal{S}_A[L]$  embeds into  $(C^{\infty}(T^*Q))^G$  and the tautological Poisson algebra  $\mathcal{S}_A[L]$  yields a Poisson algebra of continuous functions

on  $(T^*Q)/G$ . Thus, provided that the extension (6.1) splits in the category of A-modules, Theorem 4.2 applies and furnishes an explicit description of the induced Poisson structure on the algebra  $\mathcal{S}_A[L]$  of continuous functions on the orbit space  $(T^*Q)/G$ . As for the algebra  $\mathcal{S}_A[L'']$ , it is tempting to try to interpret it as an algebra of continuous functions on the total space  $T^*B$  of the cotangent bundle of B but this interpretation is only available when the orbit space B = Q/G is a smooth manifold; in general we can think of  $\mathcal{S}_A[L'']$  as a replacement for an algebra of functions on  $T^*B$ .

**Theorem 6.1.** Suppose that the orbit space B = Q/G is a smooth manifold in such a way that the projection  $\pi \colon Q \to B$  is a smooth submersion. Then the algebra A coincides with the algebra  $C^{\infty}(B)$  of ordinary smooth functions on B, the  $(\mathbb{R}, A)$ -Lie algebra L'' amounts to the  $(\mathbb{R}, A)$ -Lie algebra  $\mathfrak{X}(B)$  of ordinary smooth vector fields on B, the extension (6.1) splits in the category of A-modules, that is, admits a connection, and the identities (4.8)-(4.14) yield an explicit description of the induced Poisson algebra  $S_A[L]$  of continuous functions on the orbit space  $(T^*Q)/G$ .

*Proof.* The algebra A plainly coincides with the algebra  $C^{\infty}(B)$  of ordinary smooth functions on B, and the regularity assumption implies that the injection from L'' into the  $(\mathbb{R}, A)$ -Lie algebra  $\mathrm{Der}(A) \cong \mathfrak{X}(B)$  of ordinary smooth vector fields on B is surjective as well. Since, as an A-module,  $\mathfrak{X}(B)$  is projective, the extension (6.1) splits in the category of A-modules.

A special case of Theorem 6.1 arises when  $\pi\colon Q\to B$  is a principal G-bundle. This case has been dealt with in Theorem 5.17 above. In particular, the induced Poisson algebra  $\mathcal{S}_A[L]$  then yields a complete description of the Poisson algebra  $\left(C^\infty(\mathrm{T}^*Q)^G, \{\,\cdot\,,\,\cdot\,\}\right)$  on the quotient space  $(\mathrm{T}^*Q)/G$ . In Theorem 6.1, we do not assert that, for general G and G0, we obtain a complete description of the Poisson algebra  $\left(C^\infty(\mathrm{T}^*Q)^G, \{\,\cdot\,,\,\cdot\,\}\right)$  on the quotient space  $(\mathrm{T}^*Q)/G$ ; we recover only the subalgebra  $\mathcal{S}_A[L]$  and, in general, the relationship between  $\mathcal{S}_A[L]$  and  $C^\infty(\mathrm{T}^*Q)^G$  is more subtle than that between the functions on the total space of a cotangent bundle that are polynomial on the fibers and all smooth functions on that total space. In particular, the algebra  $\mathcal{S}_A[L]$  is not necessarily a Fréchet dense subalgebra of  $C^\infty(\mathrm{T}^*Q)^G$ , and invariants that cannot be recovered from  $\mathcal{S}_A[L]$  may show up. We shall illustrate this fact in Subsection 6.2 below.

Under the circumstances of Theorem 6.1, even though the quotient  $(\mathbb{R}, A)$ -Lie algebra L'' is the  $(\mathbb{R}, A)$ -Lie algebra of smooth vector fields on an ordinary smooth manifold and hence arises from an ordinary Lie algebroid over B, the  $(\mathbb{R}, A)$ -Lie algebra L is not necessarily projective as an A-module and hence does not necessarily arise from an ordinary Lie algebroid over B unless the G-action is principal.

For illustration, suppose that the G-action on Q is proper and has a single orbit type. Then the orbit space B = Q/G is a smooth manifold, the projection  $\pi \colon Q \to B$  is a smooth locally trivial fiber bundle (cf. [35]), and Theorem 6.1 applies. This situation has been explored in [11]. Moreover, under these circumstances, the assignment to a vector in the Lie algebra  $\mathfrak g$  of its induced fundamental vector field on Q yields an exact sequence

$$(6.2) Q \times \mathfrak{g} \longrightarrow TQ \longrightarrow Q \times_B TB \longrightarrow 0$$

of smooth G-vector bundles over Q, each of the unlabeled horizontal arrows being of constant rank.

6.2. **Homogeneous spaces.** We will now justify the claim made in the previous subsection that, when the G-action on Q is no longer free, the algebra  $\mathcal{S}_A[L]$  does not simply come down to a Fréchet dense subalgebra of  $C^{\infty}(T^*Q)^G$ .

As before, let G be a Lie group. Let H be a closed subgroup and consider the homogeneous space Q = G/H, viewed as the base of the (right) principal H-bundle  $\pi \colon G \to G/H$  having G as total space; left translation in G turns this bundle into a principal H-bundle in the category of left G-spaces. Lift the left G-action to  $\mathrm{T}(G/H)$  and  $\mathrm{T}^*(G/H)$  in the standard manner. To adjust the exposition to the standard description of homogeneous spaces, we work here with left G-manifolds rather than with right G-manifolds, and the reader will easily translate the reasoning to right G-manifolds. The subsequent reasoning is independent of our discussion of Poisson structures and there is no need to rebuild the theory with left G-actions rather than right G-actions.

We will now suppose that G/H is reductive (cf. [33]), i.e., as an H-module,  $\mathfrak{g}$  decomposes as a direct sum  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$  in such a way that  $[\mathfrak{h}, \mathfrak{q}] \subset \mathfrak{q}$ . From the exact sequence

$$(6.3) 0 \longrightarrow G \times \mathfrak{h} \longrightarrow TG \longrightarrow G \times \mathfrak{q} \longrightarrow 0$$

of H-vector bundles over G (spelled out for the total spaces) we see that the canonical morphism

$$(6.4) G \times_H \mathfrak{q} \longrightarrow T(G/H)$$

yields an isomorphism of smooth vector bundles over G/H whence the algebra  $(C^{\infty}(\mathrm{T}^*(G/H)))^G$  of smooth G-invariant functions on  $\mathrm{T}^*(G/H)$  equals  $(C^{\infty}(\mathfrak{q}^*))^H$ , the algebra of smooth H-invariant functions on  $\mathfrak{q}^*$ ; this algebra contains the algebra  $(S[\mathfrak{q}])^H$  of H-invariant polynomials in  $\mathfrak{q}$  in an obvious manner and  $(S[\mathfrak{q}])^H$  is Fréchet dense in  $(C^{\infty}(\mathfrak{q}^*))^H$ . When H is compact, in view of a result in [39], the algebra  $(C^{\infty}(\mathfrak{q}^*))^H$  of smooth H-invariant functions on  $\mathfrak{q}^*$  is the algebra of smooth functions in the generators of  $(S[\mathfrak{q}])^H$ .

To reconcile this description with that given earlier, we note first that the space B of G-orbits in Q = G/H is a single point; hence the algebra A amounts to the ground field  $\mathbb{R}$ , and the extension (6.1) becomes the identity mapping  $L \to L$  where L is the ordinary Lie algebra  $(\mathfrak{X}(Q))^G$  of smooth G-invariant vector fields on Q = G/H. However, inspection of the morphism

of vector bundles shows that, as vector spaces,  $(\mathfrak{X}(Q))^G \cong \mathfrak{q}^H$ , the vector space of H-fixed points in  $\mathfrak{q}$ . Plainly, the algebra  $(\mathfrak{S}[\mathfrak{q}])^H$  of H-invariant polynomials in  $\mathfrak{q}$  contains the algebra  $\mathfrak{S}[L] \cong \mathfrak{S}[\mathfrak{q}^H]$  of polynomials in  $\mathfrak{q}^H$  but in general the two will *not* coincide. In particular, when the H-representation  $\mathfrak{q}$  does not contain the trivial representation,  $\mathfrak{q}^H$  is zero whereas the algebra  $(\mathfrak{S}[\mathfrak{q}])^H$  of H-invariant polynomials in  $\mathfrak{q}$  is non-trivial, unless  $\mathfrak{q}$  is zero, since the space of H-orbits in  $\mathfrak{q}$  does not reduce to a point. Thus Theorem 4.2 cannot recover a Fréchet dense subalgebra of the algebra  $(C^\infty(T^*(G/H)))^G$  of smooth G-invariant functions on  $T^*(G/H)$ .

**Example 6.2.** Let G be a Lie group and view Q = G as a homogeneous space of the group  $G \times G$  in the standard manner, that is, G is viewed as a  $(G \times G)$ -space via left and right translation. In other words,  $H = \Delta G \cong G$  being the diagonal group in  $G \times G$ , G is identified with the space  $G \times G = G$  or orbits in  $G \times G$  relative to the diagonal action. The decomposition

$$\mathfrak{g} \oplus \mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}, \ \mathfrak{h} = \{(X, X); X \in \mathfrak{g}\}, \ \mathfrak{q} = \{(Y, -Y); Y \in \mathfrak{g}\}$$

is reductive, and the algebra  $(S[\mathfrak{q}])^H$  of H-invariant polynomials on  $\mathfrak{q}$  coincides with the algebra  $(S[\mathfrak{g}])^G$  of invariants on  $\mathfrak{g}$  (under the adjoint action). When  $\mathfrak{g}$  is semisimple,  $\mathfrak{g}^G$  is zero whereas  $(S[\mathfrak{g}])^G$  is never zero unless  $\mathfrak{g}$  is zero. The Poisson bracket on  $(S[\mathfrak{g}])^G$  is zero; indeed, the algebra  $(S[\mathfrak{g}])^G$  is that of Casimir elements in the Lie-Poisson algebra  $S[\mathfrak{g}]$ . Plainly, the method which works well in the case of a principal G-action does not suffice in the present situation, that is, Theorem 4.2 cannot recover the algebra under discussion.

Example 6.3. Embed  $H = \mathrm{SO}(2,\mathbb{R})$  into  $G = \mathrm{SO}(3,\mathbb{R})$  as rotations about the vertical axis. Then G/H is the standard 2-sphere  $S^2$ . In the decomposition  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$  of the Lie algebra  $\mathfrak{g} = \mathfrak{so}(3,\mathbb{R})$ ,  $\dim \mathfrak{h} = 1$ , and  $\mathfrak{q}$  is the defining (irreducible) 2-dimensional representation of  $\mathrm{SO}(2,\mathbb{R})$  via rotations in the plane  $\mathfrak{q}$ ; here we write the elements of  $\mathfrak{q}$  as  $\mathbf{x} = (x_1, x_2)$ . Since the representation  $\mathfrak{q}$  is irreducible,  $\mathfrak{q}^H$  is zero whereas the algebra  $(S[\mathfrak{q}])^H$  of H-invariant polynomials on  $\mathfrak{q}$  is the polynomial algebra in the single variable  $\mathbf{x}^2 = x_1^2 + x_2^2$ . Thus, the space  $T^*S^2$  being viewed as the total space of a vector bundle on  $S^2$  with a  $\mathrm{O}(3,\mathbb{R})$ -invariant Riemannian structure, the assignment to a covector  $\alpha_{\mathfrak{q}} \in T^*S^2$   $(\mathfrak{q} \in S^2)$  of the square  $\alpha_{\mathfrak{q}}^2$  of its length relative to the Riemannian structure induces a diffeomorphism (beware: the notion of diffeomorphism being suitably interpreted, see below) between  $(T^*S^2)/G$  and  $\mathbb{R}_{\geq 0}$ , and the Poisson bracket on the resulting algebra  $(C^\infty(T^*S^2))^G$  is zero. Here  $C^\infty(\mathbb{R}_{\geq 0})$  is not an algebra of ordinary smooth functions: the space  $\mathbb{R}_{\geq 0}$  is endowed with the algebra of Whitney functions relative to the embedding into  $\mathbb{R}$ ; these are continuous functions that are restrictions of smooth functions on some open interval of the kind  $]-\varepsilon,+\infty[$  where  $\varepsilon>0$ . Again, the method which works well in the case of a principal G-action does not suffice in the present situation, that is, Theorem 4.2 cannot recover this kind of algebra under discussion.  $\blacklozenge$ 

#### APPENDIX

We justify the claim that the left-hand S-module morphism

$$(6.6) S \otimes_A D_A \longrightarrow D_S, \ s \otimes_A da \longmapsto sda, \ s \in S, \ a \in A,$$

in (2.9) is injective.

Since V is finitely generated and projective as an A-module, there is a finitely generated projective A-module W such that  $V \oplus W$  is free. Consequently there is a free R-module U such that  $V \oplus W = A \otimes U$ . Let  $\mathcal{S}_U = \mathcal{S}_A[A \otimes U]$ . Since the R-algebra  $\mathcal{S}[U]$  admits a canonical augmentation map  $\mathcal{S}[U] \to R$ , the canonical morphism  $A \to \mathcal{S}_U$  of R-modules is injective and, since as an R-module,  $\mathcal{S}[U]$  is free, the canonical morphism  $\mathcal{S}[U] \to \mathcal{S}_U$  of R-modules is, likewise, injective. Now

$$S_U = S_A[A \otimes U] \cong A \otimes S[U] \cong S_A[V] \otimes_A S_A[W],$$

and the injections  $A \to \mathbb{S}_U$  and  $\mathbb{S}[U] \to \mathbb{S}_U$  induce a direct sum decomposition

$$D_{\mathcal{S}_U} \cong \mathcal{S}[U] \otimes D_A \oplus \mathcal{S}_U \otimes U.$$

We rewrite this decomposition in the form

This decomposition implies that the canonical morphism  $S_U \otimes_A D_A \to D_{S_U}$  of  $S_U$ -modules is injective; furthermore, the decomposition splits the resulting extension

$$(6.9) 0 \longrightarrow S_{II} \otimes_A D_A \longrightarrow D_{S_{II}} \longrightarrow D_{S_{II}|A} \longrightarrow 0$$

of  $S_U$ -modules. Notice that  $D_{S_U|A} \cong S_A[V] \otimes_A S_A[W] \otimes_A (V \oplus W)$  (canonically).

Recall that  $S = S_A[V]$  and consider the canonical surjection  $S_U \to S$  of A-algebras, viewed as a morphism of R-algebras. Since (6.9) is a split extension in the category of  $S_U$ -modules, applying the functor  $S \otimes_{S_U}$  to (6.9), we obtain the extension

$$(6.10) 0 \longrightarrow S \otimes_A D_A \longrightarrow S \otimes_{S_U} D_{S_U} \longrightarrow S \otimes_A (V \oplus W) \longrightarrow 0$$

of S-modules, necessarily split; the salient feature here is the exactness at  $S \otimes_A D_A$  (injectivity of the second unlabeled arrow).

Let  $I_W \subseteq S_A[W]$  be the kernel of the obvious algebra surjection  $S_A[W] \to A$ , and recall the fundamental exact sequence

$$(6.11) I_W/I_W^2 \longrightarrow A \otimes_{S_A[W]} D_{S_A[W]} \longrightarrow D_A \longrightarrow 0$$

in the category of A-modules. Standard homological algebra yields an isomorphism

$$\operatorname{Tor}^{\mathbb{S}_A[W]}(A,A) \longrightarrow A \otimes_{\mathbb{S}_A[W]} I_W,$$

and the canonical surjection  $I_W \to A \otimes_{\mathcal{S}_A[W]} I_W$  descends to an isomorphism  $I_W / I_W^2 \to A \otimes_{\mathcal{S}_A[W]} I_W$ . Likewise, let  $J_U$  denote the kernel of the R-algebra surjection  $\mathcal{S}_U \to \mathcal{S}$ , and consider the fundamental exact sequence

$$(6.12) J_U/J_U^2 \longrightarrow S \otimes_{S_U} D_{S_U} \longrightarrow D_S \longrightarrow 0$$

associated to the surjection  $S_U \to S$ .

The obvious map

$$S_U \otimes_{S_A[W]} I_W \longrightarrow S_U \otimes_{S_A[W]} S_A[W] \cong S_U$$

identifies  $S_U \otimes_{S_A[W]} I_W$  with the kernel  $J_U$  of  $S_U \to S$ , and the obvious morphism

$$S \otimes_A \operatorname{Tor}^{S_A[W]}(A,A) \cong S \otimes_A (I_W/I_W^2) \longrightarrow J_U/J_U^2 \cong S \otimes_{S_U} J_U \cong \operatorname{Tor}^{S_U}(S,S)$$


is an isomorphism of S-modules. Indeed, this is the canonical morphism

$$S \otimes_A \operatorname{Tor}^{S_A[W]}(A,A) \longrightarrow \operatorname{Tor}^{S \otimes_A S_A[W]}(S,S)$$

of S-modules, necessarily an isomorphism since  $S = S_A[V]$  is projective as an A-module, the A-module V being projective by assumption. Consequently the fundamental exact sequence (6.12) takes the form

$$(6.13) S \otimes_A (I_W/I_W^2) \longrightarrow S \otimes_{S_U} D_{S_U} \longrightarrow D_S \longrightarrow 0.$$

The surjection  $S_U \to S$  induces the commutative diagram



with exact rows and columns in the category of S-modules. The middle horizontal row is the (split) extension (6.10), necessarily exact as pointed out above. The right-hand vertical column is obviously exact. The middle vertical column is the exact sequence (6.13), with an arrow added emanating at 0. The A-module  $I_W/I_W^2$  is canonically isomorphic to W whence  $S \otimes_A (I_W/I_W^2) \to S \otimes_A W$  is an isomorphism of S-modules. Consequently the middle vertical column is exact at  $S \otimes_A (I_W/I_W^2)$  and hence exact.

Diagram chasing or the snake lemma implies that the induced morphism  $S \otimes_A D_A \to D_S$  of S-modules is injective.

#### ACKNOWLEDGEMENTS

We are indebted to H. Cendra, J. Marsden, R. Montgomery and J. Stasheff for various comments that helped us improve the exposition, to S. Chase, Y. Kosmann-Schwarzbach, K. Mackenzie, P. Michor, J. Stasheff and A. Weinstein for discussions about Lie-Rinehart algebras and, last but not least, to the referee for his remarks. T.S. Ratiu was partially supported by Swiss NSF grant 200021-121512. J. Huebschmann acknowledges partial support by the CNRS and by the Labex CEMPI (ANR-11-LABX-0007-01).

#### References

- [1] R. Abraham and J. E. Marsden: Foundations of Mechanics. Benjamin-Cummings Publishing Company, 1978.
- [2] R. Almeida and P. Molino: Suites d'Atiyah et feuilletages transversalement complets. C. R. Acad. Sci. Paris I 300 (1985), 13-15.
- [3] V. I. Arnold: Mathematical Methods of Classical Mechanics. Graduate Texts in Mathematics, No. 60, Springer Verlag, Berlin · Heidelberg · New York · Tokyo, 1978, 1989 (2nd edition).
- [4] M. F. Atiyah: Complex analytic connections in fibre bundles. Trans. Amer. Math. Soc. 85 (1957), 181–207.
- [5] A. Cannas da Silva and A. Weinstein: Geometric models for noncommutative algebras. Berkeley Mathematics Lecture Notes 10, American Mathematical Society, Providence, RI, 1999.
- [6] H. Cendra, J.E. Marsden, T.S. Ratiu: Lagrangian reduction by stages, Mem. Amer. Math. Soc., 152 (2001), no. 722.
- [7] H. Cendra, J.E. Marsden, S. Pekarsky, T.S. Ratiu: Variational principles for Lie-Poisson and Hamilton-Poincaré equations, Mosc. Math. J. 3, (2003), 833–867.
- [8] T. Courant: Dirac manifolds, Trans. Amer. Math. Soc., 319, (1990), 631-661.
- [9] T. Courant: Tangent Lie algebroids, J. Phys. A, 27, (1994), 4527–4536.
- [10] W. Greub, St. Halperin, and R. Vanstone: Connections, curvature, and cohomology. Vol. I: De Rham cohomology of manifolds and vector bundles. Pure and Applied Mathematics, Vol. 47, Academic Press, New York, 1972.
- [11] S. Hochgerner and A. Rainer: Singular Poisson reduction of cotangent bundles. Rev. Mat. Complut. 19 (2006), 431–466.
- [12] J. Huebschmann: Poisson cohomology and quantization. J. reine angew. Math. 408 (1990), 57–113.
- [13] J. Huebschmann: On the quantization of Poisson algebras. In: Symplectic Geometry and Mathematical Physics, Actes du colloque en l'honneur de Jean-Marie Souriau, P. Donato, C. Duval, J. Elhadad, G. M. Tuynman, eds.; Progress in Mathematics, Vol. 99, Birkhäuser-Verlag, Boston · Basel · Berlin, 204–233 (1991).
- [14] J. Huebschmann: Lie-Rinehart algebras, descent, and quantization. Fields Institute Communications 43 (2004), 295–316, math.SG/0303016.
- [15] J. Huebschmann: Lie-Rinehart algebras, Gerstenhaber algebras, and Batalin-Vilkovisky algebras. Annales de l'Institut Fourier 48 (1998), 425–440, math.DG/9704005.
- [16] J. Huebschmann: Extensions of Lie-Rinehart algebras and the Chern-Weil construction. In: Festschrift to honor the 60-th birthday of Jim Stasheff, Cont. Math. 227 (1999), 145–176, math.DG/9706002.

- [17] J. Huebschmann: Duality for Lie-Rinehart algebras and the modular class. J. reine angew. Math. 510 (1999), 103–159, math.DG/9702008.
- [18] J. Huebschmann: Differential Batalin-Vilkovisky algebras arising from twilled Lie-Rinehart algebras. Banach center publications 51 (2002), 87–102.
- [19] J. Huebschmann: Higher homotopies and Maurer-Cartan algebras: quasi-Lie-Rinehart, Gerstenhaber-, and Batalin-Vilkovisky algebras. In: The Breadth of Symplectic and Poisson Geometry, Festschrift in Honor of Alan Weinstein, J. Marsden and T. Ratiu, eds., Progress in Mathematics, Vol. 232, Birkhäuser-Verlag, Boston · Basel · Berlin, 237–302 (2004), math.DG/0311294
- [20] J. Huebschmann: Kähler spaces, nilpotent orbits, and singular reduction. Memoirs AMS 172/814 (2004), Amer. Math. Society, Providence, R. I., math.DG/0104213.
- [21] J. Huebschmann: Singular Poisson-Kähler geometry of certain adjoint quotients. In: The Mathematical Legacy of C. Ehresmann, J. Kubarski, and R. Wolak, eds., Banach Center Publications 76 (2007), 325–347, math.SG/0610614.
- [22] J. Huebschmann: Stratified Kähler structures on adjoint quotients. Differential Geometry and its Applications 26 (2008), 704-731, math.DG/0404141.
- [23] J. Huebschmann: Kähler reduction and quantization J. reine angew. Math. 591 (2006), 75-109, math. SG/0207166.
- [24] J. Huebschmann, G. Rudolph and M. Schmidt: A gauge model for quantum mechanics on a stratified space, Commun. Math. Phys. 286 (2009), 459–494, hep-th/0702017.
- [25] K. Mackenzie: General Theory of Lie groupoids and Lie algebroids. London Math. Soc. Lecture Note Series, vol. 213, Cambridge University Press, Cambridge, England, 2005.
- [26] J.E. Marsden: Lectures on Geometric Methods in Mathematical Physics, Volume 37, SIAM, Philadelphia, 1981.
- [27] J.E. Marsden, G. Misiołek, J.-P. Ortega, M. Perlmutter, T.S. Ratiu: Hamiltonian Reduction by Stages, Lecture Notes in Mathematics, 1913 (2007), Springer-Verlag.
- [28] J.E. Marsden and M. Perlmutter: The orbit bundle picture of cotangent bundle reduction, C. R. Math. Acad. Sci. Soc. R. Can., 22(2)(2000).
- [29] J.E. Marsden and T.S. Ratiu: Introduction to Mechanics and Symmetry, Texts in Applied Mathematics 17 (1994), second ed., second printing (2003), Springer-Verlag.
- [30] J. Marsden and J. Scheurle: The reduced Euler-Lagrange equations. In: M. J. Enos. ed. Dynamics and Control of mechanical systems. The falling cat and related problems. Papers from the Fields Institute Workshop held in Waterloo, Ontario, March 1992, Fields Institute Communications 1 (1993), 139–164, American Math. Society, Providence, R. I.
- [31] R. Montgomery: The Bundle Picture in Mechanics. Ph.D. Thesis, Department of Mathematics, University of California, Berkeley, 1986.
- [32] R. Montgomery, J.E. Marsden, and T.S. Ratiu: Gauged Lie-Poisson Structures, in *Fluids and Plasmas: Geometry and Dynamics* (J. Marsden, ed.) *Cont. Math.* **28** (1984), 101–114.
- [33] K. Nomizu: Invariant affine connections on homogeneous spaces. Amer. J. of Math. 76 (1954), 33-65.
- [34] J.-P. Ortega and T.S. Ratiu: Momentum Maps and Hamiltonian Reduction, Progress in Mathematics 222 (2004), Birkhäuser-Verlag, Boston.
- [35] R. S. Palais: On the existence of slices for actions of non-compact Lie groups. Ann. of Math. 73(2) (1961), 295–323.
- [36] M. Perlmutter and M. Rodríguez-Olmos: On singular Poisson Sternberg spaces. J. Symplectic Geom. 7(2) (2009), 15–49.
- [37] M. Perlmutter, M. Rodríguez-Olmos, and M. E. Sousa-Dias: On the geometry of reduced cotangent bundles at zero momentum. J. Geom. Phys. 5(2) (2007), 571–596.
- [38] G. Rinehart: Differential forms for general commutative algebras. Trans. Amer. Math. Soc. 108 (1963), 195–222.
- [39] G. W. Schwarz: Smooth functions invariant under the action of a compact Lie group. Topology 14 (1975), 63-68.
- [40] R. Sjamaar and E. Lerman: Stratified symplectic spaces and reduction. Ann. of Math. 134(2) (1991), 375–422.
- [41] S. Sternberg, On minimal coupling and the symplectic mechanics of a classical particle in the presence of a Yang-Mills field, *Proc. Nat. Acad. Sci.*, **74** (1977), 5253–5254.
- [42] I. Vaisman: Lectures on the Geometry of Poisson Manifolds. Progress in Mathematics 118, Birkhäuser-Verlag, Boston,
- [43] A. Weinstein, A universal phase space for particles in Yang-Mills fields. Lett. Math. Phys. 2(5) (1977/78), 417-420
- [44] N. Zaalani: Phase space reduction and Poisson structure. J. Math. Phys. 40 (1999), 3431–3438.

(JH)USTL, UFR DE MATHÉMATIQUES, CNRS-UMR 8524, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE E-mail address: Johannes.Huebschmann@math.univ-lille1.fr

(MP)DEPARTAMENTO DE MATEMATICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, BRASIL E-mail address: matthew@mat.ufmg.br

(TSR)SECTION DE MATHÉMATIQUES AND BERNOULLI CENTER, ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, CH-1015 LAUSANNE, SWITZERLAND

 $E ext{-}mail\ address: tudor.ratiu@epfl.ch}$