

A statistical mechanical interpretation of algorithmic information theory

Kohtaro Tadaki

Research and Development Initiative, Chuo University
1-13-27 Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan
E-mail: tadaki@kc.chuo-u.ac.jp

Abstract. We develop a statistical mechanical interpretation of algorithmic information theory by introducing the notion of thermodynamic quantities, such as free energy, energy, statistical mechanical entropy, and specific heat, into algorithmic information theory. We investigate the properties of these quantities by means of program-size complexity from the point of view of algorithmic randomness. It is then discovered that, in the interpretation, the temperature plays a role as the compression rate of the values of all these thermodynamic quantities, which include the temperature itself. Reflecting this self-referential nature of the compression rate of the temperature, we obtain fixed point theorems on compression rate.

Key words: algorithmic information theory, algorithmic randomness, Chaitin's Ω , compression rate, fixed point theorem, statistical mechanics, temperature

1 Introduction

Algorithmic information theory is a framework to apply information-theoretic and probabilistic ideas to recursive function theory. One of the primary concepts of algorithmic information theory is the *program-size complexity* (or *Kolmogorov complexity*) $H(s)$ of a finite binary string s , which is defined as the length of the shortest binary program for the universal self-delimiting Turing machine U to output s . By the definition, $H(s)$ can be thought of as the information content of the individual finite binary string s . In fact, algorithmic information theory has precisely the formal properties of classical information theory (see [4]). The concept of program-size complexity plays a crucial role in characterizing the randomness of a finite or infinite binary string. In [4] Chaitin introduced the halting probability Ω as an example of random infinite string. His Ω is defined as the probability that the universal self-delimiting Turing machine U halts, and plays a central role in the development of algorithmic information theory. The first n bits of the base-two expansion of Ω solves the halting problem for a program of size not greater than n . By this property, the base-two expansion of Ω is shown to be a random infinite binary string. In [5] Chaitin encoded this random property of Ω onto an exponential Diophantine equation in the manner that a certain property of the set of the solutions of the equation is indistinguishable from coin tosses. Moreover, based on this random property of the equation, Chaitin derived several quantitative versions of Gödel's incompleteness theorems.

In [15, 16] we generalized Chaitin's halting probability Ω to Ω^D by

$$\Omega^D = \sum_{p \in \text{dom } U} 2^{-\frac{|p|}{D}}, \quad (1)$$

so that the degree of randomness of Ω^D can be controlled by a real number D with $0 < D \leq 1$. Here, $\text{dom } U$ denotes the set of all programs p for U . As D becomes larger, the degree of randomness of Ω^D increases. When $D = 1$, Ω^D becomes a random real number, i.e., $\Omega^1 = \Omega$. The properties of Ω^D and its relations to self-similar sets were studied in [15, 16].

Recently, Calude and Stay [3] pointed out a formal correspondence between Ω^D and a partition function in statistical mechanics. In statistical mechanics, the partition function $Z(T)$ at temperature T is defined by

$$Z(T) = \sum_{x \in X} e^{-\frac{E_x}{kT}},$$

where X is a complete set of energy eigenstates of a statistical mechanical system and E_x is the energy of an energy eigenstate x . The constant k is called the Boltzmann Constant. The partition function $Z(T)$ is of particular importance in equilibrium statistical mechanics. This is because all the thermodynamic quantities of the system can be expressed by using the partition function $Z(T)$, and the knowledge of $Z(T)$ is sufficient to understand all the macroscopic properties of the system. Calude and Stay [3] pointed out that the partition function $Z(T)$ has the same form as Ω^D by performing the following replacements in $Z(T)$:

Replacements 1.1.

- (i) Replace the complete set X of energy eigenstates x by the set $\text{dom } U$ of all programs p for U .
- (ii) Replace the energy E_x of an energy eigenstate x by the length $|p|$ of a program p .
- (iii) Set the Boltzmann Constant k to $1/\ln 2$, where the \ln denotes the natural logarithm.

In this paper, inspired by their suggestion, we develop a statistical mechanical interpretation of algorithmic information theory, where Ω^D appears as a partition function.

Generally speaking, in order to give a statistical mechanical interpretation to a framework which looks unrelated to statistical mechanics at first glance, it is important to identify a microcanonical ensemble in the framework. Once we can do so, we can easily develop an equilibrium statistical mechanics on the framework according to the theoretical development of normal equilibrium statistical mechanics. Here, the microcanonical ensemble is a certain sort of uniform probability distribution. In fact, in the work [17] we developed a statistical mechanical interpretation of the noiseless source coding scheme in information theory by identifying a microcanonical ensemble in the scheme. Then, in [17] the notions in statistical mechanics such as statistical mechanical entropy, temperature, and thermal equilibrium are translated into the context of noiseless source coding.

Thus, in order to develop a statistical mechanical interpretation of algorithmic information theory, it is appropriate to identify a microcanonical ensemble in the framework of the theory. Note, however, that algorithmic information theory is not a physical theory but a purely mathematical theory. Therefore, in order to obtain significant results for the development of algorithmic information theory itself, we have to develop a statistical mechanical interpretation of algorithmic information theory in a mathematically rigorous manner, unlike in normal statistical mechanics in physics where arguments are not necessarily mathematically rigorous. A fully rigorous mathematical treatment of statistical mechanics is already developed (see [14]). At present, however, it would not as yet seem to be an easy task to merge algorithmic information theory with this mathematical treatment in a satisfactory manner.

On the other hand, if we do not stick to the mathematical strictness of an argument, we can develop a statistical mechanical interpretation of algorithmic information theory while realizing a perfect correspondence to normal statistical mechanics. In fact, in the last part of this paper (i.e., in Section 6) we develop a statistical mechanical interpretation of algorithmic information theory by making an argument on the same level of mathematical strictness as statistical mechanics in physics. There, we identify a microcanonical ensemble in algorithmic information theory in a similar manner to [17], based on the probability measure which gives Chaitin's Ω the meaning of the halting probability actually. In consequence, for example, the statistical mechanical meaning of Ω^D is clarified.

In the main part of this paper, for mathematical strictness we develop a statistical mechanical interpretation of algorithmic information theory in a different way from the physical argument in Section 6.¹ We introduce the notion of thermodynamic quantities into algorithmic information theory based on Replacements 1.1 above. Section 6 plays a role in clarifying the statistical mechanical meaning of these notion and motivating the introduction of them in the main part of this paper.

After the preliminary section on the mathematical notion needed in this paper, we prove some results on the degree of randomness of real numbers in Section 3. These results themselves and the techniques used in proving these results are frequently used throughout this paper.

Then, in Section 4 we introduce the notion of the thermodynamic quantities at any given fixed temperature T , such as partition function, free energy, energy, statistical mechanical entropy, and specific heat, into algorithmic information theory by performing Replacements 1.1 for the corresponding thermodynamic quantities in statistical mechanics. These thermodynamic quantities in algorithmic information theory are real numbers which depend only on the temperature T . We prove that if the temperature T is a computable real number with $0 < T < 1$ then, for each of these thermodynamic quantities, the compression rate by the program-size complexity H is equal to T . Thus, the temperature T plays a role as the compression rate of the thermodynamic quantities in this statistical mechanical interpretation of algorithmic information theory.

Among all thermodynamic quantities in thermodynamics, one of the most typical thermodynamic quantities is temperature itself. Thus, based on the results of Section 4, the following question naturally arises: Can the compression rate of the temperature T be equal to the temperature itself in the statistical mechanical interpretation of algorithmic information theory? This question is rather self-referential. However, in Section 5 we answer it affirmatively by proving Theorem 5.1. One consequence of Theorem 5.1 has the following form: For every $T \in (0, 1)$, if $\Omega^T = \sum_{p \in \text{dom } U} 2^{-\frac{|p|}{T}}$ is a computable real number, then

$$\lim_{n \rightarrow \infty} \frac{H(T_n)}{n} = T,$$

where T_n is the first n bits of the base-two expansion of T . This is just a fixed point theorem on compression rate, which reflects the self-referential nature of the question.

The works [15, 16] on Ω^D might be regarded as an elaboration of the technique used by Chaitin [4] to prove that Ω is random. The mathematical results of this paper, which are obtained except for in Section 6, may be regarded as further elaborations of the technique.

Finally, in Section 6, based on a physical and informal argument we develop a total statistical mechanical interpretation of algorithmic information theory which attains a perfect correspondence

¹ We make an argument in a fully mathematically rigorous manner in this paper except for Section 6. Any consequence of the argument in Section 6 is not used in any other parts of this paper.

to normal statistical mechanics. In consequence, we justify the interpretation of Ω^D as a partition function and clarify the statistical mechanical meaning of the thermodynamic quantities introduced into algorithmic information theory in Section 4.

2 Preliminaries

We start with some notation about numbers and strings which will be used in this paper. $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ is the set of natural numbers, and \mathbb{N}^+ is the set of positive integers. \mathbb{Z} is the set of integers, and \mathbb{Q} is the set of rational numbers. \mathbb{R} is the set of real numbers. $\{0, 1\}^* = \{\lambda, 0, 1, 00, 01, 10, 11, 000, 001, 010, \dots\}$ is the set of finite binary strings where λ denotes the *empty string*. For any $s \in \{0, 1\}^*$, $|s|$ is the *length* of s . A subset S of $\{0, 1\}^*$ is called a *prefix-free set* if no string in S is a prefix of another string in S . $\{0, 1\}^\infty$ is the set of infinite binary strings, where an infinite binary string is infinite to the right but finite to the left. For any $\alpha \in \{0, 1\}^\infty$, α_n is the prefix of α of length n . For any partial function f , the domain of definition of f is denoted by $\text{dom } f$. We write “r.e.” instead of “recursively enumerable.”

Normally, $o(n)$ denotes any one function $f: \mathbb{N}^+ \rightarrow \mathbb{R}$ such that $\lim_{n \rightarrow \infty} f(n)/n = 0$. On the other hand, $O(1)$ denotes any one function $g: \mathbb{N}^+ \rightarrow \mathbb{R}$ such that there is $C \in \mathbb{R}$ with the property that $|g(n)| \leq C$ for all $n \in \mathbb{N}^+$.

Let T be an arbitrary real number. $T \bmod 1$ denotes $T - \lfloor T \rfloor$, where $\lfloor T \rfloor$ is the greatest integer less than or equal to T , and $T \bmod '1$ denotes $T - \lceil T \rceil + 1$, where $\lceil T \rceil$ is the smallest integer greater than or equal to T . Hence, $T \bmod 1 \in [0, 1)$ but $T \bmod '1 \in (0, 1]$. We identify a real number T with the infinite binary string α such that $0.\alpha$ is the base-two expansion of $T \bmod 1$ with infinitely many zeros. Thus, T_n denotes the first n bits of the base-two expansion of $T \bmod 1$ with infinitely many zeros.

We say that a real number T is *computable* if there exists a total recursive function $f: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $|T - f(n)| < 1/n$ for all $n \in \mathbb{N}^+$. We say that T is *right-computable* if there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $T \leq g(n)$ for all $n \in \mathbb{N}^+$ and $\lim_{n \rightarrow \infty} g(n) = T$. We say that T is *left-computable* if $-T$ is right-computable. It is then easy to see that, for any $T \in \mathbb{R}$, T is computable if and only if T is both right-computable and left-computable. See e.g. [11, 19] for the detail of the treatment of the computability of real numbers and real functions on a discrete set.

2.1 Algorithmic information theory

In the following we concisely review some definitions and results of algorithmic information theory [4, 6]. A *computer* is a partial recursive function $C: \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that $\text{dom } C$ is a prefix-free set. For each computer C and each $s \in \{0, 1\}^*$, $H_C(s)$ is defined by $H_C(s) = \min \{ |p| \mid p \in \{0, 1\}^* \ \& \ C(p) = s \}$. A computer U is said to be *optimal* if for each computer C there exists a constant $\text{sim}(C)$ with the following property; if $C(p)$ is defined, then there is a p' for which $U(p') = C(p)$ and $|p'| \leq |p| + \text{sim}(C)$. It is then shown that there exists an optimal computer. We choose any one optimal computer U as the standard one for use, and define $H(s)$ as $H_U(s)$, which is referred to as the *program-size complexity* of s , the *information content* of s , or the *Kolmogorov complexity* of s [9, 10, 4]. Thus, $H(s)$ has the following property:

$$\forall C : \text{computer} \quad H(s) \leq H_C(s) + \text{sim}(C). \quad (2)$$

It can be shown that there is $c \in \mathbb{N}$ such that, for any $s \neq \lambda$,

$$H(s) \leq |s| + 2 \log_2 |s| + c. \quad (3)$$

For each $s \in \{0, 1\}^*$, $P(s)$ is defined as $\sum_{U(p)=s} 2^{-|p|}$. Chaitin's halting probability Ω is defined by

$$\Omega = \sum_{p \in \text{dom } U} 2^{-|p|}.$$

For any $\alpha \in \{0, 1\}^\infty$, we say that α is *weakly Chaitin random* if there exists $c \in \mathbb{N}$ such that $n - c \leq H(\alpha_n)$ for all $n \in \mathbb{N}^+$ [4, 6]. Then [4] showed that Ω is weakly Chaitin random. For any $\alpha \in \{0, 1\}^\infty$, we say that α is *Chaitin random* if $\lim_{n \rightarrow \infty} H(\alpha_n) - n = \infty$ [4, 6]. It is then shown that, for any $\alpha \in \{0, 1\}^\infty$, α is weakly Chaitin random if and only if α is Chaitin random (see [6] for the proof and historical detail). Thus Ω is Chaitin random.

The class of computers is equal to the class of functions which are computed by *self-delimiting Turing machines*. A self-delimiting Turing machine is a deterministic Turing machine which has two tapes, a program tape and a work tape. The program tape is infinite to the right, while the work tape is infinite in both directions. An input string in $\{0, 1\}^*$ is put on the program tape. See Chaitin [4] for the detail of self-delimiting Turing machines. Let M be a self-delimiting Turing machine which computes the optimal computer U . Then $P(s)$ is the probability that M halts and outputs s when M starts on the program tape filled with an infinite binary string generated by infinitely repeated tosses of a fair coin. Therefore $\Omega = \sum_{s \in \{0, 1\}^*} P(s)$ is the probability that M just halts under the same setting.

The program-size complexity $H(s)$ is originally defined using the concept of program-size, as stated above. However, it is possible to define $H(s)$ without referring to such a concept, i.e., we first introduce a *universal probability* m , and then define $H(s)$ as $-\log_2 m(s)$. A universal probability is defined through the following two definitions [20].

Definition 2.1. For any $r: \{0, 1\}^* \rightarrow [0, 1]$, we say that r is a *lower-computable semi-measure* if r satisfies the following two conditions:

(i) $\sum_{s \in \{0, 1\}^*} r(s) \leq 1$.

(ii) There exists a total recursive function $f: \mathbb{N}^+ \times \{0, 1\}^* \rightarrow \mathbb{Q}$ such that, for each $s \in \{0, 1\}^*$, $\lim_{n \rightarrow \infty} f(n, s) = r(s)$ and $\forall n \in \mathbb{N}^+ f(n, s) \leq r(s)$. □

Definition 2.2. Let m be a lower-computable semi-measure. We say that m is a *universal probability* if for any lower-computable semi-measure r , there exists a real number $c > 0$ such that, for all $s \in \{0, 1\}^*$, $cr(s) \leq m(s)$. □

Chaitin [4] showed the following theorem.

Theorem 2.3. Both $2^{-H(s)}$ and $P(s)$ are universal probabilities. □

By Theorem 2.3, we see that, for any universal probability m ,

$$H(s) = -\log_2 m(s) + O(1). \quad (4)$$

Thus it is possible to define $H(s)$ as $-\log_2 m(s)$ with any one universal probability m instead of as $H_U(s)$. Note that the difference up to an additive constant is inessential to algorithmic information theory.

In the works [15, 16], we generalized the notion of the randomness of an infinite binary string so that the degree of the randomness can be characterized by a real number D with $0 < D \leq 1$ as follows.

Definition 2.4 (weakly Chaitin D -random). *Let $D \in \mathbb{R}$ with $D \geq 0$, and let $\alpha \in \{0, 1\}^\infty$. We say that α is weakly Chaitin D -random if there exists $c \in \mathbb{R}$ such that $Dn - c \leq H(\alpha_n)$ for all $n \in \mathbb{N}^+$. \square*

Definition 2.5 (D -compressible). *Let $D \in \mathbb{R}$ with $D \geq 0$, and let $\alpha \in \{0, 1\}^\infty$. We say that α is D -compressible if $H(\alpha_n) \leq Dn + o(n)$, which is equivalent to*

$$\overline{\lim}_{n \rightarrow \infty} \frac{H(\alpha_n)}{n} \leq D.$$

\square

In the case of $D = 1$, the weak Chaitin D -randomness results in the weak Chaitin randomness. For any $D \in [0, 1]$ and any $\alpha \in \{0, 1\}^\infty$, if α is weakly Chaitin D -random and D -compressible, then

$$\lim_{n \rightarrow \infty} \frac{H(\alpha_n)}{n} = D. \quad (5)$$

Hereafter the left-hand side of (5) is referred to as the *compression rate* of an infinite binary string α in general. Note, however, that (5) does not necessarily imply that α is weakly Chaitin D -random.

In the works [15, 16], we generalized Chaitin's halting probability Ω to Ω^D by (1) for any real number $D > 0$. Thus, $\Omega = \Omega^1$. If $0 < D \leq 1$, then Ω^D converges and $0 < \Omega^D < 1$, since $\Omega^D \leq \Omega < 1$.

Theorem 2.6 (Tadaki [15, 16]). *Let $D \in \mathbb{R}$.*

(i) *If $0 < D \leq 1$ and D is computable, then Ω^D is weakly Chaitin D -random and D -compressible.*

(ii) *If $1 < D$, then Ω^D diverges to ∞ . \square*

Definition 2.7 (Chaitin D -randomness, Tadaki [15, 16]). *Let $D \in \mathbb{R}$ with $D \geq 0$, and let $\alpha \in \{0, 1\}^\infty$. We say that α is Chaitin D -random if $\lim_{n \rightarrow \infty} H(\alpha_n) - Dn = \infty$. \square*

In the case of $D = 1$, the Chaitin D -randomness results in the Chaitin randomness. Obviously, for any $D \in [0, 1]$ and any $\alpha \in \{0, 1\}^\infty$, if α is Chaitin D -random, then α is weakly Chaitin D -random. However, in 2005 Reimann and Stephan [13] showed that, in the case of $D < 1$, the converse does not necessarily hold. This contrasts with the equivalence between the weakly Chaitin randomness and the Chaitin randomness, each of which corresponds to the case of $D = 1$. In the next section, for any computable real number D with $0 < D < 1$, we give an instance of a real number which is Chaitin D -random and D -compressible.

3 Chaitin D -randomness and divergence

For each real numbers $Q > 0$ and $D > 0$, we define $W(Q, D)$ by

$$W(Q, D) = \sum_{p \in \text{dom } U} |p|^Q 2^{-\frac{|p|}{D}}.$$

As the first result of this paper, we show the following theorem.

Theorem 3.1. *Let Q and D be positive real numbers.*

(i) *If Q and D are computable and $0 < D < 1$, then $W(Q, D)$ converges to a left-computable real number which is Chaitin D -random and D -compressible.*

(ii) *If $1 \leq D$, then $W(Q, D)$ diverges to ∞ .* □

The techniques used in the proofs of Theorem 2.6 (see [16]) and Theorem 3.1 are frequently used throughout the rest of this paper as basic tools. We see that the weak Chaitin D -randomness in Theorem 2.6 is replaced by the Chaitin D -randomness in Theorem 3.1 in exchange for the divergence at $D = 1$. In order to derive this divergence we make use of Theorem 3.2 (i) below. We prove Theorem 3.2 in a more general form, and show that the *Shannon entropy* $-\sum_{s \in \{0,1\}^*} m(s) \log_2 m(s)$ of an arbitrary universal probability m diverges to ∞ . We say that a function $f: \mathbb{N}^+ \rightarrow [0, \infty)$ is *lower-computable* if there exists a total recursive function $a: \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that, for each $n \in \mathbb{N}^+$, $\lim_{k \rightarrow \infty} a(k, n) = f(n)$ and $\forall k \in \mathbb{N}^+ a(k, n) \leq f(n)$.

Theorem 3.2. *Let A be an infinite r.e. subset of $\{0, 1\}^*$ and let $f: \mathbb{N}^+ \rightarrow [0, \infty)$ be a lower-computable function such that $\lim_{n \rightarrow \infty} f(n) = \infty$. Then the following hold.*

(i) $\sum_{U(p) \in A} f(|p|)2^{-|p|}$ *diverges to ∞ .*

(ii) *If there exists $l_0 \in \mathbb{N}^+$ such that $f(l)2^{-l}$ is a nonincreasing function of l for all $l \geq l_0$, then $\sum_{s \in A} f(H(s))2^{-H(s)}$ diverges to ∞ .*

Proof. (i) Contrarily, assume that $\sum_{U(p) \in A} f(|p|)2^{-|p|}$ converges. Then, there exists $d \in \mathbb{N}^+$ such that $\sum_{U(p) \in A} f(|p|)2^{-|p|} \leq d$. We define the function $r: \{0, 1\}^* \rightarrow [0, \infty)$ by

$$r(s) = \frac{1}{d} \sum_{U(p)=s} f(|p|)2^{-|p|}$$

if $s \in A$; $r(s) = 0$ otherwise. Then we see that $\sum_{s \in \{0,1\}^*} r(s) \leq 1$ and therefore r is a lower-computable semi-measure. Since $P(s)$ is a universal probability, there exists $c \in \mathbb{N}^+$ such that $r(s) \leq cP(s)$ for all $s \in \{0, 1\}^*$. Hence we have

$$\sum_{U(p)=s} (cd - f(|p|)2^{-|p|}) \geq 0 \tag{6}$$

for all $s \in A$. On the other hand, since A is an infinite set and $\lim_{n \rightarrow \infty} f(n) = \infty$, there is $s_0 \in A$ such that $f(|p|) > cd$ for all p with $U(p) = s_0$. Therefore we have $\sum_{U(p)=s_0} (cd - f(|p|)2^{-|p|}) < 0$. However, this contradicts (6), and the proof of (i) is completed.

(ii) We first note that there is $n_0 \in \mathbb{N}$ such that $H(s) \geq l_0$ for all s with $|s| \geq n_0$. Now, let us assume contrarily that $\sum_{s \in A} f(H(s))2^{-H(s)}$ converges. Then, there exists $d \in \mathbb{N}^+$ such that $\sum_{s \in A} f(H(s))2^{-H(s)} \leq d$. We define the function $r: \{0, 1\}^* \rightarrow [0, \infty)$ by

$$r(s) = \frac{1}{d} f(H(s))2^{-H(s)}$$

if $s \in A$ and $|s| \geq n_0$; $r(s) = 0$ otherwise. Then we see that $\sum_{s \in \{0,1\}^*} r(s) \leq 1$ and therefore r is a lower-computable semi-measure. Since $2^{-H(s)}$ is a universal probability by Theorem 2.3, there

exists $c \in \mathbb{N}^+$ such that $r(s) \leq c2^{-H(s)}$ for all $s \in \{0, 1\}^*$. Hence, if $s \in A$ and $|s| \geq n_0$, then $cd \geq f(H(s))$. On the other hand, since A is an infinite set and $\lim_{n \rightarrow \infty} f(n) = \infty$, there is $s_0 \in A$ such that $|s_0| \geq n_0$ and $f(H(s_0)) > cd$. Thus, we have a contradiction, and the proof of (ii) is completed. \square

Corollary 3.3. *If m is a universal probability and A is an infinite r.e. subset of $\{0, 1\}^*$, then $-\sum_{s \in A} m(s) \log_2 m(s)$ diverges to ∞ .*

Proof. We first note that there is a real number $x_0 > 0$ such that the function $x2^{-x}$ of a real number x is decreasing for $x \geq x_0$. For this x_0 , there is $n_0 \in \mathbb{N}$ such that $-\log_2 m(s) \geq x_0$ for all s with $|s| \geq n_0$. On the other hand, by (4), there is $c \in \mathbb{N}$ such that $-\log_2 m(s) \leq H(s) + c$ for all $s \in \{0, 1\}^*$. Thus, we see that

$$\begin{aligned} -\sum_{s \in A \text{ \& } |s| \geq n_0} m(s) \log_2 m(s) &\geq \sum_{s \in A \text{ \& } |s| \geq n_0} (H(s) + c)2^{-H(s)-c} \\ &= 2^{-c} \sum_{s \in A \text{ \& } |s| \geq n_0} H(s)2^{-H(s)} + c2^{-c} \sum_{s \in A \text{ \& } |s| \geq n_0} 2^{-H(s)}. \end{aligned}$$

By Theorem 3.2 (ii), $\sum_{s \in A} H(s)2^{-H(s)}$ diverges to ∞ . Hence, we see, by the inequality above, that $-\sum_{s \in A} m(s) \log_2 m(s)$ also diverges to ∞ . \square

By Corollary 3.3, we see that the Shannon entropy of an arbitrary universal probability diverges to ∞ .

The proof of Theorem 3.1 is given as follows.

The proof of Theorem 3.1. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. Then, for every $D > 0$, $W(Q, D) = \lim_{m \rightarrow \infty} \widetilde{W}_m(Q, D)$, where

$$\widetilde{W}_m(Q, D) = \sum_{i=1}^m |p_i|^Q 2^{-\frac{|p_i|}{D}}.$$

(i) First we show that $W(Q, D)$ converges to a left-computable real number. Since $D < 1$, there is $l_0 \in \mathbb{N}^+$ such that

$$\frac{1}{D} - \frac{Q \log_2 l}{l} \geq 1$$

for all $l \geq l_0$. Then there is $m_0 \in \mathbb{N}^+$ such that $|p_i| \geq l_0$ for all $i > m_0$. Thus, we see that, for each $i > m_0$,

$$|p_i|^Q 2^{-\frac{|p_i|}{D}} = 2^{-\left(\frac{1}{D} - \frac{Q \log_2 |p_i|}{|p_i|}\right)|p_i|} \leq 2^{-|p_i|}.$$

Hence, for each $m > m_0$,

$$\widetilde{W}_m(Q, D) - \widetilde{W}_{m_0}(Q, D) = \sum_{i=m_0+1}^m |p_i|^Q 2^{-\frac{|p_i|}{D}} \leq \sum_{i=m_0}^m 2^{-|p_i|} < \Omega.$$

Thus, since $\{\widetilde{W}_m(Q, D)\}_m$ is an increasing sequence of real numbers, it converges to a real number $W(Q, D)$ as $m \rightarrow \infty$. Moreover, since Q and D are computable, $W(Q, D)$ is shown to be left-computable.

We then show that $W(Q, D)$ is Chaitin D -random. Let α be the infinite binary string such that $0.\alpha$ is the base-two expansion of $W(Q, D) \bmod '1$ with infinitely many ones. Then, since Q and D are computable real numbers and $\lceil W(Q, D) \rceil - 1 + 0.\alpha_n < W(Q, D)$ for all $n \in \mathbb{N}^+$, there exists a partial recursive function $\xi: \{0, 1\}^* \rightarrow \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$,

$$\lceil W(Q, D) \rceil - 1 + 0.\alpha_n < \widetilde{W}_{\xi(\alpha_n)}(Q, D).$$

It is then easy to see that

$$W(Q, D) - \widetilde{W}_{\xi(\alpha_n)}(Q, D) < 2^{-n}.$$

It follows that, for all $i > \xi(\alpha_n)$, $|p_i|^Q 2^{-\frac{|p_i|}{D}} < 2^{-n}$ and therefore $QD \log_2 |p_i| < |p_i| - Dn$. Thus, given α_n , by calculating the set $\{U(p_i) \mid i \leq \xi(\alpha_n)\}$ and picking any one finite binary string which is not in this set, one can obtain $s \in \{0, 1\}^*$ such that $QD \log_2 H(s) < H(s) - Dn$.

Hence, there exists a partial recursive function $\Psi: \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$QD \log_2 H(\Psi(\alpha_n)) < H(\Psi(\alpha_n)) - Dn.$$

Applying this inequality to itself, we have $QD \log_2 n < H(\Psi(\alpha_n)) - Dn + O(1)$. On the other hand, using (2) there is a natural number c_Ψ such that $H(\Psi(\alpha_n)) < H(\alpha_n) + c_\Psi$. Therefore, we have

$$QD \log_2 n < H(\alpha_n) - Dn + O(1).$$

Hence, α is Chaitin T -random. It follows that α has infinitely many zeros, which implies that $W(Q, D) \bmod 1 = W(Q, D) \bmod '1 = 0.\alpha$ and therefore $(W(Q, D))_n = \alpha_n$. Thus, $W(Q, D)$ is Chaitin D -random.

Next, we show that $W(Q, D)$ is D -compressible. Since Q and D are computable real numbers, there exists a total recursive function $g: \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow \mathbb{Z}$ such that, for all $m, n \in \mathbb{N}^+$,

$$\left| \widetilde{W}_m(Q, D) - \lceil W(Q, D) \rceil - 2^{-n}g(m, n) \right| < 2^{-n}. \quad (7)$$

Let d be any computable real number with $D < d < 1$. Then, the limit value $W(Q, d)$ exists since $d < 1$. Let β be the infinite binary string such that $0.\beta$ is the base-two expansion of $W(Q, d) \bmod '1$ with infinitely many ones.

Given n and $\beta_{\lceil Dn/d \rceil}$ (i.e., the first $\lceil Dn/d \rceil$ bits of β), one can find $m_0 \in \mathbb{N}^+$ such that

$$\lceil W(Q, d) \rceil - 1 + 0.\beta_{\lceil Dn/d \rceil} < \widetilde{W}_{m_0}(Q, d).$$

This is possible since $\lceil W(Q, d) \rceil - 1 + 0.\beta_{\lceil Dn/d \rceil} < W(Q, d)$ and $\lim_{m \rightarrow \infty} \widetilde{W}_m(Q, d) = W(Q, d)$. It is then easy to see that

$$\sum_{i=m_0+1}^{\infty} |p_i|^Q 2^{-\frac{|p_i|}{d}} < 2^{-Dn/d}.$$

Raising both sides of this inequality to the power d/D and using the inequality $a^c + b^c \leq (a+b)^c$ for real numbers $a, b > 0$ and $c \geq 1$,

$$2^{-n} > \sum_{i=m_0+1}^{\infty} |p_i|^{Qd/D} 2^{-\frac{|p_i|}{D}} > \sum_{i=m_0+1}^{\infty} |p_i|^Q 2^{-\frac{|p_i|}{D}}.$$

It follows that

$$\left| W(Q, D) - \widetilde{W}_{m_0}(Q, D) \right| < 2^{-n}. \quad (8)$$

From (7), (8), and

$$| \lfloor W(Q, D) \rfloor + 0.(W(Q, D))_n - W(Q, D) | < 2^{-n},$$

it is shown that

$$| (W(Q, D))_n - g(m_0, n) | < 3$$

and therefore

$$(W(Q, D))_n = g(m_0, n), g(m_0, n) \pm 1, g(m_0, n) \pm 2,$$

where $(W(Q, D))_n$ is regarded as a dyadic integer. Thus, there are still 5 possibilities of $(W(Q, D))_n$, so that one needs only 3 bits more in order to determine $(W(Q, D))_n$.

Thus, there exists a partial recursive function $\Phi: \mathbb{N}^+ \times \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = 3 \quad \& \quad \Phi(n, \beta_{\lceil Dn/d \rceil}, s) = (W(Q, D))_n.$$

It follows from (3) that $H((W(Q, D))_n) \leq |\beta_{\lceil Dn/d \rceil}| + o(n) \leq Dn/d + o(n)$, which implies that $W(Q, D)$ is D/d -compressible. Since d is any computable real number with $D < d < 1$, it follows that $W(Q, D)$ is D -compressible.

(ii) We choose any one computable real number Q' with $Q \geq Q' > 0$. Then, using Theorem 3.2 (i), we can show that $W(Q', 1) = \sum_{p \in \text{dom } U} |p|^{Q'} 2^{-|p|}$ diverges to ∞ . Thus, since $\widetilde{W}_m(Q, D) \geq \widetilde{W}_m(Q', 1)$, we see that $W(Q, D)$ also diverges to ∞ , and the proof is completed. \square

4 Temperature as a compression rate

In this section we introduce the notion of thermodynamic quantities such as partition function, free energy, energy, entropy, and specific heat, into algorithmic information theory by performing Replacements 1.1 for the corresponding thermodynamic quantities in statistical mechanics.² We investigate their convergence and the degree of randomness. For that purpose, we first choose any one enumeration q_1, q_2, q_3, \dots of the countably infinite set $\text{dom } U$ as the standard one for use throughout this section.³

In statistical mechanics, the partition function $Z_{\text{sm}}(T)$ at temperature T is given by

$$Z_{\text{sm}}(T) = \sum_{x \in X} e^{-\frac{E_x}{kT}}. \quad (9)$$

Motivated by the formula (9) and taking into account Replacements 1.1, we introduce the notion of partition function into algorithmic information theory as follows.

Definition 4.1 (partition function). *For each $n \in \mathbb{N}^+$ and each real number $T > 0$, we define $Z_n(T)$ by*

$$Z_n(T) = \sum_{i=1}^n 2^{-\frac{|q_i|}{T}}.$$

² For the thermodynamic quantities in statistical mechanics, see Chapter 16 of [1] and Chapter 2 of [18]. To be precise, the partition function is not a thermodynamic quantity but a statistical mechanical quantity.

³ The enumeration $\{q_i\}$ is quite arbitrary and therefore we do not, ever, require $\{q_i\}$ to be a recursive enumeration of $\text{dom } U$.

Then, for each $T > 0$, the partition function $Z(T)$ is defined by $Z(T) = \lim_{n \rightarrow \infty} Z_n(T)$. \square

Since $Z(T) = \Omega^T$, we restate Theorem 2.6 as in the following form.

Theorem 4.2 (Tadaki [15, 16]). *Let $T \in \mathbb{R}$.*

(i) *If $0 < T \leq 1$ and T is computable, then $Z(T)$ converges to a left-computable real number which is weakly Chaitin T -random and T -compressible.*

(ii) *If $1 < T$, then $Z(T)$ diverges to ∞ .* \square

In statistical mechanics, the free energy $F_{\text{sm}}(T)$ at temperature T is given by

$$F_{\text{sm}}(T) = -kT \ln Z_{\text{sm}}(T), \quad (10)$$

where $Z_{\text{sm}}(T)$ is given by (9). Motivated by the formula (10) and taking into account Replacements 1.1, we introduce the notion of free energy into algorithmic information theory as follows.

Definition 4.3 (free energy). *For each $n \in \mathbb{N}^+$ and each real number $T > 0$, we define $F_n(T)$ by*

$$F_n(T) = -T \log_2 Z_n(T).$$

Then, for each $T > 0$, the free energy $F(T)$ is defined by $F(T) = \lim_{n \rightarrow \infty} F_n(T)$. \square

Theorem 4.4. *Let $T \in \mathbb{R}$.*

(i) *If $0 < T \leq 1$ and T is computable, then $F(T)$ converges to a right-computable real number which is weakly Chaitin T -random and T -compressible.*

(ii) *If $1 < T$, then $F(T)$ diverges to $-\infty$.*

Proof. (i) Since $Z(T)$ converges by Theorem 4.2 (i) and $Z(T) > 0$, $F(T)$ also converges and

$$F(T) = -T \log_2 Z(T).$$

Note that T is a right-computable real number and $-\log_2 Z(T) > 0$. Since $Z(T)$ is a left-computable real number by Theorem 4.2 (i), $F(T)$ is a right-computable real number.

We show that $F(T)$ is weakly Chaitin T -random. By the mean value theorem, there exists $c \in \mathbb{N}^+$ such that, for any $A, B \in \mathbb{R}$, if $A \geq F(T)$ and $B \geq 1/T$ then

$$0 \leq Z(T) - 2^{-AB} \leq 2^c \max\{A - F(T), B - 1/T\}. \quad (11)$$

Since $F(T)$ is a right-computable real number, there exists a total recursive function $f: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $F(T) \leq f(m)$ for all $m \in \mathbb{N}^+$ and $\lim_{m \rightarrow \infty} f(m) = F(T)$. Since T is a computable real number, there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $0 \leq g(n) - 1/T < 2^{-n}$ for all $n \in \mathbb{N}^+$.

Given $(F(T))_n$, one can find $m_0 \in \mathbb{N}^+$ such that

$$f(m_0) < \lfloor F(T) \rfloor + 0.(F(T))_n + 2^{-n}.$$

This is possible because $F(T) < \lfloor F(T) \rfloor + 0.(F(T))_n + 2^{-n}$. It follows that $0 \leq f(m_0) - F(T) < 2^{-n}$. Therefore, by (11) it is shown that $0 \leq Z(T) - 2^{-f(m_0)g(n)} < 2^{c-n}$. Let l_n be the first n bits of the base-two expansion of $2^{-f(m_0)g(n)}$ with infinitely many zeros. It follows that

$$0 \leq 0.(Z(T))_n - 0.l_n < Z(T) - 2^{-f(m_0)g(n)} + 2^{-n} < (2^c + 1)2^{-n}.$$

Hence

$$(Z(T))_n = l_n, l_n + 1, l_n + 2, \dots, l_n + (2^c + 1),$$

where $(Z(T))_n$ and l_n are regarded as a dyadic integer. Therefore, there are still $2^c + 2$ possibilities of $(Z(T))_n$, so that one needs only $c + 1$ bits more in order to determine $(Z(T))_n$.

Thus, there exists a partial recursive function $\Phi: \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = c + 1 \quad \& \quad \Phi((F(T))_n, s) = (Z(T))_n.$$

It follows that there exists $c_\Phi \in \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$,

$$H((Z(T))_n) \leq H((F(T))_n) + c_\Phi.$$

Hence, $F(T)$ is weakly Chaitin T -random by Theorem 4.2 (i).

Next, we show that $F(T)$ is T -compressible. Since T is a computable real number, there exists a total recursive function $a: \{0, 1\}^* \times \mathbb{N}^+ \rightarrow \mathbb{Z}$ such that, for all $s \in \{0, 1\}^*$ and all $n \in \mathbb{N}^+$, if $0.s > 0$ then

$$| -T \log_2 0.s - \lfloor F(T) \rfloor - 2^{-n} a(s, n) | < 2^{-n}. \quad (12)$$

By the mean value theorem, it is also shown that there is $d \in \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$, if $0.(Z(T))_n > 0$ then

$$| -T \log_2 0.(Z(T))_n - F(T) | < 2^{d-n}. \quad (13)$$

From (12), (13), and

$$| \lfloor F(T) \rfloor + 0.(F(T))_n - F(T) | < 2^{-n},$$

it is shown that, for all $n \in \mathbb{N}^+$, if $0.(Z(T))_n > 0$ then

$$| (F(T))_n - a((Z(T))_n, n) | < 2^d + 2$$

and therefore

$$(F(T))_n = a((Z(T))_n, n), a((Z(T))_n, n) \pm 1, a((Z(T))_n, n) \pm 2, \dots, \\ a((Z(T))_n, n) \pm (2^d + 1),$$

where $(F(T))_n$ is regarded as a dyadic integer. Therefore, there are still $2^{d+1} + 3$ possibilities of $(F(T))_n$, so that one needs only $d + 2$ bits more in order to determine $(F(T))_n$.

Thus, there exists a partial recursive function $\Psi: \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = d + 2 \quad \& \quad \Psi((Z(T))_n, s) = (F(T))_n.$$

It follows that there exists $c_\Psi \in \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$,

$$H((F(T))_n) \leq H((Z(T))_n) + c_\Psi.$$

Since $Z(T)$ is T -compressible by Theorem 4.2 (i), $F(T)$ is also T -compressible.

(ii) In the case of $T > 1$, since $\lim_{n \rightarrow \infty} Z_n(T) = \infty$ by Theorem 4.2 (ii), we see that $F_n(T)$ diverges to $-\infty$ as $n \rightarrow \infty$. \square

In statistical mechanics, the energy $E_{\text{sm}}(T)$ at temperature T is given by

$$E_{\text{sm}}(T) = \frac{1}{Z_{\text{sm}}(T)} \sum_{x \in X} E_x e^{-\frac{E_x}{kT}}, \quad (14)$$

where $Z_{\text{sm}}(T)$ is given by (9). Motivated by the formula (14) and taking into account Replacements 1.1, we introduce the notion of energy into algorithmic information theory as follows.

Definition 4.5 (energy). *For each $n \in \mathbb{N}^+$ and each real number $T > 0$, we define $E_n(T)$ by*

$$E_n(T) = \frac{1}{Z_n(T)} \sum_{i=1}^n |q_i| 2^{-\frac{|q_i|}{T}}.$$

Then, for each $T > 0$, the energy $E(T)$ is defined by $E(T) = \lim_{n \rightarrow \infty} E_n(T)$. \square

Theorem 4.6. *Let $T \in \mathbb{R}$.*

(i) *If $0 < T < 1$ and T is computable, then $E(T)$ converges to a left-computable real number which is Chaitin T -random and T -compressible.*

(ii) *If $1 \leq T$, then $E(T)$ diverges to ∞ .*

Proof. (i) First we show that $E(T)$ converges. By Theorem 4.2 (i), the denominator $Z_n(T)$ of $E_n(T)$ converges to the real number $Z(T) > 0$ as $n \rightarrow \infty$. On the other hand, by Theorem 3.1 (i), the numerator $\sum_{i=1}^n |q_i| 2^{-\frac{|q_i|}{T}}$ of $E_n(T)$ converges to the real number $W(1, T)$ as $n \rightarrow \infty$. Thus, $E_n(T)$ converges to the real number $W(1, T)/Z(T)$ as $n \rightarrow \infty$.

Next, we show that $E(T)$ is a left-computable real number. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $m \in \mathbb{N}^+$, we define $\widetilde{W}_m(T)$ and $\widetilde{Z}_m(T)$ by

$$\widetilde{W}_m(T) = \sum_{i=1}^m |p_i| 2^{-\frac{|p_i|}{T}} \quad \text{and} \quad \widetilde{Z}_m(T) = \sum_{i=1}^m 2^{-\frac{|p_i|}{T}},$$

and then define $\widetilde{E}_m(T)$ by $\widetilde{E}_m(T) = \widetilde{W}_m(T)/\widetilde{Z}_m(T)$. Since the numerator and the denominator of $E_n(T)$ are positive term series which converge as $n \rightarrow \infty$, we see that $W(1, T) = \lim_{m \rightarrow \infty} \widetilde{W}_m(T)$, $Z(T) = \lim_{m \rightarrow \infty} \widetilde{Z}_m(T)$, and $E(T) = \lim_{m \rightarrow \infty} \widetilde{E}_m(T)$. We then see that

$$\widetilde{E}_{m+1}(T) - \widetilde{E}_m(T) = \frac{\widetilde{Z}_m(T) |p_{m+1}| - \widetilde{W}_m(T)}{\widetilde{Z}_{m+1}(T) \widetilde{Z}_m(T)} 2^{-\frac{|p_{m+1}|}{T}}.$$

Since $\widetilde{W}_m(T)$ and $\widetilde{Z}_m(T)$ converge as $m \rightarrow \infty$ and $\lim_{m \rightarrow \infty} |p_m| = \infty$, there exist $a \in \mathbb{N}^+$ and $m_1 \in \mathbb{N}^+$ such that, for any $m \geq m_1$,

$$\widetilde{E}_{m+1}(T) - \widetilde{E}_m(T) > |p_{m+1}| 2^{-\frac{|p_{m+1}|}{T} - a}. \quad (15)$$

In particular, $\widetilde{E}_m(T)$ is an increasing function of m for all $m \geq m_1$ by the above inequality. Thus, since T is a computable real number, $E(T)$ is shown to be a left-computable real number.

We then show that $E(T)$ is Chaitin T -random. Let α be the infinite binary string such that $0.\alpha$ is the base-two expansion of $E(T) \bmod' 1$ with infinitely many ones. Then, since T is a computable

real number and $\lceil E(T) \rceil - 1 + 0.\alpha_n < E(T)$ for all $n \in \mathbb{N}^+$, there exists a partial recursive function $\xi: \{0, 1\}^* \rightarrow \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$, $\xi(\alpha_n) \geq m_1$ and

$$\lceil E(T) \rceil - 1 + 0.\alpha_n < \tilde{E}_{\xi(\alpha_n)}(T).$$

It is then easy to see that $E(T) - \tilde{E}_{\xi(\alpha_n)}(T) < 2^{-n}$. It follows from (15) that, for all $i > \xi(\alpha_n)$, $|p_i| 2^{-\frac{|p_i|}{T}-a} < 2^{-n}$ and therefore $T \log_2 |p_i| - Ta < |p_i| - Tn$. Thus, given α_n , by calculating the set $\{U(p_i) \mid i \leq \xi(\alpha_n)\}$ and picking any one finite binary string which is not in this set, one can obtain $s \in \{0, 1\}^*$ such that $T \log_2 H(s) - Ta < H(s) - Tn$.

Hence, there exists a partial recursive function $\Psi: \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$T \log_2 H(\Psi(\alpha_n)) - Ta < H(\Psi(\alpha_n)) - Tn.$$

Applying this inequality to itself, we have $T \log_2 n < H(\Psi(\alpha_n)) - Tn + O(1)$. On the other hand, using (2) there is a natural number c_Ψ such that $H(\Psi(\alpha_n)) < H(\alpha_n) + c_\Psi$. Therefore, we have

$$T \log_2 n < H(\alpha_n) - Tn + O(1).$$

Hence, α is Chaitin T -random. It follows that α has infinitely many zeros, which implies that $E(T) \bmod 1 = E(T) \bmod '1 = 0.\alpha$ and therefore $(E(T))_n = \alpha_n$. Thus, $E(T)$ is Chaitin T -random.

Next, we show that $E(T)$ is T -compressible. Since T is a computable real number, there exists a total recursive function $g: \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow \mathbb{Z}$ such that, for all $m, n \in \mathbb{N}^+$,

$$\left| \tilde{E}_m(T) - \lceil E(T) \rceil - 2^{-n}g(m, n) \right| < 2^{-n}. \quad (16)$$

It is also shown that there is $c \in \mathbb{N}^+$ such that, for all $m \in \mathbb{N}^+$,

$$\left| E(T) - \tilde{E}_m(T) \right| < 2^c \max \left\{ \left| W(1, T) - \tilde{W}_m(T) \right|, \left| Z(T) - \tilde{Z}_m(T) \right| \right\}. \quad (17)$$

Let t be any computable real number with $T < t < 1$, Then, $W(1, t) = \lim_{m \rightarrow \infty} \tilde{W}_m(t)$, where

$$\tilde{W}_m(t) = \sum_{i=1}^m |p_i| 2^{-\frac{|p_i|}{t}}.$$

The limit value $W(1, t)$ exists since $t < 1$. Let β be the infinite binary string such that $0.\beta$ is the base-two expansion of $W(1, t) \bmod '1$ with infinitely many ones.

Given n and $\beta_{\lceil Tn/t \rceil}$ (i.e., the first $\lceil Tn/t \rceil$ bits of β), one can find $m_0 \in \mathbb{N}^+$ such that

$$\lceil W(1, t) \rceil - 1 + 0.\beta_{\lceil Tn/t \rceil} < \tilde{W}_{m_0}(t).$$

This is possible since t is a computable real number and $\lceil W(1, t) \rceil - 1 + 0.\beta_{\lceil Tn/t \rceil} < W(1, t)$. It is then easy to see that

$$\sum_{i=m_0+1}^{\infty} |p_i| 2^{-\frac{|p_i|}{t}} < 2^{-Tn/t}.$$

Raising both sides of this inequality to the power t/T and using the inequality $a^c + b^c \leq (a + b)^c$ for real numbers $a, b > 0$ and $c \geq 1$,

$$2^{-n} > \sum_{i=m_0+1}^{\infty} |p_i|^{t/T} 2^{-\frac{|p_i|}{T}} > \sum_{i=m_0+1}^{\infty} |p_i| 2^{-\frac{|p_i|}{T}}$$

and therefore

$$2^{-n} > \sum_{i=m_0+1}^{\infty} 2^{-\frac{|p_i|}{T}}.$$

It follows that

$$\left| W(1, T) - \widetilde{W}_{m_0}(T) \right| < 2^{-n} \quad \text{and} \quad \left| Z(T) - \widetilde{Z}_{m_0}(T) \right| < 2^{-n}. \quad (18)$$

From (16), (17), (18), and

$$| \lfloor E(T) \rfloor + 0.(E(T))_n - E(T) | < 2^{-n},$$

it is shown that

$$| (E(T))_n - g(m_0, n) | < 2^c + 2$$

and therefore

$$(E(T))_n = g(m_0, n), g(m_0, n) \pm 1, g(m_0, n) \pm 2, \dots, g(m_0, n) \pm (2^c + 1),$$

where $(E(T))_n$ is regarded as a dyadic integer. Thus, there are still $2^{c+1} + 3$ possibilities of $(E(T))_n$, so that one needs only $c + 2$ bits more in order to determine $(E(T))_n$.

Thus, there exists a partial recursive function $\Phi: \mathbb{N}^+ \times \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = c + 2 \quad \& \quad \Phi(n, \beta_{\lceil Tn/t \rceil}, s) = (E(T))_n.$$

It follows from (3) that $H((E(T))_n) \leq |\beta_{\lceil Tn/t \rceil}| + o(n) \leq Tn/t + o(n)$, which implies that $E(T)$ is T/t -compressible. Since t is any computable real number with $T < t < 1$, it follows that $E(T)$ is T -compressible.

(ii) In the case of $T = 1$, by Theorem 3.1 (ii), the numerator $W(1, 1) = \sum_{p \in \text{dom } U} |p| 2^{-|p|}$ of $E(1)$ diverges to ∞ . On the other hand, the denominator $Z(1)$ of $E(1)$ converges. Thus, $E(1)$ diverges to ∞ .

The case of $T > 1$ is treated as follows. We note that $\lim_{n \rightarrow \infty} |q_n| = \infty$. Given $M > 0$, there is $n_0 \in \mathbb{N}^+$ such that $|q_i| \geq 2M$ for all $i > n_0$. Since $\lim_{n \rightarrow \infty} Z_n(T) = \infty$ by Theorem 4.2 (ii), there is $n_1 \in \mathbb{N}^+$ such that

$$\frac{1}{Z_n(T)} \sum_{i=1}^{n_0} 2^{-\frac{|q_i|}{T}} \leq \frac{1}{2}$$

for all $n > n_1$. Thus, for every $n > \max\{n_0, n_1\}$,

$$\begin{aligned}
E_n(T) &= \frac{1}{Z_n(T)} \sum_{i=1}^{n_0} |q_i| 2^{-\frac{|q_i|}{T}} + \frac{1}{Z_n(T)} \sum_{i=n_0+1}^n |q_i| 2^{-\frac{|q_i|}{T}} \\
&> \frac{2M}{Z_n(T)} \sum_{i=n_0+1}^n 2^{-\frac{|q_i|}{T}} \\
&= 2M \left(1 - \frac{1}{Z_n(T)} \sum_{i=1}^{n_0} 2^{-\frac{|q_i|}{T}} \right) \\
&\geq 2M \frac{1}{2} = M.
\end{aligned}$$

Hence, $\lim_{n \rightarrow \infty} E_n(T) = \infty$, and the proof is completed. \square

In statistical mechanics, the entropy $S_{\text{sm}}(T)$ at temperature T is given by

$$S_{\text{sm}}(T) = \frac{1}{T} E_{\text{sm}}(T) + k \ln Z_{\text{sm}}(T), \quad (19)$$

where $Z_{\text{sm}}(T)$ and $E_{\text{sm}}(T)$ are given by (9) and (14), respectively. Motivated by the formula (19) and taking into account Replacements 1.1, we introduce the notion of statistical mechanical entropy into algorithmic information theory as follows.

Definition 4.7 (statistical mechanical entropy). *For each $n \in \mathbb{N}^+$ and each real number $T > 0$, we define $S_n(T)$ by*

$$S_n(T) = \frac{1}{T} E_n(T) + \log_2 Z_n(T).$$

Then, for each $T > 0$, the statistical mechanical entropy $S(T)$ is defined by $S(T) = \lim_{n \rightarrow \infty} S_n(T)$. \square

Theorem 4.8. *Let $T \in \mathbb{R}$.*

(i) *If $0 < T < 1$ and T is computable, then $S(T)$ converges to a left-computable real number which is Chaitin T -random and T -compressible.*

(ii) *If $1 \leq T$, then $S(T)$ diverges to ∞ .*

Proof. (i) Since $Z(T)$ and $E(T)$ converge by Theorem 4.2 (i) and Theorem 4.6 (i), respectively, $S(T)$ also converges and

$$S(T) = \frac{1}{T} E(T) + \log_2 Z(T).$$

Since $E(T)$ is a left-computable real number and $E(T) \notin \mathbb{N}$ by Theorem 4.6 (i), there exists a total recursive function $f: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $0 < f(m) \leq E(T)$ and $\lfloor f(m) \rfloor = \lfloor E(T) \rfloor$ for all $m \in \mathbb{N}^+$ and $\lim_{m \rightarrow \infty} f(m) = E(T)$. Since T is a computable real number, there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $0 \leq g(m) \leq 1/T$ for all $m \in \mathbb{N}^+$ and $\lim_{m \rightarrow \infty} g(m) = 1/T$. Since $Z(T)$ is a left-computable real numbers by Theorem 4.2 (i), there exists a total recursive function $h: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $h(m) \leq \log_2 Z(T)$ for all $m \in \mathbb{N}^+$ and $\lim_{m \rightarrow \infty} h(m) = \log_2 Z(T)$.

Hence, $g(m)f(m) + h(m) \leq S(T)$ for all $m \in \mathbb{N}^+$ and $\lim_{m \rightarrow \infty} g(m)f(m) + h(m) = S(T)$. Thus, $S(T)$ is a left-computable real number.

We then show that $S(T)$ is weakly Chaitin T -random. Let α be the infinite binary string such that $0.\alpha$ is the base-two expansion of $S(T) \bmod 1$ with infinitely many ones.

Given α_n , one can find $m_0 \in \mathbb{N}^+$ such that

$$\lceil S(T) \rceil - 1 + 0.\alpha_n < g(m_0)f(m_0) + h(m_0).$$

This is possible because $\lceil S(T) \rceil - 1 + 0.\alpha_n < S(T)$ and $\lim_{m \rightarrow \infty} g(m)f(m) + h(m) = S(T)$. It is shown that

$$\begin{aligned} 2^{-n} &> \left(\frac{1}{T}E(T) + \log_2 Z(T) \right) - (g(m_0)f(m_0) + h(m_0)) \\ &\geq \frac{1}{T}E(T) - g(m_0)f(m_0) \\ &\geq \frac{1}{T}(E(T) - f(m_0)) \\ &\geq E(T) - f(m_0). \end{aligned}$$

Thus, $0 \leq E(T) - f(m_0) < 2^{-n}$. Let l_n be the first n bits of the base-two expansion of the rational number $f(m_0) - \lfloor f(m_0) \rfloor$ with infinitely many zeros. It follows that

$$0 \leq 0.(E(T))_n - 0.l_n < E(T) - f(m_0) + 2^{-n} < 2 \cdot 2^{-n}.$$

Hence

$$(E(T))_n = l_n, l_n + 1,$$

where $(E(T))_n$ and l_n are regarded as a dyadic integer. Thus, there are still 2 possibilities of $(E(T))_n$, so that one needs only 1 bit more in order to determine $(E(T))_n$.

Thus, there exists a partial recursive function $\Phi: \{0, 1\}^* \times \{0, 1\} \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists b \in \{0, 1\} \quad \Phi(\alpha_n, b) = (E(T))_n.$$

It follows that there exists $c_\Phi \in \mathbb{N}^+$ such that, for all $n \in \mathbb{N}^+$,

$$H((E(T))_n) \leq H(\alpha_n) + c_\Phi.$$

Hence, α is Chaitin T -random by Theorem 4.6 (i). It follows that α has infinitely many zeros, which implies that $S(T) \bmod 1 = S(T) \bmod 1 = 0.\alpha$ and therefore $(S(T))_n = \alpha_n$. Thus, $S(T)$ is also Chaitin T -random.

Next, we show that $S(T)$ is T -compressible. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $m \in \mathbb{N}^+$, we define $\widetilde{W}_m(T)$ and $\widetilde{Z}_m(T)$ by

$$\widetilde{W}_m(T) = \sum_{i=1}^m |p_i| 2^{-\frac{|p_i|}{T}} \quad \text{and} \quad \widetilde{Z}_m(T) = \sum_{i=1}^m 2^{-\frac{|p_i|}{T}},$$

and then define $\widetilde{S}_m(T)$ by

$$\widetilde{S}_m(T) = \frac{1}{T} \frac{\widetilde{W}_m(T)}{\widetilde{Z}_m(T)} + \log_2 \widetilde{Z}_m(T).$$

Since, in the definition of $S_n(T)$, $Z_n(T)$ and the numerator and the denominator of $E_n(T)$ are positive term series which converge as $n \rightarrow \infty$, we see that $W(1, T) = \lim_{m \rightarrow \infty} \widetilde{W}_m(T)$, $Z(T) = \lim_{m \rightarrow \infty} \widetilde{Z}_m(T)$, and $S(T) = \lim_{m \rightarrow \infty} \widetilde{S}_m(T)$.

Since T is a computable real number, there exists a total recursive function $a: \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow \mathbb{Z}$ such that, for all $m, n \in \mathbb{N}^+$,

$$\left| \widetilde{S}_m(T) - \lfloor S(T) \rfloor - 2^{-n} a(m, n) \right| < 2^{-n}. \quad (20)$$

It is also shown that there is $d \in \mathbb{N}^+$ such that, for all $m \in \mathbb{N}^+$,

$$\left| S(T) - \widetilde{S}_m(T) \right| < 2^d \max \left\{ \left| W(1, T) - \widetilde{W}_m(T) \right|, \left| Z(T) - \widetilde{Z}_m(T) \right| \right\}. \quad (21)$$

Based on the inequalities (20) and (21), in the same manner as the proof of the T -compressibility of $E(T)$ in Theorem 4.6 (i), we can show that $S(T)$ is T -compressible.

(ii) In the case of $T \geq 1$, since $\lim_{n \rightarrow \infty} E_n(T) = \infty$ by Theorem 4.6 (ii) and $\log_2 Z_n(T)$ is bounded to the below by Theorem 4.2, we see that $\lim_{n \rightarrow \infty} S_n(T) = \infty$. \square

Finally, in statistical mechanics, the specific heat $C_{\text{sm}}(T)$ at temperature T is given by

$$C_{\text{sm}}(T) = \frac{d}{dT} E_{\text{sm}}(T), \quad (22)$$

where $E_{\text{sm}}(T)$ is given by (14). Motivated by the formula (22), we introduce the notion of specific heat into algorithmic information theory as follows.

Definition 4.9 (specific heat). *For each $n \in \mathbb{N}^+$ and each real number $T > 0$, we define $C_n(T)$ by*

$$C_n(T) = E'_n(T),$$

where $E'_n(T)$ is the derived function of $E_n(T)$. Then, for each $T > 0$, the specific heat $C(T)$ is defined by $C(T) = \lim_{n \rightarrow \infty} C_n(T)$. \square

Theorem 4.10. *Let $T \in \mathbb{R}$.*

(i) *If $0 < T < 1$ and T is computable, then $C(T)$ converges to a left-computable real number which is Chaitin T -random and T -compressible, and moreover $C(T) = E'(T)$ where $E'(T)$ is the derived function of $E(T)$.*

(ii) *If $T = 1$, then $C(T)$ diverges to ∞ .*

Proof. (i) First we show that $C(T)$ converges. Note that

$$C_n(T) = \frac{\ln 2}{T^2} \left\{ \frac{Y_n(T)}{Z_n(T)} - \left(\frac{W_n(T)}{Z_n(T)} \right)^2 \right\},$$

where

$$Y_n(T) = \sum_{i=1}^n |q_i|^2 2^{-\frac{|q_i|}{T}} \quad \text{and} \quad W_n(T) = \sum_{i=1}^n |q_i| 2^{-\frac{|q_i|}{T}}.$$

By Theorem 4.2 (i), $Z_n(T)$ converges to the real number $Z(T) > 0$ as $n \rightarrow \infty$. On the other hand, by Theorem 3.1 (i), $Y_n(T)$ and $W_n(T)$ converge to the real numbers $W(2, T)$ and $W(1, T)$, respectively, as $n \rightarrow \infty$. Thus, $C_n(T)$ also converges to a real number $C(T)$ as $n \rightarrow \infty$.

Next, we show that $C(T)$ is a left-computable real number. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $m \in \mathbb{N}^+$, we define $\tilde{Y}_m(T)$, $\tilde{W}_m(T)$, and $\tilde{Z}_m(T)$ by

$$\tilde{Y}_m(T) = \sum_{i=1}^m |p_i|^2 2^{-\frac{|p_i|}{T}}, \quad \tilde{W}_m(T) = \sum_{i=1}^m |p_i| 2^{-\frac{|p_i|}{T}}, \quad \text{and} \quad \tilde{Z}_m(T) = \sum_{i=1}^m 2^{-\frac{|p_i|}{T}},$$

respectively. Since $Y_n(T)$, $W_n(T)$, and $Z_n(T)$ are positive term series which converge as $n \rightarrow \infty$, we see that $W(2, T) = \lim_{m \rightarrow \infty} \tilde{Y}_m(T)$, $W(1, T) = \lim_{m \rightarrow \infty} \tilde{W}_m(T)$, $Z(T) = \lim_{m \rightarrow \infty} \tilde{Z}_m(T)$, and $C(T) = \lim_{m \rightarrow \infty} \tilde{C}_m(T)$ where

$$\tilde{C}_m(T) = \frac{\ln 2}{T^2} \left\{ \frac{\tilde{Y}_m(T)}{\tilde{Z}_m(T)} - \left(\frac{\tilde{W}_m(T)}{\tilde{Z}_m(T)} \right)^2 \right\}.$$

We then see that $\tilde{C}_{m+1}(T) - \tilde{C}_m(T)$ is calculated as

$$\begin{aligned} & \frac{\ln 2}{T^2} \frac{2^{-\frac{|p_{m+1}|}{T}}}{\tilde{Z}_{m+1}(T)} \left[|p_{m+1}|^2 - \left\{ \frac{\tilde{W}_{m+1}(T)}{\tilde{Z}_{m+1}(T)} + \frac{\tilde{W}_m(T)}{\tilde{Z}_m(T)} \right\} |p_{m+1}| \right. \\ & \left. + \left\{ \frac{\tilde{W}_{m+1}(T)}{\tilde{Z}_{m+1}(T)} + \frac{\tilde{W}_m(T)}{\tilde{Z}_m(T)} \right\} \frac{\tilde{W}_m(T)}{\tilde{Z}_m(T)} - \frac{\tilde{Y}_m(T)}{\tilde{Z}_m(T)} \right]. \end{aligned} \quad (23)$$

Since $\tilde{Y}_m(T)$, $\tilde{W}_m(T)$, and $\tilde{Z}_m(T)$ converge as $m \rightarrow \infty$ and $\lim_{m \rightarrow \infty} |p_m| = \infty$, there exists $m_1 \in \mathbb{N}^+$ such that, for any $m \geq m_1$,

$$\tilde{C}_{m+1}(T) - \tilde{C}_m(T) > |p_{m+1}| 2^{-\frac{|p_{m+1}|}{T}}. \quad (24)$$

In particular, $\tilde{C}_m(T)$ is an increasing function of m for all $m \geq m_1$ by the above inequality. Thus, since T is a computable real number, $C(T)$ is shown to be a left-computable real number.

Based on the computability of T and the inequality (24), in the same manner as the proof of Theorem 4.6 (i) we can show that $C(T)$ is Chaitin T -random.

Next, we show that $C(T)$ is T -compressible. Since T is a computable real number, there exists a total recursive function $g: \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow \mathbb{Z}$ such that, for all $m, n \in \mathbb{N}^+$,

$$\left| \tilde{C}_m(T) - \lfloor C(T) \rfloor - 2^{-n} g(m, n) \right| < 2^{-n}. \quad (25)$$

It is also shown that there is $c \in \mathbb{N}^+$ such that, for all $m \in \mathbb{N}^+$,

$$\begin{aligned} & \left| C(T) - \tilde{C}_m(T) \right| \\ & < 2^c \max \left\{ \left| Y(T) - \tilde{Y}_m(T) \right|, \left| W(T) - \tilde{W}_m(T) \right|, \left| Z(T) - \tilde{Z}_m(T) \right| \right\}. \end{aligned} \quad (26)$$

Let t be any computable real number with $T < t < 1$. Then, $W(2, t) = \lim_{m \rightarrow \infty} \tilde{Y}_m(t)$, where

$$\tilde{Y}_m(t) = \sum_{i=1}^m |p_i|^2 2^{-\frac{|p_i|}{t}}.$$

The limit value $W(2, t)$ exists since $t < 1$. Let β be the infinite binary string such that $0.\beta$ is the base-two expansion of $W(2, t) \bmod 1$ with infinitely many ones.

Given n and $\beta_{\lceil Tn/t \rceil}$ (i.e., the first $\lceil Tn/t \rceil$ bits of β), one can find $m_0 \in \mathbb{N}^+$ such that

$$\lceil W(2, t) \rceil - 1 + 0.\beta_{\lceil Tn/t \rceil} < \tilde{Y}_{m_0}(t).$$

This is possible since t is a computable real number and $\lceil W(2, t) \rceil - 1 + 0.\beta_{\lceil Tn/t \rceil} < W(2, t)$. It is then easy to see that

$$\sum_{i=m_0+1}^{\infty} |p_i|^2 2^{-\frac{|p_i|}{t}} < 2^{-Tn/t}.$$

Raising both sides of this inequality to the power t/T and using the inequality $a^c + b^c \leq (a + b)^c$ for real numbers $a, b > 0$ and $c \geq 1$,

$$2^{-n} > \sum_{i=m_0+1}^{\infty} |p_i|^{2t/T} 2^{-\frac{|p_i|}{T}} > \sum_{i=m_0+1}^{\infty} |p_i|^2 2^{-\frac{|p_i|}{T}}$$

and therefore

$$2^{-n} > \sum_{i=m_0+1}^{\infty} |p_i| 2^{-\frac{|p_i|}{T}} \quad \text{and} \quad 2^{-n} > \sum_{i=m_0+1}^{\infty} 2^{-\frac{|p_i|}{T}}.$$

It follows that

$$\max \left\{ \left| W(2, T) - \tilde{Y}_{m_0}(T) \right|, \left| W(1, T) - \tilde{W}_{m_0}(T) \right|, \left| Z(T) - \tilde{Z}_{m_0}(T) \right| \right\} < 2^{-n}. \quad (27)$$

From (25), (26), (27), and

$$| \lceil C(T) \rceil + 0.(C(T))_n - C(T) | < 2^{-n},$$

it is shown that

$$| (C(T))_n - g(m_0, n) | < 2^c + 2$$

and therefore

$$(C(T))_n = g(m_0, n), g(m_0, n) \pm 1, g(m_0, n) \pm 2, \dots, g(m_0, n) \pm (2^c + 1),$$

where $(C(T))_n$ is regarded as a dyadic integer. Thus, there are still $2^{c+1} + 3$ possibilities of $(C(T))_n$, so that one needs only $c + 2$ bits more in order to determine $(C(T))_n$.

Thus, there exists a partial recursive function $\Phi: \mathbb{N}^+ \times \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = c + 2 \quad \& \quad \Phi(n, \beta_{\lceil Tn/t \rceil}, s) = (C(T))_n.$$

It follows from (3) that $H((C(T))_n) \leq |\beta_{\lceil Tn/t \rceil}| + o(n) \leq Tn/t + o(n)$, which implies that $C(T)$ is T/t -compressible. Since t is any computable real number with $T < t < 1$, it follows that $C(T)$ is T -compressible.

By evaluating $C_{n+1}(x) - C_n(x)$ for all $x \in (0, 1)$ like (23), we can show that $C_n(x)$ converges uniformly in the wider sense on $(0, 1)$ to $C(x)$ as $n \rightarrow \infty$. Hence, we have $C(T) = E'(T)$.

(ii) It can be shown that

$$C_n(1) = \frac{\ln 2}{2} \frac{1}{Z_n(1)^2} \sum_{i=1}^n \sum_{j=1}^n (|q_i| - |q_j|)^2 2^{-|q_i|} 2^{-|q_j|}.$$

By Theorem 3.2 (i), $\sum_{j=1}^n (|q_1| - |q_j|)^2 2^{-|q_j|}$ diverges to ∞ as $n \rightarrow \infty$. Therefore $\sum_{i=1}^n \sum_{j=1}^n (|q_i| - |q_j|)^2 2^{-|q_i|} 2^{-|q_j|}$ also diverges to ∞ as $n \rightarrow \infty$. On the other hand, by Theorem 4.2 (i), $Z_n(1)$ converges as $n \rightarrow \infty$. Thus, $C(1)$ diverges to ∞ , and the proof is completed. \square

Thus, the theorems in this section show that the temperature T plays a role as the compression rate for all the thermodynamic quantities introduced into algorithmic information theory in this section. These theorems also show that the values of the thermodynamic quantities: partition function, free energy, energy, and statistical mechanical entropy diverge in the case of $T > 1$. This phenomenon might be regarded as some sort of phase transition in statistical mechanics.⁴

5 Fixed point theorems on compression rate

In this section, we prove the following theorem.

Theorem 5.1 (fixed point theorem on compression rate). *For every $T \in (0, 1)$, if $Z(T)$ is a computable real number, then the following hold:*

(i) T is right-computable and not left-computable.

(ii) T is weakly Chaitin T -random and T -compressible.

(iii) $\lim_{n \rightarrow \infty} H(T_n)/n = T$. \square

Theorem 5.1 follows immediately from Theorem 5.3, Theorem 5.5, and Theorem 5.6 below. From a purely mathematical point of view, Theorem 5.1 is just a fixed point theorem on compression rate, where the computability of the value $Z(T)$ gives a sufficient condition for T to be a fixed point on compression rate. Note that $Z(T)$ is a monotonically increasing continuous function on $(0, 1)$. In fact, [15, 16] showed that $Z(T)$ is a function of class C^∞ on $(0, 1)$. Thus, since computable real numbers are dense in \mathbb{R} , we have the following corollary of Theorem 5.1.

Corollary 5.2. *The set $\{T \in (0, 1) \mid \lim_{n \rightarrow \infty} H(T_n)/n = T\}$ is dense in $[0, 1]$.* \square

From the point of view of the statistical mechanical interpretation introduced in the previous section, Theorem 5.1 shows that the compression rate of temperature is equal to the temperature itself. Thus, Theorem 5.1 further confirms the role of temperature as the compression rate, which is observed in the previous section.

As a first step to prove Theorem 5.1, we prove the following theorem which gives the weak Chaitin T -randomness of T in Theorem 5.1.

Theorem 5.3. *For every $T \in (0, 1)$, if $Z(T)$ is a right-computable real number, then T is weakly Chaitin T -random.*

⁴ It is still open whether $C(T)$ diverges or not in the case of $T > 1$.

Proof. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $k \in \mathbb{N}^+$, we define a function $\tilde{Z}_k: (0, 1) \rightarrow \mathbb{R}$ by

$$\tilde{Z}_k(x) = \sum_{i=1}^k 2^{-\frac{|p_i|}{x}}.$$

Then, $\lim_{k \rightarrow \infty} \tilde{Z}_k(x) = Z(x)$ for every $x \in (0, 1)$. On the other hand, since $Z(T)$ is right-computable, there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $Z(T) \leq g(m)$ for all $m \in \mathbb{N}^+$, and $\lim_{m \rightarrow \infty} g(m) = Z(T)$.

We choose any one real number t with $T < t < 1$. For each $i \in \mathbb{N}^+$, using the mean value theorem we see that

$$2^{-\frac{|p_i|}{x}} - 2^{-\frac{|p_i|}{T}} < \frac{\ln 2}{T^2} |p_i| 2^{-\frac{|p_i|}{t}} (x - T)$$

for all $x \in (T, t)$. We choose any one $c \in \mathbb{N}$ with $W(1, t) \ln 2 / T^2 \leq 2^c$. Here, the limit value $W(1, t)$ exists by Theorem 3.1 (i), since $0 < t < 1$. Then, it follows that

$$\tilde{Z}_k(x) - \tilde{Z}_k(T) < 2^c (x - T) \tag{28}$$

for all $k \in \mathbb{N}^+$ and $x \in (T, t)$.

We choose any one $n_0 \in \mathbb{N}^+$ such that $T < 0.T_n + 2^{-n} < t$ for all $n \geq n_0$. Such n_0 exists since $T < t$ and $\lim_{n \rightarrow \infty} 0.T_n + 2^{-n} = T$.

Given T_n with $n \geq n_0$, one can find $k_0, m_0 \in \mathbb{N}^+$ such that

$$g(m_0) < \tilde{Z}_{k_0}(0.T_n + 2^{-n}).$$

This is possible from $Z(T) < Z(0.T_n + 2^{-n})$, $\lim_{k \rightarrow \infty} \tilde{Z}_k(0.T_n + 2^{-n}) = Z(0.T_n + 2^{-n})$, and the properties of g . It follows from $Z(T) \leq g(m_0)$ and (28) that

$$\sum_{i=k_0+1}^{\infty} 2^{-\frac{|p_i|}{T}} = Z(T) - \tilde{Z}_{k_0}(T) < \tilde{Z}_{k_0}(0.T_n + 2^{-n}) - \tilde{Z}_{k_0}(T) < 2^{c-n}.$$

Hence, for every $i > k_0$, $2^{-\frac{|p_i|}{T}} < 2^{c-n}$ and therefore $Tn - Tc < |p_i|$. Thus, by calculating the set $\{U(p_i) \mid i \leq k_0\}$ and picking any one finite binary string which is not in this set, one can then obtain an $s \in \{0, 1\}^*$ such that $Tn - Tc < H(s)$.

Hence, there exists a partial recursive function $\Psi: \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that $Tn - Tc < H(\Psi(T_n))$ for all $n \geq n_0$. Using (2), there is $c_\Psi \in \mathbb{N}^+$ such that $H(\Psi(T_n)) < H(T_n) + c_\Psi$ for all $n \geq n_0$. Therefore, $Tn - Tc - c_\Psi < H(T_n)$ for all $n \geq n_0$. It follows that T is weakly Chaitin T -random. \square

Remark 5.4. *By elaborating Theorem 4.2 (i), we can see that the left-computability of T results in the weak Chaitin T -randomness of $Z(T)$. On the other hand, by Theorem 5.3, the right-computability of $Z(T)$ results in the weak Chaitin T -randomness of T . We can integrate these two extremes into the following form: For every $T \in (0, 1]$, there exists $c \in \mathbb{N}^+$ such that, for every $n \in \mathbb{N}^+$,*

$$Tn - c \leq H(T_n, (Z(T))_n), \tag{29}$$

where $H(s, t)$ is defined as $H(\langle s, t \rangle)$ with any one computable bijection $\langle s, t \rangle$ from $(s, t) \in \{0, 1\}^* \times \{0, 1\}^*$ to $\{0, 1\}^*$ (see [4] for the detail of the notion of $H(s, t)$). In fact, if T is left-computable, then we can show that $H((Z(T))_n) = H(T_n, (Z(T))_n) + O(1)$, and therefore the inequality (29) results in the weak Chaitin T -randomness of $Z(T)$. On the other hand, if $Z(T)$ is right-computable, then we can show that $H(T_n) = H(T_n, (Z(T))_n) + O(1)$, and therefore the inequality (29) results in the weak Chaitin T -randomness of T .

Note, however, that the inequality (29) is not necessarily tight except for these two extremes, that is, the following inequality does not hold: For every $T \in (0, 1]$,

$$H(T_n, (Z(T))_n) \leq Tn + o(n), \quad (30)$$

where $o(n)$ depends on T in addition to n . To see this, contrarily assume that the inequality (30) holds. Then, by setting T to Chaitin's Ω , we have $H(\Omega_n) \leq H(\Omega_n, (Z(\Omega))_n) + O(1) \leq \Omega n + o(n)$. Since $\Omega < 1$, this contradicts the fact that Ω is weakly Chaitin random. Thus, the inequality (30) does not hold. \square

The following Theorem 5.5 and Theorem 5.6 give the T -compressibility of T in Theorem 5.1 together.

Theorem 5.5. *For every $T \in (0, 1)$, if $Z(T)$ is a right-computable real number, then T is also a right-computable real number.*

Proof. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $k \in \mathbb{N}^+$, we define a function $\tilde{Z}_k: (0, 1) \rightarrow \mathbb{R}$ by

$$\tilde{Z}_k(x) = \sum_{i=1}^k 2^{-\frac{|p_i|}{x}}.$$

Then, $\lim_{k \rightarrow \infty} \tilde{Z}_k(x) = Z(x)$ for every $x \in (0, 1)$. Since $Z(T)$ is right-computable, there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $Z(T) \leq g(m)$ for all $m \in \mathbb{N}^+$, and $\lim_{m \rightarrow \infty} g(m) = Z(T)$. Since $Z(x)$ is an increasing function of x , we see that, for every $x \in \mathbb{Q}$ with $0 < x < 1$, $T < x$ if and only if there are $m, k \in \mathbb{N}^+$ such that $g(m) < \tilde{Z}_k(x)$. Thus, T is right-computable. This is because the set $\{(m, n) \mid m \in \mathbb{Z} \ \& \ n \in \mathbb{N}^+ \ \& \ T < m/n\}$ is r.e. if and only if T is right-computable. \square

The converse of Theorem 5.5 does not hold. To see this, consider an arbitrary computable real number $T \in (0, 1)$. Then, obviously T is right-computable. On the other hand, $Z(T)$ is left-computable and weakly Chaitin T -random by Theorem 4.2 (i). Thus, $Z(T)$ is not right-computable.

Theorem 5.6. *For every $T \in (0, 1)$, if $Z(T)$ is a left-computable real number and T is a right-computable real number, then T is T -compressible.*

Proof. Let p_1, p_2, p_3, \dots be a recursive enumeration of the r.e. set $\text{dom } U$. For each $k \in \mathbb{N}^+$, we define a function $\tilde{Z}_k: (0, 1) \rightarrow \mathbb{R}$ by

$$\tilde{Z}_k(x) = \sum_{i=1}^k 2^{-\frac{|p_i|}{x}}.$$

Then, $\lim_{k \rightarrow \infty} \tilde{Z}_k(x) = Z(x)$ for every $x \in (0, 1)$.

For each $i \in \mathbb{N}^+$, using the mean value theorem we see that

$$2^{-\frac{|p_1|}{t}} - 2^{-\frac{|p_1|}{T}} > (\ln 2) |p_1| 2^{-\frac{|p_1|}{T}} (t - T)$$

for all $t \in (T, 1)$. We choose any one $c \in \mathbb{N}^+$ such that $(\ln 2) |p_1| 2^{-\frac{|p_1|}{T}} \geq 2^{-c}$. Then, it follows that

$$\tilde{Z}_k(t) - \tilde{Z}_k(T) > 2^{-c}(t - T) \quad (31)$$

for all $k \in \mathbb{N}^+$ and $t \in (T, 1)$.

Since T is a right-computable real number with $T < 1$, there exists a total recursive function $f: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $T < f(l) < 1$ for all $l \in \mathbb{N}^+$, and $\lim_{l \rightarrow \infty} f(l) = T$. On the other hand, since $Z(T)$ is left-computable, there exists a total recursive function $g: \mathbb{N}^+ \rightarrow \mathbb{Q}$ such that $g(m) \leq Z(T)$ for all $m \in \mathbb{N}^+$, and $\lim_{m \rightarrow \infty} g(m) = Z(T)$. Let β be the infinite binary sequence such that $0.\beta$ is the base-two expansion of $Z(1)$ (i.e., Chaitin's Ω).

Given n and $\beta_{\lceil Tn \rceil}$ (i.e., the first $\lceil Tn \rceil$ bits of β), one can find $k_0 \in \mathbb{N}^+$ such that

$$0.\beta_{\lceil Tn \rceil} < \sum_{i=1}^{k_0} 2^{-|p_i|}.$$

It is then easy to see that

$$\sum_{i=k_0+1}^{\infty} 2^{-|p_i|} < 2^{-Tn}.$$

Using the inequality $a^d + b^d \leq (a + b)^d$ for real numbers $a, b > 0$ and $d \geq 1$, it follows that

$$Z(T) - \tilde{Z}_{k_0}(T) < 2^{-n}. \quad (32)$$

Note that $\tilde{Z}_{k_0}(T) < \tilde{Z}_{k_0}(f(l))$ for all $l \in \mathbb{N}^+$, and $\lim_{l \rightarrow \infty} \tilde{Z}_{k_0}(f(l)) = \tilde{Z}_{k_0}(T)$. Thus, since $\tilde{Z}_{k_0}(T) < Z(T)$, one can then find $l_0, m_0 \in \mathbb{N}^+$ such that

$$\tilde{Z}_{k_0}(f(l_0)) < g(m_0).$$

It follows from (32) and (31) that

$$2^{-n} > g(m_0) - \tilde{Z}_{k_0}(T) > \tilde{Z}_{k_0}(f(l_0)) - \tilde{Z}_{k_0}(T) > 2^{-c}(f(l_0) - T).$$

Thus, $0 < f(l_0) - T < 2^{c-n}$. Let t_n be the first n bits of the base-two expansion of the rational number $f(l_0)$ with infinitely many zeros. Then, $|f(l_0) - 0.t_n| < 2^{-n}$. It follows from $|T - 0.T_n| < 2^{-n}$ that $|0.T_n - 0.t_n| < (2^c + 2)2^{-n}$. Hence

$$T_n = t_n, t_n \pm 1, t_n \pm 2, \dots, t_n \pm (2^c + 1),$$

where T_n and t_n are regarded as a dyadic integer. Thus, there are still $2^{c+1} + 3$ possibilities of T_n , so that one needs only $c + 2$ bits more in order to determine T_n .

Thus, there exists a partial recursive function $\Phi: \mathbb{N}^+ \times \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that

$$\forall n \in \mathbb{N}^+ \quad \exists s \in \{0, 1\}^* \quad |s| = c + 2 \quad \& \quad \Phi(n, \beta_{\lceil Tn \rceil}, s) = T_n.$$

It follows from (3) that $H(T_n) \leq |\beta_{\lceil Tn \rceil}| + o(n) \leq Tn + o(n)$, which implies that T is T -compressible. \square

In a similar manner to the proof of Theorem 5.1, we can prove another version of a fixed point theorem on compression rate as follows. Here, the weak Chaitin T -randomness is replaced by the Chaitin T -randomness.

Theorem 5.7 (fixed point theorem on compression rate II). *Let Q be a computable real number with $Q > 0$. For every $T \in (0, 1)$, if $W(Q, T)$ is a computable real number, then the following hold:*

(i) T is right-computable and not left-computable.

(ii) T is Chaitin T -random and T -compressible. □

Remark 5.8. *The computability of $Z(T)$ in the premise of Theorem 5.1 can be replaced by the computability of $F(T)$. On the other hand, the computability of $W(Q, T)$ in the premise of Theorem 5.7 can be replaced by the computability of $E(T)$ or $S(T)$. □*

6 Total statistical mechanical interpretation of algorithmic information theory: Physical and informal argument

In what follows, based on a physical argument we develop a total statistical mechanical interpretation of algorithmic information theory which attains a perfect correspondence to normal statistical mechanics. In consequence, we justify the interpretation of Ω^D as a partition function and clarify the statistical mechanical meaning of the thermodynamic quantities introduced into algorithmic information theory in Section 4. In the work [17], we developed a statistical mechanical interpretation of the noiseless source coding scheme based on an absolutely optimal instantaneous code by identifying a microcanonical ensemble in the scheme. In a similar manner to [17] we develop a statistical mechanical interpretation of algorithmic information theory in what follows. This can be possible because the set $\text{dom } U$ is prefix-free and therefore the action of the optimal computer U can be regarded as an instantaneous code which is extended over an infinite set. Note that, in what follows, we do not stick to the mathematical strictness of the argument and we make an argument on the same level of mathematical strictness as statistical mechanics in physics. We start with some reviews of statistical mechanics.

In statistical mechanics we consider a quantum system $\mathcal{S}_{\text{total}}$ which consists in a large number of identical quantum subsystems. Let N be the number of such subsystems. For example, $N \sim 10^{22}$ for 1 cm^3 of a gas at room temperature. We assume here that each quantum subsystem can be distinguishable from others. Thus, we deal with quantum particles which obey Maxwell-Boltzmann statistics and not Bose-Einstein statistics or Fermi-Dirac statistics. Under this assumption, we can identify the i th quantum subsystem \mathcal{S}_i for each $i = 1, \dots, N$. In quantum mechanics, any quantum system is described by a quantum state completely. In statistical mechanics, among all quantum states, energy eigenstates are of particular importance. Any energy eigenstate of each subsystem \mathcal{S}_i can be specified by a number $n = 1, 2, 3, \dots$, called a *quantum number*, where the subsystem in the energy eigenstate specified by n has the energy E_n . Then, any energy eigenstate of the system $\mathcal{S}_{\text{total}}$ can be specified by an N -tuple (n_1, n_2, \dots, n_N) of quantum numbers. If the state of the system $\mathcal{S}_{\text{total}}$ is the energy eigenstate specified by (n_1, n_2, \dots, n_N) , then the state of each subsystem \mathcal{S}_i is the energy eigenstate specified by n_i and the system $\mathcal{S}_{\text{total}}$ has the energy $E_{n_1} + E_{n_2} + \dots + E_{n_N}$.

Then, the fundamental postulate of statistical mechanics, called *the principle of equal probability*, is stated as follows.

The Principle of Equal Probability: If the energy of the system $\mathcal{S}_{\text{total}}$ is known to have a constant value in the range between E and $E + \delta E$, where δE is the indeterminacy in measurement of the energy of the system $\mathcal{S}_{\text{total}}$, then the system $\mathcal{S}_{\text{total}}$ is equally likely to be in any energy eigenstate specified by (n_1, n_2, \dots, n_N) such that $E \leq E_{n_1} + E_{n_2} + \dots + E_{n_N} \leq E + \delta E$. \square

Let $\Theta(E, N)$ be the total number of energy eigenstates of $\mathcal{S}_{\text{total}}$ specified by (n_1, n_2, \dots, n_N) such that $E \leq E_{n_1} + E_{n_2} + \dots + E_{n_N} \leq E + \delta E$. The above postulate states that any energy eigenstate of $\mathcal{S}_{\text{total}}$ whose energy lies between E and $E + \delta E$ occurs with the probability $1/\Theta(E, N)$. This uniform distribution of energy eigenstates whose energy lie between E and $E + \delta E$ is called a *microcanonical ensemble*. In statistical mechanics, the *entropy* $S(E, N)$ of the system $\mathcal{S}_{\text{total}}$ is then defined by

$$S(E, N) = k \ln \Theta(E, N),$$

where k is a positive constant, called the *Boltzmann Constant*, and the \ln denotes the natural logarithm. The average energy ε per one subsystem is given by E/N . In a normal case where ε has a finite value, the entropy $S(E, N)$ is proportional to N . On the other hand, the indeterminacy δE of the energy contributes to $S(E, N)$ through the term $k \ln \delta E$, which can be ignored compared to N unless δE is too small. Thus the magnitude of the indeterminacy δE of the energy does not matter to the value of the entropy $S(E, N)$ unless it is too small. The *temperature* $T(E, N)$ of the system $\mathcal{S}_{\text{total}}$ is defined by

$$\frac{1}{T(E, N)} = \frac{\partial S}{\partial E}(E, N).$$

Thus the temperature is a function of E and N .

Now we give a statistical mechanical interpretation to algorithmic information theory. As considered in [4], think of the optimal computer U as decoding equipment at the receiving end of a noiseless binary communication channel. Regard its programs (i.e., finite binary strings in $\text{dom } U$) as codewords and regard the result of the computation by U , which is a finite binary string, as a decoded “symbol.” Since $\text{dom } U$ is a prefix-free set, such codewords form what is called an “instantaneous code,” so that successive symbols sent through the channel in the form of concatenation of codewords can be separated.

For establishing the statistical mechanical interpretation of algorithmic information theory, we assume that the infinite binary string sent through the channel is generated by infinitely repeated tosses of a fair coin. Under this assumption, the success probability of decoding one symbol is equal to Chaitin’s halting probability Ω , and the probability of getting a finite binary string s as the first decoded symbol is equal to $P(s)$. Hereafter the infinite binary string sent through the channel is referred to as *the channel infinite string*. For each $r \in \{0, 1\}^*$, let $Q(r)$ be the probability that the channel infinite string has the prefix r . It follows that $Q(r) = 2^{-|r|}$. Thus, the channel infinite string is the random variable drawn according to Lebesgue measure on $\{0, 1\}^\infty$.

Let N be a large number, say $N \sim 10^{22}$. We relate algorithmic information theory to the statistical mechanics reviewed above in the following manner. Among all infinite binary strings, consider infinite binary strings of the form $p_1 p_2 \dots p_N \alpha$ with $p_1, p_2, \dots, p_N \in \text{dom } U$ and $\alpha \in \{0, 1\}^\infty$. For each i , the i th slot fed by p_i corresponds to the i th quantum subsystem \mathcal{S}_i . On the other hand, the ordered sequence of the 1st slot, the 2nd slot, \dots , and the N th slot corresponds to

the quantum system $\mathcal{S}_{\text{total}}$. We relate a codeword $p \in \text{dom } U$ to an energy eigenstate of a subsystem, and relate a codeword length $|p|$ to an energy E_n of the energy eigenstate of the subsystem. Then, a finite binary string $p_1 \cdots p_N$ corresponds to an energy eigenstate of $\mathcal{S}_{\text{total}}$ specified by (n_1, \dots, n_N) . Thus, $|p_1| + \cdots + |p_N| = |p_1 \cdots p_N|$ corresponds to the energy $E_{n_1} + \cdots + E_{n_N}$ of the energy eigenstate of $\mathcal{S}_{\text{total}}$.

We define a subset $C(L, N)$ of $\{0, 1\}^*$ as the set of all finite binary strings of the form $p_1 \cdots p_N$ with $p_i \in \text{dom } U$ whose total length $|p_1 \cdots p_N|$ lie between L and $L + \delta L$. Then, $\Theta(L, N)$ is defined as the cardinality of $C(L, N)$. Therefore, $\Theta(L, N)$ is the total number of all concatenations of N codewords whose total length lie between L and $L + \delta L$. We can see that if $p_1 \cdots p_N \in C(L, N)$, then $2^{-(L+\delta L)} \leq Q(p_1 \cdots p_N) \leq 2^{-L}$. Thus, all concatenations $p_1 \cdots p_N \in C(L, N)$ of N codewords occur in a prefix of the channel infinite string with the same probability 2^{-L} . Note here that we care nothing about the magnitude of δL , as in the case of statistical mechanics. Thus, the following principle, called *the principle of equal conditional probability*, holds.

The Principle of Equal Conditional Probability: Given that a concatenation of N codewords of total length L occurs in a prefix of the channel infinite string, all such concatenations occur with the same probability $1/\Theta(L, N)$. \square

We introduce a microcanonical ensemble into algorithmic information theory in this manner. Thus, we can develop a certain sort of statistical mechanics on algorithmic information theory. Note that, in statistical mechanics, the principle of equal probability is just a conjecture which is not yet proved completely in a realistic physical system. On the other hand, in our statistical mechanical interpretation of algorithmic information theory, the principle of equal conditional probability is automatically satisfied.

The *statistical mechanical entropy* $S(L, N)$ is defined by

$$S(L, N) = \log_2 \Theta(L, N). \quad (33)$$

The *temperature* $T(L, N)$ is then defined by

$$\frac{1}{T(L, N)} = \frac{\partial S}{\partial L}(L, N). \quad (34)$$

Thus, the temperature is a function of L and N .

According to the theoretical development of equilibrium statistical mechanics,⁵ we can introduce a canonical ensemble into algorithmic information theory in the following manner. We investigate the probability distribution of the left-most codeword p_1 of the channel infinite string, given that a concatenation of N codewords of total length L occurs in a prefix of the channel infinite string. For each $p \in \{0, 1\}^*$, let $R(p)$ be the probability that the left-most codeword of the channel infinite string is p , given that a concatenation of N codewords of total length L occurs in a prefix of the channel infinite string. Based on the principle of equal conditional probability, it can be shown that

$$R(p) = \frac{\Theta(L - |p|, N - 1)}{\Theta(L, N)}.$$

From the general definition (33) of statistical mechanical entropy, we have

$$R(p) = 2^{S(L-|p|, N-1) - S(L, N)}. \quad (35)$$

⁵ We follow the argument of Section 16-1 of Callen [1] in particular.

Let $E(L, N)$ be the expected length of the left-most codeword of the channel infinite string, given that a concatenation of N codewords of total length L occurs in a prefix of the channel infinite string. Then, the following equality is expected to hold:

$$S(L, N) = S(E(L, N), 1) + S(L - E(L, N), N - 1). \quad (36)$$

Here, the term $S(L, N)$ in the left-hand side denotes the statistical mechanical entropy of the whole concatenation of N codewords of total length L . On the other hand, the first term $S(E(L, N), 1)$ in the right-hand side denotes the statistical mechanical entropy of the left-most codeword of the concatenation of N codewords of total length L while the second term $S(L - E(L, N), N - 1)$ in the right-hand side denotes the statistical mechanical entropy of the remaining $N - 1$ codewords of the concatenation of N codewords of total length L . Thus, the equality (36) represents the additivity of the statistical mechanical entropy. We assume here that the equality (36) holds.

By expanding $S(L - |p|, N - 1)$ around the equilibrium point $L - E(L, N)$, we have

$$\begin{aligned} & S(L - |p|, N - 1) \\ &= S(L - E(L, N) + E(L, N) - |p|, N - 1) \\ &= S(L - E(L, N), N - 1) + \frac{\partial S}{\partial L}(L - E(L, N), N - 1)(E(L, N) - |p|). \end{aligned} \quad (37)$$

Here, we ignore the higher order terms than the first order. Since $N \gg 1$ and $L \gg E(L, N)$, using the definition (34) of temperature we have

$$\frac{\partial S}{\partial L}(L - E(L, N), N - 1) = \frac{\partial S}{\partial L}(L, N) = \frac{1}{T(L, N)}. \quad (38)$$

Hence, by (37) and (38), we have

$$S(L - |p|, N - 1) = S(L - E(L, N), N - 1) + \frac{1}{T(L, N)}(E(L, N) - |p|). \quad (39)$$

Thus, using (35), (36), and (39), we obtain

$$R(p) = 2^{\frac{E(L, N) - T(L, N)S(E(L, N), 1)}{T(L, N)}} 2^{-\frac{|p|}{T(L, N)}}.$$

Then, according to statistical mechanics we define the *free energy* $F(L, N)$ of the left-most codeword of the concatenation of N codewords of total length L by

$$F(L, N) = E(L, N) - T(L, N)S(E(L, N), 1). \quad (40)$$

It follows that

$$R(p) = 2^{\frac{F(L, N)}{T(L, N)}} 2^{-\frac{|p|}{T(L, N)}}. \quad (41)$$

Using $\sum_{p \in \text{dom } U} R(p) = 1$, we can show that, for any $p \in \text{dom } U$,

$$R(p) = \frac{1}{Z(T(L, N))} 2^{-\frac{|p|}{T(L, N)}}, \quad (42)$$

where $Z(T)$ is defined by

$$Z(T) = \sum_{p \in \text{dom } U} 2^{-\frac{|p|}{T}} \quad (T > 0). \quad (43)$$

$Z(T)$ is called the *partition function* (of the left-most codeword of the channel infinite string). Thus, in our statistical mechanical interpretation of algorithmic information theory, the partition function $Z(T)$ has exactly the same form as Ω^D . The distribution in the form of $R(p)$ is called a *canonical ensemble* in statistical mechanics.

Then, using (41) and (42), $F(L, N)$ is calculated as

$$F(L, N) = F(T(L, N)), \quad (44)$$

where $F(T)$ is defined by

$$F(T) = -T \log_2 Z(T) \quad (T > 0). \quad (45)$$

On the other hand, from the definition of $R(p)$, $E(N, L)$ is calculated as

$$E(N, L) = \sum_{p \in \text{dom } U} |p| R(p).$$

Thus, we have

$$E(L, N) = E(T(L, N)), \quad (46)$$

where $E(T)$ is defined by

$$E(T) = \frac{1}{Z(T)} \sum_{p \in \text{dom } U} |p| 2^{-\frac{|p|}{T}} \quad (T > 0). \quad (47)$$

Then, using (40), (44), (45), and (46), the statistical mechanical entropy $S(E(L, N), 1)$ of the left-most codeword of the concatenation of N codewords of total length L is calculated as

$$S(E(L, N), 1) = S(T(L, N)),$$

where $S(T)$ is defined by

$$S(T) = \frac{1}{T} E(T) + \log_2 Z(T) \quad (T > 0). \quad (48)$$

Note that the statistical mechanical entropy $S(E(L, N), 1)$ coincides with the *Shannon entropy*

$$- \sum_{p \in \text{dom } U} R(p) \log_2 R(p)$$

of the distribution $R(p)$.

Finally, the *specific heat* $C(T)$ of the left-most codeword of the channel infinite string is defined by

$$C(T) = E'(T) \quad (T > 0), \quad (49)$$

where $E'(T)$ is the derived function of $E(T)$.

Thus, a statistical mechanical interpretation of algorithmic information theory can be established, based on a physical argument. We can check that the formulas in this argument: the partition function (43), the free energy (45), the expected length of the left-most codeword (47), the statistical mechanical entropy (48), and the specific heat (49) correspond to the definitions in Section 4: Definition 4.1, Definition 4.3, Definition 4.5, Definition 4.7, and Definition 4.9, respectively. Thus, the statistical mechanical meaning of the notion of thermodynamic quantities introduced in Section 4 into algorithmic information theory is clarified by this argument.

7 Concluding remarks

In this paper, we have developed a statistical mechanical interpretation of algorithmic information theory by introducing the notion of thermodynamic quantities into algorithmic information theory and investigating their properties from the point of view of algorithmic randomness. As a result, we have discovered that, in the interpretation, the temperature plays a role as the compression rate of all these thermodynamic quantities, which include the temperature itself. Thus, in particular, we have obtained fixed point theorems on compression rate, which reflect this self-referential nature of the compression rate of the temperature. In the last part of this paper, we have also developed a total statistical mechanical interpretation of algorithmic information theory, which realizes a perfect correspondence to normal statistical mechanics and motivates the above introduction of the thermodynamic quantities into algorithmic information theory. However, the argument used in the total statistical mechanical interpretation is on the same level of mathematical strictness as statistical mechanics. Thus, we try to make the argument a mathematically rigorous form in a future study. This effort might stimulate a further unexpected development of the research of this line.

Acknowledgments

The author is grateful to Prof. Shigeo Tsujii for the financial supports.

References

- [1] H. B. Callen, *Thermodynamics and an Introduction to Thermostatistics*, 2nd ed. John Wiley & Sons, Inc., Singapore, 1985.
- [2] C. S. Calude, P. H. Hertling, B. Khoussainov, and Y. Wang, “Recursively enumerable reals and Chaitin Ω numbers,” *Theoret. Comput. Sci.*, vol. 255, pp. 125–149, 2001.
- [3] C. S. Calude and M. A. Stay, “Natural halting probabilities, partial randomness, and zeta functions,” *Inform. and Comput.*, vol. 204, pp. 1718–1739, 2006.
- [4] G. J. Chaitin, “A theory of program size formally identical to information theory,” *J. Assoc. Comput. Mach.*, vol. 22, pp. 329–340, 1975.
- [5] G. J. Chaitin, “Incompleteness theorems for random reals,” *Adv. in Appl. Math.*, vol. 8, pp. 119–146, 1987.
- [6] G. J. Chaitin, *Algorithmic Information Theory*. Cambridge University Press, Cambridge, 1987.
- [7] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*, 2nd ed. Springer, New York, 1998.
- [8] P. A. M. Dirac, *The Principles of Quantum Mechanics*, 4th ed. Oxford University Press, London, 1958.
- [9] P. Gács, “On the symmetry of algorithmic information,” *Soviet Math. Dokl.*, vol. 15, pp. 1477–1480, 1974; correction, *ibid.* vol. 15, pp. 1480, 1974.

- [10] L. A. Levin, “Laws of information conservation (non-growth) and aspects of the foundations of probability theory,” *Problems of Inform. Transmission*, vol. 10, pp. 206–210, 1974.
- [11] M. B. Pour-El and J. I. Richards, *Computability in Analysis and Physics*. Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1989.
- [12] F. Reif, *Fundamentals of Statistical and Thermal Physics*. McGraw-Hill, Inc., Singapore, 1965.
- [13] J. Reimann and F. Stephan, On hierarchies of randomness tests. Proceedings of the 9th Asian Logic Conference, World Scientific Publishing, August 16-19, 2005, Novosibirsk, Russia.
- [14] D. Ruelle, *Statistical Mechanics, Rigorous Results*, 3rd ed. Imperial College Press and World Scientific Publishing Co. Pte. Ltd., Singapore, 1999.
- [15] K. Tadaki, Algorithmic information theory and fractal sets. Proceedings of 1999 Workshop on Information-Based Induction Sciences (IBIS’99), pp. 105–110, August 26-27, 1999, Syuzenji, Shizuoka, Japan. In Japanese.
- [16] K. Tadaki, “A generalization of Chaitin’s halting probability Ω and halting self-similar sets,” *Hokkaido Math. J.*, vol. 31, pp. 219–253, 2002. Electronic Version Available: <http://arxiv.org/abs/nlin/0212001>
- [17] K. Tadaki, A statistical mechanical interpretation of instantaneous codes. Proceedings of 2007 IEEE International Symposium on Information Theory (ISIT2007), pp. 1906–1910, June 24-29, 2007, Nice, France.
- [18] M. Toda, R. Kubo, and N. Saitô, *Statistical Physics I. Equilibrium Statistical Mechanics*, 2nd ed. Springer, Berlin, 1992.
- [19] K. Weihrauch, *Computable Analysis*. Springer, Berlin, 2000.
- [20] A. K. Zvonkin and L. A. Levin, “The complexity of finite objects and the development of the concepts of information and randomness by means of the theory of algorithms,” *Russian Math. Surveys*, vol. 25, no. 6, pp. 83–124, 1970.