

Method to estimate ISCO and ring-down frequencies in binary systems and consequences for gravitational wave data analysis

Chad Hanna¹, Miguel Megevand¹, Evan Ochsner², and Carlos Palenzuela^{1,3}

¹ Department of Physics and Astronomy, Louisiana State University Baton Rouge, LA 70802, USA

² Department of Physics, University of Maryland, College Park, MD 20742

³ Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Golm, Germany

E-mail: channa@phys.lsu.edu, megevand@phys.lsu.edu, evano@umd.edu, carlos@lsu.edu

Abstract.

Recent advances in the description of compact binary systems have produced gravitational waveforms that include inspiral, merger and ring-down phases. Comparing results from numerical simulations with those of post-Newtonian (PN), and related, expansions has provided motivation for employing PN waveforms in near merger epochs when searching for gravitational waves and has encouraged the development of analytic fits to full numerical waveforms. The models and simulations do not yet cover the full binary coalescence parameter space. For these yet un-simulated regions, data analysts can still conduct separate inspiral, merger and ring-down searches. Improved knowledge about the end of the inspiral phase, the beginning of the merger, and the ring-down frequencies could increase the efficiency of both coherent inspiral-merger-ring-down (IMR) searches and searches over each phase separately. Insight can be gained for all three cases through a recently presented theoretical calculation, which, corroborated by the numerical results, provides an implicit formula for the final spin of the merged black holes, accurate to within 10% over a large parameter space. Knowledge of the final spin allows one to predict the end of the inspiral phase and the quasinormal mode ring-down frequencies, and in turn provides information about the bandwidth and duration of the merger. In this work we will discuss a few of the implications of this calculation for data analysis.

1. Introduction

Gravitational radiation emitted during the inspiral, merger and ring-down of two compact objects with a total mass below $\sim 100M_{\odot}$ is a likely detection source for interferometric gravitational wave detectors [1]. Maximal signal-to-noise ratio (SNR) is achieved in searches for compact binary coalescence with matched filtering banks of template waveforms [2, 3, 4, 5, 6] allowing faint signals to be detected reliably. The SNR depends on having a template waveform that matches the true signal correctly [2]. Because templates are weighted by the noise power spectral density of the detector with non-trivial frequency dependence it is important *when* in the binary evolution the template matches well. Numerical simulations are improving our knowledge of the full waveforms emitted during the inspiral, merger and ring-down phases of binary black hole coalescence [7, 8, 9], yet they still do not cover an adequate parameter space for constructing filter banks [10, 9]. When the full parameter space is simulated it will be useful to have fully parameterized models of these waveforms and even now many have considered hybrid waveforms that patch together PN and numerical simulations [11, 12, 7, 8, 9] as data analysis templates. Others have introduced a phenomenological fourth order correction to make effective one-body (EOB) waveforms match numerical relativity results near the merger [11, 7] known as *pseudo* 4PN (p4PN).

Before the full parameter space is simulated it is still possible to handle the three binary coalescence epochs - inspiral, merger and ring-down - separately. We will show that a recent calculation [13] which has power to predict the final spin of a binary merger, can provide both insight into when the inspiral phase ends and what the quasi-normal ring-down frequency of the merger product should be for a large, yet un-simulated, parameter space. We will discuss the following consequences of this calculation:

- The calculation of the final spin given in [13] contains an implicit reference to the innermost stable circular orbit (ISCO) of a test particle orbiting the Kerr black hole that results from the merger. We will also discuss the impact of using the new ISCO frequency as a cut off for current gravitational wave searches that are targeting coalescing compact objects including the impact on SNR and the ability to probe neutron star properties.
- Knowing the final spin gives an estimate for the quasi-normal mode ring-down frequencies. These frequencies could be used in a separate matched filter search [26].
- We can also use the final spin to calculate the light ring (smallest, unstable circular orbit for a photon orbiting a Kerr black hole) and quasinormal mode ring-down frequencies of the final black hole; we will see that the $l=2, m=2, n=0$ QNM is quite close to the light ring frequency. We will define a merger epoch as the evolution from ISCO to light ring and discuss how to obtain its time frequency content for yet un-simulated systems.

We define the merger epoch as everything that occurs between the ISCO and ring-down frequencies. We show that the duration of the radiation seems to be well

described by the infall duration of a test particle orbiting the merged black hole just below the ISCO. Knowing the duration and bandwidth gives a time-frequency volume, which is useful in data analysis for unmodeled signals.

2. Formalism

The inspiral phase of two compact objects is generally considered to end when the binary has evolved to an innermost stable circular orbit (ISCO), should it exist. The ISCO can be defined by waveform models and numerical relativity [17, 18, 7]. Unfortunately there is a difference in the numbers obtained from different methods. Some LIGO searches for non-spinning systems have taken the conservative approach to use the ISCO defined for a test particle orbiting a Schwarzschild black hole.

Intuitively the space-time of an inspiralling comparable mass binary system will not be well described by a Schwarzschild space-time, and thus the Schwarzschild ISCO will not capture the dynamics correctly. If the merger product is a black hole, it will be a Kerr-like black hole with a final spin that is a function of the masses and spins of the components. Recently [13] has proposed a simple set of assumptions that predict the final spin of the black hole merger product to well within 10% of numerical simulations based on first principle arguments. Implicitly in the derivation of [13] is the ISCO radius of a test particle orbiting a Kerr black hole having a final spin equal to the spin of the merger product. The fact that the ISCO radius used gives the correct answer for the final space-time suggests that it plays an important role in the pre-merger dynamics. We hypothesize that the test particle ISCO of the merged Kerr black hole may describe a way to define the end of the inspiral phase for binaries that produce black holes.

The ISCO solution for a test particle orbiting a Kerr black hole is [20],

$$\begin{aligned} Z_1 &\equiv 1 + \left(1 - \frac{a_f^2}{M^2}\right)^{1/3} \left[\left(1 + \frac{a_f}{M}\right)^{1/3} + \left(1 - \frac{a_f}{M}\right)^{1/3}\right] \\ Z_2 &\equiv \left(3\frac{a_f^2}{M^2} + Z_1^2\right)^{1/2} \\ r_{\text{ISCO}} &= M\{3 + Z_2 \mp [(3 - Z_1)(3 + Z_1 + 2Z_2)]^{1/2}\} \quad , \end{aligned} \quad (1)$$

where M is the total mass of the black hole and a_f is the angular momentum. The final spin is required in order to compute this ISCO. In [13], assuming the amount of mass and angular momentum radiated beyond the ISCO is small, the following implicit formula for the final angular momentum of a black hole a_f with component spins aligned with the orbit is calculated,

$$\frac{a_f}{M} = \frac{L_{\text{orb}}}{M^2} \left(q, \frac{r}{M} = \frac{r_{\text{ISCO}}}{M}, \frac{a_f}{M}\right) + \frac{q^2\chi_1 + \chi_2}{(1+q)^2} \quad , \quad (2)$$

where $\chi_i = a_i/m_i$, $q = m_1/m_2 \in [0, 1]$ and $M = m_1 + m_2$ is the total mass. The implicitly found a_f agrees well with numerical simulations [13] and the analysis can be modified to include arbitrary spin angles. L_{orb} is the orbital angular momentum

contribution calculated from the orbital angular momentum of a particle at the ISCO of a Kerr black hole with spin parameter a_f , which has the following expression [20],

$$\frac{L_{orb}}{M^2} \left(q, \frac{r}{M}, \frac{a_f}{M} \right) = \frac{q}{(1+q)^2} \frac{\pm(r^2 \mp 2a_f M^{1/2} r^{1/2} + a_f^2)}{M^{1/2} r^{3/4} (r^{3/2} - 3Mr^{1/2} \pm 2a_f M^{1/2})^{1/2}} , \quad (3)$$

where the upper/lower signs correspond to prograde/retrograde orbits. In order to agree with numerical simulations this function has to be evaluated at $r = r_{ISCO}$ given by equation (1). Alternative radii in the above prescription – like the photon-ring radius or Schwarzschild radius – give a prediction that is off the numerically obtained value. This fact, combined with the same analysis extended to elliptic binaries, again depending sensitively on the ISCO location [23], suggests the ISCO radius determined by this approach is relevant[24].

In order to get the gravitational wave frequency at ISCO we use the coordinate angular velocity of a circular orbit [20],

$$\Omega = \pm \frac{M^{1/2}}{r^{3/2} \pm a_f M^{1/2}} , \quad (4)$$

with a_f given by the implicit equation (2). The gravitational wave frequency at a given radius is then $f = \Omega/\pi$, and so we define $f_{ISCO[BKL]}$ as the frequency obtained by solving the system of equations (1-4) at the Kerr ISCO radius.

The solution space for $f_{ISCO[BKL]}$ can be written as a function of the final unknown spin a_f . For convenience, we prefer to extend it as a surface parametrized by (a_f, q, χ) , as it is shown in figure 1. This way the mass ratio dependence of the lines corresponding to different individual spins can be seen explicitly. For the equal mass case without spin the approximate expression for the Kerr test particle ISCO frequency is,

$$f_{ISCO[BKL]}(M, q) \approx (0.8q^3 - 2.6q^2 + 2.8q + 1) \times \frac{1}{\pi(6M)^{3/2}} , \quad (5)$$

which can be compared to the Schwarzschild test particle ISCO $f_{ISCO[SCH]} = \pi^{-1}(6M)^{-3/2}$.

Once the final spin is known it is possible to compute the last stable photon orbit (“light ring”) and the quasi-normal mode ring-down frequencies [16]. The light ring for a Kerr black hole is, [20]

$$r_{light} = 2M \{1 + \cos [2/3 \cos^{-1} (\mp a_f/M)]\} \quad (6)$$

$$\Omega_{light} = \pm \frac{M^{1/2}}{r_{light}^{3/2} \pm a_f M^{1/2}} , \quad (7)$$

An approximate fit to the quasinormal mode ring-down frequencies as a function of the final spin is given in [25, 26, 27]. For the $l=2, m=2, n=0$ mode we have,

$$\Omega_{QNM} \approx \frac{0.5}{M} (1 - 0.63(1 - a_f)^{0.3}) , \quad (8)$$

Others ring-down modes could be computed as necessary.

Figure 2 compares the BKL ISCO for non-spinning binaries with the ISCOs computed from the minimum energy conditions at 3PN order [19] as well as 2PN and




Figure 1. The surface of solutions of the frequencies at the ISCO as a function of the mass ratio q and the final spin a_f for components with spins that are aligned with the orbital angular momentum. Also shown, are curves corresponding to the solution of the equal spin case $\chi_1 = \chi_2 = \chi$.

p4PN EOB ISCOs [15, 16]. BKL and EOB agree exactly with the Schwarzschild ISCO at the extreme mass ratio limit, but the PN calculations are known to be inconsistent [19] in that regime. Predicting the final spin gives the $l = m = 2$ ring-down mode frequency by (8) and the light ring frequency. Both are shown in figure 2. The ISCO and light ring set a natural “merger” epoch which can be searched for by burst search techniques [28] *in between* the inspiral and ring-down matched filter searches. The ring-down frequencies shown in [16] agree well with these predictions. Although figure 2 is plotted for the non-spinning case the formalism described above can be easily generalized.

It has been shown [29, 30, 32] to be useful to understand un-modeled time-frequency evolutions in gravitational wave analysis in terms of time-frequency volumes, $\Delta f \Delta t$. This concept can provide insight into the merger epochs of various systems. We will consider three timescales for the merger 1) the Newtonian free fall time scale t_{ff} , 2) the quadrupole radiation time scale t_{quad} and 3) the time scale of a test particle following its geodesic just inside the ISCO of the Kerr space-time (corresponding to the merger product). For comparison we also plot the time-frequency volumes found from the




Figure 2. ISCO and QNM frequency estimates for non-spinning binaries as a function of mass ratio using different methods. Knowing the final spin of the black hole gives the expected ring-down frequency which agrees with the light ring frequency. The ISCO frequencies are very different depending on the method. The PN minimum energy condition gives an inconsistent result in the extreme mass ratio limit whereas the other methods (EOB, BKL) agree with the Schwarzschild ISCO at small values of q . The ISCO and QNM frequencies define a natural merger epoch which can be analyzed, albeit sub-optimally, even without knowing the numerical waveform.

figures in [16]. The Newtonian free fall time-frequency volume is,

$$\Delta f_{\text{ISCO}} \Delta t_{\text{ff}} = \frac{2^{-3/2}}{\sqrt{M\Omega_{\text{ISCO}}}} . \quad (9)$$

The quadrupole time-frequency volume is calculated by assuming that an ISCO doesn't exist and that the system continues to be driven by radiation. To calculate this we use the quadrupole approximation to the inspiral waveform given by [31], which gives the frequency evolution as,

$$M\Omega = \left[\frac{5(1+q)^2}{256q} \frac{M}{t-t_o} \right]^{3/8} \quad (10)$$

This leads to the time frequency volume,

$$\Delta f_{\text{BKLISCO}} \Delta t_{\text{quad}} = \frac{5(1+q)^2}{256\pi q} \left[(M\Omega_{\text{ISCO}})^{-8/3} - (M\Omega_{\text{light}})^{-8/3} \right] \times [(M\Omega_{\text{light}}) - (M\Omega_{\text{ISCO}})] , \quad (11)$$

where the ISCO frequencies and light ring frequencies are obtained from the BKL approximation as in figure 2.

The merger evolution will not be purely dominated by radiation by virtue that there are no circular orbits below the ISCO. The numbers obtained by this estimate should greatly over predict the time scale for extreme mass ratios where particles would take an asymptotically infinite time to fall in. The free fall time scale will under predict the time since the system may still complete some (unstable) orbits before merging. The estimate that agrees best with numerical relativity simulations is the infall time of a test particle falling from a circular orbit just below the ISCO to the light ring of the the merged Kerr black hole. Figure 3 shows the time-frequency volume for the merger epochs as a function of mass ratio for non-spinning binaries. The figure shows that the numbers taken from the simulations presented in [16] agree well with the predictions from the test particle in the Kerr space-time. By these estimates the mergers from yet un-simulated evolutions (low mass ratio) are likely to have the largest time frequency content and searches may benefit from the numerical simulations of these [30]. This information should be useful in conducting IMR searches and also in guiding the construction of analytic full waveform models. It is of course possible to repeat this analysis for arbitrary spin configurations using the above prescription and others [13, 21, 22, 33]. We will leave further discussion of these matters to future work and will spend the rest of this work considering the implications of using the ISCO found from the above set of equations as a termination frequency for PN waveforms in initial LIGO data analysis.




Figure 3. An estimate of the time-frequency volume for the merger epoch of non-spinning binary black holes. Numerical relativity results agree well with the time-frequency volume of a test particle falling into the merged Kerr black hole from a circular orbit just inside the ISCO.

3. Possible impact on data analysis for the inspiral phase of compact binary coalescence searches

As mentioned previously, there is ambiguity on how to define when the inspiral phase ends. However, what is really necessary is the characteristic frequency at which a given template waveform ceases to resemble the numerical simulations. Some searches for low mass binary systems of non-spinning component masses use the ISCO of a test particle orbiting a Schwarzschild black hole [14]. Other searches use the ISCOs calculated from explicit PN energy considerations, and some abandon the use of an ISCO altogether and use the Schwarzschild light ring as a termination frequency [15].

Recall that for low mass ratio ($q \sim 0$) systems of non-spinning objects, the Schwarzschild test particle limit is a good approximation for the expected ISCO frequency since the merger product will be a Schwarzschild-like black hole. However, for systems with comparable masses and/or spin the true ISCO frequency may be different since the non-trivial contribution from the orbital angular momentum will have a strong impact on the final black hole's spin, and the space-time in the near merger epoch. Many have addressed how well various PN approximants stay faithful to numerical relativity solutions [11, 41, 42, 43]. Most approximants stay faithful through the Schwarzschild ISCO frequency [43]. Some approximants fare extremely well beyond this point [43].

The fact that some approximants do remain faithful far beyond the Schwarzschild ISCO leads us to examine using the ISCO frequency described in the previous section, equation (5), as the termination frequency for inspiral data analysis. As shown, this frequency is consistent with some of the PN and EOB models for predicting the ISCO in the equal mass regime (fig. 2) and is consistent with exact solutions in the test particle limit. It also has the advantage of being waveform model, or fit, independent (based on first principles) and is easy to calculate. We will conservatively model the impact of possible phase errors incurred by using an approximant that doesn't quite match numerical relativity results in our assessment.

We use the stationary phase approximation waveforms, which are often employed in searches, and can be defined as,

$$\tilde{h}(f) \propto f^{-7/6} e^{i\Psi(f;M,\mu)} \quad , \quad (12)$$

where M is the total mass and μ is the reduced mass [14]. In order to ascertain the phase fidelity we turn to comparisons given in [11] which show for equal mass binary simulations the phase difference between numerical results and 3.5 PN waveforms as a function of frequency. The figures in [11] indicate a good fidelity in phase through the Schwarzschild ISCO frequency. However substantial phase errors may accumulate between the Schwarzschild and BKL ISCO frequencies. In order to accurately predict the impact of integrating PN waveforms to the BKL ISCO we decided to model the possibility that secular phase errors could be $\pm\pi$ or more radians between the Schwarzschild and BKL ISCOs. We propose calculating the SNR ratio of an ensemble of waveforms where phase errors are allowed to accumulate linearly between the

Schwarzschild ISCO $f_2(M, q)$ and the BKL ISCO $f_1(M, q)$ given by,

$$\frac{\text{SNR}_{f_1}}{\text{SNR}_{f_2}} = \left\langle \left(\int_{f_{\text{low}}}^{f_1} \frac{e^{\pi i R_n (f_1 - f_2) / f_1} \Theta(f_1 - f_2) \, df}{f^{7/3} S_n(f)} \right)^{1/2} \left(\int_{f_{\text{low}}}^{f_2} \frac{df}{f^{7/3} S_n(f)} \right)^{-1/2} \right\rangle, \quad (13)$$

where R_n is a normally distributed variable with mean 0 and variance 1. f_{low} is 40Hz [6].
 \ddagger We note that the above calculation is a conservative estimate of the SNR gain for two reasons: 1) A template with different parameters may match the numerical waveforms better, producing better phase agreement. Since search results are maximized over SNR this is an important possibility. 2) The phase errors do not accumulate linearly between these frequencies, most of the phase error occurs near the BKL ISCO. Using the $\{M, q\}$ dependent ISCO frequencies as $f_1(M, q)$ and the Schwarzschild ISCO frequencies as $f_2(M)$ we plot the expected SNR ratio for initial LIGO [35] in figure 4, which shows that for some total mass and mass ratio combinations there is an appreciable gain in SNR by integrating to the new ISCO frequencies despite modeled phase errors. This calculation is not intended to suggest that SPA templates cut at the BKL ISCO produce the optimal gain in SNR, since other approximants or frequency cut offs could do better. Instead, it aims to simply illustrate the effect of integrating to the BKL ISCO as a function of M and q for present searches. The mass range is limited to $80 M_\odot$ so that at least two cycles exist in the waveform between 40Hz and the Schwarzschild ISCO frequency.

4. Probing the tidal disruption of neutron stars

If the BKL ISCO truly marks the transition from an inspiral, radiation-dominated evolution to a dynamical one then it does have some consequence for the ability to probe the tidal disruption of neutron stars through gravitational waves. Estimation of gravitational-wave frequencies for tidal disruption of NS-BH binaries may be useful for determining various properties of the neutron star, such as the radius [36, 37]. Knowing the radius of the neutron star would in turn provide information about its equation of state [38, 39]. However, as pointed out in [36], it would be difficult to extract information about the disruption unless it occurs before the binary reaches the ISCO. If the disruption occurs afterwards, its signature in the produced gravitational waves might be too weak for extracting accurate information. Thus, the ‘‘useful’’ cases are those for which the tidal-disruption frequency is less than the frequency at the plunge ($f_{\text{td}} < f_{\text{plunge}} \equiv f_{\text{ISCO}}$).

A more accurate calculation of f_{plunge} would imply a better estimate of the range of NS-BH binaries for which the disruption is more plausible to be measured. In this section we show how that range changes when the formalism of section 2 is applied, as compared to that in [36].

\ddagger We take $S_n(f)$ to be approximated analytically as [34],

$$S_n(f) \approx 9 \times 10^{-46} \left[\left(\frac{4.49f}{150} \right)^{-56} + 0.16 \left(\frac{f}{150} \right)^{-4.52} + 0.32 \left(\frac{f}{150} \right)^2 + 0.52 \right]. \quad (14)$$




Figure 4. The SNR ratio defined by equation (13) for the non-spinning case $\chi = 0$. The solid lines include phase errors of the form (13) whereas the dashed lines assume perfect phase coherence for comparison.

The frequencies obtained in [36] correspond to the extreme mass ratio case, $q = 0$, and non-spinning neutron star, $\chi_{\text{NS}} = 0$. In figure 5 we reproduce those results (compare to figure 2 of [36]), together with the values obtained using the formalism presented in this work. We also assume in this case that the neutron star is non-spinning, but with $q = m_{\text{NS}}/m_{\text{BH}} \neq 0$. The curves of R vs f_{td} are calculated using the formula given in equation. (3) of [36], for the same parameters as in that work: $m_{\text{NS}} = 1.4M_{\odot}$; $m_{\text{BH}} = 2.5, 10, 20, 40, 80M_{\odot}$. The circles over each curve specify where the plunge occurs for different values of χ_{BH} . Filled circles correspond to the extreme mass ratio approximation. The meaningful part of each curve will be that to the left of the white circles ($f_{\text{td}} < f_{\text{plunge}}$).

The plunge frequencies calculated in [36] for the case with $m_{\text{BH}} = 2.5M_{\odot}$ (black circles in the right-hand-side curve of figure 5) correspond to retrograde orbits. However, for that mass ratio ($q = 0.56$), the test particle orbiting the merger product ISCO does so in a prograde fashion due to the orbital angular momentum dominating the final spin. Thus, the final spin is aligned with the orbital angular momentum even though its initial spin is anti-aligned.

We now compare with the NR results obtained in [40]. We calculate f_{BKL} for the six cases considered in in that work, models A-F. The BKL frequencies are shown in table 1, together with the corresponding parameters q and $M \equiv M_{\text{BH}} + M_{\text{NS}}$ (χ_{BH} and




Figure 5. Radius of the neutron star vs. the disruption frequency. The circles indicate the points at which the plunge would occur; the filled circles correspond to f_{plunge} as calculated in [36], included here for comparing. The value of χ_{BH} is indicated next to the corresponding circle where possible, otherwise a dashed line was used to connect the value to the circle. Each curve is labeled with the value of the BH mass: $m_{\text{BH}} = 2.5 - 80M_{\odot}$.

χ_{NS} are set to zero as in [40]). Although the models studied in [40] expand only a small range of parameters; we see that, at least in that range, the frequencies obtained with the BKL approach are consistent with the NR results (See figure 7 of [40]) - that is they predict in every case a higher value than the measured f_{cut} caused by the tidal effects.

Table 1. BKL frequencies for the same parameters as models A-F of [40]. These values are consistent with the NR results - that is the BKL ISCO frequency is higher than the observable f_{cut} from the simulations [40]

Model	q	$M(M_{\odot})$	$f_{\text{BKL}}(\text{kHz})$	f_{cut} from [40]
A	0.327	5.277	1.38	1.16
B	0.327	5.244	1.39	1.41
C	0.328	5.311	1.37	0.92
D	0.392	4.623	1.65	1.14
E	0.392	4.594	1.66	1.40
F	0.281	5.929	1.18	1.09

5. Conclusion

The estimate for the spin of the black hole that results from BH-BH, BH-NS, and some NS-NS mergers found in [13] leads to an estimate for the ISCO, light ring, and quasinormal mode ring-down frequencies of a general compact binary system that is waveform model independent. Both frequencies have impact on searches for the inspiral, merger and ring-down epochs of the gravitational waves emitted by these systems. We have shown that the most interesting merger-epoch time-frequency volumes are for extreme mass ratios. More work may have to be done to model merger epochs for these systems. The formalism presented here to describe inspiral, merger and ring-down epochs can be extended to arbitrary spin configurations and may help guide the construction of better analytic models.

As for inspiral epoch searches alone, we have shown that for symmetric mass, non-spinning systems with greater than $45 M_{\odot}$ total mass there can be a 30% gain in SNR over using the Schwarzschild ISCO radius as is done currently [14] even including conservative phase errors. It is worth noticing that more general fittings can be computed in order to cover the full range of parameters (q, χ_1, χ_2) within the BKL approximation. Additionally we discussed cases where neutron star tidal disruption may be observed more easily through gravitational waves. We would like to stress that this work is open ended. Here we have only considered spins that are aligned and anti-aligned with the orbital angular momentum. It is certainly possible to extend this work to make predictions about more general cases, all of which should be compared with numerical relativity results as they are produced.

6. Acknowledgements

The authors would like to acknowledge L. Lehner for suggesting this project and for invaluable discussions. The authors also thank L. Lehner and F. Pretorius for providing the test particle time of flight used in section 2. Gabriela González, Patrick Brady, Alessandra Buonanno and Jolien Creighton provided motivating discussions and insightful comments. C. Hanna would like to thank the LIGO Scientific Collaboration Compact Binary Coalescence (LIGO CBC) working group. This work was supported in part by NSF grants PHY-0605496, PHY-0653369 and PHY-0653375 to Louisiana State University and PHY-0603762 to the University of Maryland. C. Hanna would like to thank the Kavli Institute for Theoretical Physics, for their hospitality, where some of this work was completed. The Kavli Institute is supported by NSF grant PHY05-51164.

7. References

- [1] “The last three minutes: Issues in gravitational-wave measurements of coalescing compact binaries.” C. Cutler et al., *Phys. Rev. Lett.* 70, 2984 (1993).
- [2] “Search templates for gravitational waves from inspiraling binaries: Choice of template spacing.” B. J. Owen. *Phys. Rev. D* 53, 6749 (1996).

- [3] “Analysis of LIGO data for gravitational waves from binary neutron stars.” The LIGO Scientific Collaboration. Phys. Rev. D 69, 122001 (2004).
- [4] “Search for gravitational waves from galactic and extra-galactic binary neutron stars.” The LIGO Scientific Collaboration. Phys. Rev. D 72, 082001 (2005).
- [5] “Search for gravitational waves from binary black hole inspirals in LIGO data.” The LIGO Scientific Collaboration. Phys. Rev. D 73, 062001 (2006).
- [6] “Search for gravitational waves from binary inspirals in S3 and S4 LIGO data.” The LIGO Scientific Collaboration <http://arxiv.org/abs/0704.3368> (2007).
- [7] “Inspirals, merger and ring-down of equal-mass black-hole binaries.” Alessandra Buonanno, Gregory B. Cook, Frans Pretorius <http://lanl.arxiv.org/abs/gr-qc/0610122> (2006).
- [8] “Inspirals, merger and ring-down of unequal mass black hole binaries: a multipolar analysis.” Emanuele Berti, Vitor Cardoso, Jose A. Gonzalez, Ulrich Sperhake, Mark Hannam, Sascha Husa, Bernd Bruegmann <http://arxiv.org/abs/gr-qc/0703053> (2007).
- [9] “Matched filtering of numerical relativity templates of spinning binary black holes.” Birjoo Vaishnav, Ian Hinder, Frank Herrmann, and Deirdre Shoemaker Phys. Rev. D 76, 084020 (2007).
- [10] “Binary black hole late inspiral: Simulations for gravitational wave observations.” J. G. Baker, S.T. McWilliams, J. R. van Meter, J. Centrella, D. Choi, B. J. Kelly, and M. Koppitz. Phys. Rev. D 75, 124024 (2007).
- [11] “A data-analysis driven comparison of analytic and numerical coalescing binary waveforms: non-spinning case.” Yi Pan et al arXiv:0704.1964v2 [gr-qc] (2007).
- [12] “Gravitational-wave data analysis using binary black-hole waveforms.” P. Ajith arXiv:0712.0343v1 [gr-qc]
- [13] “Estimating the final spin of a binary black hole coalescence.” Alessandra Buonanno, Lawrence E. Kidder, Luis Lehner <http://arxiv.org/abs/0709.3839> (2007).
- [14] “FINDCHIRP: an algorithm for detection of gravitational waves from inspiraling compact binaries” Allen, B.; Anderson, W. G.; Brady, P. R.; Brown, D. A.; Creighton, J. D. E. <http://arxiv.org/abs/gr-qc/0509116> (2005).
- [15] “Effective one-body approach to general relativistic two-body dynamics.” A. Buonanno and T Damour. Phys. Rev. D 59, 084006 (1999)
- [16] “Toward faithful templates for non-spinning binary black holes using the effective-one-body approach.” Buonanno, Alessandra; Pan, Yi; Baker, John G.; Centrella, Joan; Kelly, Bernard J.; McWilliams, Sean T.; van Meter, James R. <http://arxiv.org/abs/0706.3732> (2007)
- [17] “Innermost stable orbits for coalescing binary systems of compact objects” L E Kidder, C M Will and A G Wiseman. Class. Quantum Grav. 9 L125-L131 (1992)
- [18] “Innermost circular orbit of binary black holes at the third post-Newtonian approximation” Luc Blanchet. Phys.Rev. D65 (2002).
- [19] “Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries.” Luc Blanchet Living Rev. Relativity 9, 4. URL (cited on 2007-12-20): <http://www.livingreviews.org/lrr-2006-4> (2006)
- [20] “Rotating Black Holes: Locally Nonrotating Frames, Energy Extraction, and Scalar Synchrotron Radiation.” Bardeen, J. M., Press, W. H., and Teukolsky, S. A. ApJ, **178** 347 (1972).
- [21] “On the final spin from the coalescence of two black holes” Luciano Rezzolla, Enrico Barausse, Ernst Nils Dorband, Denis Pollney, Christian Reisswig, Jennifer Seiler, Sascha Husa. <http://xxx.arxiv.org/abs/0712.3541>
- [22] “The final spin from the coalescence of aligned-spin black-hole binaries.” L. Rezzolla, P. Diener, E. N. Dorband, D. Pollney, C. Reisswig, E. Schnetter, J. Seiler. <http://www.arxiv.org/abs/0710.3345> (2007)
- [23] “Eccentric binary black-hole mergers: The transition from inspiral to plunge in general relativity.” U. Sperhake, E. Berti, V. Cardoso, J. A. Gonzalez, B. Bruegmann, M. Ansorg <http://arxiv.org/abs/0710.3823> (2007).
- [24] Luis Lehner *private communication November 2007*.

- [25] “Gravitational-wave measurements of the mass and angular momentum of a black hole” Fernando Echeverria. *Phys. Rev. D* 40, 3194 - 3203 (1989)
- [26] “Search techniques for gravitational waves from black-hole ringdowns” Jolien D. E. Creighton. *Phys.Rev. D*60 022001 (1999)
- [27] “On gravitational-wave spectroscopy of massive black holes with the space interferometer LISA” Emanuele Berti, Vitor Cardoso, Clifford M. Will *Phys.Rev. D*73 064030 (2006)
- [28] “Search for gravitational-wave bursts in LIGO data from the fourth science run” B Abbott et al 2007 *Class. Quantum Grav.* 24 5343-5369 (2007)
- [29] “An excess power statistic for detection of burst sources of gravitational radiation” Warren G. Anderson, Patrick R. Brady, Jolien D. E. Creighton, Eanna E. Flanagan. *Phys.Rev. D*63 (2001)
- [30] “Learning about compact binary merger: the interplay between numerical relativity and gravitational-wave astronomy” Thomas Baumgarte, Patrick Brady, Jolien D E Creighton, Luis Lehner, Frans Pretorius, Ricky DeVoe. <http://arxiv.org/abs/gr-qc/0612100>
- [31] “Observing binary inspiral in gravitational radiation: One interferometer” Lee Samuel Finn and David Chernoff. *Phys. Rev. D* 47, 2198 - 2219 (1993)
- [32] “Measuring gravitational waves from binary black hole coalescences. I. Signal to noise for inspiral, merger, and ring-down” Éanna É. Flanagan and Scott A. Hughes *Phys. Rev. D* 57, 4535 - 4565 (1998)
- [33] “Binary black hole merger: symmetry and the spin expansion” Latham Boyle, Michael Kesden, Samaya Nissanke. <http://arxiv.org/abs/0709.0299>
- [34] “Comparison of search templates for gravitational waves from binary inspiral” Thibault Damour, Bala R. Iyer and B. S. Sathyaprakash *Phys. Rev. D* 63, 044023 (2001)
- [35] LIGO Science Requirement Document noise curve http://www.ligo.caltech.edu/~jzweizig/distribution/LSC_Data/srd.html.
- [36] “Prospects for gravitational-wave observation of neutron-star tidal disruption in neutron-star/black-hole binaries.” Vallisneri, M. *Phys.Rev.Lett.* **84** (2000) 3519
- [37] “Possible direct method to determine the radius of a star from the spectrum of gravitational wave signals II : Spectra for various cases.” Motoyuki Saijo, Takashi Nakamura *Phys.Rev. D*63 (2001) 064004.
- [38] “Determining the nuclear equation of state from neutron-star masses and radii.” Lindblom, Lee *Astrophysical Journal, Part 1* (ISSN 0004-637X), vol. 398, no. 2, p. 569-573 (1992).
- [39] “Reconstructing the equation of state for cold nuclear matter from the relationship of any two properties of neutron stars.” Tomohiro Harada *Phys.Rev. D*64 (2001) 048801
- [40] “Merger of black hole and neutron star in general relativity: Tidal disruption, torus mass, and gravitational waves.” Masaru Shibata and Keisuke Taniguchi [arXiv:0711.1410v1](https://arxiv.org/abs/0711.1410v1) [gr-qc]
- [41] “Where post-Newtonian and numerical-relativity waveforms meet.” Mark Hannam, Sascha Husa, Jose A. Gonzalez, Ulrich Sperhake, Bernd Bruegmann <http://arxiv.org/abs/0706.1305> (2007).
- [42] “Consistency of post-Newtonian waveforms with numerical relativity.” John G. Baker, James R. van Meter, Sean T. McWilliams, Joan Centrella, Bernard J. Kelly *Phys. Rev. Lett.* vol. 99, 181101 (2007).
- [43] “High-accuracy comparison of numerical relativity simulations with post-Newtonian expansions.” Boyle et al [arXiv:0710.0158v1](https://arxiv.org/abs/0710.0158v1) [gr-qc].