

Necessary and sufficient optimality conditions for relaxed and strict control problems of forward-backward systems*

Seid BAHALALI

Laboratory of Applied Mathematics, University Med Khider,
Po. Box 145, Biskra 07000, Algeria.
sbahlali@yahoo.fr

December 3, 2018

Abstract

Stochastic maximum principle of nonlinear controlled forward-backward systems, where the set of strict (classical) controls need not be convex and the coefficients depend explicitly on the variable control, is an open problem impossible to solve by the classical method of spike variation. In this paper, we introduce a new approach to solve this open problem and we establish necessary as well as sufficient conditions of optimality, in the form of global stochastic maximum principle, for two models. The first concerns the relaxed controls, who are a measure-valued processes. The second is a restriction of the first to strict control problems.

AMS Subject Classification. 93 Exx

Keywords. Forward-backward stochastic differential equations, Stochastic maximum principle, Strict control, Relaxed control, Adjoint equations, Variational inequality.

1 Introduction

We study a stochastic control problem where the system is governed by a non-linear forward-backward stochastic differential equation (FBSDE for short) of the type

$$\begin{cases} dx_t^v = b(t, x_t^v, v_t) dt + \sigma(t, x_t^v, v_t) dW_t, \\ x_0^v = \xi, \\ dy_t = -f(t, x_t^v, y_t^v, z_t^v, v_t) dt + z_t^v dW_t, \\ y_T = \varphi(x_T^v), \end{cases}$$

*This work is partially supported by Algerian-French cooperation, Tassili 07 MDU 705.

where b , σ , f and φ are given maps, $W = (W_t)_{t \geq 0}$ is a standard Brownian motion, defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$, satisfying the usual conditions.

The control variable $v = (v_t)$, called strict (classical) control, is an \mathcal{F}_t adapted process with values in some set U of \mathbb{R}^k . We denote by \mathcal{U} the class of all strict controls.

The criteria to be minimized, over the set \mathcal{U} , has the form

$$J(v) = \mathbb{E} \left[g(x_T^v) + h(y_0^v) + \int_0^T l(t, x_t^v, y_t^v, z_t^v, v_t) dt \right],$$

where g , h and l are given functions and (x_t^v, y_t^v, z_t^v) is the trajectory of the system controlled by v .

A control $u \in \mathcal{U}$ is called optimal if it satisfies

$$J(u) = \inf_{v \in \mathcal{U}} J(v).$$

Stochastic control problems for backward and forward-backward systems have been studied by many authors including Peng [37], Xu [41], El-Karoui et al [14, 15], Wu [40], Dokuchaev and Zhou [10], Peng and Wu [38], Bahlali and Labed [4], Bahlali [7], Shi and Wu [39], Ji and Zhou [26]. The dynamic programming approaches have been studied by Fuhrman and Tessitore [19].

All the previous works on stochastic maximum principle of forward-backward systems are established in the cases where the control domain is convex or uncontrolled diffusion coefficient. The general case, where the set of controls need not be convex and the diffusion coefficient depends explicitly on the control variable, is an open problem unsolved until now. There is no result in the literature concerning this problem, because the classical way which consists to use the spike variation method on the strict controls does not lead to any result. In this paper, we introduce a new approach to solve this open problem. The main idea is to use a bigger new class \mathcal{R} of processes by replacing the U -valued process (v_t) by a $\mathbb{P}(U)$ -valued process (q_t) , where $\mathbb{P}(U)$ is the space of probability measures on U equipped with the topology of weak convergence. This new class of processes is called relaxed controls and have a richer structure of compacity and convexity, for which the control problem becomes solvable.

In the relaxed model, the system is governed by the FBSDE

$$\begin{cases} dx_t^q = \int_U b(t, x_t^q, a) q_t(da) dt + \int_U \sigma(t, x_t^q, a) q_t(da) dW_t, \\ x_0^q = \xi, \\ dy_t^q = - \int_U f(t, x_t^q, y_t^q, z_t^q, a) q_t(da) dt + z_t^q dW_t, \\ y_T^q = \varphi(x_T^q). \end{cases}$$

The functional cost to be minimized, over the class \mathcal{R} of relaxed controls, is defined by

$$J(q) = \mathbb{E} \left[g(x_T^q) + h(y_0^q) + \int_0^T \int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) dt \right].$$

A relaxed control μ is called optimal if it solves

$$J(\mu) = \inf_{q \in \mathcal{R}} J(q).$$

The relaxed control problem is a generalization of the problem of strict controls. Indeed, if $q_t(da) = \delta_{v_t}(da)$ is a Dirac measure concentrated at a single point $v_t \in U$, then we get a strict control problem as a particular case of the relaxed one.

Our aim in this paper, is to establish necessary as well as sufficient conditions of optimality in the form of global stochastic maximum principle, for both relaxed and strict controls. To achieve this goal, we derive these results as follows.

Firstly, we give the optimality conditions for relaxed controls. The idea is to use the fact that the set of relaxed controls is convex. Then, we establish necessary optimality conditions by using the classical way of the convex perturbation method. More precisely, if we denote by μ an optimal relaxed control and q is an arbitrary element of \mathcal{R} , then with a sufficiently small $\theta > 0$ and for each $t \in [0, T]$, we can define a perturbed control as follows

$$\mu_t^\theta = \mu_t + \theta(q_t - \mu_t).$$

We derive the variational equation from the state equation, and the variational inequality from the inequality

$$0 \leq J(\mu^\theta) - J(\mu).$$

By using the fact that the coefficients b, σ, f and l are linear with respect to the relaxed control variable, necessary optimality conditions are obtained directly in the global form.

To achieve this part of the paper, we prove under minimal additional hypothesis, that these necessary optimality conditions for relaxed controls are also sufficient.

The second main result in the paper characterizes the optimality for strict control processes. It is directly derived from the above result by restricting from relaxed to strict controls. The idea is to replace the relaxed controls by a Dirac measures charging a strict controls. Thus, we reduce the set \mathcal{R} of relaxed controls and we minimize the cost J over the subset $\delta(\mathcal{U}) = \{q \in \mathcal{R} \mid q = \delta_v ; v \in \mathcal{U}\}$. Necessary optimality conditions for strict controls are then obtained directly from those of relaxed one. Finally, we prove that these necessary conditions becomes sufficient, without imposing neither the convexity of U nor that of the Hamiltonian H in v .

The paper is organized as follows. In Section 2, we formulate the strict and relaxed control problems and give the various assumptions used throughout the paper. Section 3 is devoted to study the relaxed control problems and we establish necessary as well as sufficient conditions of optimality for relaxed controls. In the last section, we derive directly from the results of Section 3, the optimality conditions for strict controls.

Along this paper, we denote by C some positive constant and we need the following matrix notations. We denote by $\mathcal{M}_{n \times d}(\mathbb{R})$ the space of $n \times d$ real matrices and by $\mathcal{M}_{n \times n}^d(\mathbb{R})$ the linear space of vectors $M = (M_1, \dots, M_d)$ where $M_i \in \mathcal{M}_{n \times n}(\mathbb{R})$.

For any $M, N \in \mathcal{M}_{n \times n}^d(\mathbb{R})$, $L, S \in \mathcal{M}_{n \times d}(\mathbb{R})$, $Q \in \mathcal{M}_{n \times n}(\mathbb{R})$, $\alpha, \beta \in \mathbb{R}^n$ and $\gamma \in \mathbb{R}^d$, we use the following notations

$$\alpha\beta = \sum_{i=1}^n \alpha_i \beta_i \in \mathbb{R} \text{ is the product scalar in } \mathbb{R}^n;$$

$$LS = \sum_{i=1}^d L_i S_i \in \mathbb{R}, \text{ where } L_i \text{ and } S_i \text{ are the } i^{th} \text{ columns of } L \text{ and } S;$$

$$ML = \sum_{i=1}^d M_i L_i \in \mathbb{R}^n;$$

$$M\alpha\gamma = \sum_{i=1}^d (M_i \alpha) \gamma_i \in \mathbb{R}^n;$$

$$MN = \sum_{i=1}^d M_i N_i \in \mathcal{M}_{n \times n}(\mathbb{R});$$

$$MQN = \sum_{i=1}^d M_i Q N_i \in \mathcal{M}_{n \times n}(\mathbb{R});$$

$$MQ\gamma = \sum_{i=1}^d M_i Q \gamma_i \in \mathcal{M}_{n \times n}(\mathbb{R}).$$

We denote by L^* the transpose of the matrix L and $M^* = (M_1^*, \dots, M_d^*)$.

2 Formulation of the problem

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$ be a filtered probability space satisfying the usual conditions, on which a d -dimensional Brownian motion $W = (W_t)_{t \geq 0}$ is defined. We assume that (\mathcal{F}_t) is the \mathcal{P} -augmentation of the natural filtration of W .

Let T be a strictly positive real number and U a non-empty set of \mathbb{R}^k .

2.1 The strict control problem

Definition 1 *An admissible strict control is an \mathcal{F}_t -adapted process $v = (v_t)$ with values in U such that*

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |v_t|^2 \right] < \infty.$$

We denote by \mathcal{U} the set of all admissible strict controls.

For any $v \in \mathcal{U}$, we consider the following controlled FBSDE

$$\begin{cases} dx_t^v = b(t, x_t^v, v_t) dt + \sigma(t, x_t^v, v_t) dW_t, \\ x_0^v = \xi, \\ dy_t^v = -f(t, x_t^v, y_t^v, z_t^v, v_t) dt + z_t^v dW_t, \\ y_T^v = \varphi(x_T^v), \end{cases} \quad (1)$$

where,

$$\begin{aligned} b &: [0, T] \times \mathbb{R}^n \times U \longrightarrow \mathbb{R}^n, \\ \sigma &: [0, T] \times \mathbb{R}^n \times U \longrightarrow \mathcal{M}_{n \times d}(\mathbb{R}), \\ f &: [0, T] \times \mathbb{R}^n \times \mathbb{R}^m \times \mathcal{M}_{m \times d}(\mathbb{R}) \times U \longrightarrow \mathbb{R}^m, \\ \varphi &: \mathbb{R}^n \longrightarrow \mathbb{R}^m, \end{aligned}$$

and ξ is an n -dimensional \mathcal{F}_0 -measurable random variable such that

$$\mathbb{E} |\xi|^2 < \infty.$$

The criteria to be minimized is defined from \mathcal{U} into \mathbb{R} by

$$J(v) = \mathbb{E} \left[g(x_T^v) + h(y_0^v) + \int_0^T l(t, x_t^v, y_t^v, z_t^v, v_t) dt \right], \quad (2)$$

where,

$$\begin{aligned} g &: \mathbb{R}^n \longrightarrow \mathbb{R}, \\ h &: \mathbb{R}^m \longrightarrow \mathbb{R}, \\ l &: [0, T] \times \mathbb{R}^n \times \mathbb{R}^m \times \mathcal{M}_{m \times d}(\mathbb{R}) \times U \longrightarrow \mathbb{R}. \end{aligned}$$

A strict control u is called optimal if it satisfies

$$J(u) = \inf_{v \in \mathcal{U}} J(v). \quad (3)$$

We assume that

$$\begin{aligned} b, \sigma, f, g, h, l, \varphi &\text{ are continuously differentiable with respect to } (x, y, z), \\ &\text{they are bounded by } C(1 + |x| + |y| + |z| + |v|) \text{ and} \\ &\text{their derivatives with respect to } (x, y, z) \text{ are continuous and bounded.} \end{aligned} \quad (4)$$

Under the above hypothesis, for every $v \in U$, equation (1) has a unique strong solution and the functional cost J is well defined from \mathcal{U} into \mathbb{R} .

2.2 The relaxed model

Definition 2 A relaxed control $(q_t)_t$ is a $\mathbb{P}(U)$ -valued process, progressively measurable with respect to $(\mathcal{F}_t)_t$ and such that for each t , $1_{[0, t]} \cdot q$ is \mathcal{F}_t -measurable. We denote by \mathcal{R} the set of all relaxed controls.

Remark 3 *The set of strict controls is embedded into the set of relaxed controls by the mapping*

$$f : v \longmapsto f_v(dt, da) = dt\delta_{v_t}(da),$$

where δ_v is the atomic measure concentrated at a single point v .

For more details on relaxed controls, see [3], [5], [6], [12], [17], [29], [32], [33].

For any $q \in \mathcal{R}$, we consider the following relaxed FBSDE

$$\begin{cases} dx_t^q = \int_U b(t, x_t^q, a) q_t(da) dt + \int_U \sigma(t, x_t^q, a) q_t(da) dW_t, \\ x_0^q = \xi, \\ dy_t^q = - \int_U f(t, x_t^q, y_t^q, z_t^q, a) q_t(da) dt + z_t^q dW_t, \\ y_T^q = \varphi(x_T^q). \end{cases} \quad (5)$$

The expected cost to be minimized, in the relaxed model, is defined from \mathcal{R} into \mathbb{R} by

$$J(q) = \mathbb{E} \left[g(x_T^q) + h(y_0^q) + \int_0^T \int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) dt \right]. \quad (6)$$

A relaxed control μ is called optimal if it solves

$$J(\mu) = \inf_{q \in \mathcal{R}} J(q). \quad (7)$$

Remark 4 *If we put*

$$\begin{aligned} \bar{b}(t, x_t^q, q_t) &= \int_U b(t, x_t^q, a) q_t(da), \\ \bar{\sigma}(t, x_t^q, q_t) &= \int_U \sigma(t, x_t^q, a) q_t(da), \\ \bar{f}(t, x_t^q, q_t) &= \int_U f(t, x_t^q, y_t^q, z_t^q, a) q_t(da), \\ \bar{l}(t, x_t^q, q_t) &= \int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da). \end{aligned}$$

Then, equation (5) becomes

$$\begin{cases} dx_t^q = \bar{b}(t, x_t^q, q_t) dt + \bar{\sigma}(t, x_t^q, q_t) dW_t, \\ x_0^q = \xi, \\ dy_t^q = -\bar{f}(t, x_t^q, y_t^q, z_t^q, q_t) dt + z_t^q dW_t, \\ y_T^q = \varphi(x_T^q). \end{cases} \quad (5')$$

With a functional cost given by

$$J(q) = \mathbb{E} \left[g(x_T^q) + h(y_0^q) + \int_0^T \bar{l}(t, x_t^q, y_t^q, z_t^q, q_t) dt \right]. \quad (8)$$

Hence, by introducing relaxed controls, we have replaced U by a larger space $\mathbb{P}(U)$. We have gained the advantage that $\mathbb{P}(U)$ is both compact and convex. Furthermore, the new coefficients of equation (5) and the running cost are linear with respect to the relaxed control variable.

Remark 5 The coefficients $\bar{b}, \bar{\sigma}$ and \bar{f} (defined in the above remark) check respectively the same assumptions as b, σ and f . Then, under assumptions (4), $\bar{b}, \bar{\sigma}$ and \bar{f} are uniformly Lipschitz and with linear growth. Then by classical results on FBSDEs, for every $q \in \mathcal{R}$ equation (5') admits a unique strong solution. Consequently, for every $q \in \mathcal{R}$ equation (5) has a unique strong solution.

On the other hand, It is easy to see that \bar{l} checks the same assumptions as l . Then, the functional cost J is well defined from \mathcal{R} into \mathbb{R} .

Remark 6 If $q_t = \delta_{v_t}$ is an atomic measure concentrated at a single point $v_t \in U$, then for each $t \in [0, T]$ we have

$$\begin{aligned} \int_U b(t, x_t^q, a) q_t(da) &= \int_U b(t, x_t^q, a) \delta_{v_t}(da) = b(t, x_t^q, v_t), \\ \int_U \sigma(t, x_t^q, a) q_t(da) &= \int_U \sigma(t, x_t^q, a) \delta_{v_t}(da) = \sigma(t, x_t^q, v_t), \\ \int_U f(t, x_t^q, y_t^q, z_t^q, a) q_t(da) &= \int_U f(t, x_t^q, y_t^q, z_t^q, a) \delta_{v_t}(da) = f(t, x_t^q, y_t^q, z_t^q, v_t), \\ \int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) &= \int_U l(t, x_t^q, y_t^q, z_t^q, a) \delta_{v_t}(da) = l(t, x_t^q, y_t^q, z_t^q, v_t). \end{aligned}$$

In this case $(x^q, y^q, z^q) = (x^v, y^v, z^v)$, $J(q) = J(v)$ and we get a strict control problem. So the problem of strict controls $\{(1), (2), (3)\}$ is a particular case of relaxed control problem $\{(5), (6), (7)\}$.

Remark 7 The relaxed control problems studied e.g. in El Karoui et al [12] and Bahlali-Mezzerdi-Djehiche [3] is different to ours, in that they relax the corresponding infinitesimal generator of the state process, which leads to a martingale problem for which the state process driven by an orthogonal martingale measure. In our setting the driving martingale measure $q_t(da) dW_t$ is however not orthogonal. See Ma-Yong [29] for more details.

3 Necessary and sufficient optimality conditions for relaxed controls

In this section, we study the problem $\{(5), (6), (7)\}$ and we establish necessary as well as sufficient conditions of optimality for relaxed controls.

3.1 Preliminary results

Since the set \mathcal{R} is convex, then the classical way to derive necessary optimality conditions for relaxed controls is to use the convex perturbation method. More

precisely, let μ be an optimal relaxed control and $(x_t^\mu, y_t^\mu, z_t^\mu)$ the solution of (5) controlled by μ . Then, we can define a perturbed relaxed control as follows

$$\mu_t^\theta = \mu_t + \theta (q_t - \mu_t), \quad (9)$$

where, $\theta > 0$ is sufficiently small and q is an arbitrary element of \mathcal{R} .

Denote by $(x_t^\theta, y_t^\theta, z_t^\theta)$ the solution of (5) associated with μ^θ .

From optimality of μ , the variational inequality will be derived from the fact that

$$0 \leq J(\mu^\theta) - J(\mu). \quad (10)$$

For this end, we need the following classical lemmas.

Lemma 8 *Under assumptions (4), we have*

$$\lim_{\theta \rightarrow 0} \left[\sup_{0 \leq t \leq T} \mathbb{E} |x_t^\theta - x_t^\mu|^2 \right] = 0, \quad (11)$$

$$\lim_{\theta \rightarrow 0} \left[\sup_{0 \leq t \leq T} \mathbb{E} |y_t^\theta - y_t^\mu|^2 \right] = 0, \quad (12)$$

$$\lim_{\theta \rightarrow 0} \mathbb{E} \int_0^T |z_t^\theta - z_t^\mu|^2 dt = 0. \quad (13)$$

Proof. We have

$$\begin{aligned} x_t^\theta - x_t^\mu &= \int_0^t \left[\int_U b(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U b(s, x_s^\mu, a) \mu_s(da) \right] ds \\ &\quad + \int_0^t \left[\int_U \sigma(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s(da) \right] dW_s \\ &= \int_0^t \left[\int_U b(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U b(s, x_s^\mu, a) \mu_s^\theta(da) \right] ds \\ &\quad + \int_0^t \left[\int_U b(s, x_s^\mu, a) \mu_s^\theta(da) - \int_U b(s, x_s^\mu, a) \mu_s(da) \right] ds \\ &\quad + \int_0^t \left[\int_U \sigma(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s^\theta(da) \right] dW_s \\ &\quad + \int_0^t \left[\int_U \sigma(s, x_s^\mu, a) \mu_s^\theta(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s(da) \right] dW_s. \end{aligned}$$

By using the definition of μ_t^θ and taking expectations, we have

$$\begin{aligned}\mathbb{E} |x_t^\theta - x_t^\mu|^2 &\leq C\mathbb{E} \int_0^t \left| \int_U b(s, x_s^\theta, a) \mu_s(da) - \int_U b(s, x_s^\mu, a) \mu_s(da) \right|^2 ds \\ &\quad + C\theta^2 \mathbb{E} \int_0^t \left| \int_U b(s, x_s^\theta, a) q_s(da) - \int_U b(s, x_s^\theta, a) \mu_s(da) \right|^2 ds \\ &\quad + C\mathbb{E} \int_0^t \left| \int_U \sigma(s, x_s^\theta, a) \mu_s(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s(da) \right|^2 ds \\ &\quad + C\theta^2 \mathbb{E} \int_0^t \left| \int_U \sigma(s, x_s^\theta, a) q_s(da) - \int_U \sigma(s, x_s^\theta, a) \mu_s(da) \right|^2 ds.\end{aligned}$$

By (4), b and σ are uniformly Lipschitz with respect to x . Hence,

$$\mathbb{E} |x_t^\theta - x_t^\mu|^2 \leq C\mathbb{E} \int_0^t |x_s^\theta - x_s^\mu|^2 ds + C\theta^2.$$

By using Gronwall's lemma and Buckholder-Davis-Gundy inequality, we obtain (11).

Let us now prove (12) and (13).

Applying Itô's formula to $(y_t^\theta - y_t^\mu)^2$, we have

$$\begin{aligned}&\mathbb{E} |y_t^\theta - y_t^\mu|^2 + \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds = \mathbb{E} |\varphi(x_T^\theta) - \varphi(x_T^\mu)|^2 \\ &+ 2\mathbb{E} \int_t^T \left| (y_s^\theta - y_s^\mu) \left[\int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s^\theta(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\mu, a) \mu_s(da) \right] \right| ds.\end{aligned}$$

From the Young formula, for every $\varepsilon > 0$, we have

$$\begin{aligned}&\mathbb{E} |y_t^\theta - y_t^\mu|^2 + \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds \\ &\leq \mathbb{E} |\varphi(x_T^\theta) - \varphi(x_T^\mu)|^2 + \frac{1}{\varepsilon} \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds \\ &+ \varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s^\theta(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\mu, a) \mu_s(da) \right|^2 ds.\end{aligned}$$

Then,

$$\begin{aligned}
& \mathbb{E} |y_t^\theta - y_t^\mu|^2 + \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds \\
& \leq \mathbb{E} |\varphi(x_T^\theta) - \varphi(x_T^\mu)|^2 + \frac{1}{\varepsilon} \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s^\theta(da) - \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\theta, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\mu, y_s^\theta, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\mu, y_s^\mu, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\mu, a) \mu_s(da) \right|^2 ds.
\end{aligned}$$

By the definition of μ_t^θ , we have

$$\begin{aligned}
& \mathbb{E} |y_t^\theta - y_t^\mu|^2 + \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds \\
& \leq \mathbb{E} |\varphi(x_T^\theta) - \varphi(x_T)|^2 + \frac{1}{\varepsilon} \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds \\
& + C\varepsilon \theta^2 \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) q_s(da) - \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\theta, y_s^\theta, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\theta, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\mu, y_s^\theta, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\theta, a) \mu_s(da) \right|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T \left| \int_U f(s, x_s^\mu, y_s^\mu, z_s^\theta, a) \mu_s(da) - \int_U f(s, x_s^\mu, y_s^\mu, z_s^\mu, a) \mu_s(da) \right|^2 ds.
\end{aligned}$$

Since φ and f are uniformly Lipschitz with respect to x, y, z , then

$$\begin{aligned}
\mathbb{E} |y_t^\theta - y_t^\mu|^2 + \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds & \leq \left(\frac{1}{\varepsilon} + C\varepsilon \right) \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds \\
& + C\varepsilon \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds + \alpha_t^\theta,
\end{aligned} \tag{14}$$

where α_t^θ is given by

$$\alpha_t^\theta = \mathbb{E} |x_T^\theta - x_T^\mu|^2 + C\varepsilon \mathbb{E} \int_t^T |x_s^\theta - x_s^\mu|^2 ds + C\varepsilon \theta^2.$$

By (11), we have

$$\lim_{\theta \rightarrow 0} \alpha_t^\theta = 0. \quad (15)$$

Choose $\varepsilon = \frac{1}{2C}$, then (14) becomes

$$\mathbb{E} |y_t^\theta - y_t^\mu|^2 + \frac{1}{2} \mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds \leq \left(2C + \frac{1}{2}\right) \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds + \alpha_t^\theta.$$

From the above inequality, we derive two inequalities

$$\mathbb{E} |y_t^\theta - y_t^\mu|^2 \leq \left(2C + \frac{1}{2}\right) \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds + \alpha_t^\theta, \quad (16)$$

$$\mathbb{E} \int_t^T |z_s^\theta - z_s^\mu|^2 ds \leq (4C + 1) \mathbb{E} \int_t^T |y_s^\theta - y_s^\mu|^2 ds + 2\alpha_t^\theta. \quad (17)$$

By using (15), (16), Gronwall's lemma and Burkholder-Davis-Gundy inequality, we obtain (12). Finally, (13) is derived from (15) and (12). ■

Lemma 9 *Let \tilde{x}_t and \tilde{y}_t are respectively the solutions of the following linear equations (called variational equations)*

$$\begin{cases} d\tilde{x}_t = \int_U b_x(t, x_t^\mu, a) \mu_t(da) \tilde{x}_t dt + \int_U \sigma_x(t, x_t^\mu, a) \mu_t(da) \tilde{x}_t dW_t \\ \quad + \left[\int_U b(t, x_t^\mu, a) \mu_t(da) - \int_U b(t, x_t^\mu, a) q_t(da) \right] dt \\ \quad + \left[\int_U \sigma(t, x_t^\mu, a) \mu_t(da) - \int_U \sigma(t, x_t^\mu, a) q_t(da) \right] dW_t, \\ \tilde{x}_0 = 0. \end{cases} \quad (18)$$

$$\begin{cases} d\tilde{y}_t = - \int_U [f_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{x}_t + f_y(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{y}_t + f_z(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{z}_t] \mu_t(da) dt \\ \quad + \left[\int_U f(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) - \int_U f(t, x_t^\mu, y_t^\mu, z_t^\mu, a) q_t(da) \right] dt + \tilde{z}_t dW_t, \\ \tilde{y}_T = \varphi_x(x_T^\mu) \tilde{x}_T. \end{cases} \quad (19)$$

Then, the following estimations hold

$$\lim_{\theta \rightarrow 0} \mathbb{E} \left| \frac{x_t^\theta - x_t^\mu}{\theta} - \tilde{x}_t \right|^2 = 0, \quad (20)$$

$$\lim_{\theta \rightarrow 0} \mathbb{E} \left| \frac{y_t^\theta - y_t^\mu}{\theta} - \tilde{y}_t \right|^2 = 0, \quad (21)$$

$$\lim_{\theta \rightarrow 0} \mathbb{E} \int_0^T \left| \frac{z_t^\theta - z_t^\mu}{\theta} - \tilde{z}_t \right|^2 dt = 0. \quad (22)$$

Proof. For simplicity, we put

$$X_t^\theta = \frac{x_t^\theta - x_t^\mu}{\theta} - \tilde{x}_t, \quad (23)$$

$$Y_t^\theta = \frac{y_t^\theta - y_t^\mu}{\theta} - \tilde{y}_t, \quad (24)$$

$$Z_t^\theta = \frac{z_t^\theta - z_t^\mu}{\theta} - \tilde{z}_t. \quad (25)$$

i) Proof of (20).

We have

$$\begin{aligned} X_t^\theta &= \frac{1}{\theta} \int_0^t \int_U b(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U b(s, x_s^\mu, a) \mu_s^\theta(da) ds \\ &\quad + \frac{1}{\theta} \int_0^t \left[\int_U b(s, x_s^\mu, a) \mu_s^\theta(da) - \int_U b(s, x_s^\mu, a) \mu_s(da) \right] ds \\ &\quad + \frac{1}{\theta} \int_0^t \left[\int_U \sigma(s, x_s^\theta, a) \mu_s^\theta(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s^\theta(da) \right] dW_s \\ &\quad + \frac{1}{\theta} \int_0^t \left[\int_U \sigma(s, x_s^\mu, a) \mu_s^\theta(da) - \int_U \sigma(s, x_s^\mu, a) \mu_s(da) \right] dW_s \\ &\quad - \int_0^t \int_U b_x(s, x_s^\mu, a) \mu_s(da) \tilde{x}_s ds - \int_0^t \int_U \sigma_x(s, x_s^\mu, a) \mu_s(da) \tilde{x}_s dW_s \\ &\quad - \int_0^t \left[\int_U b(s, x_s^\mu, a) \mu_s(da) - \int_U b(s, x_s^\mu, a) q_s(da) \right] ds \\ &\quad - \int_0^t \left[\int_U \sigma(s, x_s^\mu, a) \mu_s(da) - \int_U \sigma(s, x_s^\mu, a) q_s(da) \right] dW_s. \end{aligned}$$

By using the definition of μ^θ and taking expectations, we get

$$\begin{aligned} \mathbb{E} |X_t^\theta|^2 &\leq C \mathbb{E} \int_0^t \int_0^1 \int_U |b_x(s, x_s^\mu + \lambda \theta (X_s^\theta + \tilde{x}_s), a) X_s^\theta|^2 \mu_s(da) d\lambda ds \\ &\quad + C \mathbb{E} \int_0^t \int_0^1 \int_U |\sigma_x(s, x_s^\mu + \lambda \theta (X_s^\theta + \tilde{x}_s), a) X_s^\theta|^2 \mu_s(da) d\lambda ds \\ &\quad + C \mathbb{E} |\beta_t^\theta|^2, \end{aligned}$$

where, β_t^θ is given by

$$\begin{aligned}
\beta_t^\theta = & \int_0^t \int_0^1 \int_U b_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a)(x_s^\theta - x_s^\mu) q_s(da) d\lambda ds \\
& - \int_0^t \int_0^1 \int_U b_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a)(x_s^\theta - x_s^\mu) \mu_s(da) d\lambda ds \\
& + \int_0^t \int_0^1 \int_U \sigma_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a)(x_s^\theta - x_s^\mu) q_s(da) d\lambda dW_s \\
& - \int_0^t \int_0^1 \int_U \sigma_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a)(x_s^\theta - x_s^\mu) \mu_s(da) d\lambda dW_s \\
& + \int_0^t \int_0^1 \int_U b_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a) \tilde{x}_s \mu_s(da) d\lambda ds \\
& + \int_0^t \int_0^1 \int_U \sigma_x(s, x_s^\mu + \lambda\theta(X_s^\theta + \tilde{x}_s), a) \tilde{x}_s \mu_s(da) d\lambda dW_s \\
& - \int_0^t \int_U b_x(s, x_s^\mu, a) \tilde{x}_s \mu_s(da) ds - \int_0^t \int_U \sigma_x(s, x_s^\mu, a) \tilde{x}_s \mu_s(da) dW_s.
\end{aligned}$$

Since b_x and σ_x are continuous and bounded, then

$$\mathbb{E} |X_t^\theta|^2 \leq C \mathbb{E} \int_0^t |X_s^\theta|^2 ds + C \mathbb{E} |\beta_t^\theta|^2,$$

$$\lim_{\theta \rightarrow 0} \mathbb{E} |\beta_t^\theta|^2 = 0.$$

We obtain (20) by using Gronwall's lemma in the above inequality.

ii) Proof of (21) and (22).

For simplicity, we put

$$\Lambda_t^\theta(a) = (t, x_t^\mu + \lambda\theta(X_t^\theta + \tilde{x}_t), y_t^\mu + \lambda\theta(Y_t^\theta + \tilde{y}_t), z_t^\mu + \lambda\theta(Z_t^\theta + \tilde{z}_t), a).$$

By (24), we have the following FBSDE

$$\begin{cases} dY_t^\theta = (F_t^y Y_t^\theta dt + F_t^y Z_t^\theta - \gamma_t^\theta) dt + Z_t^\theta dW_t, \\ Y_T^\theta = \frac{\varphi(x_T^\theta) - \varphi(x_T^\mu)}{\theta} - \varphi_x(x_T^\mu) \tilde{x}_T, \end{cases}$$

where,

$$\begin{aligned}
F_t^y &= - \int_0^1 \int_U f_y(\Lambda_t^\theta(a)) \mu_t(da) d\lambda, \\
F_t^z &= - \int_0^1 \int_U f_z(\Lambda_t^\theta(a)) \mu_t(da) d\lambda,
\end{aligned}$$

and γ_t^θ is given by

$$\begin{aligned}\gamma_t^\theta &= \int_t^T \int_U f_x(\Lambda_s^\theta(a)) X_s^\theta \mu_s(da) ds \\ &+ \int_t^T \int_U [f_x(\Lambda_s^\theta(a)) (x_s^\theta - x_s^\mu) + f_y(\Lambda_s^\theta(a)) (y_s^\theta - y_s^\mu) + f_z(\Lambda_s^\theta(a)) (z_s^\theta - z_s^\mu)] q_s(da) ds \\ &- \int_t^T \int_U [f_x(\Lambda_s^\theta(a)) (x_s^\theta - x_s^\mu) + f_y(\Lambda_s^\theta(a)) (y_s^\theta - y_s^\mu) + f_z(\Lambda_s^\theta(a)) (z_s^\theta - z_s^\mu)] \mu_s(da) ds.\end{aligned}$$

Since f_x, f_y and f_z are continuous and bounded, then from (20), (11), (12) and (13), we have

$$\lim_{\theta \rightarrow 0} \mathbb{E} |\gamma_t^\theta|^2 = 0. \quad (26)$$

Applying Itô's formula to $(Y_t^\theta)^2$, we get

$$\mathbb{E} |Y_t^\theta|^2 + \mathbb{E} \int_t^T |Z_s^\theta|^2 ds = \mathbb{E} |Y_T^\theta|^2 + 2\mathbb{E} \int_t^T |Y_s^\theta (F_s^y Y_s^\theta + F_s^z Z_s^\theta - \gamma_s^\theta)| ds.$$

By using the Young formula, for every $\varepsilon > 0$, we have

$$\begin{aligned}\mathbb{E} |Y_t^\theta|^2 + \mathbb{E} \int_t^T |Z_s^\theta|^2 ds &\leq \mathbb{E} |Y_T^\theta|^2 + \frac{1}{\varepsilon} \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + \varepsilon \mathbb{E} \int_t^T |(F_s^y Y_s^\theta + F_s^z Z_s^\theta - \gamma_s^\theta)|^2 ds \\ &\leq \mathbb{E} |Y_T^\theta|^2 + \frac{1}{\varepsilon} \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + C\varepsilon \mathbb{E} \int_t^T |F_s^y Y_s^\theta|^2 ds \\ &\quad + C\varepsilon \mathbb{E} \int_t^T |F_s^z Z_s^\theta|^2 ds + C\varepsilon \mathbb{E} \int_t^T |\gamma_s^\theta|^2 ds.\end{aligned}$$

Since F_t^y and F_t^z are bounded, then

$$\mathbb{E} |Y_t^\theta|^2 + \mathbb{E} \int_t^T |Z_s^\theta|^2 ds \leq \left(\frac{1}{\varepsilon} + C\varepsilon \right) \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + C\varepsilon \mathbb{E} \int_t^T |Z_s^\theta|^2 ds + \eta_t^\theta,$$

where

$$\eta_t^\theta = \mathbb{E} |Y_T^\theta|^2 + C\varepsilon \mathbb{E} \int_t^T |\gamma_s^\theta|^2 ds.$$

Choose $\varepsilon = \frac{1}{2C}$, then we have

$$\mathbb{E} |Y_t^\theta|^2 + \frac{1}{2} \mathbb{E} \int_t^T |Z_s^\theta|^2 ds \leq \left(2C + \frac{1}{2} \right) \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + \eta_t^\theta.$$

From the above inequality, we deduce two inequalities

$$\mathbb{E} |Y_t^\theta|^2 \leq \left(2C + \frac{1}{2} \right) \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + \eta_t^\theta, \quad (27)$$

$$\mathbb{E} \int_t^T |Z_s^\theta|^2 ds \leq (4C + 1) \mathbb{E} \int_t^T |Y_s^\theta|^2 ds + 2\eta_t^\theta. \quad (28)$$

On the other hand, we have

$$\begin{aligned} \mathbb{E} |Y_T^\theta|^2 &= \mathbb{E} \left| \tilde{y}_T - \frac{y_T^\theta - y_T^\mu}{\theta} \right|^2 \\ &= \mathbb{E} \left| \varphi_x(x_T^\mu) \tilde{x}_T - \frac{\varphi(x_T^\theta) - \varphi(x_T^\mu)}{\theta} \right|^2 \\ &\leq 2\mathbb{E} \int_0^1 |[\varphi_x(x_T^\mu) - \varphi_x(x_T^\mu + \lambda\theta(\tilde{x}_T + X_T^\theta))] \tilde{x}_T|^2 d\lambda \\ &\quad + 2\mathbb{E} \int_0^1 |\varphi_x(x_T^\mu + \lambda\theta(\tilde{x}_T + X_T^\theta)) X_T^\theta|^2 d\lambda. \end{aligned}$$

By using (20) and the continuity and boundeness of φ_x , we get

$$\lim_{\theta \rightarrow 0} \mathbb{E} |Y_T^\theta|^2 = 0. \quad (29)$$

From (26) and (29), we deduce that

$$\lim_{\theta \rightarrow 0} \eta_t^\theta = 0. \quad (30)$$

By using (27), (30), Gronwall's lemma and Burkholder-Davis-Gundy inequality, we obtain (21). Finally (22) is derived from (27), (30), (21). ■

Lemma 10 *Let μ be an optimal control minimizing the functional J over \mathcal{R} and $(x_t^\mu, y_t^\mu, z_t^\mu)$ the solution of (1) associated with μ . Then for any $q \in \mathcal{R}$, we have*

$$\begin{aligned} 0 &\leq \mathbb{E} [g_x(x_T^\mu) \tilde{x}_T] + \mathbb{E} [h_y(y_0^\mu) \tilde{y}_0] \\ &\quad + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt \\ &\quad + \mathbb{E} \int_0^T \left[\int_U l_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{x}_t + \int_U l_y(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{y}_t \right] dt \\ &\quad + \mathbb{E} \int_0^T \int_U l_z(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{z}_t dt. \end{aligned} \quad (31)$$

Proof. Let μ be an optimal relaxed control minimizing the cost J over \mathcal{R} , then

from (10) we have

$$\begin{aligned}
0 &\leq \mathbb{E} [g(x_T^\theta) - g(x_T^\mu)] + \mathbb{E} [h(y_0^\theta) - h(y_0^\mu)] \\
&+ \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) \mu_t^\theta(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt \\
&= \mathbb{E} [g(x_T^\theta) - g(x_T^\mu)] + \mathbb{E} [h(y_0^\theta) - h(y_0^\mu)] \\
&+ \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) \mu_t^\theta(da) - \int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) \mu_t(da) \right] dt \\
&+ \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) \mu_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt.
\end{aligned}$$

From the definition of μ^θ , we get

$$\begin{aligned}
0 &\leq \mathbb{E} [g(x_T^\theta) - g(x_T^\mu)] + \mathbb{E} [h(y_0^\theta) - h(y_0^\mu)] \\
&+ \theta \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) q_t(da) - \int_U l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) \mu_t(da) \right] dt \\
&+ \mathbb{E} \int_0^T \int_U [l(t, x_t^\theta, y_t^\theta, z_t^\theta, a) - l(t, x_t^\mu, y_t^\mu, z_t^\mu, a)] \mu_t(da) dt.
\end{aligned}$$

Then,

$$\begin{aligned}
0 &\leq \mathbb{E} \int_0^1 [g_x(x_T^\mu + \lambda \theta (\tilde{x}_T + X_T^\theta)) \tilde{x}_T] d\lambda \\
&+ \mathbb{E} \int_0^1 [h_y(y_0^\mu + \lambda \theta (\tilde{y}_0 + Y_0^\theta)) \tilde{y}_0] d\lambda \\
&+ \mathbb{E} \int_0^T \int_0^1 \int_U [l_x(\Lambda_t^\theta(a)) \tilde{x}_t + l_y(\Lambda_t^\theta(a)) \tilde{y}_t + l_z(\Lambda_t^\theta(a)) \tilde{z}_t] \mu_t(da) d\lambda dt \\
&+ \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt + \rho_t^\theta,
\end{aligned} \tag{32}$$

where ρ_t^θ is given by

$$\begin{aligned}
\rho_t^\theta &= \mathbb{E} \int_0^1 [g_x(x_T^\mu + \lambda \theta (\tilde{x}_T + X_T^\theta)) X_T^\theta] d\lambda \\
&+ \mathbb{E} \int_0^1 [h_y(y_0^\mu + \lambda \theta (\tilde{y}_0 + Y_0^\theta)) Y_0^\theta] d\lambda \\
&+ \mathbb{E} \int_0^T \int_0^1 \int_U [l_x(\Lambda_t^\theta(a)) (x_t^\theta - x_t^\mu) + l_y(\Lambda_t^\theta(a)) (y_t^\theta - y_t^\mu) + l_z(\Lambda_t^\theta(a)) (z_t^\theta - z_t^\mu)] q_t(da) d\lambda dt \\
&+ \mathbb{E} \int_0^T \int_0^1 \int_U [l_x(\Lambda_t^\theta(a)) (x_t^\theta - x_t^\mu) + l_y(\Lambda_t^\theta(a)) (y_t^\theta - y_t^\mu) + l_z(\Lambda_t^\theta(a)) (z_t^\theta - z_t^\mu)] \mu_t(da) d\lambda dt \\
&+ \mathbb{E} \int_0^T \int_0^1 \int_U [l_x(\Lambda_t^\theta(a)) X_T^\theta + l_y(\Lambda_t^\theta(a)) Y_t + l_z(\Lambda_t^\theta(a)) Z_t^\theta] \mu_t(da) d\lambda dt.
\end{aligned}$$

Since the derivatives g_x, h_y, l_x, l_y, l_z are continuous and bounded, then by using (11), (12), (13), (20), (21), (22) and the Cauchy-Schwartz inequality, we have

$$\lim_{\theta \rightarrow 0} \rho_t^\theta = 0.$$

By letting θ go to 0 in (32), the proof is completed. ■

3.2 Adjoint equations and variational inequality

Introduce the following system of stochastic differential equations, called adjoint equations

$$\begin{cases} dp_t^\mu = -\mathcal{H}_x(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) dt + P_t^\mu dW_t, \\ p_T^\mu = g_x(x_T^\mu) + \varphi_x(x_T^\mu) k_T^\mu. \end{cases} \quad (33)$$

$$\begin{cases} dk_t^\mu = \mathcal{H}_y(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) dt \\ \quad + \mathcal{H}_z(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) dW_t, \\ k_0^\mu = h_y(y_0^\mu). \end{cases} \quad (34)$$

Where the Hamiltonian \mathcal{H} is defined from $[0, T] \times \mathbb{R}^n \times \mathbb{R}^m \times \mathcal{M}_{m \times d}(\mathbb{R}) \times \mathbb{P}(U) \times \mathbb{R}^n \times \mathbb{R}^m$ into \mathbb{R} by

$$\begin{aligned} \mathcal{H}(t, x, y, z, q, p, k) = & \int_U l(t, x, y, z, a) q_t(da) + p \int_U b(t, x, a) q_t(da) \\ & + P \int_U \sigma(t, x, a) q_t(da) + k \int_U f(t, x, y, z, a) q_t(da). \end{aligned}$$

Since $p_T^\mu = g_x(x_T^\mu) + \varphi_x(x_T^\mu) k_T^\mu$ and $k_0^\mu = h_y(y_0^\mu)$, then (31) becomes

$$\begin{aligned} 0 \leq & \mathbb{E}[p_T^\mu \tilde{x}_T] + \mathbb{E}[k_0^\mu \tilde{y}_0] - \mathbb{E}[\varphi_x(x_T^\mu) k_T^\mu] + \mathbb{E} \int_0^T \int_U l_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{x}_t \mu_t(da) dt \\ & + \mathbb{E} \int_0^T \int_U [l_y(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{y}_t + l_z(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \tilde{z}_t] \mu_t(da) dt \\ & + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt. \end{aligned} \quad (35)$$

By applying Itô's formula to $(p_t^\mu \tilde{x}_t)$ and $(k_t^\mu \tilde{y}_t)$, we have

$$\begin{aligned} \mathbb{E}[p_T^\mu \tilde{x}_T] = & -\mathbb{E} \int_0^T \left[\int_U f_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) k_t^\mu + \int_U l_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] \tilde{x}_t dt \\ & + \mathbb{E} \int_0^T p_t^\mu \left[\int_U b(t, x_t^\mu, a) q_t(da) - \int_U b(t, x_t^\mu, a) \mu_t(da) \right] dt \\ & + \mathbb{E} \int_0^T P_t^\mu \left[\int_U \sigma(t, x_t^\mu, a) q_t(da) - \int_U \sigma(t, x_t^\mu, a) \mu_t(da) \right] dt. \end{aligned}$$

$$\begin{aligned}
\mathbb{E}[k_0^\mu \tilde{y}_0] &= \mathbb{E}[k_T^\mu \tilde{y}_T] - \mathbb{E} \left[\int_0^T \int_U l_y(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{y}_t + \int_U f_x(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{x}_t k_t^\mu \right] dt \\
&\quad + \mathbb{E} \int_0^T k_t^\mu \left[\int_U f(t, x_t^\mu, y_t^\mu, z_t^\mu, a) q_t(da) - \int_U f(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt \\
&\quad - \mathbb{E} \int_0^T \int_U l_z(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \tilde{z}_t dt.
\end{aligned}$$

Then for every $q \in \mathcal{R}$, (35) becomes

$$0 \leq \mathbb{E} \int_0^T [\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu) - \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu)] dt. \quad (36)$$

3.3 Necessary optimality conditions for relaxed controls

Starting from the variational inequality (36), we can now state necessary optimality conditions for the relaxed control problem $\{(5), (6), (7)\}$ in the global form.

Theorem 11 (*Necessary optimality conditions for relaxed controls*) *Let μ be an optimal relaxed control minimizing the functional J over \mathcal{R} and $(x_t^\mu, y_t^\mu, z_t^\mu)$ the solution of (5) associated with μ . Then, there exist two unique adapted processes p^μ and k^μ , which are respectively solutions of (33) and (34), such that*

$$\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \leq \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu); \quad \forall q_t \in \mathbb{P}(U); \quad ae, \quad as. \quad (37)$$

Proof. The result follows immediately from (36). ■

3.4 Sufficient optimality conditions for relaxed controls

In this subsection, we study when necessary optimality conditions (37) becomes sufficient. We recall assumptions (4) and the system of adjoints equations $\{(33), (34)\}$. For any $q \in \mathcal{R}$, we denote by (x^q, y^q, z^q) the solution of equation (5) controlled by q .

Theorem 12 (*Sufficient optimality conditions for relaxed controls*). *Assume that the functions g , h , φ and $(x, y, z) \mapsto \mathcal{H}(t, x, y, z, q, p, k)$ are convex. Then, μ is an optimal solution of the relaxed control problem $\{(5), (6), (7)\}$, if it satisfies (37).*

Proof. Let μ be an arbitrary element of \mathcal{R} (candidate to be optimal). For any $q \in \mathcal{R}$, we have

$$\begin{aligned}
J(q) - J(\mu) &= \mathbb{E}[g(x_T^q) - g(x_T^\mu)] + \mathbb{E}[h(y_0^q) - h(y_0^\mu)] \\
&\quad + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt.
\end{aligned}$$

Since g and h are convex, then

$$\begin{aligned} g(x_T^q) - g(x_T^\mu) &\geq g_x(x_T^\mu)(x_T^q - x_T^\mu), \\ h(y_0^q) - h(y_0^\mu) &\geq h_y(y_0^\mu)(y_0^q - y_0^\mu). \end{aligned}$$

Thus,

$$\begin{aligned} J(q) - J(\mu) &\geq \mathbb{E}[g_x(x_T^\mu)(x_T^q - x_T^\mu)] + \mathbb{E}[h_y(y_0^\mu)(y_0^q - y_0^\mu)] \\ &\quad + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt. \end{aligned}$$

We remark from (33) and (34), that

$$\begin{aligned} p_T^\mu &= g_x(x_T^\mu) + \varphi_x(x_T^\mu) k_T^\mu, \\ k_0^\mu &= h_y(y_0^\mu). \end{aligned}$$

Then, we have

$$\begin{aligned} J(q) - J(\mu) &\geq \mathbb{E}[p_T^\mu(x_T^q - x_T^\mu)] - \mathbb{E}[k_T^\mu \varphi_x(x_T^\mu)(x_T^q - x_T^\mu)] + \mathbb{E}[k_0^\mu(y_0^q - y_0^\mu)] \\ &\quad + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt. \end{aligned}$$

Since φ is convex, then

$$\varphi_x(x_T^\mu)(x_T^q - x_T^\mu) \leq \varphi(x_T^q) - \varphi(x_T^\mu) = y_T^q - y_T^\mu.$$

Thus,

$$\begin{aligned} J(q) - J(\mu) &\geq \mathbb{E}[p_T^\mu(x_T^q - x_T^\mu)] - \mathbb{E}[k_T^\mu(y_T^q - y_T^\mu)] + \mathbb{E}[k_0^\mu(y_0^q - y_0^\mu)] \\ &\quad + \mathbb{E} \int_0^T \left[\int_U l(t, x_t^q, y_t^q, z_t^q, a) q_t(da) - \int_U l(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt. \end{aligned}$$

By applying Itô's formula respectively to $p_t^\mu(x_t^q - x_t^\mu)$ and $k_t^\mu(y_t^q - y_t^\mu)$, we obtain

$$\begin{aligned} \mathbb{E}[p_T^\mu(x_T^q - x_T^\mu)] &= -\mathbb{E} \int_0^T \mathcal{H}_x(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu)(x_t^q - x_t^\mu) dt \\ &\quad + \mathbb{E} \int_0^T p_t^\mu \left[\int_U b(t, x_t^q, a) q_t(da) - \int_U b(t, x_t^\mu, a) \mu_t(da) \right] dt \\ &\quad + \mathbb{E} \int_0^T P_t^\mu \left[\int_U \sigma(t, x_t^q, a) q_t(da) - \int_U \sigma(t, x_t^\mu, a) \mu_t(da) \right] dt, \end{aligned}$$

$$\begin{aligned} \mathbb{E}[k_0^\mu(y_0^q - y_0^\mu)] &= \mathbb{E}[k_T^\mu(y_T^q - y_T^\mu)] - \mathbb{E} \int_0^T \mathcal{H}_y(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu)(y_t^q - y_t^\mu) dt \\ &\quad + \mathbb{E} \int_0^T k_t^\mu \left[\int_U f(t, x_t^q, y_t^q, z_t^q, a) q_t(da) - \int_U f(t, x_t^\mu, y_t^\mu, z_t^\mu, a) \mu_t(da) \right] dt \\ &\quad - \mathbb{E} \int_0^T \mathcal{H}_z(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu)(z_t^q - z_t^\mu) dt. \end{aligned}$$

Then,

$$\begin{aligned}
& J(q) - J(\mu) \\
& \geq \mathbb{E} \int_0^T [\mathcal{H}(t, x_t^q, y_t^q, z_t^q, q_t, p_t^\mu, k_t^\mu) - \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu)] dt \\
& - \mathbb{E} \int_0^T \mathcal{H}_x(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (x_t^q - x_t^\mu) dt \\
& - \mathbb{E} \int_0^T \mathcal{H}_y(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (y_t^q - y_t^\mu) dt \\
& - \mathbb{E} \int_0^T \mathcal{H}_z(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (z_t^q - z_t^\mu) dt.
\end{aligned} \tag{38}$$

Since \mathcal{H} is convex in (x, y, z) and linear in μ , then by using the Clarke generalized gradient of \mathcal{H} evaluated at (x_t, y_t, z_t, μ_t) and the necessary optimality conditions (37), it follows by [44, Lemmas 2.2 (4) and 2.3] that

$$\begin{aligned}
& \mathcal{H}(t, x_t^q, y_t^q, z_t^q, q_t, p_t^\mu, k_t^\mu) - \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \\
& \geq \mathcal{H}_x(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (x_t^q - x_t^\mu) + \mathcal{H}_y(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (y_t^q - y_t^\mu) \\
& + \mathcal{H}_z(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (z_t^q - z_t^\mu).
\end{aligned}$$

Or equivalently,

$$\begin{aligned}
0 & \leq \mathcal{H}(t, x_t^q, y_t^q, z_t^q, q_t, p_t^\mu, k_t^\mu) - \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \\
& - \mathcal{H}_x(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (x_t^q - x_t^\mu) \\
& - \mathcal{H}_y(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (y_t^q - y_t^\mu) \\
& - \mathcal{H}_z(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) (z_t^q - z_t^\mu).
\end{aligned}$$

Then from (38), we get

$$J(q) - J(\mu) \geq 0.$$

The theorem is proved. ■

4 Necessary and sufficient optimality conditions for strict controls

In this section, we study the strict control problem $\{(1), (2), (3)\}$ and from the results of section 3, we derive the optimality conditions for strict controls. For this end, consider the following subset of \mathcal{R}

$$\delta(\mathcal{U}) = \{q \in \mathcal{R} \mid q = \delta_v \ ; \ v \in \mathcal{U}\}. \tag{39}$$

The set $\delta(\mathcal{U})$ is the collection of all relaxed controls in the form of Dirac measure charging a strict control.

Denote by $\delta(U)$ the action set of all relaxed controls in $\delta(\mathcal{U})$.

If $q \in \delta(\mathcal{U})$, then $q = \delta_v$ with $v \in \mathcal{U}$. In this case we have for each t , $q_t \in \delta(U)$ and $q_t = \delta_{v_t}$.

To establish necessary optimality conditions for strict controls, we need the following lemma

Lemma 13 *The strict control u minimizes J over \mathcal{U} if and only if the relaxed control $\mu = \delta_u$ minimizes J over \mathcal{R} .*

Proof. Suppose that u minimizes the cost J over \mathcal{U} , then

$$J(u) = \inf_{v \in \mathcal{U}} J(v).$$

We know that the value function of the ordinary control problem coincides with that of the relaxed control problem (see Ma and Yong [29], page 16, line 7 and Theorem 36). Then, we have

$$\inf_{v \in \mathcal{U}} J(v) = \inf_{q \in \mathcal{R}} J(q). \quad (40)$$

Consequently, we get

$$J(u) = \inf_{q \in \mathcal{R}} J(q).$$

Since $\mu = \delta_u$, then

$$\begin{cases} (x^\mu, y^\mu, z^\mu) = (x^u, y^u, z^u), \\ J(\mu) = J(u), \end{cases} \quad (41)$$

This implies that

$$J(\mu) = \inf_{q \in \mathcal{R}} J(q).$$

Conversely, if $\mu = \delta_u$ minimize J over \mathcal{R} , then

$$J(\mu) = \inf_{q \in \mathcal{R}} J(q).$$

From (40), we get

$$J(\mu) = \inf_{v \in \mathcal{U}} J(v).$$

And from (41), we obtain

$$J(u) = \inf_{v \in \mathcal{U}} J(v).$$

The proof is completed. ■

The following lemma, who will be used to establish sufficient optimality conditions for strict controls, shows that we get the results of the above lemma if we replace \mathcal{R} by $\delta(\mathcal{U})$.

Lemma 14 *The strict control u minimizes J over \mathcal{U} if and only if the relaxed control $\mu = \delta_u$ minimizes J over $\delta(\mathcal{U})$.*

Proof. Let $\mu = \delta_u$ be an optimal relaxed control minimizing the cost J over $\delta(\mathcal{U})$, we have then

$$J(\mu) \leq J(q), \quad \forall q \in \delta(\mathcal{U}).$$

Since $q \in \delta(\mathcal{U})$, then there exists $v \in \mathcal{U}$ such that $q = \delta_v$.

It is easy to see that

$$\begin{cases} (x^\mu, y^\mu, z^\mu) = (x^u, y^u, z^u), \\ (x^q, y^q, z^q) = (x^v, y^v, z^v), \\ J(\mu) = J(u), \\ J(q) = J(v). \end{cases} \quad (42)$$

Then, we get

$$J(u) \leq J(v), \quad \forall v \in \mathcal{U}.$$

Conversely, let u be a strict control minimizing the cost J over \mathcal{U} . Then

$$J(u) \leq J(v), \quad \forall v \in \mathcal{U}.$$

Since the controls $u, v \in \mathcal{U}$, then there exist $\mu, q \in \delta(\mathcal{U})$ such that

$$\mu = \delta_u, \quad q = \delta_v.$$

This implies that relations (42) hold. Consequently, we get

$$J(\mu) \leq J(q), \quad \forall q \in \delta(\mathcal{U}).$$

The proof is completed. ■

4.1 Necessary optimality conditions for strict controls

Define the Hamiltonian H in the strict case from $[0, T] \times \mathbb{R}^n \times \mathbb{R}^m \times \mathcal{M}_{m \times d}(\mathbb{R}) \times U \times \mathbb{R}^n \times \mathbb{R}^m$ into \mathbb{R} by

$$H(t, x, y, z, v, p, k) = l(t, x, y, z, v) + pb(t, x, v) + P\sigma(t, x, v) + kf(t, x, y, z, v).$$

Theorem 15 *(Necessary optimality conditions for strict controls) Let u be an optimal control minimizing the functional J over \mathcal{U} and (x_t^u, y_t^u, z_t^u) the solution of (1) associated with u . Then there exist two unique adapted processes p^u and k^u , respectively solutions of*

$$\begin{cases} dp_t^u = -H_x(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u) dt + P_t^u dW_t, \\ p_T^u = g_x(x_T^u) + \varphi_x(x_T^u) k_T^u. \end{cases} \quad (43)$$

$$\begin{cases} dk_t^u = H_y(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u) dt \\ \quad + H_z(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u) dW_t, \\ k_0^u = h_y(y_0^u). \end{cases} \quad (44)$$

Such that

$$H(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u) \leq H(t, x_t^u, y_t^u, z_t^u, v_t, p_t^u, k_t^u); \forall v_t \in U; \text{ ae, as.} \quad (45)$$

Proof. Let u be an optimal solution of the strict control problem $\{(1), (2), (3)\}$. Then, there exist $\mu \in \delta(\mathcal{U})$ such that

$$\mu = \delta_u.$$

Since u minimizes the cost J over \mathcal{U} , then by lemma 13, μ minimizes J over \mathcal{R} . Hence, by the necessary optimality conditions for relaxed controls (Theorem 11), there exist two unique adapted processes p^μ and k^μ , respectively solutions of (33) and (34), such that

$$\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \leq \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu); \quad \forall q_t \in \mathbb{P}(U); \quad a.e., \quad a.s. \quad (46)$$

Since $\delta(U) \subset \mathbb{P}(U)$, then we get

$$\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \leq \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu); \quad \forall q_t \in \delta(U); \quad a.e., \quad a.s. \quad (47)$$

Since $q \in \delta(\mathcal{U})$, then there exist $v \in \mathcal{U}$ such that $q = \delta_v$.

We note that v is an arbitrary element of \mathcal{U} since q is arbitrary.

We can easily see that

$$\left\{ \begin{array}{l} (x^\mu, y^\mu, z^\mu) = (x^u, y^u, z^u), \\ (x^q, y^q, z^q) = (x^v, y^v, z^v), \\ (p^\mu, P^\mu, k^\mu) = (p^u, P^u, k^u), \\ \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) = H(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u), \\ \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu) = H(t, x_t^u, y_t^u, z_t^u, v_t, p_t^u, k_t^u), \end{array} \right. \quad (48)$$

where, the pair (p^u, P^u) and k^u are respectively the unique solutions of (43) and (44).

By using (46), (47) and (48), we can easily deduce (45). The proof is completed. ■

4.2 Sufficient optimality conditions for strict controls

We recall assumptions (4) on the system of adjoint equations $\{(43), (44)\}$.

We need the following corollary of theorem 12 whose shows that the sufficient optimality conditions for relaxed controls hold if we replace \mathcal{R} by $\delta(\mathcal{U})$ and $\mathbb{P}(U)$ by $\delta(U)$.

Corollary 16 *Assume that the functions g, h, φ and $(x, y, z) \mapsto \mathcal{H}(t, x, y, z, q, p, k)$ are convex. Let $\mu \in \delta(\mathcal{U})$ such that*

$$\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \leq \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu); \quad \forall q_t \in \delta(U); \quad a.e., \quad a.s.$$

Then

$$J(\mu) = \inf_{q \in \delta(\mathcal{U})} J(q).$$

Proof. We have the same proof that in theorem 12. ■

Theorem 17 (*Sufficient optimality conditions for strict controls*) Assume that the functions g , h , φ and $(x, y, z) \mapsto H(t, x, y, z, q, p, k)$ are convex. Then, u is an optimal solution of the control problem $\{(1), (2), (3)\}$, if it satisfies (45).

Proof. Let u be a strict control (candidate to be optimal) such that necessary optimality conditions for strict controls hold. That is

$$H(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u) \leq H(t, x_t^u, y_t^u, z_t^u, v_t, p_t^u, k_t^u); \forall v_t \in U; \text{ a.e., a.s.} \quad (49)$$

The controls u, v are elements of \mathcal{U} , then there exist $\mu, q \in \delta(\mathcal{U})$ such that

$$\mu = \delta_u, \quad q = \delta_v.$$

This implies that

$$\begin{cases} (x^\mu, y^\mu, z^\mu) = (x^u, y^u, z^u), \\ \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) = H(t, x_t^u, y_t^u, z_t^u, u_t, p_t^u, k_t^u), \\ \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu) = H(t, x_t^u, y_t^u, z_t^u, v_t, p_t^u, k_t^u). \end{cases}$$

By the above equalities and (49), we deduce that

$$\mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, \mu_t, p_t^\mu, k_t^\mu) \leq \mathcal{H}(t, x_t^\mu, y_t^\mu, z_t^\mu, q_t, p_t^\mu, k_t^\mu); \forall q_t \in \delta(U); \text{ a.e., a.s.}$$

Since H is convex in (x, y, z) , it is easy to see that \mathcal{H} is convex in (x, y, z) , and since g , h and φ are convex, then from the corollary 16, μ minimizes the cost J over $\delta(\mathcal{U})$.

Finally by lemma 14, we deduce that u minimizes the cost J over \mathcal{U} . The proof is completed. ■

Remark 18 *The sufficient optimality conditions for strict controls are proved without assuming neither the convexity of U nor that of H in v .*

5 The case of fully coupled forward-backward systems

In the case where the system is governed by a fully coupled FBSDE, the proofs of necessary as well as sufficient conditions of optimality, for relaxed and strict controls, are practically the same as in the preceding sections. But, it is necessary to put additional assumptions on the coefficients, to ensure existence and uniqueness of the solution of the fully coupled FBSDE (for that, we propose to refer to the paper by Peng and Wu [38]). These additional hypothesis are not in contradiction with our assumptions (4).

References

- [1] F. Antonelli, *Backward-forward stochastic differential equations*. Annals of Applied Probability, 1993, 3, pp. 777–793.
- [2] F. Antonelli and J. Ma, *Weak solution of Forward-Backward SDE's*. Stochastic Analysis and Application, 2003, 21, n.3, pp. 493–514
- [3] S. Bahlali, B. Mezerdi and B. Djehiche, *Approximation and optimality necessary conditions in relaxed stochastic control problems*, Journal of Applied Mathematics and Stochastic Analysis, Volume 2006, pp 1-23.
- [4] S. Bahlali and B. Labed, *Necessary and sufficient conditions of optimality for optimal control problem with initial and terminal costs*, Rand. Operat. and Stoch. Equ, 2006, Vol 14, No3, pp 291-301.
- [5] S. Bahlali, B. Djehiche and B. Mezerdi, *The relaxed maximum principle in singular control of diffusions*, SIAM J. Control and Optim, 2007, Vol 46, Issue 2, pp 427-444.
- [6] S. Bahlali, *Necessary and sufficient conditions of optimality for relaxed and strict control problems*, SIAM J. Control and Optim, To appear.
- [7] S. Bahlali, *Necessary and sufficient condition of optimality for optimal control problem of forward and backward systems*, Theory of Probability and Its Applications (TVP), In revision.
- [8] A. Bensoussan, *Non linear filtering and stochastic control*. Proc. Cortona 1981, Lect. notes in Math. 972, Springer Verlag.
- [9] F. Delarue, *On the existence and uniqueness of solutions to FBSDEs in a non-degenerate case*. Stochastic Process. Appl., 2002, 99, pp. 209–286.
- [10] N. Dokuchaev and X. Y. Zhou, *Stochastic controls with terminal contingent conditions*, Journal Of Mathematical Analysis And Applications, 1999, 238, pp 143-165.
- [11] J. Douglas, J. Ma and P. Protter, *Numerical methods for forward-backward stochastic differential equations*, 1996, Ann. Appl. Probab., 6(3), pp 940-968.
- [12] N. El Karoui, N. Huu Nguyen and M. Jeanblanc Piqué, *Compactification methods in the control of degenerate diffusions*. Stochastics, Vol. 20, 1987, pp 169-219.
- [13] N. El Karoui and L. Mazliak, *Backward stochastic differential equations*, 1997, Addison Wesley, Longman.
- [14] N. El-Karoui, S. Peng, and M. C. Quenez, *Backward stochastic differential equations in finance*, 1997, Math. finance 7.

- [15] N. El-Karoui, S. Peng and M.C. Quenez, *A dynamic maximum principle for the optimization of recursive utilities under constraints*, Annals of Applied Probability, 11(2001), pp 664-693.
- [16] R.J. Elliott and M. Kohlmann, *The variational principle and stochastic optimal control*. Stochastics 3, 1980, pp 229-241.
- [17] W.H. Fleming, *Generalized solutions in optimal stochastic control*, Differential games and control theory 2, (Kingston conference 1976), Lect. Notes in Pure and Appl. Math.30, 1978.
- [18] N.F. Framstad, B. Oksendal and A. Sulem, *A sufficient stochastic maximum principle for optimal control of jump diffusions and applications to finance*, J. Optim. Theory and applications. 121, 2004, pp 77-98.
- [19] M. Fuhrman and G. Tessitore, *Existence of optimal stochastic controls and global solutions of forward-backward stochastic differential equations*, SIAM J. Control and Optim, 2004, Vol 43, N° 3, pp 813-830.
- [20] U.G. Haussmann, *General necessary conditions for optimal control of stochastic systems*, Math. Programming Studies 6, 1976, pp 30-48.
- [21] U.G. Haussmann, *A Stochastic maximum principle for optimal control of diffusions*, Pitman Research Notes in Math, 1986, Series 151.
- [22] Y. Hu, *On the solution of Forward-backward SDEs with monotone and continuous coefficients*, Nonlinear Anal., 1999, 42, pp 1-12.
- [23] Y. Hu and S. Peng, *Solution of forward-backward stochastic differential equations*. Probab.Theory Rel. Fields, 1995,103, pp. 273–283.
- [24] Y. Hu and J. Yong, *Forward-backward stochastic differential equations with nonsmooth coefficients*. Stochastic Process. Appl., 2000, 87, pp. 93–106.
- [25] J. Jacod and J. Mémin, *Sur un type de convergence intermédiaire entre la convergence en loi et la convergence en probabilité*. Sem. Proba.XV, Lect. Notes in Math. 851, Springer Verlag, 1980.
- [26] S.Ji and X. Y. Zhou, *A maximum principle for stochastic optimal control with terminal state constraints, and its applications*. Commun. Inf. Syst, 2006, 6(4), pp 321-338.
- [27] H.J. Kushner, *Necessary conditions for continuous parameter stochastic optimization problems*, SIAM J. Control Optim, Vol. 10, 1973, pp 550-565.
- [28] J. Ma, P. Protter and J. Yong, *Solving forward-backward stochastic differential equations explicitly - a four step scheme*, Probab. Theory Rel. Fields, 1994, 98, pp 339–359.

- [29] J. Ma and J. Yong, *Solvability of forward-backward SDEs and the nodal set of Hamilton-Jacobi-Bellman equations*. A Chinese summary appears in Chinese Ann. Math. Ser. A 16 (1995), no. 4, 532. Chinese Ann. Math. Ser. B 16, 1995, no. 3, pp 279–298.
- [30] J. Ma and J. Yong, *Forward-backward stochastic differential equations and their applications*, In Lecture Notes Math., 1999, volume 1702. Springer, Berlin.
- [31] J. Ma and J. Zhang, *Representation theorems for backward stochastic differential equations*, Ann. Appl. Probab., 2002, 12(4), pp 1390-1418, 2002.
- [32] B. Mezerdi and S. Bahlali, *Approximation in optimal control of diffusion processes*, Rand. Operat. and Stoch. Equ, 2000, Vol.8, No 4, pp 365-372.
- [33] B. Mezerdi and S. Bahlali, *Necessary conditions for optimality in relaxed stochastic control problems*, Stochastics And Stoch. Reports, 2002, Vol 73 (3-4), pp 201-218.
- [34] E. Pardoux and S. Peng, *Adapted solutions of backward stochastic differential equations*, Sys. Control Letters, 1990, Vol. 14, pp 55-61.
- [35] E. Pardoux and S. Tang, *Forward-Backward stochastic differential equations and quasilinear parabolic PDEs*. Probab. Theory Rel. Fields, 1999, 114, pp 123–150.
- [36] S. Peng, *A general stochastic maximum principle for optimal control problems*. SIAM J. Control and Optim. 1990, 28, N° 4, pp 966-979.
- [37] S. Peng, *Backward stochastic differential equations and application to optimal control*, Appl. Math. Optim. 1993, 27, pp 125-144.
- [38] S. Peng and Z. Wu, *Fully coupled forward-backward stochastic differential equations and applications to optimal control*. SIAM J. Control Optim., 1999, 37, no. 3, pp. 825–843.
- [39] J.T. Shi and Z. Wu, *The maximum principle for fully coupled forward-backward stochastic control system*, Acta Automatica Sinica, Vol 32, No 2, 2006, pp 161-169.
- [40] Z. Wu, *Maximum Principle for Optimal Control Problem of Fully Coupled Forward-Backward Stochastic Systems*, Systems Sci. Math. Sci, 1998, 11, No.3, pp 249-259.
- [41] W. Xu, *Stochastic maximum principle for optimal control problem of forward and backward system*, J. Austral. Math. Soc. Ser. B 37, 1995, pp 172-185.
- [42] J. Yong, *Finding adapted solutions of forward-backward stochastic differential equations - method of continuation*, Probablity Theory Related Fields, 1997, 107, pp. 537–572.

- [43] J. Yong and X.Y. Zhou, *Stochastic controls : Hamilton systems and HJB equations*, vol 43, Springer, New York, 1999.
- [44] X.Y. Zhou, *Sufficient conditions of optimality for stochastic systems with controllable diffusions*. IEEE Trans. on Automatic Control, 1996, 41, pp 1176-1179.