# Some properties of the k-dimensional Lyness' map

Anna Cima<sup>(1)</sup>, Armengol Gasull<sup>(1)</sup> and Víctor Mañosa <sup>(2)</sup>

<sup>(1)</sup> Dept. de Matemàtiques, Facultat de Ciències, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain cima@mat.uab.cat, gasull@mat.uab.cat

<sup>(2)</sup> Dept. de Matemàtica Aplicada III (MA3), Control, Dynamics and Applications Group (CoDALab) Universitat Politècnica de Catalunya (UPC) Colom 1, 08222 Terrassa, Spain victor.manosa@upc.edu

September 4, 2021

#### Abstract

This paper is devoted to study some properties of the k-dimensional Lyness' map  $F(x_1, \ldots, x_k) = (x_2, \ldots, x_k, (a + \sum_{i=2}^k x_i)/x_1)$ . Our main result presentes a rational vector field that gives a Lie symmetry for F. This vector field is used, for  $k \leq 5$ , to give information about the nature of the invariant sets under F. When k is odd, we also present a new (as far as we know) first integral for  $F \circ F$  which allows to deduce in a very simple way several properties of the dynamical system generated by F. In particular for this case we prove that, except on a given codimension one algebraic set, none of the positive initial conditions can be a periodic point of odd period.

2000 Mathematics Subject Classification 39A20, 37E35.

PACS numbers 02.30.Ik, 05.45.-a.

*Keywords:* Lyness' difference equations, Lie symmetry, first integral, integrable and non– integrable discrete dynamical system.

## 1 Introduction and main results

The second and more specially the third order Lyness' difference equations

$$y_{n+2} = \frac{a+y_{n+1}}{y_n}$$
 and  $y_{n+3} = \frac{a+y_{n+1}+y_{n+2}}{y_n}$ , with  $a \ge 0$ ,

have been considered as emblematic examples of integrable discrete systems, see for instance [15, 20, 22, 23]. The dynamics of the above equations, or their associated maps, has been

the objective of recent intensive investigation. Nowadays the behaviour of the orbits is well known when positive initial conditions are considered, see [2, 5, 7, 10, 24]. However few results have been obtained for negative initial conditions, see [6, 10].

On the other hand, the study of the 2 and 3-dimensional Lyness' maps is the starting point for the study of other birational integrable maps, see for instance [3, 8, 10, 13, 16, 17, 18, 23], and [9] for a general paper on this topic.

For  $k \ge 4$ , very few results, apart of the ones obtained recently by Bastien and Rogalsky [4], are known for the k-th order Lyness' equation

$$y_{n+k} = \frac{a + \sum_{i=1}^{k-1} y_{n+i}}{y_n}.$$
(1)

The main difference between the k = 2, 3 and the  $k \ge 4$  scenarios is that the first cases are *integrable* in the sense that the associated maps have 1 and 2 functionally independent first integrals, respectively (in this paper we say that a map F is integrable if it has k - 1functionally independent first integrals). It seems that this property is not shared for the Lyness' equations when  $k \ge 4$ .

Indeed, consider the k-dimensional Lyness' map associated to the difference equation (1),

$$F(x_1, \dots, x_k) = \left(x_2, \dots, x_k, \frac{a + \sum_{i=2}^k x_i}{x_1}\right), \text{ with } a \ge 0,$$
 (2)

which is a diffeomorphism from  $Q^+ := \{ \mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k : x_1 > 0, x_2 > 0, \dots, x_k > 0 \}$ into itself. It is well known that it has the following couple of functionally independent first integrals

$$V_1(\mathbf{x}) = \left(a + \sum_{i=1}^k x_i\right) \left(\prod_{i=1}^k (x_i + 1)\right) / (x_1 \cdots x_k) \tag{3}$$

and

$$V_2(\mathbf{x}) = \left(a + \sum_{i=1}^k x_i + x_1 x_k\right) \left(\prod_{i=1}^{k-1} (1 + x_i + x_{i+1})\right) / (x_1 \cdots x_k).$$
(4)

A third functionally independent first integral for  $k \ge 5$  has recently been given in [11]. Moreover in that paper it is conjectured that for any k, the Lyness' map has up to  $E\left(\frac{k+1}{2}\right)$  functionally independent first integrals, where  $E(\cdot)$  denotes the integer part function. The conjecture seems to be true for  $k \le 6$ , see again [11].

The integrable structure for k = 2, 3 implies that the dynamics of the maps studied in the above references is in fact one-dimensional. In any case, although for  $k \ge 4$  the above assertion seems not to be true, the existence of several first integrals reduces the dimension of the space where the dynamics takes place. If the above conjecture would be true, then  $k - E\left(\frac{k+1}{2}\right)$ , would be, generically, the dimension of the invariant manifold given by the level sets of the first integrals.

One geometrical object that has played a key role to understand the dynamics of a large class of 2 and 3-dimensional maps is the *Lie symmetry* of the map, [7, 8]. Recall, that a vector field  $\mathbf{X}$  is said to be a Lie symmetry of a map G if it satisfies the condition

$$\mathbf{X}(G(\mathbf{x})) = (DG(\mathbf{x})) \, \mathbf{X}(\mathbf{x}). \tag{5}$$

The vector field  $\mathbf{X}$  is related with the dynamics of G in the following sense: G maps any orbit of the differential system determined by the vector field, to another orbit of this system, see [8]. In the integrable case, where the dynamics are in fact one dimensional, the existence of a Lie symmetry fully characterizes the dynamics. In [8, Thm. 1] it is proved that if  $G: \mathcal{U} \to \mathcal{U}$  is a diffeomorphism having a Lie symmetry  $\mathbf{X}$ , and such G preserves  $\gamma$ , a solution of the differential equation  $\dot{x} = \mathbf{X}(x)$ , then the dynamics of  $G|_{\gamma}$  is either conjugated to a rotation, conjugated to a translation of the line, or constant, according whether  $\gamma$  is homeomorphic to  $\mathbb{S}^1$ ,  $\mathbb{R}$ , or a point, respectively. Other properties of the Lie symmetries of discrete systems are studied in [14].

One of the main results of this paper, is the following theorem where the expression of a Lie symmetry for the k-dimensional Lyness' map is given.

**Theorem 1.** For  $k \geq 3$ , the vector field  $\mathbf{X}_k = \sum_{i=1}^k X_i \frac{\partial}{\partial x_i}$ , is a Lie symmetry for the k-dimensional Lyness' map (2), where

$$X_{1}(\mathbf{x}) = \frac{(x_{1}+1)\left[\prod_{i=2}^{k-1}(1+x_{i}+x_{i+1})\right](a+\sum_{i=1}^{k-1}x_{i}-x_{2}x_{k})}{\prod_{i=2}^{k}x_{i}},$$

$$(6)$$

$$X_{m}(\mathbf{x}) = \frac{(x_{m}+1)\left[\prod_{i=1,i\neq m-1,m}^{k-1}(1+x_{i}+x_{i+1})\right](a+\sum_{i=1}^{k}x_{i}+x_{1}x_{k})(x_{m-1}-x_{m+1})}{k},$$

 $\prod_{i=1,i\neq m}^k x_i$ 

for all  $2 \leq m \leq k-1$ , and

$$X_k(\mathbf{x}) = -\frac{(x_k+1)\left[\prod_{i=1}^{k-2}(1+x_i+x_{i+1})\right](a+\sum_{i=2}^k x_i - x_1 x_{k-1})}{\prod_{i=1}^{k-1} x_i}.$$
(8)

(7)

Once we have the candidate  $\mathbf{X}_k$  to be a Lie symmetry of the Lyness' map F the proof of Theorem 1 only will consist in checking that (5) holds. We give here some hints of how we have found the above  $\mathbf{X}_k$ . Observe that if there exists a vector field  $\mathbf{X}_k$ , satisfying equation (5) for the k-dimensional Lyness' map (2), then

$$\begin{pmatrix} X_1(F) \\ X_2(F) \\ \vdots \\ X_k(F) \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ \vdots & & & & \\ -\frac{a+\sum_{i=2}^k x_i}{x_1^2} & \frac{1}{x_1} & \cdots & \frac{1}{x_1} \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_k \end{pmatrix}.$$

Hence it is necessary that

$$X_{i+1} = X_i(F), \text{ for } i = 1, \dots, k-1,$$
 (9)

and the "compatibility condition":

$$X_k(F) = -\left(\frac{a + \sum_{i=2}^k x_i}{x_1^2}\right) X_1 + \frac{1}{x_1} \left[\sum_{i=2}^k X_i\right].$$

Thus, the construction of a Lie symmetry is straightforward once the right expression of  $X_1$ , as a seed of equations (9), is obtained. In [5, 7] and [8] the expressions of  $\mathbf{X}_2$  and  $\mathbf{X}_3$  are given. The idea of these papers is that these vector fields have to be multiples of  $\nabla V_1$  and  $\nabla V_1 \times \nabla V_2$ , respectively. These constructions are used to force F and  $\mathbf{X}_k$  to share the same set of first integrals. When  $k \geq 4$  we can not use anymore this idea because there are no enough first integrals. Nevertheless we observe that the first components of  $\mathbf{X}_2$  and  $\mathbf{X}_3$ , are

$$\frac{(x_1+1)(a+x_1-x_2^2)}{x_2} \quad \text{and} \quad \frac{(x_1+1)(1+x_2+x_3)(a+x_1+x_2-x_2x_3)}{x_2x_3}$$

respectively. Thus it seems natural to try with  $X_1$  as the expression given in (6) and, indeed, it works!. From this starting point, the proof for a given small k is only a matter of computations, while the proof for a general k is long and tedious, but straightforward. It is done in Section 2. We suggest to skip this section in a first reading of the paper.

Another result that helps for understanding the dynamics generated by F, when k is odd, is given in next proposition. In this result, the key point is the existence of a new (as far as we know) first integral for  $F^2 = F \circ F$  for any odd  $k \ge 3$ . As we will see, our proof of the existence of this function is inspired in the paper [22], where this first integral is given for k = 3.

**Theorem 2.** Set  $k = 2\ell + 1$ ,  $\mathbf{x} = (x_1, \dots, x_{2\ell+1})$  and consider F from  $\mathcal{Q}^+$  into itself. Then (a) The function

$$W(\mathbf{x}) = \frac{\prod_{j=0}^{\ell} (x_{2j+1} + 1)}{\prod_{j=1}^{\ell} x_{2j}},$$
(10)

is a first integral of  $F^2$ .

(b) For any  $\ell \geq 2$ , the function  $V_3 := W + W(F)$ , which is,

$$V_{3}(\mathbf{x}) = \frac{\prod_{j=0}^{\ell} x_{2j+1}(x_{2j+1}+1) + (a + \sum_{j=1}^{2\ell+1} x_j) \prod_{j=1}^{\ell} x_{2j}(x_{2j}+1)}{\prod_{i=1}^{2\ell+1} x_i}, \quad (11)$$

is a first integral of F which is functionally independent with the first integrals  $V_1$  and  $V_2$  given in (3) and (4), respectively.

- (c) The function  $W \cdot W(F)$  coincides with the first integral of F,  $V_1$  given in (3). In other words,  $V_1 = W \cdot W(F)$ .
- (d) The algebraic set  $\mathcal{G} := \{ \mathbf{x} \in \mathcal{Q}^+ : W(\mathbf{x}) W(F(\mathbf{x})) = 0 \}$  is invariant by F.
- (e) If the map F has some periodic point of odd period then it has to be contained in  $\mathcal{G}$ .
- (f) Setting  $\mathcal{G}^{\pm} := \{\mathbf{x} \in \mathcal{Q}^{+} : \pm(W(\mathbf{x}) W(F(\mathbf{x}))) > 0\}$ , the map F sends  $\mathcal{G}^{+}$  into  $\mathcal{G}^{-}$ and viceversa, and both sets are invariant by  $F^{2}$ . Furthermore, the dynamics of  $F^{2}$ on each of these sets are conjugated, being the map F itself the conjugation.
- (g) The measure

$$m_1(B) := \int_B \frac{\pm 1}{\Pi(\mathbf{x})(W(\mathbf{x}) - W(F(\mathbf{x})))} \, d\mathbf{x},$$

where  $\Pi(\mathbf{x}) = \prod_{i=1}^{k} x_i$ , is an invariant measure for  $F^2$ , i.e.  $m_1(F^2(B)) = m_1(B)$ , where B is any measurable set in  $\mathcal{G}^{\pm}$ .

(h) The measure

$$m_2(B) := \int_B \frac{1}{\Pi(\mathbf{x})} \, d\mathbf{x},$$

is an invariant measure for  $F^2$ , i.e.  $m_2(F^2(B)) = m_2(B)$ , where B is any measurable set in  $Q^+$ .

We remark that, when k is odd, the first integral  $V_3$  given above coincides with the one given recently in [11]. Observe also that the invariant algebraic surface  $\mathcal{G}$  was already found in [7], but only for k = 3. Also, as we will see in Subsection 3.1, the function W is useful to make an explicit simple order reduction when we study the dynamics of F for k odd.

Although, by using both theorems we have not been able to present a complete study of the higher dimensional Lyness' map, in next results we give some information about the invariant sets in the phase space when k = 4, 5. We prove:

**Proposition 3.** The vector field  $\mathbf{X}_4$  given by equations (6)–(8) for k = 4, is a Lie symmetry for the 4-dimensional Lyness' map. Moreover,  $\mathbf{X}_4(V_i) = 0$ , for i = 1, 2, and then the sets  $I_{h,k} := \{V_1 = h\} \cap \{V_2 = k\} \cap \mathcal{Q}^+$  are invariant by F and by the flow of  $\mathbf{X}_4$ .

Furthermore, if we assume that both first integrals intersect transversally on  $C_{h,k}$ , a connected component of  $I_{h,k}$ , then  $C_{h,k}$  is diffeomorphic to a torus.

**Proposition 4.** The vector field  $\mathbf{X}_5$  given by equations (6)–(8) for k = 5 is a Lie symmetry for the 5-dimensional Lyness' map. Moreover,  $\mathbf{X}_5(V_i) = 0$ , for i = 1, 2, 3, where

$$V_{3}(\mathbf{x}) = \frac{1}{x_{1}x_{2}x_{3}x_{4}x_{5}} \left( \begin{array}{c} x_{1}x_{3}x_{5} (1+x_{1}) (1+x_{3}) (1+x_{5}) \\ +x_{2}x_{4} (1+x_{2}) (1+x_{4}) (a+x_{1}+x_{2}+x_{3}+x_{4}+x_{5}) \end{array} \right),$$

and the sets  $I_{h,k,l} := \{V_1 = h\} \cap \{V_2 = k\} \cap \{V_3 = \ell\} \cap \mathcal{Q}^+$ , which generically have at least two connected components, are invariant by F and by the flow of  $\mathbf{X}_5$ .

Furthermore, if we assume that the three first integrals intersect transversally on  $C_{h,k,\ell}$ , a connected component of  $I_{h,k,\ell}$ , then  $C_{h,k,\ell}$  is diffeomorphic to a (two-dimensional) torus.

It is important to notice that in the above results we do not assert that most of the connected components  $C_{h,k}$  and  $C_{h,k,\ell}$  are tori, since we did not succeed to prove that over them the intersection of the energy levels at the whole sets  $C_{h,k}$  and  $C_{h,k,\ell}$  are transversal. Hence it remains open to decide whether they are two-dimensional differentiable manifolds or not, and in the case that they are not differentiable manifolds to decide which are their topology. In any case, our result reduces the problem to a computational question.



Figure 1. Projections into  $\mathbb{R}^3$  of the flow associated to the Lie symmetry  $\mathbf{X}_4$ , and the orbit of the Lyness' map, for k = 4 and a = 4, both with initial condition (1,2,3,4).

Our numerical simulations seem to indicate that for k = 4, all the generic level curves  $I_{h,k}$ , are connected. On the other hand, for k = 5, generically the sets  $I_{h,k,l}$  seem to have exactly two connected components. In Figures 1 and 2 we give a projection in  $\mathbb{R}^3$  of these surfaces. Indeed, in Figure 1 we represent, both an orbit of F and an orbit of  $\mathbf{X}_4$  starting with the same initial condition, and in Figure 2 an orbit of F for k = 5. Notice that in both cases the behavior of the orbits seems to indicate that two, (respectively three) is the maximum number of independent first integrals for F when k = 4 (resp. k = 5), as it is suggested in [11].



Figure 2. Projection into  $\mathbb{R}^3$  of the first 5000 iterates of the Lyness' map for k = 5 and a = 1, staring at (1,2,3,4,5). Odd and even iterates are in different connected components.

A remarkable fact that Figure 1 shows is that, although for k = 4 the manifold  $I_{h,k}$  is invariant for both the map F and the flow of  $\mathbf{X}_4$ , the map F seems to send an orbit of  $\mathbf{X}_4$ to a different orbit of the vector field. Further numerical experiments seems to confirm this fact. Under this situation, we can not use the techniques developed in [8]. This fact makes more difficult the knowledge of the behavior of F restricted to each  $I_{h,k}$  and is one of the important differences between the cases k = 2, 3 and k = 4.

## 2 Proof of Theorem 1

As we have already explained in the introduction, the existence of a vector field  $\mathbf{X}_k = \sum_{i=i}^k X_i \frac{\partial}{\partial x_i}$  satisfying equation (5) for the k-dimensional Lyness' map (2), is equivalent to the set of equations

$$X_{i+1} = X_i(F), \text{ for } i = 1, \dots, k-1,$$
 (12)

together with the compatibility condition:

$$X_k(F) = -\left(\frac{a + \sum_{i=2}^k x_i}{x_1^2}\right) X_1 + \frac{1}{x_1} \left[\sum_{i=2}^k X_i\right].$$
 (13)

The proof will consist in checking that the choice of  $\mathbf{X}_k$  given in the statement satisfies equations (12) and (13). The result is straightforward for k = 3, 4, 5 and we omit the details. So, from now on, we assume that  $k \ge 6$ .

We proceed in two steps:

First step: We will show that from the expression (6) of  $X_1$  as a seed of equations (12) we obtain the expressions of  $X_m$  for m = 2, ..., k - 1 and  $X_k$  given by equations (7) and (8), respectively.

Second step: We will prove that the compatibility condition (13) is satisfied.

**First step:** We start with some preliminary observations:

Observation 1. Set  $K_i := x_i + 1$  for i = 1, ..., k - 1. Then  $K_i(F) = x_{i+1} + 1$ . Observation 2. If we set  $L_i := 1 + x_i + x_{i+1}$ , then for all  $1 \leq i \leq k - 1$ ,  $L_i(F) = 1$  $1 + x_{i+1} + x_{i+2} = L_{i+1}$ , and

$$L_{k-1}(F) = 1 + x_k + x_{k+1} = 1 + x_k + \frac{a + \sum_{i=2}^k x_i}{x_1} = \frac{a + \sum_{i=1}^k x_i + x_1 x_k}{x_1}.$$

For this reason

(a) If  $M_1 := \prod_{i=1}^{k-1} (1 + x_i + x_{i+1})$ , then  $M_1(F) = \frac{(a + \sum_{i=1}^k x_i + x_1 x_k) \left( \prod_{i=2}^{k-1} (1 + x_i + x_{i+1}) \right)}{x_1}.$ 

(b) Setting  $M_m := \prod_{i=1, i \neq m-1, m}^{k-1} (1 + x_i + x_{i+1})$  for  $2 \le m \le k-2$ , we obtain

$$M_m(F) = \left(\prod_{i=2, i \neq m, m+1}^{k-1} (1+x_i+x_{i+1})\right) (a + \sum_{i=1}^k x_i + x_1 x_k) / x_1.$$

(c) If 
$$M_{k-1} := \prod_{i=1}^{k-3} (1 + x_i + x_{i+1})$$
, then  $M_{k-1}(F) = \prod_{i=2}^{k-2} (1 + x_i + x_{i+1})$ .

Observation 3. Set  $N = a + \sum_{i=1}^{k-1} x_i - x_2 x_k$ , then

$$N(F) = a + \sum_{i=2}^{k} x_i - x_3 x_{k+1} = a + \sum_{i=2}^{k} x_i - x_3 \frac{a + \sum_{i=2}^{k} x_i}{x_1}$$
$$= \frac{a + \sum_{i=2}^{k} x_i}{x_1} (x_1 - x_3) = x_{k+1} (x_1 - x_3).$$

Observation 4. Set  $R = a + \sum_{i=1}^{k} x_i + x_1 x_k$ , then

$$R(F) = a + \sum_{i=2}^{k} x_i + x_{k+1} + x_2 x_{k+1} = \frac{a + \sum_{i=2}^{k} x_i}{x_1} (1 + x_1 + x_2) = x_{k+1} (1 + x_1 + x_2).$$

Observation 5. (a) For all  $2 \le i \le k - 2$  set  $S_i = x_{i-1} - x_{i+1}$ , then  $S_i(F) = x_i - x_{i+2}$ .

(b) Set  $S_{k-1} = x_{k-2} - x_k$ , then

$$S_{k-1}(F) = x_{k-1} - x_{k+1} = x_{k-1} - \frac{a + \sum_{i=2}^{k} x_i}{x_1} = -\frac{a + \sum_{i=2}^{k} x_i - x_1 x_{k-1}}{x_1}.$$

If we now consider the seed  $X_1$  given by equation (6), using Observations 1, 2a, and 3 we obtain that

$$X_2 = \frac{(x_2+1)\left[\prod_{i=3}^{k-1}(1+x_i+x_{i+1})\right](a+\sum_{i=1}^k x_i+x_1x_k)(x_1-x_3)}{\prod_{i=1,i\neq 2}^k x_i}.$$

Now applying systematically Observations 1, 2b, 4 and 5a, we obtain that for  $2 \le m \le k-1$  the component  $X_m = X_{m-1}(F)$  is given by equation (7).

Observe that in particular

$$X_{k-1} = X_{k-2}(F) = \frac{(x_{k-1}+1)\left[\prod_{i=1}^{k-3}(1+x_i+x_{i+1})\right](a+\sum_{i=1}^{k}x_i+x_1x_k)(x_{k-2}-x_k)}{\prod_{i=1}^{k-2}x_i},$$

hence the term  $L_{k-1} = 1 + x_{k-1} + x_k$  does not appear and so, in order to compute  $X_k$  we need to use Observations 1, 2c, 4 and 5b, obtaining the expression of  $X_k$  given by (8).

Second step (compatibility condition (13)). A simple computations shows that

$$X_k(F) = \frac{\mathbf{A}}{x_1\left(\prod_{i=1}^k x_i\right)},$$

where

$$\mathbf{A} = -\left(a + \sum_{i=1}^{k} x_i\right) \left[\prod_{i=2}^{k-1} L_i\right] \left(a + \sum_{i=2}^{k} x_i + x_1 \left(a + \sum_{i=3}^{k} x_i - x_2 x_k\right)\right).$$

Another computation gives that

$$-\left(\frac{a+\sum_{i=2}^{k} x_i}{x_1^2}\right)X_1 + \frac{1}{x_1}\left[\sum_{i=2}^{k} X_i\right] = \frac{\mathbf{B}}{x_1\left(\prod_{i=1}^{k} x_i\right)}$$

where

$$\mathbf{B} = -(x_1+1)\left(a + \sum_{i=2}^k x_i\right) \left[\prod_{i=2}^{k-1} L_i\right] \left(a + \sum_{i=1}^{k-1} x_i - x_2 x_k\right) + \left(a + \sum_{i=1}^k x_i + x_1 x_k\right) \mathbf{C} - x_k (x_k+1) \left(a + \sum_{i=2}^k x_i - x_1 x_{k-1}\right) \left[\prod_{i=1}^{k-2} L_i\right],$$

and

$$\mathbf{C} = \sum_{m=2}^{k-1} x_m (x_m + 1) S_m M_m.$$

Recall that  $L_i = 1 + x_i + x_{i+1}$  for all i = 1, ..., k - 1,  $S_m = x_{m-1} - x_{m+1}$  and  $M_m = \prod_{i=1, i \neq m-1, m}^{k-1} (1 + x_i + x_{i+1})$  for  $2 \le m \le k - 1$ . Therefore we want to prove that  $\mathbf{A} = \mathbf{B}$ .

Step 2a. First we show that **B** contains  $L_2$  and  $L_3$  as a factors. To see this, it suffices to check that  $L_2$  and  $L_3$  are factors of **C**. Observe that if  $L_2 = 0$ , then we can write  $L_3 = x_4 - x_2$ , and  $x_3 = -(1 + x_2)$  hence  $x_1 - x_3 = x_1 + x_2 + 1$ . Therefore

$$\begin{aligned} \mathbf{C}|_{\{L_2=0\}} &= \sum_{m=2}^{3} x_m (x_m+1) S_m M_m \\ &= \sum_{m=2}^{3} x_m (x_m+1) (x_{m-1} - x_{m+1}) \left( \prod_{i=1, i \neq m-1, m}^{k-1} (1 + x_i + x_{i+1}) \right) \\ &= x_2 (1 + x_2) (x_1 - x_3) (x_4 - x_2) \left( \prod_{i=4}^{k-1} (1 + x_i + x_{i+1}) \right) \\ &+ x_2 (1 + x_2) (x_2 - x_4) (1 + x_1 + x_2) \left( \prod_{i=4}^{k-1} (1 + x_i + x_{i+1}) \right) = 0. \end{aligned}$$

So it is a factor of **C**.

If  $L_3 = 0$ , then  $L_2 = x_2 - x_4$ , and  $x_3 - x_5 = -(1 + x_4 + x_5)$ . Hence

$$\begin{aligned} \mathbf{C}|_{\{L_3=0\}} &= \sum_{m=3}^{4} x_m (x_m+1) S_m M_m = \\ &= \sum_{m=3}^{4} x_m (x_m+1) (x_{m-1} - x_{m+1}) \left( \prod_{i=1, i \neq m-1, m}^{k-1} (1 + x_i + x_{i+1}) \right) \\ &= x_4 (1 + x_4) (x_2 - x_4) (1 + x_1 + x_2) \left( \prod_{i=4}^{k-1} (1 + x_i + x_{i+1}) \right) \\ &- x_4 (1 + x_4) (1 + x_4 + x_5) (1 + x_2 + x_3) (1 + x_1 + x_2) \times \\ &\times \left( \prod_{i=5}^{k-1} (1 + x_i + x_{i+1}) \right) = 0. \end{aligned}$$

Hence  $L_3$  is a factor of **C**.

The above results imply that

$$\mathbf{C} = L_2 L_3 Q_k(x_1, x_2, x_4, \dots, x_k),$$

(observe that  $x_3$  does not appear in the expression of  $Q_k$ ). Hence  $L_2$  and  $L_3$  are factors in the expression of **B**, and then

$$\mathbf{B} = L_2 L_3 \left[ -(x_1+1) \left( a + \sum_{i=2}^k x_i \right) \left[ \prod_{i=4}^{k-1} L_i \right] \left( a + \sum_{i=1}^{k-1} x_i - x_2 x_k \right) \right. \\ \left. + \left( a + \sum_{i=1}^k x_i + x_1 x_k \right) Q_k - x_k (x_k+1) \left( a + \sum_{i=2}^k x_i - x_1 x_{k-1} \right) L_1 \left[ \prod_{i=4}^{k-2} L_i \right] \right].$$

Furthermore

$$Q_{k} = -x_{2}(1+x_{2})\left(\prod_{i=4}^{k-1}L_{i}\right) + (x_{2}-x_{4})L_{1}\left(\prod_{i=4}^{k-1}L_{i}\right) + x_{4}(1+x_{4})L_{1}\left(\prod_{i=5}^{k-1}L_{i}\right) + \sum_{m=5}^{k-1}x_{m}(x_{m}+1)(x_{m-1}-x_{m+1})L_{1}\left(\prod_{i=4, j\neq m-1, m}^{k-1}L_{j}\right).$$

Step 2b. Now we state the following claim, that will we proved at the end of the proof.

**Claim:** For  $k \ge 6$ ,  $Q_k(x_1, x_2, x_4, \dots, x_k) = \left(\prod_{i=4}^{k-2} L_i\right) [x_1 x_2 L_{k-1} - x_{k-1} x_k L_1].$ 

By using the claim,  $\mathbf{B} = \left[\prod_{i=2}^{k-2} L_i\right] \mathbf{D}$ , where

$$\mathbf{D} = -(x_1+1)\left(a + \sum_{i=2}^{k} x_i\right)\left(a + \sum_{i=1}^{k-1} x_i - x_2 x_k\right) L_{k-1} \\ + \left(a + \sum_{i=1}^{k} x_i + x_1 x_k\right) [x_1 x_2 L_{k-1} - x_{k-1} x_k L_1] \\ - x_k (x_k+1)\left(a + \sum_{i=2}^{k} x_i - x_1 x_{k-1}\right) L_1.$$

Step 2c. Observe that

$$\begin{aligned} &-x_k(x_k+1)\left(a+\sum_{i=2}^k x_i-x_1x_{k-1}\right)L_1-x_{k-1}x_kL_1\left(a+\sum_{i=1}^k x_i+x_1x_k\right)\\ &= -L_1x_k\left[(x_k+1)\left(a+\sum_{i=2}^k x_i-x_1x_{k-1}\right)-x_{k-1}\left(a+\sum_{i=1}^k x_i+x_1x_k\right)\right]\\ &= -L_1x_k\left[x_{k-1}\left(a+\sum_{i=1}^k x_i+x_1x_k-x_1-x_1x_k\right)+(1+x_k)\left(a+\sum_{i=2}^k x_i\right)\right]\\ &= -L_1x_k\left[x_{k-1}\left(a+\sum_{i=2}^k x_i\right)+(1+x_k)\left(a+\sum_{i=2}^k x_i\right)\right]\\ &= -L_1x_kL_{k-1}\left(a+\sum_{i=2}^k x_i\right).\end{aligned}$$

Thus **D** also contains the factor  $L_{k-1}$ , and therefore  $\mathbf{B} = \left[\prod_{i=2}^{k-1} L_i\right] \mathbf{E}$ , where

$$\mathbf{E} = \begin{bmatrix} -(x_1+1)\left(a + \sum_{i=2}^{k} x_i\right)\left(a + \sum_{i=1}^{k-1} x_i - x_2 x_k\right) \\ + \left(a + \sum_{i=1}^{k} x_i + x_1 x_k\right)x_1 x_2 - x_k L_1\left(a + \sum_{i=2}^{k} x_i\right)\end{bmatrix}.$$

Step 2d. Is not difficult to check that **E** is a quadratic polynomial in  $x_3$ . Now we see that **E** vanishes either when  $x_3 = x_{3,1} := -a - \sum_{i=1, i \neq 3}^k x_i$  (that is when  $a + \sum_{i=1}^k x_i = 0$ ), and when

$$x_3 = x_{3,2} := -\left(a + \sum_{i=4}^{k-1} x_i + \frac{x_2(1 - x_1 x_k)}{1 + x_1}\right).$$

In the first case we have

$$\begin{aligned} \mathbf{E}|_{\{a+\sum_{i=1}^{k} x_i=0\}} &= x_1(x_1+1)(-x_k-x_2x_k) + x_1^2x_2x_k + L_1x_1x_k \\ &= x_1x_k \left[-(x_1+1)(x_2+1) + x_1x_2 + 1 + x_1 + x_2\right] \\ &= x_1x_k \left[-1-x_1-x_2-x_1x_2 + x_1x_2 + 1 + x_1 + x_2\right] = 0. \end{aligned}$$

In the second case, since  $a + \sum_{i=3}^{k-1} x_i = x_2(x_1x_k - 1)/(x_1 + 1)$ , we have

$$\mathbf{E}|_{\{x_3=x_{3,2}\}} = -\left(x_2 + \frac{x_2(x_1x_k - 1)}{x_1 + 1}\right)(x_1 + 1)\left(\frac{x_2(x_1x_k - 1)}{x_1 + 1} + x_1 + x_2 - x_k - x_2x_k\right) \\ + x_1x_2\left(x_1 + x_2 + \frac{x_2(x_1x_k - 1)}{x_1 + 1} + x_1x_k\right) \\ - L_1x_k\left(x_2 + \frac{x_2(x_1x_k - 1)}{x_1 + 1}\right) =: \frac{1}{x_1 + 1}\mathbf{F},$$

where (after some computations)

$$\mathbf{F} = -x_2 x_1 (x_k + 1) L_1 (x_1 - x_k) + x_1^2 x_2 L_1 (x_k + 1) - L_1 x_1 x_2 x_k (x_k + 1)$$
  
=  $x_1 x_2 L_1 (x_k + 1) [-(x_1 - x_k) + x_1 - x_k] = 0.$ 

Therefore  $\mathbf{E}|_{\{x_3=x_{3,2}\}} = 0.$ 

In summary, as a quadratic polynomial in  $x_3$ , **E** factorizes as

$$\mathbf{E} = -(x_1+1)\left(x_3+a+\sum_{i=1,i\neq 3}^k x_i\right)\left(x_3+a+\sum_{i=4}^{k-1} x_i+(x_2(1-x_1x_k))/(x_1+1)\right) = -\left(a+\sum_{i=1}^k x_i\right)\left(\left(x_3+a+\sum_{i=4}^{k-1} x_i\right)(x_1+1)+x_2(1-x_1x_k)\right) = -\left(a+\sum_{i=1}^k x_i\right)\left((x_1+1)\left(a+\sum_{i=3}^{k-1} x_i\right)+x_2-x_1x_2x_k\right) = -\left(a+\sum_{i=1}^k x_i\right)\left(a+\sum_{i=2}^{k-1} x_i+x_1\left(a+\sum_{i=3}^{k-1} x_i-x_2x_k\right)\right).$$

So, finally, we get that

$$\mathbf{B} = -\left[\prod_{i=2}^{k-1} L_i\right] \left(a + \sum_{i=1}^k x_i\right) \left(a + \sum_{i=2}^{k-1} x_i + x_1 \left(a + \sum_{i=3}^{k-1} x_i - x_2 x_k\right)\right) = \mathbf{A},$$

as we wanted to show.

To end the proof it only remains to prove the claim. We proceed by induction. That it is true when k = 6 is straightforward. Assume now that the claim is true for  $Q_k$ , then

$$Q_{k+1} = -x_2(1+x_2) \left(\prod_{i=4}^k L_i\right) + (x_2 - x_4) L_1 \left(\prod_{i=4}^k L_i\right) + x_4(1+x_4) L_1 \left(\prod_{i=5}^k L_i\right) + \sum_{m=5}^{k-1} x_m(1+x_m)(x_{m-1} - x_{m+1}) L_1 \left(\prod_{i=4,i\neq m-1,m}^k L_i\right)$$
  
=  $L_k Q_k + L_1 \left(\prod_{i=4}^{k-2} L_i\right) (x_k(1+x_k)(x_{k-1} - x_{k+1})).$ 

By using the hypothesis of induction, we obtain that

$$Q_{k+1} = L_k \left( \prod_{i=4}^{k-2} L_i \right) [x_1 x_2 L_{k-1} - x_{k-1} x_k L_1] + L_1 \left( \prod_{i=4}^{k-2} L_i \right) (x_k (1+x_k) (x_{k-1} - x_{k+1})) = \left( \prod_{i=4}^{k-2} L_i \right) (x_1 x_2 L_k L_{k-1} + L_1 [x_k (1+x_k) (x_{k-1} - x_{k+1}) - x_{k-1} x_k L_k]).$$

Thus

$$Q_{k+1} = L_k \left(\prod_{i=4}^{k-2} L_i\right) \left[x_1 x_2 L_{k-1} - x_{k-1} x_k L_1\right] + L_1 \left(\prod_{i=4}^{k-2} L_i\right) \left(x_k (1+x_k) (x_{k-1} - x_{k+1})\right) \\ = \left(\prod_{i=4}^{k-2} L_i\right) \left(x_1 x_2 L_k L_{k-1} + L_1 \left[x_k (1+x_k) (x_{k-1} - x_{k+1}) - x_{k-1} x_k L_k\right]\right).$$

An easy computation shows that  $x_k(1+x_k)(x_{k-1}-x_{k+1})-x_{k-1}x_kL_k = -L_{k-1}x_kx_{k+1}$ , and the result follows. Therefore, Theorem 1 is proved.

## 3 Geometrical issues in the odd case.

Before proving Theorem 2 and Propositions 3 and 4 we need some preliminary results. Recall that a map H which is a first integral for  $G^p := G \circ \stackrel{p}{\cdots} \circ G$  is also called sometimes  $a \ p-first \ integral$ , or simply, for short, a p-integral, of G, see [23].

The following lemma, which is very easy to prove, gives light on one utility of p-first integrals, specially if they are not symmetric functions of their arguments.

**Lemma 5.** Let H be a p-integral of a map G. Then then for any symmetric function of p variables S, the function  $V_S := S(H, H \circ G, H \circ G^2, \ldots, H \circ G^{p-1})$  is a first integral of G.

**Lemma 6.** Set  $k = 2\ell + 1$ ,  $\mathbf{x} = (x_1, x_2, \dots, x_k) \in \mathcal{Q}^+$  and let W be the function given in (10). Then, if we define the polynomial

$$Z(\mathbf{x}): = (\prod_{i=1}^{k} x_i)[W(\mathbf{x}) - W(F(\mathbf{x}))] =$$
  
=  $\prod_{j=0}^{\ell} x_{2j+1}(x_{2j+1}+1) - \left(a + \sum_{i=1}^{2\ell+1} x_i\right) \prod_{j=1}^{\ell} x_{2j}(x_{2j}+1),$ 

it holds that  $Z(F(\mathbf{x})) = \det(DF(\mathbf{x}))Z(\mathbf{x})$ .

**Proof.** In Theorem 2 (a) we will prove that W is a 2-integral of F. Thus, if we define  $\tilde{Z} := W - W(F)$ , then

$$\tilde{Z}(F(\mathbf{x})) = -\tilde{Z}(\mathbf{x}). \tag{14}$$

From the above equality it is clear that  $\{\mathbf{x} \in \mathcal{Q}^+ : \tilde{Z}(\mathbf{x}) = 0\} = \{\mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) = 0\}$  is an invariant hypersurface by *F*. Notice also that if we define  $\Pi(\mathbf{x}) := \prod_{j=1}^k x_j$ , it holds that

$$\Pi(F(\mathbf{x})) = \frac{a + \sum_{i=2}^{k} x_i}{x_1^2} \Pi(\mathbf{x}) = -\det(DF(\mathbf{x})) \Pi(\mathbf{x}).$$
(15)

Since  $Z(\mathbf{x}) = \Pi(\mathbf{x})\tilde{Z}(\mathbf{x})$ , by using equalities (14) and (15), we obtain that

$$Z(F(\mathbf{x})) = \Pi(F(\mathbf{x}))\tilde{Z}(F(\mathbf{x})) = \det(DF(\mathbf{x}))\Pi(\mathbf{x})\tilde{Z}(\mathbf{x}) = \det(DF(\mathbf{x}))Z(\mathbf{x}),$$

as we wanted to see.

**Proof of Theorem 2.** (a) The proof of the equality  $W(F^2(\mathbf{x})) = W(\mathbf{x})$  is straightforward. We have obtained the expression (10) inspired in the results of [22]. In that paper it is proved that

$$\frac{(y_n+1)(y_{n+2}+1)}{y_{n+1}} = \frac{(y_{n+2}+1)(y_{n+4}+1)}{y_{n+3}}$$

where  $\{y_n\}$  is the sequence given by the 3-rd order Lyness' recurrence  $y_{n+3} = (a + y_{n+1} + y_{n+2})/y_n$ . Notice that this property is equivalent to say that for k = 3, W is a 2-integral for F.

(b-c) By applying Lemma 5 with S(u, v) = u + v and S(u, v) = uv, we obtain the first integrals  $V_3$  and  $V_1$ , respectively. The functionally independence of  $V_1, V_2$  and  $V_3$ , for  $\ell \ge 2$ , follows from straightforward computations and it is already established in [11].

(d-f) From Lemma 6 we know that  $Z(F(\mathbf{x})) = \det(DF(\mathbf{x}))Z(\mathbf{x})$ , where recall that  $Z(\mathbf{x}) = (\prod_{i=1}^{k} x_i)[W(\mathbf{x}) - W(F(\mathbf{x}))]$ . Note also that

$$\det(DF(\mathbf{x})) = (-1)^k \frac{a + x_2 + \dots + x_{k-1}}{x_k^2}.$$

Since when k is odd  $\det(DF) < 0$  on  $\mathcal{Q}^+$ , equation  $Z(F) = \det(DF)Z$  means that

$$\mathcal{G} = \{ \mathbf{x} \in \mathcal{Q}^+ : W(\mathbf{x}) = W(F(\mathbf{x})) \} = \{ \mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) = 0 \}$$

is invariant by F and that F maps the region  $\{\mathbf{x} : Z(\mathbf{x}) > 0\}$  into the region  $\{\mathbf{x} : Z(\mathbf{x}) < 0\}$ and viceversa. Furthermore it implies that the dynamics of  $F^2$  on each of these sets are conjugated, being the map F itself the conjugation. Moreover, any periodic orbit with odd period must lie in  $\mathcal{G}$ , as we wanted to see. (g-h) By using the Change of Variables Theorem it is easy to see that if G is a diffeomorphism of  $\mathcal{U}$ , and on this region  $\mu$  is a positive function that satisfies  $\mu(G(\mathbf{x})) = \det(DG(\mathbf{x}))\mu(\mathbf{x})$ , then

$$m(B) = \int_B \frac{1}{\mu(\mathbf{x})} \, d\mathbf{x}$$

is an invariant measure for G. By Lemma 6 we know that  $Z(F(\mathbf{x})) = \det(DF(\mathbf{x}))Z(\mathbf{x})$  and by equality (15), that  $\Pi(F(\mathbf{x})) = -\det(DF(\mathbf{x}))\Pi(\mathbf{x})$ . By using these results we have that  $Z(F^2(\mathbf{x})) = \det(DF^2(\mathbf{x}))Z(\mathbf{x})$  and  $\Pi(F^2(\mathbf{x})) = \det(DF^2(\mathbf{x}))\Pi(\mathbf{x})$ , being both equalities in the corresponding domains, which are invariant by  $F^2$ . Hence (f) and (g) follow.

### 3.1 Order reduction

Recall that in the previous section we have seen that when  $k = 2\ell + 1$ , the regions  $\{\mathbf{x} : Z(\mathbf{x}) > 0\}$  and  $\{\mathbf{x} : Z(\mathbf{x}) < 0\}$  are invariant by  $F^2$ , and the dynamics on both region are conjugated. This observation allows us to give a new application of the invariant W. We can reduce the study of the dynamics of F on  $\{\mathbf{x} : Z(\mathbf{x}) \neq 0\}$  to the study of a new (k-1)-dimensional map, having one more parameter.

For instance for n = 3, we get that W(x, y, z) = (x + 1)(z + 1)/y, and hence any admissible level surface  $\{\mathbf{x} : W(\mathbf{x}) = w\}, w \neq 0$ , can be described as y = k(x + 1)(z + 1), where k = 1/w. Therefore

$$F^{2}|_{\{W=w\}}(x,y,z) = \left(z, \frac{a+z+k(x+1)(z+1)}{x}, \frac{a+k+z(k+1)}{kx(z+1)}\right),$$

and we can reduce the study of the dynamics of  $F^2$  to the study of the reduced map

$$\tilde{F}_2(x,z) = \left(z, \frac{a+k+z(k+1)}{kx(z+1)}\right),\,$$

or, equivalently, the study of the second order difference equation

$$y_{n+2} = \frac{a+k+y_{n+1}(k+1)}{ky_n(y_{n+1}+1)},$$
(16)

as in [22, Eqs. (8) and (9)]. The dynamics of this difference equation is studied in [8, Ex. 3]. Equation (16) is sometimes called *generalized Lyness' recurrence*, see [20].

For n = 5, the integral of  $F^2$  is W(x, y, z, t, s) = (x + 1)(z + 1)/(yt). Again any admissible level surface  $\{\mathbf{x} : W(\mathbf{x}) = w\}, w \neq 0$ , can be described as t = k(x + 1)(z + 1)(s + 1)/y, where k = 1/w. Therefore proceeding as before we can reduce the study of the dynamics of  $F^2$  to the study of the reduced map

$$\tilde{F}_2(x, y, z, s) = \left(y, z, t, \frac{p_2(z, s; k)x^2 + p_1(y, z, s; a, k)x + p_0(y, z, s; a, k)}{xy^2}\right)$$

where  $p_2(z, s; k) = k (s + 1) (z + 1)$ ,  $p_1(y, z, s; a, k) = 2 k (s + 1) (z + 1) + y (a + s + z)$ , and  $p_0(y, z, s; a, k) = k (s + 1) (z + 1) + y (z + s + a + y)$ , or equivalently to the difference equation

$$y_{n+4} = \frac{p_2(y_{n+2}, y_{n+3}; k)y_n^2 + p_1(y_{n+1}, y_{n+2}, y_{n+3}; a, k)y_n + p_0(y_{n+1}, y_{n+2}, y_{n+3}; a, k)}{y_n y_{n+1}^2}.$$

Clearly, the described procedure can be generalized to higher dimensions.

## 4 Dynamics of the low dimensional cases

This subsection is devoted to prove Propositions 3 and 4.

Along the section we will use a straightforward consequence of the following result:

**Theorem 7** (Bastien and Rogalsky, [4]). Let  $\bar{\mathbf{x}}$  be the fixed point of F in  $\mathcal{Q}^+$ . For any  $h > V_1(\bar{\mathbf{x}})$ , the level sets  $\{V_1 = h\} \cap \mathcal{Q}^+$  are homeomorphic to  $\mathbb{S}^{k-1}$ .

**Corollary 8.** Let  $K \neq \emptyset$  be by the intersection of some level sets of different first integrals of F, including  $V_1$  among them. Then  $K \cap Q^+$  is a compact set, invariant by F.

For the sake of completeness and to compare with the cases k = 4, 5, we start by recalling some results for the case k = 3, most of them already proved in [7].

### 4.1 The 3-dimensional map

For k = 3,

$$F(x, y, z) = \left(y, z, \frac{a+y+z}{x}\right),$$

the Lie symmetry given in Theorem 1, is

$$\mathbf{X}_3 := \begin{bmatrix} x(x+1)(1+y+z)(a+x+y-yz)\frac{\partial}{\partial x} + y(y+1)(x-z)(a+x+y+z+xz)\frac{\partial}{\partial y} \\ + z(z+1)(1+x+y)(a+y+z-xy)\frac{\partial}{\partial z} \end{bmatrix} / (xyz)$$

and since  $\mathbf{X}_3(V_i) = 0$ , for i = 1, 2, the functions  $V_1$  and  $V_2$  given in (3) and (4) are first integrals for  $\mathbf{X}_3$  and F. Also, by Theorem 2, W(x, y, z) := (x + 1) (z + 1) / y is a 2-integral of  $F; \mathcal{G} = \{\mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) = 0\}$  is invariant by F, where

$$Z(\mathbf{x}) = x(x+1)z(z+1) - (a+x+y+z)y(y+1);$$

and F maps  $\mathcal{G}^+ := \{ \mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) > 0 \}$  into  $\mathcal{G}^- := \{ \mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) < 0 \}$ , and viceversa.

Let  $\bar{\mathbf{x}}$  be the fixed point in  $\mathcal{Q}^+$ , of F. Set  $h > V_1(\bar{\mathbf{x}})$ ,  $k > V_2(\bar{\mathbf{x}})$ . In [7] it is proved that  $\{V_1 = h\} \cap \mathcal{Q}^+$  and  $\{V_2 = k\} \cap \mathcal{Q}^+$  are diffeomorphic to two dimensional spheres, and their

transversal intersections are formed by exactly two disjoints curves, both diffeomorphic to circles. Their non transversal intersections correspond to:

(a) The 2-periodic points of F (which are equilibrium points of  $\mathbf{X}_3$ ) given by the curve  $\mathcal{L} := \{(x, (x+a)/(x-1), x) | x > 1\}.$ 

(b) The levels placed at  $\mathcal{G}$ . Those placed at  $\mathcal{G} \setminus \{\bar{\mathbf{x}}\}\$  are formed by exactly one curve, diffeomorphic to a circle.

Finally, since  $\mathbf{X}_3$  is also a Lie symmetry of  $F^2$ , as a consequence of [7, Thm. 18] or [8, Thm. 1], we know that the map  $F^2$  restricted to each of the sets  $\{V_1 = h\} \cap \{V_2 = k\}$ , which is not simply a fixed point of  $F^2$ , is conjugated to a rotation. Further discussion about the possible rotation numbers can be found in [7].

#### 4.2 The 4-dimensional map.

**Proof of Proposition 3.** Equations (6)–(8) give the following Lie symmetry for the 4-dimensional Lyness' map:

$$\mathbf{X}_{4} = \begin{bmatrix} x (x+1) (1+y+z) (1+z+t) (a+x+y+z-yt) \frac{\partial}{\partial x} \\ +y (y+1) (1+z+t) (a+x+y+z+t+xt) (x-z) \frac{\partial}{\partial y} \\ +z (z+1) (1+x+y) (a+x+y+z+t+xt) (y-t) \frac{\partial}{\partial z} \\ -t (t+1) (1+x+y) (1+y+z) (a+y+z+t-xz) \frac{\partial}{\partial t} \end{bmatrix} / (xyzt).$$

A straightforward computation shows that the Lie symmetry satisfies  $\mathbf{X}_4(V_i) = 0$ , for i = 1, 2. Hence, the orbits of both F and  $\mathbf{X}_4$  generically lie in a two dimensional surface of the form  $I_{h,k} := \{V_1 = h\} \cap \{V_2 = k\} \cap \mathcal{Q}^+$ .

Let  $C_{h,k}$  be a connected component of  $I_{h,k}$ . From Corollary 8 we know that  $C_{h,k}$  is compact. If  $\{V_1 = h\}$  and  $\{V_2 = k\}$  intersect transversally on  $C_{h,k}$ , then for all points

Rank 
$$\begin{pmatrix} (V_1)_x & (V_1)_y & (V_1)_z & (V_1)_t \\ (V_2)_x & (V_2)_y & (V_2)_z & (V_2)_t \end{pmatrix} = 2.$$

This fact implies that the dual 2-form associated to the 2-field

$$\nabla V_1 \wedge \nabla V_2 = [(V_1)_x (V_2)_y - (V_1)_y (V_2)_x] \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y} + [(V_1)_x (V_2)_z - (V_1)_z (V_2)_x] \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial z}$$
  
+ [(V\_1)\_x (V\_2)\_t - (V\_1)\_t (V\_2)\_x] \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial t} + [(V\_1)\_y (V\_2)\_z - (V\_1)\_z (V\_2)\_y] \frac{\partial}{\partial y} \wedge \frac{\partial}{\partial z}   
+ [(V\_1)\_y (V\_2)\_t - (V\_1)\_t (V\_2)\_y] \frac{\partial}{\partial y} \wedge \frac{\partial}{\partial t} + [(V\_1)\_z (V\_2)\_t - (V\_1)\_t (V\_2)\_z] \frac{\partial}{\partial z} \wedge \frac{\partial}{\partial t},

given by

$$\begin{split} \omega &= & [(V_1)_z (V_2)_t - (V_1)_t (V_2)_z] dx dy + [(V_1)_z (V_2)_x - (V_1)_x (V_2)_z] dx dz \\ &+ [(V_1)_z (V_2)_y - (V_1)_y (V_2)_z] dx dt + [(V_1)_t (V_2)_x - (V_1)_x (V_2)_t] dy dz \\ &+ [(V_1)_t (V_2)_y - (V_1)_y (V_2)_t] dy dt + [(V_1)_x (V_2)_y - (V_1)_y (V_2)_x] dz dt \end{split}$$

is nonzero at every point of  $C_{h,k}$ , and therefore it is orientable, see [1, Sec. 2.5] or [21, Sec. X, §1.].

Let  $\mathbf{X}_{4}^{h,k}$  denote the restriction of the vector field  $\mathbf{X}_{4}$  to the invariant surface  $C_{h,k}$ . Some computations show that the unique equilibrium point of  $\mathbf{X}_{4}$  in  $\mathcal{Q}^{+}$  is the fixed point of F. Hence  $\mathbf{X}_{4}^{h,k}$  has no equilibrium points, and therefore the Poincaré–Hopf formula gives:

$$0 = i(\mathbf{X}_4^{h,k}) = \chi(C_{h,k}) = 2 - 2g$$

where  $i(\mathbf{X}_{4}^{h,k})$  denotes the sum of the indices of the equilibrium points of  $\mathbf{X}_{4}^{h,k}$  in  $C_{h,k}$  and  $\chi(C_{h,k})$  and g are the Euler characteristic and the genus of the surface  $C_{h,k}$ , respectively. Hence the genus of  $C_{h,k}$  is one. An orientable, compact, connected surface of genus one is a torus, as we wanted to prove.

#### 4.3 The 5-dimensional map

**Proof of Proposition 4.** By Theorem 2 we know that

$$W(x, y, z, t, s) := \frac{(x+1)(z+1)(s+1)}{yt}$$

is a 2-integral of F. Moreover, the 3 functionally independent first integrals of F given in (3), (4) and (11), are

$$\begin{split} V_1(x,y,z,t,s) &= \frac{(a+x+y+z+t+s)\left(x+1\right)\left(y+1\right)\left(z+1\right)\left(t+1\right)\left(s+1\right)}{xyzts},\\ V_2(x,y,z,t,s) &= \frac{(a+x+y+z+t+s+xs)\left(1+x+y\right)\left(1+y+z\right)\left(1+z+t\right)\left(1+t+s\right)}{xyzts},\\ V_3(x,y,z,t,s) &= \frac{x\left(x+1\right)z\left(z+1\right)s\left(s+1\right)+\left(a+x+y+z+t+s\right)y\left(y+1\right)t\left(t+1\right)}{xyzts}. \end{split}$$

Some tedious computations show that the hypersurface  $\mathcal{G}$  is in the locus of non-transversality of the three level sets of the integrals  $V_i$ , i = 1, 2, 3 in  $Q^+$ . Recall that precisely,  $\mathcal{G} = \{\mathbf{x} \in Q^+ : Z(\mathbf{x}) = 0\}$  is invariant by F, where

$$Z(\mathbf{x}) = x(x+1)z(z+1)s(s+1) - (a+x+y+z+t+s)y(y+1)t(t+1),$$

and that F maps  $\mathcal{G}^+ = \{ \mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) > 0 \}$  into  $\mathcal{G}^- = \{ \mathbf{x} \in \mathcal{Q}^+ : Z(\mathbf{x}) < 0 \}$ , and viceversa.

Equations (6)-(8) give the following Lie symmetry for the 5-dimensional Lyness' map:

$$\begin{aligned} \mathbf{X}_{5} &= \left[ x \left( x+1 \right) \left( 1+y+z \right) \left( 1+z+t \right) \left( 1+t+s \right) \left( a+x+y+z+t-ys \right) \frac{\partial}{\partial x} \right. \\ &+ y \left( y+1 \right) \left( 1+t+s \right) \left( 1+z+t \right) \left( a+x+y+z+t+s+xs \right) \left( x-z \right) \frac{\partial}{\partial y} \right. \\ &+ z \left( z+1 \right) \left( 1+x+y \right) \left( 1+t+s \right) \left( a+x+y+z+t+s+xs \right) \left( y-t \right) \frac{\partial}{\partial z} \\ &+ t \left( t+1 \right) \left( 1+x+y \right) \left( 1+y+z \right) \left( a+x+y+z+t+s+xs \right) \left( z-s \right) \frac{\partial}{\partial t} \\ &- s \left( s+1 \right) \left( 1+x+y \right) \left( 1+y+z \right) \left( 1+z+t \right) \left( a+y+z+t+s-tx \right) \frac{\partial}{\partial s} \right] / (xyzts). \end{aligned}$$

Again, direct computations show that  $\mathbf{X}_5(V_i) = 0$ , for i = 1, 2, 3. Hence the orbits of both F, and  $\mathbf{X}_5$  lie in a two dimensional surface of the form  $I_{h,k,\ell} := \{V_1 = h\} \cap \{V_2 = k\} \cap \{V_3 = \ell\} \cap \mathcal{Q}^+$ .

Let  $C_{h,k,\ell}$  be a connected component of  $I_{h,k,\ell}$ . From Corollary 8 we know that  $C_{h,k,\ell}$  is compact. If  $\{V_1 = h\}$ ,  $\{V_2 = k\}$  and  $\{V_3 = k\}$  intersect transversally on  $C_{h,k,\ell}$ , then for all points

Rank 
$$\begin{pmatrix} (V_1)_x & (V_1)_y & (V_1)_z & (V_1)_t & (V_1)_s \\ (V_2)_x & (V_2)_y & (V_2)_z & (V_2)_t & (V_2)_s \\ (V_3)_x & (V_3)_y & (V_3)_z & (V_3)_t & (V_3)_s \end{pmatrix} = 3.$$

Similarly than in the case k = 4, this fact implies that the dual 2-form associated to the 3-field  $\nabla V_1 \wedge \nabla V_2 \wedge \nabla V_3$  is nonzero at every point of  $C_{h,k,\ell}$ , and therefore this set is a two-dimensional orientable manifold.

It is not difficult to check that all the equilibrium points of  $\mathbf{X}_5$  in  $\mathcal{Q}^+$  are the points of the curve

$$\mathcal{L} = \left\{ \mathbf{x} = \left( x, \frac{2x+a}{x-2}, x, \frac{2x+a}{x-2}, x \right) \quad \text{with} \quad x > 2 \right\},\$$

which contains a continuum of two periodic points and the fixed point of F. Moreover  $\mathcal{L}$  belongs to the locus of non-transversality of the integrals  $V_1$ , and  $V_2$  in  $\mathcal{Q}^+$ .

The above observation implies that  $\mathbf{X}_5$  has no equilibrium points in any level set  $I_{h,k,\ell}$ where the three first integrals intersect transversally. Therefore the Poincaré–Hopf formula gives

$$0 = i(\mathbf{X}_5^{h,k,\ell}) = \chi(C_{h,k,\ell}) = 2 - 2g_5$$

for each connected component  $C_{h,k,\ell}$  of such a level set (where  $\mathbf{X}_5^{h,k,\ell}$  is the restriction of  $\mathbf{X}_5$  to  $C_{h,k,\ell}$ ).

Hence g = 1, which implies that  $C_{h,k,\ell}$  is a torus (since it is two-dimensional, orientable, compact, manifold of genus one), as we wanted to proof.

Finally, observe that the fact that F maps  $\mathcal{G}^+$  into  $\mathcal{G}^-$ , and viceversa, implies that most  $I_{h,k,\ell}$  have at least one connected component on each set. In fact, it seems that each  $I_{h,k,\ell}$ , not included in  $\Gamma$ , has exactly two connected components in  $\mathcal{Q}^+$ , as it happens when k = 3.

See Figures 2 and 3 for two illustrations of this assertion.



Figure 3. Projection into  $\mathbb{R}^3$  of the first  $10^4$  iterates of the Lyness' map, for k = 5 and a = 4, starting at (1, 2, 3, 4, 5). Odd and even iterates are in different connected components.

From the above proof it is clear that if some  $I_{h,k,\ell}$  cuts  $\mathcal{L}$  then the first three integrals do not cut transversally on it. Let us see that in general  $I_{h,k,\ell} \cap \mathcal{L} = \emptyset$ . This will be a consequence of the shape of the function

$$v_1(x) := V_1\left(x, \frac{2x+a}{x-2}, x, \frac{2x+a}{x-2}, x\right), \quad x > 2, \ a \ge 0.$$

This function has a global minimum at the coordinate given by the fixed point x = 2 + $\sqrt{4+a}$ , and  $\lim_{x \to 2^+} v_1(x) = \lim_{x \to +\infty} v_1(x) = +\infty$ . Thus, given  $h > v_1(2+\sqrt{4+a})$  there are only two solutions  $x_1(h) < 2+\sqrt{4+a} < x_2(h)$ 

of the equation  $v_1(x) = h$ . Now set

$$v_i(x) := V_i\left(x, \frac{2x+a}{x-2}, x, \frac{2x+a}{x-2}, x\right), \quad x > 2, \ a \ge 0,$$

for i = 2, 3. Then, for any value of k and  $\ell$  satisfying  $k \notin \{v_2(x_1(h)), v_2(x_2(h))\}$  or  $\ell \notin$  $\{v_3(x_1(h)), v_3(x_2(h))\},$  we obtain that  $I_{h,k,\ell}$ , does not intersect  $\mathcal{L}$ .

#### Conclusions 5

Several properties for the k-dimensional Lyness' map F have been given, like the existence of a Lie symmetry for F and of a new and simple invariant for  $F^2$ . This Lie symmetry together with the new invariant give information for k = 4 and 5 about the dynamics and the topology of the level surfaces where the dynamics of F is confined. Some general results for k odd have been also presented. However, on the contrary that happens in the cases k = 2 and 3, the numerical explorations indicate that for k = 4, 5 the orbits of the map are not contained in the orbits of the flow of the Lie symmetry with the same initial condition, although they are placed in the same manifold, which has dimension  $k - E(\frac{k+1}{2})$ . This is an obstruction to apply the theoretical tools developed in [8].

Numerical simulations seem to show that for some initial conditions the projection in  $\mathbb{R}^3$  of the iterates of F when k = 6, 7 fill densely a 2-dimensional manifold, indicating that probably they live in a 3-dimensional manifold of  $\mathbb{R}^6$  and  $\mathbb{R}^7$ . These facts are coherent with the conjecture of [11] about the number of independent first integrals of the Lyness' maps, and show that for  $k \ge 6$  the dynamics are much more complicated.

When  $k = 2\ell$ , the simplest scenario that we imagine for the dynamics of the k-dimensional Lyness' map is that most of the orbits lie on invariant manifolds which are diffeomorphic to  $\ell$ -dimensional tori,  $S^1 \times \stackrel{\ell}{\cdots} \times S^1$ . On the other hand, when  $k = 2\ell + 1$ , most of them lie on two diffeomorphic copies of  $S^1 \times \stackrel{\ell}{\cdots} \times S^1$ , separated by the invariant set  $\mathcal{G}$ . Moreover these orbits jump from one of these tori to the other one and viceversa.

In any case, much more research must be done in order to have a total understanding of the dynamics and the geometrical structure of high dimensional Lyness' maps.

Acknowledgements. We want to thank Guy Bastien and Marc Rogalski for communicating their results of [4] prior to publication. The third author is grateful to Immaculada Gálvez for her kind help.

GSD-UAB and CoDALab Groups are supported by the Government of Catalonia through the SGR program. They are also supported by DGICYT through grants MTM2005-06098-C02-01 (first and second authors) and DPI2005-08-668-C03-1 (third author).

## References

- R. Abraham, J.E. Marsden. "Foundations of Mechanics" 2nd Ed. Addison–Wesley. Redwood City, California, 1987.
- [2] G. Bastien and M. Rogalski. Global behavior of the solutions of Lyness difference equation  $u_{n+2}u_n = u_{n+1} + a$ , J. Difference Equations and Appl. 10, no. 11 (2004), 977–1003.
- [3] G. Bastien and M. Rogalski. On algebraic difference equations  $u_{n+2} + u_n = \psi(u_{n+1})$  in  $\mathbb{R}$  related to a family of elliptic quartics in the plane, J. Math. Anal. Appl. 326 (2007), 822–844.

- [4] G. Bastien, M. Rogalsky. Results and conjectures about global behavior of the solutions of the order q Lyness' difference equation in ℝ<sup>+</sup><sub>\*</sub>, work in progress. Private communication. December 13<sup>th</sup>, 2006.
- [5] F. Beukers and R. Cushman. Zeeman's monotonicity conjecture, J. Differential Equations 143 (1998), 191–200.
- [6] A. Cima, A. Gasull, V. Mañosa. Dynamics of rational discrete dynamical systems via first integrals, Internat. J. Bifur. Chaos Appl. Sci. Engrg. 16 (2006), 631-645.
- [7] A. Cima, A. Gasull, V. Mañosa. Dynamics of the third order Lyness' difference Equation, J. Difference Equations & Appl. 13 (2007), 855–844.
- [8] A. Cima, A. Gasull, V. Mañosa. Studying discrete dynamical systems through differential equations, J. Differential Equations 244 (2008), 630–648.
- [9] A. Cima, F. Mañosas. Real dynamics of integrable birational maps. In preparation, 2008.
- [10] L. Gardini, G.I. Bischi, C. Mira. Invariant curves and focal points in a Lyness iterative process, Internat. J. Bifur. Chaos Appl. Sci. Engrg. 13 (2001), 1841–1852.
- M. Gao, Y Kato, M. Ito. Some invariants for k<sup>th</sup>-order Lyness equation, Appl. Math. Lett. 17 (2004), 1183-1189.
- [12] V. Guillemin, A. Pollack. "Differential Topology". Prentice Hall. Englewood Cliffs, New Jersey, 1974.
- [13] I. Gumovski, C. Mira. "Recurrences and discrete dynamic systems". Lecture Notes in Mathematics 809. Springer Verlag, Berlin, 1980.
- [14] F.A. Haggar, G.B. Byrnes, G.R. Quispel and H.W. Cappel. k-integrals and k-Lie symmetries in discrete dynamical systems, Phys. A 233 (1996), 379–394.
- [15] R. Hirota, K. Kimura and H. Yahagi. How to find the conserved quantities of nonlinear discrete equations J. Phys. A: Math. Gen. 34 (2001), 10377–10386.
- [16] A. Iatrou. Three dimensional integrable mappings, arXiv:nlin.SI/0306052vl (2003)
- [17] A. Iatrou. Higher dimensional integrable mappings, Phys. D: 179 (2003), 229–253.
- [18] A. Iatrou and , J.A.G. Roberts. Integrable mappings of the plane preserving biquadratic invariant curves, J. Phys. A: Math. Gen. 34 (2001), 6617–6636.

- [19] K. Kimura, H. Yahagi, R. Hirota, A. Ramani, B. Grammaticos and Y. Ohta. A new class of integrable discrete systems, J. Phys. A: Math. Gen. 35 (2002), 9205–9212.
- [20] M.R.S. Kulenović. Invariants and related Liapunov functions for difference equations, Appl. Math. Lett. 13 (2000), 1–8.
- [21] S. Lang. "Differential and Riemannian Manifolds". Springer. New York, 1995.
- [22] J. Matsukidaira, D. Takahashi. Third-order integrable difference equations generated by a pair of second order equations, J. Phys. A: Math. Gen. 39 (2006), 1151–1161.
- [23] J.A.G. Roberts, G.R.W. Quispel. Creating and relating three dimensional integrable maps, J. Phys. A: Math. Gen. 39 (2006), L605–L615.
- [24] E. C. Zeeman. Geometric unfolding of a difference equation. Unpublished paper. Hertford College, Oxford(1996), 1–42.